



HAL
open science

Formaldehyde around 3.5 and 5.7- μm : measurement and calculation of broadening coefficients

David Jacquemart, Anne Laraia, Fridolin Kwabia Tchana, Robert R. Gamache, Agnes Perrin, Nelly Lacombe

► To cite this version:

David Jacquemart, Anne Laraia, Fridolin Kwabia Tchana, Robert R. Gamache, Agnes Perrin, et al.. Formaldehyde around 3.5 and 5.7- μm : measurement and calculation of broadening coefficients. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 2010, 111 (9), pp.1209-1222. 10.1016/j.jqsrt.2010.02.004 . hal-00744701

HAL Id: hal-00744701

<https://hal.sorbonne-universite.fr/hal-00744701>

Submitted on 24 Oct 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Formaldehyde around 3.5 and 5.7- μm : measurement and calculation of broadening coefficients

D. JACQUEMART^(1,2,*), A. LARAIA⁽³⁾, F. KWABIA TCHANA^(1,2,†), R.R. GAMACHE⁽³⁾, A.
PERRIN⁽⁴⁾ and N. LACOME^(1,2)

¹ Université Pierre et Marie Curie-Paris 6, Laboratoire de Dynamique, Interactions et Réactivité, UMR 7075, Case Courrier 49, 4 Place Jussieu, 75252 Paris Cedex 05, France

² CNRS, UMR 7075, Laboratoire de Dynamique, Interactions et Réactivité, Case Courrier 49, 4 Place Jussieu, 75252 Paris Cedex 05, France

³ University of Mass. Lowell, Department of Environmental, Earth & Atmospheric Sciences
Lowell MA 01854, USA

⁴ Laboratoire Interuniversitaire des Systèmes Atmosphériques (LISA), CNRS/Univ Paris Est & Paris7, 61 avenue du Général de Gaulle, 94010 Créteil cedex, France

Manuscript Pages : 16

No. of Figures : 8+4

No. of Tables : 3

* Corresponding author. Tel.: + 33-1-44-27-36-82; fax: + 33-1-44-27-30-21.
E-mail address: jacquemart@spmol.jussieu.fr (D. Jacquemart).

† Current address: Laboratoire Interuniversitaire des Systèmes Atmosphériques (LISA), CNRS/Univ Paris Est & Paris 7, 61 avenue du Général de Gaulle, 94010 Créteil cedex, France.

ABSTRACT

Self- and N₂-broadening coefficients of H₂CO have been retrieved in both the 3.5 and 5.7- μ m spectral regions. These coefficients have been measured in FT spectra for transitions with various J (from 0 to 25) and K values (from 0 to 10), showing a clear dependence with both rotational quantum numbers J and K . First, an empirical model is presented to reproduce the rotational dependence of the measured self- and N₂-broadening coefficients. Then, calculations of N₂-broadening of H₂CO were made for some for 3296 ν_2 transitions using the semi-classical Robert-Bonamy formalism. These calculations have been done for various temperatures in order to obtain the temperature dependence of the line widths. Finally, self- and N₂-broadening coefficients, as well as temperature dependence of the N₂-widths has been generated to complete the whole HITRAN 2008 version of formaldehyde (available as supplementary materials).

Keywords: Formaldehyde; broadening coefficients; widths; H₂CO; Fourier transform spectroscopy; Robert Bonamy formalism

1. Introduction

Both the 3.5 and 5.7 μm spectral regions of formaldehyde are used for the optical detection of this molecule in the atmosphere [1-11]. The importance of the 3.5 and 5.7 μm regions of formaldehyde (H_2CO) has been previously presented in our previous work [12], in which line intensities measurement and calculation have been performed in both spectral regions. These new calculations [12] have been used to update and complete the HITRAN 2008 edition [13] of formaldehyde. For a better detection of formaldehyde in the atmosphere, the knowledge of accurate N_2 -broadening coefficients is necessary. Spectra of formaldehyde broadened by N_2 have recorded in Paris (LADIR) using a Bruker HR-120 spectrometer in order to perform measurements of N_2 -broadening coefficients. Both self- and N_2 -broadening coefficients have measured for numerous transitions in both spectral regions, allowing us to observe rotational dependences with respect to both J and K rotational quantum numbers. A polynomial expansion has been used to reproduce the measurements and their rotational dependences. Also, calculations of N_2 -broadening of H_2CO were made for some for 3296 ν_2 transitions using the semi-classical Robert-Bonamy formalism. These calculations have been done for various temperatures in order to obtain the temperature dependence of the line widths. The goal of this work is to complete the complete HITRAN line list of formaldehyde with both self- and N_2 broadening coefficients, as well as with the N_2 -broadening temperature dependence.

The experimental conditions of the spectra recorded in this work are detailed in Section 2. The measurements are presented in Section 3 together with the polynomial expansion used to model them, as well as the theoretical calculation of N_2 -broadening coefficients. Comparison with the measurements obtained in this work and others works is also presented. Finally, the generation of a complete line list for formaldehyde at 3.6 and 5.7 μm in HITRAN format [13] is presented in Section 4 and available as supplementary materials.

2. Experimental details

All the experimental spectra used in this work have been recorded using the Fourier transform spectrometer Bruker IFS 120 HR of LADIR in Paris. For all the spectra, the instrument was equipped with a MCT photovoltaic detector, a Ge/KBr beamsplitter, and a Globar source. The whole optical path was under vacuum, and a 0.8 mm entrance aperture

diameter was used. No optical filter has been used in order to measure both the 3.6 and 5.7 μm spectral regions of formaldehyde absorption. The recorded spectral domain is between 1400 and 4000 cm^{-1} . Each spectrum was the result of the co-addition of 100 interferograms (an over sampling ratio of 8 has been used by post-zero filling the interferograms, but no numerical apodization has been performed). A 30 cm stainless steel cell closed by two KBr windows has been used. More details on the experimental set-up can be found in Ref. [12]. The experimental conditions of spectra used in this work are summarized in Table 1.

3. Measurements, calculations, and comparisons

3.1. Measurements

Based on the 4 experimental spectra recorded with pure H_2CO gas (see conditions in Table 1 for spectra 1-4), the multispectrum fitting procedure described in Ref. [14] has been used to retrieve the line positions, the intensities, and the self-broadening coefficients [12]. In order to avoid polymerization, low pressures of H_2CO have been used. Consequently even with quite large self-broadening coefficients (around $0.5 \text{ cm}^{-1}/\text{atm}$), the determination of these coefficients can not be very accurate. For spectra 5-8 recorded with a mixture of $\text{H}_2\text{CO}-\text{N}_2$, the contribution of the self-widths is not completely negligible. In a first step the self-broadening coefficients have been obtained with spectra 1-4 and modelled using a polynomial expansion that allowed reproducing the rotational dependence with respect in J and K_a of the lower state of the transitions (see Section 3.2). In the following text, the notations J , K_a and K_c are the quantum numbers associated to the lower state of the transition. When performing the fit of spectra 5-8, the self-broadening coefficients have been fixed to the empirical polynomial expansion described latter in section 3.2.

Finally, 284 and 368 self-broadening coefficients have been retrieved respectively in the 5.7 and 3.6- μm spectral regions, and 280 and 456 N_2 -broadening coefficients respectively in the 5.7 and 3.6- μm spectral regions. All these measurements have been gathered in Table A1 and A2 given as supplementary materials and are plotted versus J in Fig. 1 for the self-broadening coefficients and in Fig.2 for the N_2 -broadening coefficients. These figures show a clear rotational dependence with respect to J . The large dispersion of the measurements in these figures hides partially the rotational dependence with respect to K_a . Indeed, when measurements for same J values are plotted versus K_a , it reveals a K_a rotational dependence

for both self- and N₂-broadening coefficients. For a set of broadening coefficients with same value of J , the widths decrease with K_a increasing. Such dependence has already been pointed out for other molecules [15-22]. As an example, this dependence is plotted in Figs. 3 and 4, respectively for the self- and N₂-broadening coefficients having same J value ($J = 8$). Let us recall that the spectra have been recorded with low pressures of formaldehyde in order to avoid polymerization. As a consequence, the determination of the self-broadening coefficients is not very accurate, which could explain that the dispersion for self-broadening coefficients is more important than for N₂-broadening coefficients. The accuracy for the self-broadening coefficients is estimated to be between 10 and 20%, whereas for the N₂-broadening coefficients the accuracy has been estimated to be between 5 and 10%. For both self- and N₂- broadening coefficients, no rotational dependence has been observed with respect to the K_c quantum numbers or with the type of transition (various values of ΔJ , ΔK_a , or ΔK_c).

3.2. Empirical polynomial expansion

Empirical polynomial expansion has been used to reproduce both the J and K_a rotational dependence for the self- and N₂-broadening coefficients measured in this work. Such a model has already been presented previously for CH₃Br [22]. Each set of same value of J was fitted by a polynomial expansion of order two in K_a (fixing the first-order term to zero):

$$\gamma_J(K_a) = a_J^0 + a_J^2 K_a^2 \quad (1)$$

Example of the rotational dependence versus K_a (for $J=8$) is plotted in Fig.3 for the self-broadening coefficients and in Fig. 4 for the N₂-broadening coefficients. The two parameters a_J^0 and a_J^2 obtained for each set of same values of J have then been plotted versus J in Figs.5 and 6 for the self- and N₂-broadening coefficients respectively. The continuous line in Fig. 5 and 6 to the manual smooth of the values deduced from experimental measurements. The smoothed values of a_J^0 and a_J^2 are given in Table 2 and can be used to generate broadening parameters for J and K_a values ranging from 0 to 30 and 0 to 7 respectively. In Figs. 7 and 8 are plotted, for K_a equal to 0 to 7, both experimental broadening coefficients and calculated one using Eq. (1) and parameters of Table 2.

The a_j^0 parameters represent the J dependence of the broadening parameters for $K^a=0$, whereas the a_j^2 parameters describe the influence of the rotational quantum number K_a on broadening parameters. Despite the important dispersion of the measurements, especially for the self-broadening coefficients, the a_j^0 and a_j^2 parameters are quite smooth and reproduced quite well the rotational dependences in Figs. 3,4,7 and 8. A part of the dispersion of the broadening coefficients in Figs. 1 and 2 is due to the K rotational dependence of the widths modelled by the a_j^2 parameters.

3.3. Theoretical calculation for N₂-broadening coefficients

3.3.1 The Complex Robert-Bonamy Formalism

The calculations made here are a complex implementation [23-25] of the semi-classical theory of Robert and Bonamy [26]. The complex Robert-Bonamy (CRB) formalism produces the half-width and line shift from a single complex calculation. In this formalism the half-width, γ , and line shift, δ , of a ro-vibrational transition $f \leftarrow i$ are given by minus the imaginary part and the real part, respectively, of the diagonal elements of the complex relaxation matrix [27,28]. In computational form γ and δ are expressed in terms of the Liouville scattering matrix

$$(\gamma - i\delta) = \frac{n_2}{2 \pi c} \left\langle v \left[1 - e^{-R S_2(f,i,J_2,v,b)} e^{-i I S_2(f,i,J_2,v,b)} \right] \right\rangle_{v,b,J} \quad (2)$$

where n_2 is the number density of perturbers and $\langle \rangle_{v,b,J_2}$ represents an average over all trajectories (impact parameter b and initial relative velocity v) and initial rotational state J_2 of the collision partner. $S_2 = {}^R S_2 + i I S_2$ is the second order terms in the successive expansion of the scattering matrix, which depends on the ro-vibrational states involved and associated collision induced jumps from these levels, on the intermolecular potential and characteristics of the collision dynamics. Note, Eq. (2) generally contains the vibrational dephasing term, S_1 , which arises only for transitions where there is a change in the vibrational state. The potential leading to S_1 is written in terms of the isotropic induction and London dispersion interactions which depend on the vibrational dependence of the dipole moment and polarizability of the radiating molecule. These parameters are not available for H₂CO and the S_1 term has been omitted from the calculation. Note also, for collision systems with a strong electrostatic

component, like H₂CO-N₂, the effect of the S_1 term on the half-width often tends to be small. The exact form of the S_2 term is given in Refs. [23-26].

The intermolecular potential used in the calculations is comprised of an electrostatic component (dipole and quadrupole moments of H₂CO with the quadrupole moment of N₂ or with the dipole and quadrupole moments of H₂CO) and an atom-atom component. The initial heteronuclear Lennard-Jones parameters for the atomic pairs are determined using the "combination rules" of Hirschfelder et al. [29]. The atom-atom distance, r_{ij} is expressed in terms of the center of mass separation, R , via the expansion in $1/R$ of Sack [30] using the formulation of Neshyba and Gamache [31] expanded to 8th order.

The dynamics of the collision process use Robert and Bonamy's [26] second order in time approximation to the true trajectories, which are based on the isotropic part of the intermolecular potential. These curved trajectories have been shown to accurately model the true trajectories [32].

3.3.2 Parameters for the H₂CO-N₂ collision system

The wavefunctions used to evaluate the reduced matrix elements are obtained by diagonalizing the Watson Hamiltonian [33] in a symmetric top basis. The wavefunctions for the ground vibrational state of H₂CO are determined using the Watson-Hamiltonian constants of Muller *et al.* [34] and those for the ν_2 vibrational state use the Watson constants of Perrin [35]. The molecular constants for N₂ are from Huber and Herzberg [36].

The molecular constants used in the calculations are as follows: The dipole moment is from the work of Fabricant *et al.* [37], $\mu=2.33$ D. The quadrupole moments are from Kukolich [38] here reported with H₂CO in the I^R representation: $Q_{xx}=-0.269\pm 0.20$ 10⁻²⁶ esu, $Q_{yy}=+0.3295\pm 0.12$ 10⁻²⁶ esu, $Q_{zz}=-0.0605\pm 0.16$ 10⁻²⁶ esu. The quadrupole moment of nitrogen is from Mulder et al. [39], $Q_{zz}=-1.4\pm 0.1$ 10⁻²⁶ esu. The starting atom-atom parameters were obtained using the standard combination rules with the atom-atom parameters for homonuclear diatomics determined by Bouanich [40] by fitting to second virial coefficient data. In the calculations, the atom-atom potential is expanded to eighth-order in the molecular centers of mass separation.

The input Lennard-Jones atom-atom parameters are not as well known as the other parameters; the ϵ values can vary by 30% and the σ values by 5% depending on the source and the ϵ values can vary by 69% and the σ values by 9% depending on whether they were derived using viscosity data or virial data [29]. Depending on how the values were derived it

is possible to find examples in the literature where the parameters for the same interaction pair differ by factors of 2. Thus it appears reasonable to adjust the atom-atom parameters and/or the resulting trajectory parameters provided there are reliable experimental data to fit to. To accomplish this, the measured spectra were studied to identify transitions for which we had high confidence in the measurement. This analysis yielded 39 transitions for the H₂CO-N₂ system.

For H₂CO-N₂ there are 6 atom-atom parameters (ϵ_{HN} , σ_{HN} , ϵ_{CN} , σ_{CN} , ϵ_{ON} , and σ_{ON}) that need to be adjusted. Starting from the combination rule values, see Table 3, individually each epsilon or sigma value was increased by 5% and calculations made. These results were compared to the values calculated using the combination rule values yielding a sense of how the half-width varies with the atom-atom parameters. Using these changes as a guide the parameters were varied and calculations made and compared with the 39 measured half-widths. After 14 iterations the percent difference went from the initial value of -11.1 to a final value of 0.1. The final atom-atom parameters for the H₂CO-N₂ system are given in Table 3. Note, the adjustment of the atom-atom parameters was done without the use of a least-squares technique due to the complexity of the atom-atom potential and the time required for a single calculation (~ 4 hours).

3.3.3 Calculation of half widths at various temperatures

A file of 3713 transitions in the ν_2 , ν_3 , and ν_5 bands of H₂CO [35] in the frequency range 1620-1840 cm⁻¹ atm⁻¹ was taken and 3296 ν_2 transitions extracted for this study. Complex Robert-Bonamy calculations of the half-width were made at 7 temperatures (200., 225., 275., 296., 350., 500., and 700. K) by solving Eq. 2. The calculations are for transitions with $J=0$ to 41 and $K_a=0$ to 16. The half-widths at 296 K range from roughly 0.02 to 0.128, a factor of 6.4.

The power law model for the temperature dependence of the half-width was given by Birnbaum [41] by considering a one term intermolecular potential and all on-resonance collisions giving

$$\gamma(T) = \gamma(T_0) \left[\frac{T_0}{T} \right]^n \quad (3)$$

where n is called the temperature exponent. While modern calculations use hundreds of terms in the intermolecular potential, Eq. (3) remains a reliable model for many systems. However, the correctness of Eq. (3) does need to be tested as discussed below.

The temperature exponent does depend on the temperature range of the fit [42,43]. Here, the temperature exponent was determined for each transition by a least-squares fit of $\ln[\gamma(T)/\gamma(T_0)]$ vs. $\ln[T_0/T]$ using five (200K-350K) and seven (200K-700K) temperatures of the study. The error in the temperature exponent was determined as follows: The temperature exponents were calculated using the half-width values at any two of the temperatures studied. For example, with seven temperatures this yields twenty-one 2-point temperature exponents. The difference between each 2-point temperature exponent and the least-squares fit value is calculated. The error is taken as the largest of these differences. While this procedure tends to yield the maximum error in the temperature exponent, given the nature of the data and other uncertainties it is thought to be more reasonable than a statistical value taken from the fit.

Discussion

Half-width as a function of the rotational quantum numbers

The half-widths at 296 K are plotted versus an energy ordered index $(J(J+1)+K_a-K_c+1)$ in Figure 7 where the plot symbols are the K_a values. The roughly vertical columns in the figure are data with the same J . There appears to be some structure with respect to the K_a with the largest half-widths belonging to transitions with the smallest K_a values. However, no simple propensity rule to predict half-widths can be derived from the data.

3.3.4 Temperature dependence of the half-width

The temperature dependence of the N₂-broadened half-widths was determined for the 3296 ν_2 band transitions studied in this work using the power law formula, Eq. (3). The “rule-of-thumb” expression for the temperature exponent given by Birnbaum [41] for a “dipole-quadrupole” system, such as H₂CO-N₂ states that $n=5/6$.

Some recent studies have shown that for certain types of radiator-perturber interactions the power law model is questionable. Wagner *et al.* [44] have observed that for certain transitions of water vapor perturbed by air, N₂ or O₂ the power law does not correctly model the temperature dependence of the half-width. This fact was later demonstrated by Toth *et al.* [45] in a study of air-broadening of water vapor transitions in the region from 696-2163 cm⁻¹. In both studies it was found that the temperature exponent, n , can be negative for many transitions. In such cases the power law model, Eq. (3), is not completely correct. The

mechanism leading to negative temperature exponents is called the resonance overtaking effect and was discussed by Hartmann *et al.* [46], Wagner *et al.* [44], and Antony *et al.* [47]. Thus, in this work the applicability of the power law model was tested.

The values for the temperature dependence of the N₂-broadened half-width go from roughly 0.83 to 0.10. While the range in values is large, there are no negative temperature exponents for this collision system. As stated above the temperature exponents were calculated for 2 temperature ranges; 200-350K, and 200-700K. Figure 8 shows the fits for the $19_{155} \leftarrow 18_{154}$ transition for the two temperature ranges. Plotted are $\ln[\gamma(T)/\gamma(T_0)]$ versus $\ln[T_0/T]$ where the slope of the fitted line is the value of n . The temperature exponent for the 7 data points is 0.40 (top panel) and that for the 5-point fit is 0.54 (bottom panel). For both cases the correlation coefficient, R , indicates the power law fit is not perfect and the error associated with the 7-point fit is almost 3 times larger than that for the 5-point fit. (Note, the scale of the abscissa is different in each panel.) For atmospheric applications the temperature range of the 5-point fit, 200-350K, is more appropriate and the temperature exponents reported here are those of the 5-point fit. In Fig. 9 $\ln[\gamma(T)/\gamma(T_0)]$ versus $\ln[T_0/T]$ is plotted for the $2_{11} \leftarrow 2_{12}$ transition (top panel) and the $20_{146} \leftarrow 19_{145}$ transition (lower panel). In the top panel the fit is quite good ($R=0.99995$) and the slope, n , is 0.761(0.038). In the lower panels is an example of a high J and K_a transition where n is 0.360(0.240) the fit is not as good ($R=0.9831$), which is reflected in the reported error. It is cautioned that there are many transitions for which the power law model does not work well.

In Fig. 10 the temperature exponents for the 3713 N₂-broadened transitions of H₂CO are plotted versus the energy ordered index. The dashed line in the plot is the “rule of thumb” value, 5/6. Plots (not shown here) were made with K_a as the plot symbol and different coordinates for the abscissa but no structure was evident. Figure 10 shows there are large variations in n as a function of the rotational quantum numbers.

3.4. Comparisons

Figure 11 shows the measured half-widths from Refs [48-51] and this work and the calculated half-widths versus $J+0.9(K_a/J)$. The star symbols are the measurements of the ν_3 band transitions by Cline and Varghese [48], The square symbol is the measurement of Burkart and Schramm on the $4_{14} \leftarrow 5_{15}$ transition in the ν_1 band [49], the open circle with

the plus sign symbol is the measurement of the $3_{13} \leftarrow 3_{12}$ transition in the rotation band [50], the open plus sign symbol is the measurement of the the $3_{03} \leftarrow 3_{13}$ transition in the ν_4 band by Nadler *et al.* [51], the solid triangles are the measurements of this work. All the measurements have the associated error bars plotted as well. The solid circles are the CRB calculations of the half-widths for ν_2 transitions. The plot shows the data of refs. [49] and [51] are low compared with the other measurements. It is not certain if this is an effect of vibrational dependence of the half-widths for these two transitions. The data of Ref. 48 agree somewhat with the measurements made here at low J but the data above $J=8$ have smaller values than the measurements made here. The calculated half-widths are lower than the measurements made here for low J . For $J \geq 5$ the agreement is good. Overall the calculations have a 4.5 % difference when compared to all 258 measurements.

4. Creation of a complete line list

The last edition of HITRAN [13] contain the recent line list of A. Perrin *et al.* [12] concerning both the 3.6 and 5.7 μm spectral regions (35958 lines), and quite all data for the 0-100 cm^{-1} spectral region (163 lines). For all these transitions, constant values have been used for both the self- and N_2 -broadening coefficients, as well as for the temperature dependence of the N_2 -broadening coefficients. Table A3 given as supplementary materials is a copy of the formaldehyde $^{12}\text{H}_2^{12}\text{C}^{16}\text{O}_2$ line list from HITRAN 2008 [13], in which self- and N_2 broadening coefficients generated with the empirical polynomial expansion together with the parameters of Table 2 (see Section 3.2). The N_2 -broadening coefficients and its temperature dependence, calculated using the complex Robert Bonamy formalism for the ν_2 band (see Section 3.3), have also been used to complete the HITRAN line list. Then, Table 3 has not really the same format as the one of the HITRAN database since two columns of N_2 -broadening coefficients are given instead of one.

ACKNOWLEDGEMENTS

This work was supported by the Programme National de Chimie Atmosphérique (PNCA).

REFERENCES

- [1] Yokelson RJ, Goode JG, Ward DE, Susott RA, Babbitt RE, Wade DD, Bertschi I, Griffith DWT, Hao WM. Emissions of formaldehyde, acetic acid, methanol, and other trace gases from biomass fires in North Carolina measured by airborne Fourier transform infrared spectroscopy. *J of Geophys Res* 1999;D104:30109-30125.
- [2] Fried A, Sewell S, Henry B, Wert BP, Gilpin T. Tunable diode laser absorption spectrometer for ground-based measurements of formaldehyde. *J of Geophys Res* 1997;D102:6253-6266.
- [3] Fried A, Lee YN, Frost G, Wert B, Henry B, Drummond JR, Hubler G, Jobson T. Airborne CH₂O measurements over the North Atlantic during the 1997 NARE campaign: Instrument comparisons and distributions. *J Geophys Res* 2002;D107:ACH14-1-21.
- [4] Wert BP, Fried A, Rauenbuehler S, Walega J, Henry B. Design and performance of a tunable diode laser absorption spectrometer for airborne formaldehyde measurements. *Journal of Geophysical Research* 2003;108(D12):ACH1-1-15.
- [5] Richter D, Fried A, Wert BP, Walega JG, Tittel FK. Development of a tunable mid-IR difference frequency laser source for highly sensitive airborne trace gas detection *Appl Phys B75* 2002 281-288
- [6] Dahnke H, von Basum G, Kleinermanns K, Hering P, Murts M. Rapid formaldehyde monitoring in ambient air by means of mid-infrared cavity leak-out spectroscopy. *Appl Phys* 2002;B75:311-316.
- [7] Coheur PF, Herbin H, Clerboux C, Hurtmans D, Wespes C, Carleer M, Turquety S, Rinsland CP, Remedios J, Hauglustaine D, Boone CD, Bernath PF. ACE-FTS observation of a young biomass burning plume: first reported measurements of C₂H₄, C₃H₆O, H₂CO and PAN by infrared occultation from space. *Atmos Chem Phys* 2007;7:5437-5446.
- [8] Wagner V, Schiller C, Fischer H. Formaldehyde measurements in the marine boundary layer of the Indian Ocean during the 1999 INDOEX cruise of the R/V Ronald H. Brown. *J Geophys Res* 2001;D106:28528-28538.
- [9] Steck T, Glatthor N, von Clarmann T, Fischer H, Flaud JM, Funke B, Grabowski U, Höpfner M, Kellmann S, Linden A, Perrin A, Stiller GP. Retrieval of global upper tropospheric and stratospheric formaldehyde (H₂CO) distributions from high-resolution MIPAS-Envisat spectra. *Atmos Chem Phys* 2008;8:463-470.
- [10] Herndon SC, Zahniser MS, Nelson DD Jnr, Shorter J, McManus JB, Jimenez R, Warneke C, de Gouw JA. Airborne measurements of HCHO and HCOOH during the New England Air Quality Study 2004 using a pulsed quantum cascade laser spectrometer. *J of Geophys Res* 2007;112(D10): D10S03-1-15.
- [11] Rinsland CP, Goldman A. Infrared spectroscopic measurements of tropospheric trace gases. *Appl Opt* 1992; 31:6969-6971.
- [12] A. Perrin, D. Jacquemart, F. Kwabia Tchana, and N. Lacombe. Absolute line intensities and new linelists for the 5.7 and 3.6 μm bands of formaldehyde. *JQSRT* 2009;110:700-16.
- [13] Rothman LS, Gordon IE, Barbe A, Chris Benner D, Bernath PF, Birk M, Brown LR, Boudon V, Champion J-P, Chance K, Coudert LH, Dana V, Fally S, Flaud J-M, Gamache RR, Goldman A, Jacquemart D, Lacombe N, Mandin J-Y, Massie ST, Mikhailenko S, Orphal J, Perevalov V, Perrin A, Rinsland CP, Šimečková M, Smith MAH, Tashkun

S, Tennyson J, Toth RA, Vandaele AC, Vander Auwera J. The HITRAN 2008 Molecular Spectroscopic Database. *J. Quant. Spectrosc. Radiat. Transfer* 110,533-572 (2009).

[14] Jacquemart D, Mandin JY, Dana V, Picqué N, Guelachvili G. A multispectrum fitting procedure to deduce molecular line parameters. Application to the 3-0 band of $^{12}\text{C}^{16}\text{O}$. *Eur Phys J D* 2001;14:55-69.

[15] Nemtchinov V, Sung, K, Varanasi P. Measurements of line intensities and half-widths in the 10- μm bands of $^{14}\text{NH}_3$. *JQSRT* 2004;83:243-65.

[16] Predoi-Cross A, Hambrook K, Brawley-Tremblay S, Bouanich JP, Malathy Devi V, Smith MAH. Room-temperature broadening and pressure-shift coefficients in the ν_2 band of $\text{CH}_3\text{D}-\text{O}_2$: Measurements and semi-classical calculations. *J Mol Spectrosc* 2006;236:75-90.

[17] Predoi-Cross A, Hambrook K, Brawley-Tremblay S, Bouanich JP, Malathy Devi V, Smith MAH. Measurements and theoretical calculations of N_2 -broadening and N_2 -shifting coefficients in the ν_2 band of CH_3D . *J Mol Spectrosc* 2006;235:35-53.

[18] Lepère M, Blanquet G, Walrand J, Bouanich JP. K -dependence of broadening coefficients for $\text{CH}_3\text{F}-\text{N}_2$ and for other systems involving a symmetric top molecule. *J Mol Spectrosc* 2000;517-518:493-502.

[19] Chackerian C, Brown LR, Lacombe N, Tarrago G. Methyl Chloride ν_5 Region Lineshape Parameters and Rotational Constants for the ν_2 , ν_5 , and $2\nu_3$ Vibrational Bands. *J Mol Spectrosc* 1998;191:148-57.

[20] Bouanich JP, Blanquet G, Walrand J. Theoretical O_2 - and N_2 -Broadening Coefficients of CH_3Cl Spectral Lines. *J Mol Spectrosc* 1993;161:416-26.

[21] Levy A, Lacombe N, Tarrago G. Hydrogen- and Helium-Broadening Phosphine Lines. *J Mol Spectrosc* 1993;157:172-81.

[22] Jacquemart D, Kwabia Tchana F, Lacombe N, Kleiner I. A complete set of line parameters for CH_3Br in the 10- μm spectral region. *JQSRT* 2007;105:264-302.

[23] Lynch R. Half-widths and line shifts of water vapor perturbed by both nitrogen and oxygen. Ph.D. dissertation, University of Massachusetts Lowell, June, 1995.

[24] Lynch R, Gamache RR, Neshyba SP. Fully Complex Implementation of the Robert-Bonamy Formalism: Halfwidths and Line Shifts of H_2O Broadened by N_2 , *J. Chem. Phys.* 1996; **105**: 5711-21.

[25] amache RR, Lynch R, Neshyba SP. New Developments in the Theory of Pressure-Broadening and Pressure-Shifting of Spectral Lines of H_2O : The Complex Robert-Bonamy Formalism, *J. Quant. Spectrosc. Radiat. Transfer* 1998; **59**: 319-35

[26] Robert D, Bonamy J. Short range force effects in semiclassical molecular line broadening calculations, *Journal De Physique* 1979; **20**: 923-43.

[27] Baranger M. General impact theory of pressure broadening, *Phys Rev* 1958; **112**: 855-65.

[28] Ben-Reuven A, Spectral Line Shapes in Gases in the Binary-Collision Approximation, *Adv. Chem. Phys.*, Prigogine I and Rice SA, Editors. 1975, Academic Press: New York. p. 235.

[29] Hirschfelder JO, Curtiss CF, Bird RB. *Molecular Theory of Gases and Liquids*, Wiley, New York 1964.

- [30] Sack RA. Two-Center Expansion for the Powers of the Distance Between Two Points, *J. Math. Phys.* 1964; **5**: 260-8.
- [31] Neshyba SP, Gamache RR. Improved Line Broadening Coefficients for Asymmetric Rotor Molecules: Application to Ozone Perturbed by Nitrogen., *J. Quant. Spectrosc. Radiat. Transfer* 1993; **50**: 443-53.
- [32] Neshyba SP, Lynch R, Gamache RR, Gabard T, Champion J-P. Pressure Induced Widths and Shifts for the ν_3 band of Methane, *J. Chem. Phys.* 1994; **101**: 9412-21.
- [33] Watson JKG. Determination of centrifugal Distortion Coefficients of Asymmetric- Top Molecules, *J. Chem. Phys.* 1967; **46**: 1935-49.
- [34] Muller HSP, Winnewisser G, Demasion J, Perrin A, Valentin A. The Ground State Spectroscopic Constants of Formaldehyde, *J. Mol. Spectrosc.* 2000; **200**: 143-4.
- [35] Perrin A. Watson Hamiltonian constants for the ν_2 band of H_2CO and H_2CO linelist, private communication, Laboratoire Inter-Universitaire des Systemes Atmospheriques (LISA), CNRS, Universite Paris XII, Paris, France, 2007.
- [36] Huber KP, Herzberg G. *Molecular Spectra and Molecular Structure: Constants of Diatomic Molecules*, Van Nostrand, New York 1979.
- [37] Fabricant B, Krieger D, Muebnter JS. Molecular beam electric resonance study of formaldehyde, thioformaldehyde, and ketene, *J. Chem. Phys.* 1977; **67**: 1576-86.
- [38] Kukolich SG. Molecular Beam Measurements of the Magnetic Susceptibility Anisotropies and Molecular Quadrupole Moment in H_2CO , *J. Chem. Phys.* 1971; **54**: 8-11.
- [39] Mulder F, van Dijk G, van der Avoird A. Multipole moments, polarizabilities and anisotropic long range interaction coefficients for N_2 , *Mol. Phys.* 1980; **39**: 407-25.
- [40] Bouanich J-P. Site-Site Lennard- Jones Potential Parameters for N_2 , O_2 , H_2 , CO and CO_2 , *J. Quant Spectrosc. Radiat. Transfer* 1992; **47**: 243-50.
- [41] Birnbaum G. Microwave Pressure Broadening and its Application to Intermolecular Forces, *Advances in Chem. Phys.* 1967; **12**: 487-548.
- [42] L. R. Brown, C. M. Humphrey and R. R. Gamache, "CO₂-broadened water in the pure rotation and ν_2 fundamental regions," *J. Mol. Spectrosc.* **246**, 1-21, 2007.
- [43] R. R. Gamache and A. L. Laraia, "N₂-, O₂-, and air-broadened half-widths, their temperature dependence, and line shifts for the rotation band of H_2^{16}O ," in press *J. Mol. Spectrosc.*, 2009.
- [44] Wagner G, Birk M, Gamache RR, Hartmann J-M. Collisional parameters of H_2O lines: effects of temperature, *J. Quant. Spectrosc. Radiat. Transfer* 2005; **92**: 211-30.
- [45] Toth RA, Brown LR, Smith MAH, Malathy Devi V, Chris Benner D, Dulick M. Air-broadening of H_2O as a function of temperature: $696\text{-}2163\text{ cm}^{-1}$, *J. Quant. Spectrosc. Radiat. Transfer* 2006; **101**: 339-66.
- [46] Hartmann JM, Taine J, Bonamy J, Labani B, Robert D. Collisional broadening of rotation-vibration lines for asymmetric-top molecules II. H_2O diode laser measurements in the 400-900 K range; calculations in the 300-2000 K range, *J. Chem. Phys.* 1987; **86**: 144-56.
- [47] Antony BK, Neshyba S, Gamache RR. Self-broadening of water vapor transitions via the complex Robert-Bonamy theory, *J. Quant. Spectrosc. Radiat. Transfer* 2006; **105**: 148-63.

- [48] Cline DS, Varghese PL. High resolution spectral measurements in the ν_3 band of formaldehyde using a tunable IR diode laser, *Applied Optics* 1988; **27**: 3219-24.
- [49] Burkart M, Schramm B. Foreign gasbroadening of an IR absorption line of formaldehyde, *J. Mol. Spectrosc.* 2003; **217**: 153-6.
- [50] Srivastava GP, Gautam HO, Kumar A. Microwave pressure broadening studies of some molecules, *J. Phys. B: Atom. Molec. Phys.* 1973; **6**: 743-55.
- [51] Nadler S, Daunt SJ, Reuter DC. Tunable diode laser measurements of formaldehyde foreign gas broadening parameters and line strengths in the 9-11 μm region, *Appl. Opt.* 1987; **26**: 1641-6

Table 1.

*Absorbing sample*Natural H₂CO (98.624 % of H₂C¹⁶O)

Estimated purity ~98.9 %

Experimental conditions

SNR 75-100

Absorption path 30 cm

Spectrum No	Pressure of H ₂ CO (hPa)	Pressure of N ₂ (hPa)	Temperature (K)
1	0.480		297.7
2	0.536		297.2
3	0.577		297.8
4	0.657		298.3
5	0.491	147.6	298.6
6	0.556	197.6	298.7
7	0.587	286.5	298.3
8	0.658	379.4	298.6

Table 2. Parameters a_J^0 and a_J^2 retained for the calculation of the self- and N₂-broadening coefficients using Eq. (1).

J	Self-broadening coefficients		N ₂ -broadening coefficients	
	a_J^0	a_J^2	a_J^0	a_J^2
0	0.670	0.0	0.1390	0.0
1	0.670	0.0	0.1440	0.0
2	0.600	0.0	0.1388	-0.00040
3	0.530	0.0	0.1340	-0.00170
4	0.500	0.0	0.1297	-0.00130
5	0.510	-0.00654	0.1259	-0.00100
6	0.570	-0.00921	0.1225	-0.00088
7	0.623	-0.00648	0.1195	-0.00070
8	0.650	-0.00485	0.1168	-0.00060
9	0.666	-0.00400	0.1145	-0.00048
10	0.665	-0.00300	0.1126	-0.00041
11	0.650	-0.00230	0.1109	-0.00032
12	0.625	-0.00178	0.1095	-0.00026
13	0.598	-0.00120	0.1084	-0.00020
14	0.560	-0.00075	0.1075	-0.00016
15	0.520	-0.00040	0.1067	-0.00001
16	0.482	-0.00013	0.1062	-0.00005
17	0.450	0.0	0.1058	0.0
18	0.410	0.0	0.1055	0.0
19	0.380	0.0	0.1054	0.0
20	0.340	0.0	0.1053	0.0
21	0.300	0.0	0.1053	0.0
22	0.250	0.0	0.1053	0.0
23	0.209	0.0	0.1053	0.0
24	0.190	0.0	0.1052	0.0
25	0.190	0.0	0.1052	0.0
26	0.190	0.0	0.1050	0.0
27	0.188	0.0	0.1048	0.0
28	0.188	0.0	0.1040	0.0
29	0.188	0.0	0.1040	0.0
30	0.188	0.0	0.1040	0.0

Table 3. Initial and final atom-atom parameters for the H₂CO-N₂ collision systems

Interacting pair	$\epsilon_{\text{initial}}/k_B$ (K)	$\epsilon_{\text{final}}/k_B$ (K)	σ_{initial} (Å)	σ_{final} (Å)
H-N	20.45	13.2925	2.99	1.9435
C-N	33.75	21.9375	3.42	2.223
O-N	43.88	28.5220	3.148	2.0462



Fig 1: Self-broadening coefficients versus J .Fig 2: N_2 -broadening coefficients versus J .



Fig.3: Rotational dependence with respect to K_a^2 for self-broadening coefficients with $J=8$.Fig.4: Rotational dependence with respect to K_a^2 for N_2 -broadening coefficients with $J=8$.

Fig.5: Parameters a_j^0 and a_j^2 deduced from the fit of the measured self-broadening coefficients using Eq. (1). The error bars are 1SD. The continuous line symbolizes a manual smooth of these parameters.




Fig.6: Parameters a_J^0 and a_J^2 deduced from the fit of the measured N₂-broadening coefficients using Eq. (1). The error bars are 1SD. The continuous line symbolizes a manual fit of these parameters.




Fig. 7: Self-broadening coefficients measurements are plotted for each value of K_a values from 0 up to 7. The error bars are 1SD. The continuous line represents calculation based on an empirical expansion (see Eq. (1)) and parameters of Table 2. Open triangles are the theoretical calculation using the Complex Robert-Bonamy Formalism for $\Delta K_a = 0$ transitions (see Section 3.3).




Fig. 8: N_2 -broadening coefficients measurements are plotted for each value of K_a values from 0 up to 7. The error bars are 1SD. The continuous line represents calculation based on an empirical expansion (see Eq. (1)) and parameters of Table 2. Open triangles are the theoretical calculation using the Complex Robert-Bonamy Formalism for $\Delta K_a = 0$ transitions (see Section 3.3).

