



HAL
open science

Derivation of a Hele-Shaw type system from a cell model with active motion

Benoît Perthame, Fernando Quirós, Min Tang, Nicolas Vauchelet

► To cite this version:

Benoît Perthame, Fernando Quirós, Min Tang, Nicolas Vauchelet. Derivation of a Hele-Shaw type system from a cell model with active motion. *Interfaces and Free Boundaries: Mathematical Analysis, Computation and Applications*, 2014, 14 (4), pp.489-508. 10.4171/IFB/327 . hal-00906168

HAL Id: hal-00906168

<https://hal.sorbonne-universite.fr/hal-00906168v1>

Submitted on 19 Nov 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Derivation of a Hele-Shaw type system from a cell model with active motion

Benoît Perthame^{*†} Fernando Quirós[‡] Min Tang[§] Nicolas Vauchelet^{*†}

July 4, 2013

Abstract

We formulate a Hele-Shaw type free boundary problem for a tumor growing under the combined effects of pressure forces, cell multiplication and active motion, the latter being the novelty of the present paper. This new ingredient is considered here as a standard diffusion process. The free boundary model is derived from a description at the cell level using the asymptotic of a stiff pressure limit.

Compared to the case when active motion is neglected, the pressure satisfies the same complementarity Hele-Shaw type formula. However, the cell density is smoother (Lipschitz continuous), while there is a deep change in the free boundary velocity, which is no longer given by the gradient of the pressure, because some kind of ‘mushy region’ prepares the tumor invasion.

Key-words: Tumor growth; Hele-Shaw equation; porous medium equation; free boundary problems.
Mathematics Subject Classification 35K55; 35B25; 76D27; 92C50.

1 Introduction

Among the several models now available to deal with cancer development, there is a class, initiated in the 70’s by Greenspan [15], that considers that cancerous cells multiplication is limited by nutrients (glucosis, oxygen) brought by blood vessels. Models of this class rely on two kinds of descriptions; either they describe the dynamics of cell population density [5] or they consider the ‘geometric’ motion of the tumor through a free boundary problem; see [11, 10, 13, 16] and the references therein. In the latter kind of models the stability or instability of the free boundary is an important issue that has attracted attention, [9, 13].

The first stage, where growth is limited by nutrients, lasts until the tumor reaches the size of $\approx 1\text{mm}$; then, lack of food leads to cell necrosis which triggers neovasculatures development [8] that supply the tumor with enough nourishment. This has motivated a new generation of models where growth

^{*}UPMC Univ Paris 06 and CNRS UMR 7598, Laboratoire Jacques-Louis Lions, F-75005, Paris, France. Email B. P.: benoit.perthame@ljl.math.upmc.fr, Email N. V.: vauchelet@ann.jussieu.fr

[†]INRIA Paris Rocquencourt EPC BANG. France.

[‡]Departamento de Matemáticas, Universidad Autónoma de Madrid, 28049-Madrid, Spain. Supported by Spanish project MTM2011-24696. Email: fernando.quirós@uam.es

[§]Department of mathematics, Institute of Natural Sciences and MOE-LSC. Shanghai Jiao Tong University, China. Email: tangmin1002@gmail.com

is limited by the competition for space [4], turning the modeling effort towards mechanical concepts, considering tissues as multiphase fluids (the phases could be interstitial water, healthy and tumor cells, extra-cellular matrix ...) [6, 7, 18, 1, 20]. This point of view is now sustained by experimental evidence [19]. The term ‘homeostatic pressure’, coined recently, denotes the lower pressure that prevents cell multiplication by contact inhibition.

In a recent paper [17] the authors explain how asymptotic analysis can link the two main approaches, cell density models and free boundary models, in the context of fluid mechanics for the simplest cell population density model, proposed in [6], in which the cell population density evolves under pressure forces and cell multiplication. The principle of the derivation is to use the stiff limit in the pressure law of state, as treated in several papers; see for instance [2, 14] and the references therein.

Besides mechanical motion induced by pressure, for some types of cancer cells it is important to take into account active motion; see [3, 12, 21]. In the present paper we extend the asymptotic analysis of [17] to a model that includes such an ingredient. We examine the specific form of the Hele-Shaw limit and draw qualitative conclusions on the behaviour of the solutions in terms of regularity and free boundary velocity.

2 Notations and main result

Our model of tumor growth incorporates active motion of cells thanks to a diffusion term,

$$\partial_t n_k - \operatorname{div}(n_k \nabla p_k) - \nu \Delta n_k = n_k G(p_k), \quad (x, t) \in Q := \mathbb{R}^d \times (0, \infty). \quad (2.1)$$

The variable n_k represents the density of tumor cells, and the variable p_k the pressure, which is considered to be given by a homogeneous law (written with a specific coefficient so as to simplify notations later on)

$$p_k(n) = \frac{k}{k-1} n^{k-1}. \quad (2.2)$$

Hence, we are dealing with a porous medium type equation; see [23] for a general reference on such problems. We complement this system with an initial condition that is supposed to satisfy

$$\begin{cases} n_k(x, 0) = n^{ini}(x) > 0, & n^{ini} \in L^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d), \\ p_k^{ini} := \frac{k}{k-1} (n^{ini})^{k-1} \leq P_M. \end{cases} \quad (2.3)$$

In a purely mechanical view, the pressure-limited growth is described by the function G , which satisfies

$$G'(\cdot) < 0 \quad \text{and} \quad G(P_M) = 0, \quad (2.4)$$

for some $P_M > 0$, usually called the homeostatic pressure; see [6, 19].

Many authors use another type of models, namely free boundary problems on the tumor region $\Omega(t)$. Our purpose is to make a rigorous derivation of one of such models from (2.1), (2.2). As it is wellknown, for $\nu = 0$ this is possible in the asymptotics k large. This is connected, in fluid mechanics, to the Hele-Shaw equations; a complete proof of the derivation is provided in [17]. Typically the limit of the cell density is an indicator function for each time $t > 0$, $n_\infty = \mathbb{1}_{\Omega(t)}$, if this is initially true, and the problem is reduced to describing the velocity of the boundary $\partial\Omega(t)$.

Our aim is thus to understand what is the effect of including active motion, that is, $\nu > 0$. We will show that both the density and the pressures have limits, n_∞ and p_∞ , as $k \rightarrow \infty$ that satisfy

$$\partial_t n_\infty - \operatorname{div}(n_\infty \nabla p_\infty) - \nu \Delta n_\infty = n_\infty G(p_\infty). \quad (2.5)$$

Compared with the case $\nu = 0$ considered in [17], a first major difference is that now the cell density n_k is smooth, since equation (2.1) is non-degenerate when $\nu > 0$. Is that translated into more regularity for the limit density? We will show that this is indeed the case. Though the limit density satisfies

$$0 \leq n_\infty \leq 1,$$

it is not an indicator function any more, and its time derivate $\partial_t n_\infty$ is a function, while it is only a measure when $\nu = 0$. As for the pressure, we will establish that we still have

$$n_\infty = 1 \text{ in } \Omega(t) = \{p_\infty(t) > 0\},$$

or in other words $p_\infty \in P_\infty(n_\infty)$, with P_∞ the limiting monotone graph

$$P_\infty(n) = \begin{cases} 0, & 0 \leq n < 1, \\ [0, \infty), & n = 1. \end{cases} \quad (2.6)$$

Furthermore, multiplying equation (2.1) by $p'_k(n_k)$ leads to

$$\partial_t p_k - n_k p'_k(n_k) \Delta p_k - |\nabla p_k|^2 - \nu \Delta p_k = n_k p'_k(n_k) G(p_k) - \nu p''_k(n_k) |\nabla n_k|^2,$$

and for the special case $p_k = \frac{k}{k-1} n_k^{k-1}$ at hand we find

$$\partial_t p_k - (k-1) p_k \Delta p_k - |\nabla p_k|^2 - \nu \Delta p_k = (k-1) p_k G(p_k) - \nu \frac{(k-2) \nabla p_k \cdot \nabla n_k}{n_k}. \quad (2.7)$$

Therefore, the ‘complementary relation’

$$-p_\infty \Delta p_\infty = p_\infty G(p_\infty) - \nu \frac{\nabla p_\infty \cdot \nabla n_\infty}{n_\infty}, \quad (2.8)$$

is expected in the limit. However, ∇p_∞ vanishes unless $p_\infty > 0$, in which case $n_\infty = 1$, therefore $\nabla n_\infty = 0$. Thus, the equation on p_∞ ignores the additional term coming from active motion and reduces to the same Hele-Shaw equation for the pressure that holds when $\nu = 0$, namely

$$p_\infty [\Delta p_\infty + G(p_\infty)] = 0. \quad (2.9)$$

Let us remark that this does not mean that active motion has no effect in the limit. Though the pressure equation is the same one as for the case $\nu = 0$, the free boundary $\partial\Omega(t)$ is not expected to move with the usual Hele-Shaw rule $V = -\nabla p_\infty$, but with a faster one; see Section 7 for a discussion on the speed of the free boundary.

The above heuristic discussion can be made rigorous.

Theorem 2.1 *Let $T > 0$ and $Q_T = \mathbb{R}^d \times (0, T)$. Assume (2.3), (2.4) and that the initial data satisfies $\partial_t n^{ini} \geq 0$. Consider a weak solution (n_k, p_k) of (2.1)–(2.2). Up to extraction of a subsequence, $(n_k, p_k)_k$ converges strongly in $L^p(Q_T)$, $1 \leq p < \infty$, to limits*

$$n_\infty \in C([0, \infty); L^1(\mathbb{R}^d)) \cap L^\infty((0, T); H^1(\mathbb{R}^d)), \quad p_\infty \in L^\infty((0, T); H^1(\mathbb{R}^d)),$$




Figure 1: First steps of the initiation of the free boundary. Results obtained thanks to a discretization of the system (2.1)–(2.2) with $k = 100$ and $\nu = 0.5$. The density n is plotted in solid line whereas the pressure p is represented in dashed line. The pressure p has the same shape as in the classical Hele-Shaw system with growth. However the density n is smoother.

such that $0 \leq n_\infty \leq 1$, $n_\infty(0) = n^{ini}$, $0 \leq p_\infty \leq P_M$, $p_\infty \in P_\infty(n_\infty)$, where P_∞ is the Hele-Shaw monotone graph given in (2.6). Moreover, the pair (n_∞, p_∞) satisfies on the one hand (2.5), and on the other hand the Hele-Shaw type equation

$$\partial_t n_\infty - \Delta p_\infty - \nu \Delta n_\infty = n_\infty G(p_\infty), \quad (2.10)$$

and the complementarity relation (2.9) for almost every $t > 0$, all three equations in the weak sense. The time derivatives of the limit functions satisfy

$$\partial_t n_\infty, \partial_t p_\infty \in \mathcal{M}^1(Q_T), \quad \partial_t n_\infty, \partial_t p_\infty \geq 0.$$

To illustrate this behaviour, we present numerical results obtained thanks to a discretization with finite volume of system (2.1)–(2.2) in the case $k = 100$, $\nu = 0.5$ and with $G(p) = 1 - p$. We display in Figure 1 the first steps of the formation of a tumor which is initially given by a small bump. As expected, we notice that the density n is smooth. The shape of the pressure p at the place where $n = 1$ is similar to the one observed for the classical Hele-Shaw system (see e.g. [17]).

The rest of the paper is organized as follows. We begin in Section 3 with some uniform (in k) a priori estimates which are necessary for strong compactness. Then, in Section 4 we prove the main statements in Theorem 2.1. The most delicate part, establishing (2.9), is postponed to Section 5. After proving uniqueness for the limit problem in Section 6, we end with a final section devoted to discuss further regularity issues and speed of the boundary of the tumor zone.

3 Estimates

To begin with, we gather in the following statement all the a priori estimates that we need later on.

Lemma 3.1 *With the assumptions and notations in Theorem 2.1, the weak solution (n_k, p_k) of (2.1)–(2.2) satisfies*

$$0 \leq n_k \leq \left(\frac{k-1}{k} P_M\right)^{1/(k-1)} \xrightarrow{k \rightarrow \infty} 1, \quad 0 \leq p_k \leq P_M,$$

$$\int_{\mathbb{R}^d} n_k(t) \leq e^{G(0)t} \int_{\mathbb{R}^d} n^{ini}, \quad \int_{\mathbb{R}^d} p_k(t) \leq C e^{G(0)t} \int_{\mathbb{R}^d} n^{ini}.$$

with C a constant independent of k . Furthermore, there exists a uniform (with respect to k) nonnegative constant such that

$$\int_{\mathbb{R}^d} \left(\nu |\nabla n_k|^2 + kn_k^{k-1} |\nabla n_k|^2 + |\nabla p_k|^2 \right) (t) \leq C \left(T, \|n^{ini}\|_{L^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)} \right) \quad \text{for all } t \in (0, T). \quad (3.1)$$

Finally,

$$\partial_t n_k, \partial_t p_k \geq 0, \quad \partial_t n_k \text{ is bounded in } L^\infty((0, T); L^1(\mathbb{R}^d)), \quad \partial_t p_k \text{ is bounded in } L^1(Q_T).$$

Proof. Estimates on n_k and p_k . The $L^\infty(Q)$ bounds are a consequence of standard comparison arguments for (2.1) and (2.7). The $L^\infty((0, T); L^1(\mathbb{R}^d))$ bound for n_k can be obtained by integrating (2.1) over \mathbb{R}^d and then using (2.4). The $L^\infty((0, T); L^1(\mathbb{R}^d))$ bound for p_k now follows from the relation between p_k and n_k .

Estimates on the time derivatives. We introduce the quantity

$$\Sigma(n_k) = n_k^k + \nu n_k, \quad \Sigma'(n_k) = kn_k^{k-1} + \nu. \quad (3.2)$$

The density equation (2.1) is rewritten in terms of this new variable as

$$\partial_t n_k - \Delta \Sigma(n_k) = n_k G(p_k). \quad (3.3)$$

Using the notation $\Sigma_k = \Sigma(n_k)$ and multiplying the above equation by $\Sigma'(n_k)$, we get

$$\partial_t \Sigma_k - \Sigma'_k \Delta \Sigma_k = n_k \Sigma'_k G(p_k). \quad (3.4)$$

Let $w_k = \partial_t \Sigma(n_k)$. Notice that $\text{sign}(\partial_t n_k) = \text{sign}(w_k)$. A straightforward computation yields

$$\partial_t w_k - \Sigma'_k \Delta w_k = \partial_t n_k \Sigma''_k (\Delta \Sigma_k + n_k G(p_k)) + \partial_t n_k \Sigma'_k G(p_k) + \partial_t n_k \Sigma'_k kn_k^{k-1} G'(p_k).$$

By using that $w_k = \Sigma'_k \partial_t n_k$ and $\Sigma'(n_k) = kn_k^{k-1} + \nu \geq \nu > 0$, the right hand side of the above equation can be written in a more handfull way as

$$\partial_t w_k - \Sigma'_k \Delta w_k = w_k \left(\frac{\Sigma''_k}{\Sigma'_k} (\Delta \Sigma_k + n_k G(p_k)) + G(p_k) + kn_k^{k-1} G'(p_k) \right).$$

Since this equation preserves positivity and $\text{sign}(w_k(0)) = \text{sign}(\partial_t n_k^{ini}) \geq 0$, we conclude that $w_k \geq 0$, that is, $\partial_t n_k \geq 0$. The relation between p_k and n_k then immediately yields $\partial_t p_k \geq 0$.

Now that we know that the time derivatives have a sign, bounds for them follow easily. Indeed, using (2.1), we get

$$\|\partial_t n_k(t)\|_{L^1(\mathbb{R}^d)} = \frac{d}{dt} \int_{\mathbb{R}^d} n_k(t) \leq G(0) \|n_k(t)\|_{L^1(\mathbb{R}^d)}.$$

This gives the bound on $\partial_t n_k$ in $L^\infty([0, T]; L^1(\mathbb{R}^d))$. For $\partial_t p_k$ we write

$$\|\partial_t p_k\|_{L^1(Q_T)} = \int_0^T \frac{d}{dt} \left(\int_{\mathbb{R}^d} p_k(t) \right) dt \leq \int_{\mathbb{R}^d} p_k(T).$$

This last expression is uniformly bounded in k .

Estimates on the gradients. We multiply equation (2.1) by n_k , integrate over \mathbb{R}^d and use integration by parts for the diffusion terms,

$$\int_{\mathbb{R}^d} (n_k \partial_t n_k)(t) + \int_{\mathbb{R}^d} (kn_k^{k-1} |\nabla n_k|^2 + \nu |\nabla n_k|^2)(t) = \int_{\mathbb{R}^d} (n_k^2 G(p_k))(t) \leq G(0) \int_{\mathbb{R}^d} n_k^2(t).$$

Since both n_k and $\partial_t n_k$ are nonnegative, we immediately obtain the estimate on the first two terms in (3.1). On the other hand, integrating equation (2.7), we deduce

$$\begin{aligned} \int_{\mathbb{R}^d} \partial_t p_k(t) + (k-2) \int_{\mathbb{R}^d} (|\nabla p_k|^2 + \nu kn_k^{k-3} |\nabla n_k|^2)(t) &= (k-1) \int_{\mathbb{R}^d} (p_k G(p_k))(t) \\ &\leq (k-1) G(0) \int_{\mathbb{R}^d} p_k(t). \end{aligned}$$

Since $\partial_t p_k \geq 0$, we easily obtain the L^2 bound on ∇p_k in (3.1). \square

4 Proof of Theorem 2.1

In this section we prove all the statements in Theorem 2.1 except the one concerning the complementarity relation for the pressure, equation (2.9), whose proof is postponed to the next section.

Strong convergence and bounds. Since the families n_k and p_k are bounded in $W_{\text{loc}}^{1,1}(Q)$, we have strong convergence in L_{loc}^1 both for n_k and p_k . To pass from local convergence to convergence in $L^1(Q_T)$, we need to prove that the mass in an initial strip $t \in [0, 1/R]$ and in the tails $|x| > R$ are uniformly (in k) small if R is large enough. The control on the initial strip is immediate using our uniform, in k and t , bounds for $\|n_k(t)\|_{L^1(\mathbb{R}^d)}$ and $\|p_k(t)\|_{L^1(\mathbb{R}^d)}$. The tails for the densities n_k are controlled using the equation, pretty in the same way as it was done for the case $\nu = 0$; see [17] for the details. The control on the tails of the pressures p_k then follows from the relation between p_k and n_k . Strong convergence in $L^p(Q_T)$ for $1 < p < \infty$ is now a consequence of the uniform bounds for n_k and p_k .

Thanks to the a priori estimates proved above, we also have that $(\nabla n_k)_k$ and $(\nabla p_k)_k$ converge weakly in $L^2(Q_T)$, and

$$0 \leq n_\infty \leq 1, \quad n_\infty, p_\infty \in L^\infty((0, T); H^1(\mathbb{R}^d)), \quad \partial_t n_\infty, \partial_t p_\infty \in \mathcal{M}^1(Q_T), \quad \partial_t n_\infty, \partial_t p_\infty \geq 0.$$

Identification of the limit. To establish equation (2.5) in the distributional sense, we just pass to the limit, by weak-strong convergence, in equation (2.1). On the other hand, using the definition of p_k in (2.2), we have

$$n_k p_k = \frac{k}{k-1} n_k^k = \left(1 - \frac{1}{k}\right)^{1/(k-1)} p_k^{k/(k-1)} \xrightarrow[k \rightarrow \infty]{} p_\infty.$$

Taking the limit $k \rightarrow \infty$, we deduce the monotone graph property

$$p_\infty(1 - n_\infty) = 0. \tag{4.1}$$

In order to show the equivalence of (2.10) and (2.5), we need to prove that $\nabla p_\infty = n_\infty \nabla p_\infty$. This is seen to be equivalent to $p_\infty \nabla n_\infty = 0$ by using the Leibnitz rule in $H^1(\mathbb{R}^d)$ for (4.1). To prove the latter identity, we first write

$$p_k \nabla n_k = \frac{k}{k-1} n_k^k \nabla n_k = \frac{\sqrt{k}}{k-1} n_k^{(k+1)/2} (\sqrt{k} n_k^{(k-1)/2} \nabla n_k).$$

From estimate (3.1), the term between parentheses is uniformly bounded in $L^2(Q_T)$ and since $(n_k)_k$ is uniformly (in k) bounded in $L^\infty(Q_T)$, we conclude that

$$\lim_{k \rightarrow \infty} \|p_k \nabla n_k\|_{L^2(Q_T)} = 0.$$

We deduce then from the strong convergence of $(p_k)_k$ and the weak convergence of $(\nabla n_k)_k$ that

$$p_\infty \nabla n_\infty = 0, \tag{4.2}$$

as desired.

Time continuity and initial trace. Time continuity for the limit density n_∞ follows from the monotonicity and the equation, as in the case $\nu = 0$. Once we have continuity, the identification of the initial trace will follow from the equation for n_k , letting first $k \rightarrow \infty$ and then $t \rightarrow 0$; see [17] for the details.

Remark. Since $p_\infty \geq 0$, (4.2) implies that

$$\nabla p_\infty \cdot \nabla n_\infty = 0. \tag{4.3}$$

5 The equation on p_∞

In this section we give a rigorous derivation of equation (2.9), which is the most delicate point in the proof of Theorem 2.1.

(i) Our first goal is to establish that, in the weak sense,

$$p_\infty \Delta p_\infty + p_\infty G(p_\infty) \leq 0. \tag{5.1}$$

Thanks to (4.2) and (4.3), this is equivalent to proving that

$$p_\infty \Delta (p_\infty + \nu n_\infty) + p_\infty G(p_\infty) \leq 0. \tag{5.2}$$

In order to prove the latter inequality, we follow an idea of [17] and use a time regularization method *à la Steklov*. To this aim, we introduce a regularizing kernel $\omega_\varepsilon(t) \geq 0$ with compact support of length ε .

Let $n_{k,\varepsilon} = n_k * \omega_\varepsilon$. From equation (2.1), we deduce

$$\partial_t n_{k,\varepsilon} - \Delta \omega_\varepsilon * (n_k^k + \nu n_k) = (n_k G(p_k)) * \omega_\varepsilon. \tag{5.3}$$

Then, for fixed $\varepsilon > 0$, $\Delta \omega_\varepsilon * (n_k^k + \nu n_k)$ is bounded in $L^q(Q_T)$ for all $q \geq 1$. Thus, we can extract a subsequence such that $(\nabla \omega_\varepsilon * (n_k^k + \nu n_k))_k$ converges strongly in $L^2(Q_T)$. Since we have strong convergence of $(n_k^k + \nu n_k)_k$ towards $p_\infty + \nu n_\infty$, we deduce that the strong limit of $(\nabla \omega_\varepsilon * (n_k^k + \nu n_k))_k$ is equal to $\nabla \omega_\varepsilon * (p_\infty + \nu n_\infty)$.

Multiplying equation (5.3) by p_k , we have

$$p_k \partial_t n_{k,\varepsilon} = p_k \Delta (n_k^k * \omega_\varepsilon + \nu n_{k,\varepsilon}) + p_k ((n_k G(p_k)) * \omega_\varepsilon).$$

We can pass to the limit $k \rightarrow \infty$ to get

$$\lim_{k \rightarrow \infty} p_k \partial_t n_{k,\varepsilon} = p_\infty \Delta (\omega_\varepsilon * (p_\infty + \nu n_\infty)) + p_\infty ((n_\infty G(p_\infty)) * \omega_\varepsilon).$$

To determine the sign, we decompose the left hand side term, divided by the harmless factor $k/(k-1)$, as

$$\int_{\mathbb{R}} n_k^{k-1}(t) \partial_t n_k(s) \omega_\varepsilon(t-s) ds = \underbrace{\int_{\mathbb{R}} n_k^{k-1}(s) \partial_t n_k(s) \omega_\varepsilon(t-s) ds}_{\mathcal{A}_k} + \underbrace{\int_{\mathbb{R}} (n_k^{k-1}(t) - n_k^{k-1}(s)) \partial_t n_k(s) \omega_\varepsilon(t-s) ds}_{\mathcal{B}_k}.$$

On the one hand we have

$$\mathcal{A}_k = \frac{1}{k} \int_{\mathbb{R}} \partial_t n^k(s) \omega_\varepsilon(t-s) ds \rightarrow 0 \quad \text{when } k \rightarrow \infty.$$

As for \mathcal{B}_k , we recall that $\partial_t n_k \geq 0$ provided $\partial_t n^{ini} \geq 0$; see Lemma 3.1. Thus, for $s > t$ we have $n_k^{k-1}(t) - n_k^{k-1}(s) \leq 0$. Then, choosing ω_ε such that $\text{supp } \omega_\varepsilon \subset \mathbb{R}_-$, we deduce that $\mathcal{B}_k \leq 0$, which yields

$$p_\infty \Delta(\omega_\varepsilon * (p_\infty + \nu n_\infty)) + p_\infty (n_\infty G(p_\infty) * \omega_\varepsilon) \leq 0.$$

It remains to pass to the limit $\varepsilon \rightarrow 0$ in the regularization process. We can pass to the limit in the weak formulation since we already know that $\nabla p_\infty \in L^2(Q_T)$. Then, using (4.1), we get the inequality (5.2) and thus (5.1).

(ii) Our second purpose is to establish the other inequality, namely

$$p_\infty \Delta p_\infty + p_\infty G(p_\infty) \geq 0. \tag{5.4}$$

To prove it, we multiply equation (2.7) by a nonnegative test function $\phi(x, t)$ and integrate, and obtain

$$\begin{aligned} \iint_{Q_T} \phi (p_k \Delta p_k + p_k G(p_k) - \nu \frac{k-2}{k-1} \frac{\nabla p_k \cdot \nabla n_k}{n_k}) \\ = \frac{1}{k-1} \iint_{Q_T} [\phi (\partial_t p_k - |\nabla p_k|^2) + \nu \nabla \phi \cdot \nabla p_k]. \end{aligned}$$

From the proved bounds, the right hand side of the above equation converges to 0 as $k \rightarrow \infty$. We can use integration by parts and rewrite the left hand side as

$$\iint_{Q_T} \left(\phi p_k G(p_k) - p_k \nabla \phi \cdot \nabla p_k - \phi |\nabla p_k|^2 - \phi \nu \frac{k(k-2)}{k-1} n_k^{k-3} |\nabla n_k|^2 \right).$$

Since the last term is nonpositive, we obtain that

$$\liminf_{k \rightarrow \infty} \iint_{Q_T} (\phi p_k G(p_k) - p_k \nabla \phi \cdot \nabla p_k - \phi |\nabla p_k|^2) \geq 0.$$

From weak-strong convergence in products, or convexity inequalities in the weak limit, we finally conclude

$$\iint_{Q_T} (\phi p_\infty G(p_\infty) - p_\infty \nabla \phi \cdot \nabla p_\infty - \phi |\nabla p_\infty|^2) \geq 0.$$

This is the weak formulation of (5.4).

Remark. A careful inspection of the proof of (5.4) shows that (2.8) holds if and only if ∇p_k converges strongly in $L^2(Q_T)$ and $kn_k^{k-3} |\nabla n_k|^2$ converges weakly to 0 locally in $L^1(Q)$. Since we have proved (2.8), we conclude that we have the two mentioned convergence results.

6 Uniqueness for the limit model

In this section we prove that the limit problem (2.10) admits at most one solution. We will adapt Hilbert's duality method in the spirit of [17].

Theorem 6.1 *Let $T > 0$, $\nu > 0$. There is a unique pair (n, p) of functions in $L^\infty([0, T]; L^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d))$, $n \in C([0, T]; L^1(\mathbb{R}^d))$, $n(0) = n^{ini}$, $p \in P_\infty(n)$, satisfying (2.10) in the sense of distributions and such that $\nabla n, \nabla p \in L^2(Q_T)$, $\partial_t n, \partial_t p \in \mathcal{M}^1(Q_T)$.*

Proof. Let us consider two solutions (n_1, p_1) and (n_2, p_2) . Then for any test function ϕ with $\phi \in W^{2,2}(Q_T)$ and $\partial_t \phi \in L^2(Q_T)$, we have

$$\iint_{Q_T} \left((n_1 - n_2) \partial_t \phi + (p_1 - p_2 + \nu(n_1 - n_2)) \Delta \phi + (n_1 G(p_1) - n_2 G(p_2)) \phi \right) = 0, \quad (6.1)$$

which can be rewritten as

$$\iint_{Q_T} (\nu(n_1 - n_2) + p_1 - p_2) (A \partial_t \phi + \Delta \phi + AG(p_1) \phi - B \phi) = 0, \quad (6.2)$$

where

$$\begin{aligned} 0 \leq A &= \frac{n_1 - n_2}{\nu(n_1 - n_2) + p_1 - p_2} \leq \frac{1}{\nu}, \\ 0 \leq B &= -n_2 \frac{G(p_1) - G(p_2)}{\nu(n_1 - n_2) + p_1 - p_2} \leq \kappa, \end{aligned}$$

for some nonnegative constant κ . To arrive to these bounds on A we set $A = 0$ when $n_1 = n_2$, even if $p_1 = p_2$. Since A can vanish, we use a smoothing argument by introducing the regularizing sequences $(A_n)_n$, $(B_n)_n$ and $(G_{1,n})_n$ such that

$$\begin{aligned} \|A - A_n\|_{L^2(Q_T)} &< \alpha/n, & 1/n < A_n \leq 1, \\ \|B - B_n\|_{L^2(Q_T)} &< \beta/n, & 0 \leq B_n \leq \beta_2, & \|\partial_t B_n\|_{L^1(Q_T)} \leq \beta_3, \\ \|G_{1,n} - G(p_1)\|_{L^2(Q_T)} &\leq \delta/n, & |G_{1,n}| < \delta_2, & \|\nabla G_{1,n}\|_{L^2(Q_T)} \leq \delta_3, \end{aligned}$$

for some nonnegative constants $\alpha, \beta, \beta_2, \beta_3, \delta, \delta_2, \delta_3$.

Given any arbitrary smooth function ψ compactly supported, we consider the solution ϕ_n of the backward heat equation

$$\begin{cases} \partial_t \phi_n + \frac{1}{A_n} \Delta \phi_n + G_{1,n} \phi_n - \frac{B_n}{A_n} \phi_n = \psi & \text{in } Q_T, \\ \phi_n(T) = 0. \end{cases} \quad (6.3)$$

The coefficient $1/A_n$ is continuous, positive and bounded below away from zero. Then the equation satisfied by ϕ_n is parabolic. Hence ϕ_n is smooth and since ψ is compactly supported, we have that $\phi_n, \Delta \phi_n$ and therefore $\partial_t \phi_n$ are L^2 -integrable. Therefore, we can use ϕ_n as a test function in (6.2). Then, by the definition of A , we have

$$\iint_{Q_T} (n_1 - n_2) \psi = \iint_{Q_T} (\nu(n_1 - n_2) + p_1 - p_2) A \psi.$$

Inserting (6.3) and subtracting (6.2), we obtain

$$\iint_{Q_T} (n_1 - n_2)\psi = I_{1n} + I_{2n} + I_{3n},$$

where

$$\begin{aligned} I_{1n} &= \iint_{Q_T} (\nu(n_1 - n_2) + p_1 - p_2) \left(\left(\frac{A}{A_n} - 1 \right) (\Delta\phi_n - C_n\phi_n) \right), \\ I_{2n} &= \iint_{Q_T} (\nu(n_1 - n_2) + p_1 - p_2) (B - B_n)\phi_n, \\ I_{3n} &= \iint_{Q_T} (n_1 - n_2) (G_{1,n} - G(p_1))\phi_n. \end{aligned}$$

The convergence towards 0 of the terms I_{in} , $i = 1, 2, 3$ is now a consequence on some estimates on the test functions ϕ_n which are gathered in Lemma 6.2 below. Indeed, applying the mentioned estimates and Cauchy-Schwarz inequality we have

$$\begin{aligned} I_{1n} &\leq K \|(A - A_n)/\sqrt{A_n}\|_{L^2(Q_T)} \leq K\sqrt{n}\|A - A_n\|_{L^2(Q_T)} \leq K\alpha/\sqrt{n}, \\ I_{2n} &\leq K\|B - B_n\|_{L^2(Q_T)} \leq K\gamma/n, \\ I_{3n} &\leq K\delta/n, \end{aligned}$$

(in all the computations, K denotes various nonnegative constants). Then letting $n \rightarrow \infty$, we conclude that

$$\iint_{Q_T} (n_1 - n_2)\psi = 0,$$

for any smooth function ψ compactly supported, hence $n_1 = n_2$. It is then obvious, thanks to (6.1), that $p_1 = p_2$. \square

Lemma 6.2 *Under the assumptions of Theorem 6.1, we have the uniform bounds, only depending on T and ψ ,*

$$\|\phi_n\|_{L^\infty(Q_T)} \leq \kappa_1, \quad \sup_{0 \leq t \leq T} \|\nabla\phi_n(t)\|_{L^2(\mathbb{R}^d)} \leq \kappa_2, \quad \|1/\sqrt{A_n}(\Delta\phi_n - B_n\phi_n)\|_{L^2(Q_T)} \leq \kappa_3.$$

Proof. The first bound is a consequence of the maximum principle on (6.3). Then multiplying (6.3) by $\Delta\phi_n - B_n\phi_n$ and integrating on \mathbb{R}^d , we get

$$\begin{aligned} & -\frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^d} |\nabla\phi_n(t)|^2 - \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^d} B_n\phi_n^2(t) + \int_{\mathbb{R}^d} \frac{1}{A_n} |\Delta\phi_n - B_n\phi_n|^2(t) + \frac{1}{2} \int_{\mathbb{R}^d} (\partial_t B_n\phi_n^2)(t), \\ & = \int_{\mathbb{R}^d} \left(G_{1,n} |\nabla\phi_n|^2 + \phi_n \nabla\phi_n \cdot \nabla G_{1,n} + B_n G_{1,n} \phi_n^2 + (\Delta\psi - B_n\psi)\phi_n \right)(t). \end{aligned}$$

After an integration in time on $[t, T]$, we deduce

$$\frac{1}{2} \|\nabla\phi_n(t)\|_{L^2(\mathbb{R}^d)}^2 + \int_t^T \int_{\mathbb{R}^d} \frac{1}{A_n} |\Delta\phi_n - B_n\phi_n|^2 \leq K \left(1 - t + \int_t^T \|\nabla\phi_n(s)\|_{L^2(\mathbb{R}^d)} ds \right),$$

where we use the bounds on $\nabla G_{1,n}$ and $\partial_t B_n$ by construction of the regularization. We conclude by applying Gronwall's Lemma. \square

7 Further regularity and velocity of the free boundary

Remember that both p_∞ and n_∞ belong to $H^1(\mathbb{R}^d)$ for almost every $t > 0$. This regularity cannot be improved, because there are jumps in the gradients of both p_∞ and n_∞ at the free boundary. As a consequence, their laplacians are not functions, but measures. However, these singularities cancel in the combination $\Sigma_\infty = p_\infty + \nu n_\infty$, as we will see now.

Lemma 7.1 *With the assumptions of Theorem 2.1, the quantity Σ_∞ belongs to $L^2((0, T); H^2(\mathbb{R}^d))$ for all $T > 0$ and we have the estimate*

$$\iint_{Q_T} (\Delta \Sigma_\infty)^2 \leq C(T).$$

Proof. We recall the definition of Σ_k in (3.2). Since $\nabla \Sigma_k = n_k \nabla p_k + \nu \nabla n_k$, estimate (3.1) yields that for all $0 < t \leq T$,

$$\int_{\mathbb{R}^d} |\nabla \Sigma_k(t)|^2 \leq C(T).$$

We now multiply the equation (3.4) by $\Delta \Sigma_k$, and integrate in Q_T , $0 < T < \infty$, to obtain, using that $\Sigma'_k > \nu$ and the fact that both n_k and $G(p_k)$ are nonnegative,

$$\iint_{Q_T} (\Delta \Sigma_k)^2 \leq \frac{1}{2} \int_{\mathbb{R}^d} |\nabla \Sigma_k|^2(0) + C(T).$$

The result follows directly. □

This implies in particular that in the limit $\Sigma_\infty(\cdot, t) \in H^2(\mathbb{R}^d)$ for almost every $t > 0$. Hence, the size of the jump (downwards) of ∇p_∞ at the free boundary coincides with the size of the jump (upwards) of $\nu \nabla n_\infty$ there.

Concerning the time regularity, the limit equation for the density (3.3), now tells us that $\partial_t n_\infty \in L^2(Q_T)$. Hence $n_\infty \in H^1(Q_T)$. We do not have a similar property for the pressure (think of the situation when two tumors meet).

Our last goal is to derive formally an asymptotic value for the free boundary speed in a particular example. Let $\Omega(t)$ denote, as before, the space filled by the tumor at time t . We notice that n_∞ solves

$$\partial_t n_\infty = \nu \Delta n_\infty + G(0)n_\infty, \quad x \in \mathbb{R}^d \setminus \Omega(t), \quad t > 0,$$

with boundary conditions

$$n_\infty = 1, \quad \nu \partial_n n_\infty = \partial_n p_\infty, \quad x \in \partial \Omega(t), \quad t > 0.$$

If $\Omega(t)$ were known, the problem would be overdetermined. This is precisely what fixes the dynamics of the free boundary. Let us assume that the tumor is a ball centered at the origin,

$$\Omega(t) = \{x : p_\infty(x, t) > 0\} = \{x : n_\infty(x, t) = 1\} = B_{R(t)}(0).$$

We look for a solution which is spherically symmetric $n_\infty(r, t)$, $p_\infty(r, t)$. We set $\sigma = R'(t)$. In opposition to other models of tumor growth (see [22] for instance), here there are no radial solutions




Figure 2: Shape of the traveling waves obtained thanks to a numerical discretization of the system (2.1)–(2.2) with $k = 100$ and $\nu = 0.5$ (left) or $\nu = 0$ (right) for the same initial data and the same final time. The density n is plotted in line whereas the pressure is represented in dashed line. We notice the regularity of n in the case $\nu = 0.5$, whereas it has a jump at the interface when $\nu = 0$. Also the free boundary moves faster when active motion is present.

with constant speed. However, following [17] Appendix A, we expect our solution to behave for large times as a one dimensional traveling wave (with constant speed).

In order to analyze the expected asymptotic constant speed, we set $n_R(r - \sigma t) = n_\infty(r, t)$ and $p_R(r - \sigma t) = p_\infty(r, t)$. Introducing this ansatz in equation (2.10), we obtain

$$-\sigma n'_R = p''_R + \frac{d-1}{r} p'_R + \nu n''_R + \nu \frac{d-1}{r} n'_R + n_R G(p_R). \quad (7.1)$$

On $\mathbb{R}^d \setminus \Omega(t)$, we have $p_\infty = 0$, then integrating (7.1) in $(R(0), \infty)$, we get

$$\sigma n_R(R(0)) = -\nu n'_R(R(0)^+) + \nu(d-1) \int_{R(0)}^{\infty} \frac{n'_R}{r} dr + G(0) \int_{R(0)}^{\infty} n_R dr.$$

In a one dimensional setting ($d = 1$) and using the boundary relation at the interface of $\Omega(0)$, we deduce

$$\sigma = -p'_R(R(0)^-) + G(0) \int_{R(0)}^{\infty} n_R(r) dr. \quad (7.2)$$

We recall that for the classical Hele-Shaw model without active motion (i.e. $\nu = 0$), the traveling velocity is $\sigma_0 = -p'_R(R(0)^-)$. Since $n_R(R(0)) = 1$ and n_R is continuous and nonnegative, we have $\int_{R(0)}^{\infty} n_R(r) dr > 0$. Then we conclude from equation (7.2) that $\sigma > \sigma_0$.

We can do a more precise computation confirming the above statement for the one-dimensional case. From the complementarity relation (2.9), we have $-p''_R = G(p_R)$ on $\Omega(0)$. Multiplying this latter equation by p'_R and integrating on $(0, R(0))$, we deduce

$$p'_R(R(0)^-)^2 = 2 \int_0^{R(0)} p'_R G(p_R) dr.$$

In the center of the tumor, we expect a maximal packing of the cells. Therefore, we have the boundary conditions

$$\lim_{r \rightarrow 0} p_R(r) = P_M, \quad \lim_{r \rightarrow 0} p'_R(r) = 0.$$

Since $p''_R = -G(p_R) \leq 0$, we deduce that $p'_R < 0$ and we can make the change of variable

$$p'_R(R(0)^-)^2 = 2 \int_0^{R(0)} p'_R G(p_R) dr = 2 \int_0^{P_M} G(q) dq.$$

The quantity $\sigma_0 = \sqrt{2 \int_0^{P_M} G(q) dq}$ is the traveling velocity for a tumor spheroid in the case $\nu = 0$; see Appendix A.1 of [17]. Combining this with (7.2), we deduce that the growth of the tumor is faster with active motion than in the case $\nu = 0$.

In Figure 2, we display numerical simulations obtained from a discretization with a finite volume scheme of system (2.1)–(2.2) for $k = 100$. The left picture presents the result for $\nu = 0.5$, and the right for $\nu = 0$ (i.e. without active motion). We use the growth function $G(p) = 1 - p$ and the results in both cases with the same initial data and at final time $t = 10$. We notice that in the case $\nu = 0.5$ the density function is smooth and the domain occupied by the tumor is larger than in the case without active motion, which suggests as explained above a faster invasion speed.

References

- [1] Bellomo, N.; Li, N. K.; Maini, P. K. *On the foundations of cancer modelling: selected topics, speculations, and perspectives*. Math. Models Methods Appl. Sci. 18 (2008), no. 4, 593–646.
- [2] Bénilan, Ph.; Igbida, N. *La limite de la solution de $u_t = \Delta_p u^m$ lorsque $m \rightarrow \infty$* . C. R. Acad. Sci. Paris Sér. I Math. 321 (1995), no. 10, 1323–1328.
- [3] Betteridge, R.; Owen, M. R.; Byrne, H. M.; Alarcón, T.; Maini, P. K. *The impact of cell crowding and active cell movement on vascular tumour growth*. Netw. Heterog. Media 1 (2006), no. 4, 515–535.
- [4] Brú, A.; Albertos, S.; Subiza, J. L.; Asenjo, J. A.; Brœ, I. *The universal dynamics of tumor growth*. Biophys. J. 85 (2003), no. 5, 2948–2961.
- [5] Byrne, H. M.; Chaplain, M. A. *Growth of necrotic tumors in the presence and absence of inhibitors*. Math. Biosci. 135 (1996), no. 15, 187–216.
- [6] Byrne, H. M.; Drasdo, D. *Individual-based and continuum models of growing cell populations: a comparison*. J. Math. Biol. 58 (2009), no. 4-5, 657–687.
- [7] Byrne, H. M.; Preziosi, L. *Modelling solid tumour growth using the theory of mixtures*. Math. Med. Biol. 20 (2003), no. 4, 341–366.
- [8] Chaplain, M. A. J. *Avascular growth, angiogenesis and vascular growth in solid tumours: the mathematical modeling of the stages of tumor development*. Math. Comput. Modeling 23 (1996), no. 6, 47–87.

- [9] Ciarletta, P.; Foret, L.; Ben Amar, M. *The radial growth phase of malignant melanoma: multiphase modelling, numerical simulations and linear stability analysis*. J. R. Soc. Interface 8 (2011) no. 56, 345–368.
- [10] Cui, S. *Formation of necrotic cores in the growth of tumors: analytic results*. Acta Math. Sci. Ser. B Engl. Ed. (2006), no. 4, 781–796.
- [11] Cui, S.; Escher, J. *Asymptotic behaviour of solutions of a multidimensional moving boundary problem modeling tumor growth*. Comm. Partial Differential Equations 33 (2008), no. 4–6, 636–655.
- [12] Drasdo, D.; Hoehme, S. *Modeling the impact of granular embedding media, and pulling versus pushing cells on growing cell clones*. New J. Phys. 14 (2012) 055025 (37pp).
- [13] Friedman, A.; Hu, B. *Stability and instability of Liapunov-Schmidt and Hopf bifurcation for a free boundary problem arising in a tumor model*. Trans. Am. Math. Soc. 360 (2008), no. 10, 5291–5342.
- [14] Gil, O.; Quirós, F. *Boundary layer formation in the transition from the porous media equation to a Hele-Shaw flow*. Ann. Inst. H. Poincaré Anal. Non Linéaire 20 (2003), no. 1, 13–36.
- [15] Greenspan, H. P. *Models for the growth of a solid tumor by diffusion*. Stud. Appl. Math. 51 (1972), no. 4, 317–340.
- [16] Lowengrub, J. S.; Frieboes H. B.; Jin, F.; Chuang, Y.-L.; Li, X.; Macklin, P.; Wise, S. M.; Cristini, V. *Nonlinear modelling of cancer: bridging the gap between cells and tumours*. Nonlinearity 23 (2010), no. 1, R1–R91.
- [17] Perthame, B.; Quirós, F.; Vázquez, J. L. *The Hele-Shaw asymptotics for mechanical models of tumor growth*. Arch. Ration. Mech. Anal., to appear.
- [18] Preziosi, L.; Tosin, A. *Multiphase modelling of tumour growth and extracellular matrix interaction: mathematical tools and applications*. J. Math. Biol. 58 (2009), no. 4-5, 625–656.
- [19] Ranft, J.; Basana, M.; Elgeti, J.; Joanny, J.-F.; Prost, J.; Jülicher, F. *Fluidization of tissues by cell division and apoptosis*. Proc. Natl. Acad. Sci. USA (2010), no. 49, 20863–20868.
- [20] Roose, T.; Chapman, S. J.; Maini, P. K. *Mathematical models of avascular tumor growth*. SIAM Rev. 49 (2007), no. 2, 179–208.
- [21] Saut, O.; Lagaert, J.-B.; Colin, T.; Fathallah-Shaykh, H. M. *A multilayer grow-or-go model for GBM: effects of invasive cells and anti-angiogenesis on growth*. Preprint 2012.
- [22] Tang, M.; Vauchelet, N.; Cheddadi, I.; Vignon-Clementel, I.; Drasdo, D.; Perthame, B. *Composite waves for a cell population system modelling tumor growth and invasion*. Chin. Ann. Math. Ser. B 34 (2013), no. 2, 295–318.
- [23] Vázquez, J. L. “The porous medium equation. Mathematical theory”. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, Oxford, 2007. ISBN: 978-0-19-856903-9.