



# **In situ activity of NAC11-7 roseobacters in coastal waters off the Chesapeake Bay based on ftsZ expression.**

Daohong Yao, Alison Buchan, Marcelino T Suzuki

## **► To cite this version:**

Daohong Yao, Alison Buchan, Marcelino T Suzuki. In situ activity of NAC11-7 roseobacters in coastal waters off the Chesapeake Bay based on ftsZ expression.. *Environmental Microbiology*, 2011, 13 (4), pp.1032-41. 10.1111/j.1462-2920.2010.02408.x . hal-01062916v1

**HAL Id: hal-01062916**

**<https://hal.sorbonne-universite.fr/hal-01062916v1>**

Submitted on 11 Sep 2014 (v1), last revised 13 Sep 2014 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

3        ***In situ* Activity of NAC11-7 Roseobacters in Coastal Waters off the**  
4        **Chesapeake Bay based on *ftsZ* Expression**

5        Daohong Yao<sup>1</sup>, Alison Buchan<sup>2</sup> and Marcelino T. Suzuki<sup>1,3\*</sup>

6        <sup>1</sup>Chesapeake Biological Laboratory, University of Maryland Center for Environmental Science.  
7        Solomons, MD, USA.

8        <sup>2</sup> Department of Microbiology, University of Tennessee Knoxville, Knoxville, TN, USA.

9        **\* to whom correspondence should be addressed:**

10        Université Pierre et Marie Curie (Paris VI) Laboratoire Arago and CNRS UMR 7621,  
11        Laboratoire d’Oceanographie Microbienne. Avenue du Fontaulé, Banyuls-sur-Mer 66650 France

12        **Electronic mail address:** suzuki@obs-banyuls.fr

13        **Tel:** +33(0)4 3019 2401 **FAX :** +33(0)4 6888 7398

14        **Running Title:** *In situ* expression of roseobacters *ftsZ*

---

<sup>3</sup> Current Address: Université Pierre et Marie Curie (Paris 6) Laboratoire Arago and CNRS UMR 7621, Laboratoire  
d’Oceanographie Microbienne, Banyuls-sur-Mer, France

## Summary

Determining *in situ* growth rates for specific bacterioplankton is of critical importance to understanding their contributions to energy and matter flow in the Ocean. Quantifying expression of genes central to cell division is a plausible approach for obtaining these measurements. In order to test this approach's assumptions, a quantitative PCR assay targeting the cell division gene *ftsZ* in the ubiquitous, but not yet cultivated, NAC11-7 group of the Rhodobacterales order of marine bacteria was developed. *ftsZ* genes and their corresponding mRNAs were measured in diel *in situ* samples and in parallel on-deck incubations. Strong correlations between *ftsZ* expression and gene abundance ( $R^2=0.62$ ), and between *ftsZ* expression and water temperature ( $R^2=0.73$ ) were observed *in situ*. Rapid and cyclical changes of NAC11-7 *ftsZ* gene copies suggested that populations from two different end-member water types were sampled. This was supported by distinct *ftsZ* gene expression levels in populations associated with these water types. An outlier to this trend occurred at a single time point (9:00), and was remarkably consistent with a concomitant peak in *ftsZ* expression in on-deck incubations, suggesting synchronous population growth. To our knowledge this is the first evidence for synchronous populations for a heterotrophic bacterioplankton group in the ocean.

## Introduction

Marine bacterioplankton play pivotal roles in oceanic food webs and the cycling of organic matter, which in turn influence global carbon pools (Azam and Malfatti, 2007; Wohlers et al., 2009). In recent years, the application of molecular, genomic, and transcriptomic techniques has provided us with vast datasets, greatly expanding our knowledge of bacterioplankton phylogenetic diversity, their environmental distributions, and metabolic capabilities (for reviews see Rappe and Giovannoni, 2003; Delong, 2009). Despite this knowledge, challenges still exist in understanding the ecological functions of **specific** bacterioplankton. Among these challenges is a lack of measurements of *in situ* growth rates of bacterioplankton at high phylogenetic resolution. This information is essential for elucidating the contributions of these ecotypes to biogeochemical cycles critical to the ecosystem.

The most commonly used approaches for determining growth rates of heterotrophic marine bacterioplankton are indirect and rely on measurements of bacterial production. Growth rates are often calculated from bacterioplankton production, which are usually estimated from the incorporation of  $^3\text{H}$ -thymidine or  $^3\text{H}$ -leucine (Fuhrman and Azam, 1982; Kirchman et al., 1985; Simon and Azam, 1989), or non-radioactive bromodeoxyuridine (Steward and Azam, 1999). These methods regard the entire bacterial community as a “black box”, provide community-level activity measurements, and therefore do not provide information on the production and growth rates of specific bacterioplankton. While the combination of microautoradiography with fluorescence *in situ* hybridization (i.e. Micro-FISH) allows the assessment of specific bacterioplankton activities at relatively narrow phylogenetic levels (i.e. Cottrell and Kirchman, 2003; Alonso and Pernthaler, 2005; Margolin, 2005), these experiments suffer from issues inherent to confinement and tracer-based studies (i.e. bottle effects).

A plausible approach to measure *in situ* activity, and perhaps growth rates of specific bacterioplankton, involves quantifying the expression (mRNAs) of genes encoding proteins involved in cell division. The idea behind such approach would be to determine direct relationships between *in situ* expression of protein-encoding genes and growth rates of specific bacterioplankton, without additions or incubations. The rationale is analogous to that of the measurement of the frequency of dividing cells (Hagstrom et al., 1979), except that the measurement could be made at specific phylogenetic levels, and earlier in the cell cycle (i.e. before a clear septal ring is formed), thus decreasing problems associated with preferential protistan grazing of dividing cells (Sherr et al., 1992). Requirements for successfully using a gene transcript tracking approach for **instantaneous** (i.e. at any time of the day) measurements of growth would be that populations of targeted organisms do not divide synchronously, and that the regulation of cell division protein synthesis occurs at the transcriptional level. In cases where the population divides synchronously, growth rates estimates are possible, but would not be instantaneous and would require measurements throughout a diel cycle (i.e. Carpenter and Chang, 1988; Chang and Carpenter, 1988; Holtzendorff et al., 2002).

FtsZ initiates cell division of most prokaryotes by self-assembling into a membrane-associated Z-ring structure and by recruiting other proteins to form the cell division septum (Margolin, 2005; Dajkovic and Lutkenhaus, 2006; Osawa and Erickson, 2006). In synchronized populations of *Caulobacter crescentus*, a model organism for the study of bacterial cell division, the transcription of *ftsZ* is controlled by two regulators: DnaA promotes *ftsZ* transcription, while CtrA is its repressor, ensures that cell division is coordinated with DNA replication (Hottes et al., 2005; McAdams and Shapiro, 2009). Levels of expression of *ftsZ* genes was also found to vary during the cell cycle in synchronized *Prochlorococcus* populations in the Gulf of Aqaba, with

the maxima of expression at the replication phase (S phase; Holtzendorff et al., 2001; Holtzendorff et al., 2002). These authors suggested that transcriptional regulation of *ftsZ* could be a major factor triggering the observed synchronized cell division of *Prochlorococcus* populations. Finally, the sequence of FtsZ protein is highly conserved in most bacteria and its phylogeny has good congruence to 16S rRNA phylogeny (Vaughan et al., 2004), allowing putative identification of genes uncovered directly from the environment.

In this study, we investigated *in situ* expression of *ftsZ* genes putatively belonging to the NAC11-7 subclade of the *Roseobacter* group of marine bacteria. Members of the NAC11-7 clade have been implicated in DMSP degradation (Zubkov et al., 2001; Buchan et al., 2005) and the genome sequence of a NAC11-7 representative (HTCC 2255) suggests members of this subclade might be aerobic anoxygenic phototrophs, since the genome of HTCC 2255 contains a *roseobacter* type *pufM* gene. We quantified *ftsZ* expression of this group over a diel cycle to determine whether evidence exists for synchronicity in cell division. Gene expression was measured as the ratio of *ftsZ* mRNA to *ftsZ* gene copies using quantitative real time PCR (qPCR) and reverse transcription qPCR (RT-qPCR), respectively, in samples collected from a station off the Chesapeake Bay and in a parallel on-board incubation.

## Results

### ***Rhodobacterales ftsZ genes from coastal waters off the Chesapeake Bay.***

Degenerate primers targeting *ftsZ* from the order Rhodobacterales were designed based on conserved regions identified in alignments of representative FtsZ sequences obtained from GenBank. A clone library was generated from fragments amplified from coastal Atlantic Ocean surface waters and 74 *ftsZ* clones were analyzed. Phylogenetic reconstruction of these sequenced clones as well as representative reference sequences revealed three interesting groups (Figure 1):

a prevalent, yet unidentified group (designated Group I) containing 38 sequences (51% of the *ftsZ* clones); a second unidentified group (Group II) with 31 sequences (42 % of the clones); and a group represented by 5 sequences that were most closely related to the *FtsZ* sequence from strain HTCC 2255 (Figure 1). 16S rRNA gene phylogeny places strain HTCC 2255 in the NAC11-7 group of the roseobacters (Buchan et al., 2005). Given the previously reported global distributions and abundances of NAC11-7 representatives, and their putative photoheterotrophic metabolism this group was selected for further investigation.

***In situ diel expression of NAC11-7 ftsZ genes.***

NAC11-7 *ftsZ* mRNA and DNA copies were monitored in surface waters over a 24 h period at a single station located at the mouth of the Chesapeake Bay. Gene copies, and more remarkably, gene expression, showed rapid and sometimes large (c.a. 4-fold) changes with time (i.e. from 9:00 to 12:00 and from 18:00 to 21:00), suggesting distinct populations (i.e. patches) of NAC11-7 may have been sampled at different times, as a result of water advection (Figure 2). Interestingly, *ftsZ* gene expression followed closely *ftsZ* gene abundance trends, except at the last two sampling times. A regression analysis between *ftsZ* gene copies and gene expression using individual replicates showed significant correlation ( $R^2 = 0.6214$ ,  $P = 0.0003$ ) for the first seven sampling times, covering 18 hours from 6:00 to midnight. However, for the remaining two time points (3:00 and 6:00 on Sept 4, 2005) this trend no longer held. As a consequence, a correlation analysis of *ftsZ* gene to gene expression inclusive of all sampling points, was not significant ( $R^2 = 0.0304$ ,  $P = 0.45$ ). These results indicate that at least for some of the samples a correlation existed between gene abundance and per-cell gene expression activity.

In the first 12 hours, *ftsZ* expression and to some degree *ftsZ* gene copies appeared to be somewhat correlated with the tidal cycle, with peaks of expression at 6:00 and 18:00 and lowest

expression at 12:00, lagging the onshore high and low tides by about 3 hours respectively (Figure 2). Temperature and salinity also showed fluctuations that loosely followed the tidal cycle with the same 3-hour lag (data not shown). After 18:00 a steady drop in temperature and simultaneous increase in salinity was observed. Overall, these trends in salinity and temperature indicate that tidal currents influenced our sampling site and that different waters with different NAC11-7 populations (or patches), were sampled.

In order to further examine the relationships between *ftsZ* gene abundance and expression with water masses, we overlaid *ftsZ* gene abundances and expression to gene ratios onto temperature-salinity (T-S) plots (Figure 3). These plots indicate a cyclical variation between warmer, less saline waters and cooler, more saline waters. While, gene copies did not show clear trends in the T-S plot (data not shown), *ftsZ* expression showed an interesting relationship. With the single exception of the 9:00 sample, samples below 25.8°C showed significantly lower *ftsZ* expression (two-sample t-test  $P=0.003$ ) than those above 25.8°C (Figure 3). A regression analysis excluding the 9:00 measurements showed a significant correlation ( $R^2=0.725$ ,  $P=0.007$ ) between *ftsZ* expression and temperature. These data strongly indicate that populations from two contrasting water types were sampled throughout the diel study. Furthermore, when the data are separated between samples above and below 25.8 °C, an outlier in *ftsZ* transcript abundance is evident at the 9:00 sampling time, suggestive of a peak in *ftsZ* gene expression at that time (Figure 3).

#### ***ARISA analysis of microbial communities.***

In order to examine the dynamics of bacterioplankton communities and populations at the different sampling times, these communities were examined using automated ribosomal intergenic spacer analysis (ARISA). We observed that a number of ARISA peaks were present

at all time points analyzed, (i.e. 456, 485, 510 and 556 bp), while several peaks showed a more ephemeral occurrence (i.e. 793, 799, 900 bp), suggesting that levels of patchiness differed for different phylotypes (Supplemental Fig 1). At least in the case of these fluctuating phylotypes, different populations (patches) appear to have been sampled at different times. Unfortunately ARISA peaks sizes corresponding to the NAC11-7 group were not readily detected, as these organisms were likely present in numbers below the resolution of ARISA.

#### ***Enclosed incubation diel expression of NAC11-7 *ftsZ* genes.***

A parallel on-deck incubation experiment in which *ftsZ* gene and gene expression were quantified in incubated samples was conducted for comparative purposes. In these samples, *ftsZ* gene abundance dropped from 6:00 to 15:00 and stabilized at ca.  $1 \times 10^4$  copies/ml (a 2.5-fold decrease). Contrastingly, *ftsZ* expression showed a very remarkable 4-fold increase in expression at 9:00 relative to the 6:00 time point. However, by 12:00 expression was 5-fold less (Figure 4). Considering that the measurements were made in three separate, incubated bottles and that the  $C_T$ s of the 9:00 cDNA samples were nearly two units lower than those of the remaining samples, we are very confident that this observation was not due to an experimental artifact.

## **Discussion**

In order to test assumptions related to the measurement of *in situ* growth rates of specific roseobacters based on cell division genes, a qPCR *ftsZ* assay was developed for the NAC11-7 clade. To the best of our knowledge these are the first quantitative data of relative expression of a functional gene for a specific roseobacter subclade in natural samples. Analysis of the NAC11-7 sequences reveals that although all the sequences were highly similar at the amino acid level, one sequence (Clone *ftsZ*05F3 was different at the DNA level (Supplemental Figure 2). Based

on the ecotype theory (Cohan, 2001), clone *ftsZ5F03* was assumed to belong to a different ecotype that was not subjected to a theoretical selective sweep, as were the remaining 5 phylotypes. In order to maximize the chance that our measurements reflected populations most coherent in their response to environmental factors, the sequence of clone *ftsZ05F03* was not targeted by our qPCR assay.

We observed that levels of normalized transcript-to-gene ratios were low. This was not completely unexpected as mRNAs exist in a complex RNA pool composed primarily of rRNA and have relatively short half-lives (Belasco, 1993). Thus, the efficiency of mRNA recovery is expected to be considerably lower than that for genomic DNA. While we did not measure reverse transcription efficiencies here, we have previously shown RT efficiencies to be constant between different samples for proteorhodopsin mRNAs using an analogous method with the same reverse transcriptase and a specific primer (Shi, 2005).

It is important to emphasize that our measurements of gene expression are based on a *per cell* average for the entire population. Thus, depending on how *ftsZ* is expressed in a single cell during the cell cycle, the mRNA to gene ratio may reflect: 1) the fraction of the population that is actively transcribing *ftsZ* in preparation for cell division or 2) the average per cell expression level [i.e. if *ftsZ* is constitutively transcribed, and *FtsZ* regulation occurs post-transcriptionally in preparation for cell division]. If a population displays synchronous growth and *ftsZ* expression varies during the cell cycle (Case 1), we expect to observe peaks in the population-level *ftsZ* expression. Conversely, if *ftsZ* transcription were constitutive (Case 2), one would not expect *ftsZ* expression peaks even in synchronous populations.

The only other study to examine *ftsZ* gene expression in natural bacterial assemblages was carried out with Red Sea populations of the cyanobacterium *Prochlorococcus*. In that study,

*ftsZ* gene expression was found to be highly synchronous, with greatest expression occurring in the twilight hours before nightfall (Holtzendorff et al., 2002). These expression profiles were consistent with the known photoautotrophic physiology of the organism; these populations are naturally synchronized and undergo cell division at night (Vaulot and Partensky, 1992 ; Vaulot et al., 1995). *Caulobacter crescentus*, an aquatic alphaproteobacterium, which is largely used as a model for cell cycle and differentiation studies (see review by McAdams and Shapiro, 2009), is the closest roseobacter relative for which extensive *ftsZ* gene expression studies have been undertaken. In this organism, *ftsZ* gene expression is cell cycle-dependent and is time constrained (Quardokus et al., 1996; Kelly et al., 1998; Sackett et al., 1998; Martin and Brun, 2000; Brun, 2001). In contrast, FtsZ protein concentration per cell has been found to be constant regardless of growth rate in the copiotrophic *E. coli* (*Gammaproteobacteria*) and *B. subtilis* (*Firmicutes*) (Rueda et al., 2003; Weart and Levin, 2003; Haeusser and Levin, 2008). Several arguments support time-constrained *ftsZ* expression in NAC11-7 members. As members of the *Alphaproteobacteria*, these bacteria are more likely to share conserved features of central physiology with *C. crescentus* than with the more distantly related *E. coli* or *B. subtilis*. In addition, the genome of NAC 11-7 isolate HTCC2255 contains the genes encoding for DnaA and CtrA (blastp e-value  $<10^{-20}$  using the *Caulobacter crescentus* amino acid sequences), the central regulators of *ftsZ* expression and cell cycle in this organism (McAdams and Shapiro, 2009). Finally, it is difficult to rationalize advantages to NAC11-7, organisms adapted to somewhat oligotrophic conditions, to synthesize a constant amount of FtsZ through the entire cell cycle unless FtsZ has an alternative unknown function.

Different lines of evidence indicate that different NAC11-7 populations (or patches) with different growth rates were sampled at different time points during the *in situ* experiment. First,

215 remarkable fluctuations of NAC11-7 *ftsZ* gene copy numbers (nearly 4-fold in 3 h) were  
216 observed *in situ* (Figure 2), implying changes in populations sizes resulting from physical  
217 processes. Second, ARISA analysis of the total bacterioplankton community structure showed  
218 evidence for patchiness, as certain phylotypes were only detected in specific time points  
219 (Supplemental Figure 1). While transient populations were likely sampled throughout the diel  
220 cycle, some trends in *ftsZ* gene and gene expression did emerge that revealed insight into *ftsZ*  
221 gene expression profiles and synchronicity of these populations. During the first seven data  
222 points (18 hrs from 6:00 to 0:00), a strong correlation ( $R^2 = 0.6214$ ) was observed between  
223 NAC11-7 abundance *in situ* (inferred from *ftsZ* copies/ml), and the average *per cell ftsZ*  
224 transcript abundance, indicating that levels of expression are not constant in this ecotype. This  
225 was suggestive that more active populations yielded, or were associated, with higher abundances  
226 of these organisms. Collectively, these results support the case for non-synchronous cell division  
227 since *ftsZ* expression was not constrained to specific times (i.e. *ftsZ* was expressed at all  
228 measured time points). Finally, the plot of gene expression over a temperature-salinity diagram,  
229 and a regression analysis showed that, with the exception of the 9:00 replicates, there is a high  
230 correlation between water temperature and *ftsZ* expression (Figure 3).

231         We found a remarkable peak of gene-normalized *ftsZ* transcript abundance at 9:00 in the  
232 on-deck incubation experiment, which could be interpreted in several ways. First, and in  
233 contrast to conclusions drawn from the *in situ* samples, *ftsZ* expression (and by extension cell  
234 division) might be synchronous in the putatively photoheterotrophic NAC11-7 clade. An  
235 alternative explanation would be that the 9:00 peak of expression resulted from stimulation  
236 caused by confinement. The observations that 1) the response was fast and time-constrained; 2)  
237 gene abundances steadily decreased with time, and 3) there is no evidence that members of the

NAC11-7 group respond positively to enrichment conditions (i.e. this group is not known as copiotrophic), all argue against this notion. The slow decline in *ftsZ* gene abundance in the bottles might be explained by a lack of growth, and continuous mortality by viral lysis and/or protistan bacterivory. The peak of *ftsZ* expression at 9:00 might be explained by a previous “commitment” of the circadian rhythm. As different NAC11-7 populations (patches) were sampled over the course of the *in situ* diel it is difficult to directly compare *in situ* *ftsZ* gene expression levels with the parallel on-deck experiment. However, the fact that the *in situ* sample from 9:00 was higher than expected based on the correlation between gene expression and temperature (Figure 3) also agrees with the incubation results.

An intriguing interpretation of these seemingly contradictory results is that a certain fraction of the population is synchronously dividing, with a peak in *ftsZ* expression at 9:00, while a background of asynchronously dividing cells also exist, some of which are expressing *ftsZ* at any given time. This "background" expression might be in fact, better correlated to temperature or other parameters influencing growth. The fact that a certain level of *ftsZ* expression was measured throughout the diel cycle in synchronously dividing *Prochlorococcus* populations (Holzendorf et al 2001) seems to support this idea. However, in the case of *Prochlorococcus*, *ftsZ* expression is at the end of the light period and cell division occurs at the dark (Holtzendorff et al., 2001). Assuming that the NAC11-7 ecotype is an anaerobic anoxygenic phototroph (AAP) as HTCC 2255 appears to be, one possible explanation for a peak of expression in the morning with subsequent cell division might be related to the fact that in the Rhodobacterales AAP studied to date (i.e. Biebl and Wagner-Dobler, 2006; Koblizek et al., 2010) bacteriochlorophyll synthesis occurs at night. Division in the early morning could assure a maximum amount of this pigment is available to daughter cells.

To our knowledge, this is the first indication that specific heterotrophic bacterioplankton grow synchronously in the environment, and if proven true, this will have very important implications to the study of bacterioplankton ecology. For instance, bacterioplankton production is in most cases measured for the bulk community at short-time incubations. Thus, understanding whether specific populations are synchronous will be paramount to the full interpretation of these bulk measurements. Since there is putative evidence that some representatives of NAC11-7 are photoheterotrophic, understanding the relationships between light-driven metabolism and synchronous growth in nature is also important. Also, future *in situ* studies should include Lagrangian sampling to minimize advective effects, and allow measurement in coherent populations.

## Experimental Procedures

### *Sampling*

Surface water samples were collected every 3 hours from 6:00 local time on September 3 to 6:00 on September 4, 2005 at station RM6 (37°05.61N, 75°42.35 W) aboard the RV Cape Henlopen. Temperature and salinity were measured using a SBE9 CTD (Seabird, Bellevue, WA), and tidal effects were estimated from the height of the tide at Ship Shoal Inlet calculated using the data at the NOAA tides and current prediction page <http://tidesandcurrents.noaa.gov>. Triplicate 590 ml water subsamples were collected pre-filtered through GF/A filters (1.6  $\mu$ m nominal pore; Whatman, Maidstone, UK). Ninety ml of pre-filtered sample was filtered through 13 mm diameter 0.2 $\mu$ m Supor200<sup>®</sup> polysulfone filters (Pall Corp., East Hills, NY) and the filters transferred to a tube containing 130 $\mu$ l lysis buffer (2 mM NaEDTA (pH 8.0), 20 mM Tris•Cl (pH 8.0), 1.2% v/v Triton X100). The remaining 500 ml were filtered through a 25 mm diameter 0.2  $\mu$ m Supor200<sup>®</sup> filters (Pall Gelman Inc.) and transferred to screw cap tubes containing 250  $\mu$ l

RNA Later (Ambion, Austin, TX). All samples were frozen at  $-20^{\circ}\text{C}$  aboard and within a week stored at  $-70^{\circ}\text{C}$  until nucleic acid extraction.

On-deck incubation experiments were conducted using water collected at the first sampling time (6:00, 03 September, 2005). Fifteen l of surface water was used to fill 24 x 500 ml (600 ml total volume) polycarbonate bottles (Nalge Nunc International Corp., Rochester, NY), and incubated at *in situ* temperatures in an on-deck incubator. At every *in situ* sampling point, three of the incubation bottles were taken and sampled for nucleic acids as described for cast samples.

#### ***Nucleic acid extraction and quantification***

Total DNA was extracted from Supor200<sup>®</sup> 0.2  $\mu\text{m}$  filters as previously described, except that the DNeasy 96 Tissue Kit was used, and therefore the DNeasy 96 plates were sealed and centrifugations performed in a Sorvall<sup>®</sup> Legend<sup>™</sup> T Centrifuge with a Highplate<sup>®</sup> rotor (Kendro, Osterode, Germany) at 5250 rpm for 10 min for binding and 5250 rpm for 5 min for washes. The DNeasy 96 plate was incubated at  $70^{\circ}\text{C}$  for 15 min evaporate traces of ethanol and DNA was eluted by centrifugation at 5250 rpm, for 2 min.

Total RNA was extracted from Supor200<sup>®</sup> 0.2  $\mu\text{m}$  filters following an optimized protocol adapted from the Qiagen RNeasy<sup>®</sup> 96 manual (Shi 2005). Filters in screw cap tubes were thawed on the ice and 320  $\mu\text{l}$  low protein binding zirconium oxide beads (200  $\mu\text{m}$ , OPS Diagnostics, Lebanon, NJ) were added into each tube.  $\beta$ -Mercaptoethanol was mixed in a 1:100 ratio into RLT buffer and 875  $\mu\text{l}$  of the mixture added to screw cap tubes. The tubes were beaten in a MM301 mixer mill (Retsch GmbH Inc., Haan, Germany) at 30.0 HZ for 2 min and incubated for 5 min at  $70^{\circ}\text{C}$ . 800  $\mu\text{l}$  of the liquid phase was transferred into a new low-RNA-binding 2 ml microcentrifuge tube (Ambion, Inc.), avoiding the beads. 800  $\mu\text{l}$  of 100% ethanol was added into

each tube and well mixed well. 800  $\mu$ l of the mixture was loaded into wells of RNeasy 96 plate, sealed and centrifuged at 5000 rcf for 5 min, and the procedure repeated. The columns were washed once with 800  $\mu$ l RW1 buffer and twice with 800  $\mu$ l RPE buffer at 5000 rcf for 5 min, with the last spin for 15 min. 35  $\mu$ l Diethylpyrocarbonate (DEPC) treated water (Ambion) were added into each column, incubation for 1 min at room temperature and 5 min centrifugation at 5000 rcf. This procedure was repeated with another 35  $\mu$ l DEPC treated water. RNA was treated with the DNA-free™ kit (Ambion) to remove co-extracted DNA.

DNA and RNA concentrations were quantified fluorometrically by PicoGreen® and RiboGreen® staining respectively (Molecular Probes, Invitrogen Corp., Carlsbad, CA) on a Spectra MAX Gemini microplate spectrofluorometer (Molecular Devices, Sunnyvale, CA).

#### ***Construction of ftsZ DNA PCR clone library***

An initial *ftsZ* and FtsZ database was developed using *arb* (Ludwig *et al.*, 2004) by importing gene sequences listed by Vaughan and coworkers (Vaughan *et al.*, 2004). 163 representative FtsZ sequences were exported and used to retrieve additional amino acid, and coding gene sequences using blastp searches against the March 2005 NCBI *nt*, *env\_nt* and *wgs* databases. All amino acid sequences were aligned using ClustalW (Thompson *et al.*, 1994) and a bayesian tree containing 507 sequences (144 homologous positions) was constructed using MrBayes version 3.0 (Ronquist and Huelsenbeck, 2003) with the following parameters: 800,000 generations, mixed models of amino acid substitution, and a burn in of 6000 trees. In this tree the *Alphaproteobacteria* formed a monophyletic clade with high confidence values and several orders including *Rhizobiales* and *Rhodobacterales* also formed monophyletic clades (data not shown). The database was updated with public sequences belonging to the *Rhodobacterales*, and was used for the design of PCR primers and probes. Degenerate primers targeting *ftsZ* from

330 *Rhodobacterales* (ftsZrb2F: 5'-AAY GCN GTS AAY AAY AT-3', ftsZrb2R: 5'-YTT NCC  
331 CAT YTC RT-3') were designed and used to retrieve *ftsZ* gene sequences by PCR and cloning  
332 from a DNA sample collected at time zero.

333 1 $\mu$ l of extracted genomic DNA was used as template in a 10  $\mu$ l-volume PCR reaction,  
334 which also included 10X PCR buffer, 0.2mM of each dNTP, 3 mM MgCl<sub>2</sub>, 500 nM forward  
335 primer ftsZ2rbF, 500 nM reverse primer ftsZ2rbR and 0.025 U/ $\mu$ l of Platinum® Taq DNA  
336 Polymerase (Invitrogen, Carlsbad, CA). Reactions were performed on a GeneAmp 9700 PCR  
337 system (Applied Biosystems) and cycling conditions were as follows: 2 min at 94 °C and 37  
338 cycles of 30 sec at 94 °C, 30 sec at 55 °C and 2 min at 72 °C. PCR was followed by a  
339 reconditioning (Thompson et al., 2002) of 5 steps. Reconditioned PCR products were loaded on  
340 a 1% modified TAE (40 mM Tris-acetate, pH 8.0, 0.1 mM Na<sub>2</sub>EDTA) agarose gel and separated  
341 by electrophoresis. Target fragments were cut, recovered by Ultrafree®-DA (Millipore) gel  
342 extraction and used to build a *ftsZ* gene clone library using the TOPO™ TA cloning kit  
343 (Invitrogen) following manufacture's instructions. 96 clones were bidirectionally sequenced  
344 using BigDye V3.1 chemistry and capillary electrophoresis on an AB3100 genetic analyzer  
345 (Applied Biosystems Inc, Foster City, CA). These sequence data have been submitted to the  
346 GenBank database under accession number HM035149-HM035222. 74 sequences identified as  
347 *ftsZ* were imported into the ARB database described above and added to the tree described above  
348 using the ADD\_BY\_PARSIMONY tool. Finally, a bayesian tree of *Rhodobacterales ftsZ* was  
349 constructed using the MPI version of Mr Bayes V3.1 (Ronquist and Huelsenbeck, 2003; Altekar  
350 et al., 2004). 188 homologous amino acid positions from 123 sequences were exported and used  
351 in this phylogenetic analysis. Two four parallel chains of 2,000,000 generations were run with  
352 mixed models of amino acid substitution; trees were sampled every 100 generations, and 11000

“burnin” trees were excluded to generate the consensus tree. The average standard deviation of split frequencies was below 0.05 after 1,100,000 generations.

### ***Real Time PCR Assays***

PCR primers and a Taqman probe were designed to target selected members of the NAC11-7 group. Results of *ftsZ* cloning and sequencing indicated that based on amino acid sequences five clone sequences were affiliated with strain HTCC 2255. However, the DNA sequence of one of these clones (F3) contained obviously higher variation to the remainder clones and HTCC 2255 (Supplemental Figure 1). Based on ecotype theory (Cohan, 2001), it appeared that the distinct clone could belong to a different ecotype, which we assumed was not subject to a periodic selection event as the remaining sequences. Thus we designed a primer and probe to exclusively target the remaining four sequences and HTCC 2255. The primer and the probe were manually designed with aid of probe match functions in the arb\_edit module of the ARB package.  $T_M$ , secondary structure and possible dimers were checked using Primer Express (Applied Biosystems) and the Oligo Analyzer online tool ([www.idtdna.com](http://www.idtdna.com)).

To test specificity of primers to the target sequences, 3 target clones and 15 non-target clones were purified using QuickLyse Miniprep kit (Qiagen), and diluted to  $10^7$  *ftsZ* copies/ $\mu$ l with nuclease-free TE buffer (Ambion). Ten  $\mu$ l PCR reaction contained 5  $\mu$ l TaqMan® Universal PCR Master Mix (Applied Biosystems), 0.5 $\mu$ M Forward primer *ftsZrbA03-2qF* (GTG AAA AAG CTA CTG AGG GTC T) and Reverse primer *ftsZrbA03-2qR* (GCT TCC TGC CAG ATG ATC), and 1  $\mu$ l plasmid template. The cycling parameters were as follows: 2 min at 50 °C, 10 min at 95 °C and 30 cycles of 15 sec at 95 °C and 1min at 57 °C. All PCR products were loaded in a 2% NuSieve® (3:1) Agarose Gel (Cambrex, Rockland, ME), electrophoresed, and

post-stained with 1:10,000 SYBR® Gold (Invitrogen) for 30 min. The gel was visualized with a FluoroChem 8900 (Alpha Innotech, San Leandro, CA).

In order to get the highest amplification efficiency in real time PCR, a primer concentration matrix was performed. In each 25  $\mu$ l reaction, the following reagents were kept constant: 1X PCR buffer, 0.2 mM of dATP, dGTP and dCTP, 0.4 mM of dUTP, 5 mM MgCl<sub>2</sub>, 200 nM probe NAC11-7 (AAC CAACAGTAGGAGCATTAGCCGCT), 1.2  $\mu$ M SuperROX™ (Biosearch Technologies, Novato, CA), 0.01 U/ $\mu$ l AmpErase® Uracil N-glycosylase (UNG) (Applied Biosystems), 0.025 U/ $\mu$ l of Platinum® Taq DNA Polymerase (Invitrogen), 2.5  $\mu$ l NAC11-7 standard (10<sup>4</sup> copies/ $\mu$ l) and a matrix of forward and reverse primer concentrations of 0.1  $\mu$ M, 0.5  $\mu$ M, 1.0  $\mu$ M, 1.5  $\mu$ M. Reactions were set in a MicroAmp® Optical 96-Well Reaction Plate (Applied Biosystems), which was sealed with an optical adhesive cover (Applied Biosystems) reactions and ran in an ABI Prism 7000 Sequence Detection system, following the cycling parameters: 2 min at 50 °C, 10 min at 95 °C and 40 cycles of 15 sec at 95 °C and 1 min at 57 °C. The primer combination yielding the lowest C<sub>T</sub> (1.5  $\mu$ M *ftsZ*rbA03-2qF; 0.5  $\mu$ M *ftsZ*rbA03-2qR) was used in all subsequent measurements.

A plasmid containing cloned DNA that was purified and linearized as previously described (Suzuki et al., 2000) was used to prepare standards for real time PCR for quantification of *ftsZ* gene and mRNA. 2.5  $\mu$ l of DNA extracts were used in 25  $\mu$ l reactions and standards ranged from 10<sup>2</sup> to 10<sup>7</sup> copies/ $\mu$ l. Reactions were run in duplicate along with non-template controls using the same cycling parameters described for the primer matrix. *ftsZ* cDNA was quantified in the same manner as *ftsZ* gene except that 5  $\mu$ l of Reverse transcription products were used as template. All real time PCR measurements were calculated as copy numbers per volume of seawater, assuming that nucleic acid extraction efficiencies were constant as shown by

Shi (Shi, 2005). mRNAs copies were assumed equal to cDNA copies. Since DNA and mRNA were measured in triplicate biological samples, *ftsZ* genes and mRNA copy numbers that were >2X or <X/2 (X is the average of the remainder two replicates) were treated as outliers and removed from the analyses

### **Optimization of Reverse Transcription**

Since preliminary results showed relatively high  $C_T$  values from *in situ* sample cDNAs, we attempted to increase *ftsZ* cDNA copy numbers by: (1). Increasing reverse transcription efficiency, by varying template amount, RT temperature and RT duration and (2). Adding increased amounts of cDNA products in real-time PCR reactions. In optimized conditions 4  $\mu$ l purified RNA (ca. 10 ng) was reverse transcribed to cDNA using ThermoScript™ Reverse Transcriptase. Template and 10 pmol of the NAC11-7 specific primer *ftsZrbA03-2qR* were heated to 65°C for 5 min and chilled on ice immediately, followed by the addition of 5X cDNA synthesis buffer, 5mM DTT, 2 U/ $\mu$ l RNaseOUT™ Inhibitor (Invitrogen), 0.75 U/ $\mu$ l of reverse transcriptase and DEPC-treated water to a final volume of 10  $\mu$ l. Reactions were incubated at 55°C for 60 min to synthesize cDNA and at 85°C for 5 min to inactivate the transcriptase. No-RT controls were performed for one of the triplicate samples using the same method except substituting reverse transcriptase with DEPC-treated water.

### **ARISA (Automated rRNA intergenic spacer analysis)**

One  $\mu$ l environmental genomic DNA was used to perform a 10  $\mu$ l ARISA reactions, containing 1X PCR buffer, 1.2 mM  $MgCl_2$ , 0.08 mM dNTPs (Promega Corp., Madison, WI), 0.5  $\mu$ M primer 1406F-FAM (Fisher and Triplett, 1999), 1.5  $\mu$ M primer 23S-Y (Dyda et al., 2009), and 0.01 U/ $\mu$ l of Platinum® Taq DNA Polymerase (Invitrogen). Reactions were run on a GeneAmp 9700 (Applied Biosystems) under the following conditions: Initial denaturation and

enzyme activation at 94 °C for 2 min, followed by 35 cycles of 94 °C for 30 sec, 55 °C for 30 sec and 65 °C for 2min. 1  $\mu$ l of each PCR reaction was mixed with 9  $\mu$ l of 1:0.06 Formamide and GS2500 size standard (Applied Biosystems), denatured at 94 °C for 2 min and separated by capillary electrophoresis using an Applied Biosystems 3100 genetic analyzer. Sizes of the fragments were analyzed by the Peak Scanner™ Software v1.0 (Applied Biosystems).

#### **Acknowledgements**

We acknowledge funding support from the National Science Foundation Grants OCE 0550547 (to M.S.) and OCE 0550485 (to A.B.), the CBL Graduate Education Committee and the crew of RV Cape Henlopen.

## References

- Alonso, C., and Pernthaler, J. (2005) Incorporation of glucose under anoxic conditions by bacterioplankton from coastal North Sea surface waters. *Appl Environ Microbiol* **71**: 1709-1716.
- Altekar, G., Dwarkadas, S., Huelsenbeck, J.P., and Ronquist, F. (2004) Parallel Metropolis coupled Markov chain Monte Carlo for Bayesian phylogenetic inference. *Bioinformatics* **20**: 407-415.
- Azam, F., and Malfatti, F. (2007) Microbial structuring of marine ecosystems. *Nat Rev Microbiol* **5**: 782-791.
- Belasco, J.G. (1993) mRNA degradation in prokaryotic cells: an overview. . In *Control of Messenger RNA Stability*. Belasco, J.G., Brawerman, G. (ed). San Diego CA: Academic Press,, pp. 3-11.
- Biebl, H., and Wagner-Dobler, I. (2006) Growth and bacteriochlorophyll a formation in taxonomically diverse aerobic anoxygenic phototrophic bacteria in chemostat culture: Influence of light regimen and starvation. *Process Biochemistry* **41**: 2153-2159.
- Brun, Y.V. (2001) Global analysis of a bacterial cell cycle: tracking down necessary functions and their regulators. *Trends Microbiol* **9**: 405-407.
- Buchan, A., Gonzalez, J.M., and Moran, M.A. (2005) Overview of the marine *Roseobacter* lineage. *Appl Environ Microbiol* **71**: 5665-5677.
- Carpenter, E.J., and Chang, J. (1988) Species-Specific Phytoplankton Growth-Rates Via Diel DNA-Synthesis Cycles .1. Concept of the Method. *Marine Ecology-Progress Series* **43**: 105-111.

452 Chang, J., and Carpenter, E.J. (1988) Species-Specific Phytoplankton Growth-Rates Via Diel  
 453 DNA-Synthesis Cycles .2. DNA Quantification and Model Verification in the  
 454 Dinoflagellate *Heterocapsa-Triquetra*. *Marine Ecology-Progress Series* **44**: 287-296.  
 455 Cohan, F.M. (2001) Bacterial species and speciation. *Syst Biol* **50**: 513-524.  
 456 Cottrell, M.T., and Kirchman, D.L. (2003) Contribution of major bacterial groups to bacterial  
 457 biomass production (thymidine and leucine incorporation) in the Delaware estuary. *Limnol*  
 458 *Oceanogr* **48**: 168-178.  
 459 Dajkovic, A., and Lutkenhaus, J. (2006) Z ring as executor of bacterial cell division. . *J Mol*  
 460 *Microbiol Biotechnol* **11**: 140-151.  
 461 Delong, E.F. (2009) The microbial ocean from genomes to biomes. *Nature* **459**: 200-206.  
 462 Dyda, R.Y., Suzuki, M.T., Yoshinaga, M.Y., and Harvey, H.R. (2009) The response of microbial  
 463 communities to diverse organic matter sources in the Arctic Ocean. *Deep-Sea Research*  
 464 *Part II-Topical Studies in Oceanography* **56**: 1249-1263.  
 465 Fisher, M.M., and Triplett, E.W. (1999) Automated Approach for Ribosomal Intergenic Spacer  
 466 Analysis of Microbial Diversity and Its Application to Freshwater Bacterial Communities.  
 467 *Appl Environ Microbiol* **65**: 4630-4636.  
 468 Fuhrman, J.A., and Azam, F. (1982) Thymidine incorporation as a measure of heterotrophic  
 469 bacterioplankton production in marine surface waters: evaluation and field results *Mar Biol*  
 470 **66**: 109-120.  
 471 Haeusser, D.P., and Levin, P.A. (2008) The great divide: coordinating cell cycle events during  
 472 bacterial growth and division. *Curr Opin Microbiol* **11**: 94-99.

473 Hagstrom, A., Larsson, U., Horstedt, P., and Normark, S. (1979) Frequency of dividing cells, a  
 474 new approach to the determination of bacterial growth rates in aquatic environments.  
 475 *Applied and Environmental Microbiology* **37**: 805-812.

476 Holtzendorff, J., Marie, D., Post, A.F., Partensky, F., Rivlin, A., and Hess, W.R. (2002)  
 477 Synchronized expression of *ftsZ* in natural *Prochlorococcus* populations of the Red Sea.  
 478 *Environ Microbiol* **4**: 644-653.

479 Holtzendorff, J., Partensky, F., Jacquet, S., Bruyant, F., Marie, D., Garczarek, L. et al. (2001)  
 480 Diel Expression of Cell Cycle-Related Genes in Synchronized Cultures of *Prochlorococcus*  
 481 sp. Strain PCC 9511. *J Bacteriol* **183**: 915-920.

482 Hottes, A.K., Shapiro, L., and McAdams, H.H. (2005) DnaA coordinates replication initiation  
 483 and cell cycle transcription in *Caulobacter crescentus*. *Mol Microbiol* **58**: 1340-1353.

484 Kelly, A.J., Sackett, M.J., Din, N., Quardokus, E., and Brun, Y.V. (1998) Cell cycle-dependent  
 485 transcriptional and proteolytic regulation of FtsZ in *Caulobacter*. *Genes Dev* **12**: 880-893.

486 Kirchman, D.L., K'Neas, E., and Hodson, R. (1985) Leucine incorporation and its potential as a  
 487 measure of protein synthesis by bacteria in natural aquatic systems. *Appl Environ*  
 488 *Microbiol* **49**: 599-607.

489 Koblizek, M., Mlcouskova, J., Kolber, Z., and Kopecky, J. (2010) On the photosynthetic  
 490 properties of marine bacterium COL2P belonging to *Roseobacter* clade. *Arch Microbiol*  
 491 **192**: 41-49.

492 Margolin, W. (2005) FtsZ and the division of prokaryotic cells and organelles. *Nat Rev Mol Cell*  
 493 *Biol* **6**: 862-871.

494 Martin, M.E., and Brun, Y.V. (2000) Coordinating development with the cell cycle in  
 495 *Caulobacter*. *Curr Opin Microbiol* **3**: 589-595.

496 McAdams, H.H., and Shapiro, L. (2009) System-level design of bacterial cell cycle control.  
 497 *FEBS Lett* **583**: 3984-3991.

498 Osawa, M., and Erickson, H.P. (2006) FtsZ from divergent foreign bacteria can function for cell  
 499 division in *Escherichia coli*. *J Bacteriol* **188**: 7132-7140.

500 Quardokus, E.M., Din, N., and Brun, Y.V. (1996) Cell cycle regulation and cell type-specific  
 501 localization of the FtsZ division initiation protein in *Caulobacter*. *Proc Natl Acad Sci* **93**:  
 502 6314-6319.

503 Rappe, M.S., and Giovannoni, S.J. (2003) The uncultured microbial majority. *Annu Rev*  
 504 *Microbial* **57**: 369-394.

505 Ronquist, F., and Huelsenbeck, J.P. (2003) MrBayes 3: Bayesian phylogenetic inference under  
 506 mixed models. *Bioinformatics* **19**: 1572-1574.

507 Rueda, S., Vicente, M., and Mingorance, J. (2003) Concentration and assembly of the division  
 508 ring proteins FtsZ, FtsA, and ZipA during the *Escherichia coli* cell cycle. *J Bacteriol* **185**:  
 509 3344-3351.

510 Sackett, M.J., Kelly, A.J., and Brun, Y.V. (1998) Ordered expression of *ftsQA* and *ftsZ* during  
 511 the *Caulobacter crescentus* cell cycle. *Mol Microbiol* **28**: 421-434.

512 Sherr, B.F., Sherr, E.B., and McDaniel, J. (1992) Effect of protistan grazing on the frequency of  
 513 dividing cells in bacterioplankton assemblages. *Appl Environ Microbiol* **58**: 2381-2385.

514 Shi, Y. (2005) Measurement of in situ expression of proteorhodopsin genes at the North Pacific  
 515 Central Gyre Station ALOHA. In *Marine-Estuarine-Environmental Sciences*. College Park:  
 516 MSc. Thesis, University of Maryland College Park, p. 127.

517 Simon, M., and Azam, F. (1989) Protein content and protein synthesis rates of planktonic marine  
 518 bacteria. *Mar Ecol Prog* **51**: 201-213.

519 Steward, G.F., and Azam, F. (1999) Bromodeoxyuridine as an alternative to H-3-thymidine for  
 520 measuring bacterial productivity in aquatic samples. *Aquat Microb Ecol* **19**: 57-66.  
 521 Suzuki, M.T., Taylor, L.T., and Delong, E.F. (2000) Quantitative analysis of small-subunit  
 522 rRNA genes in mixed microbial populations via 5'-nuclease assays. *Appl Environ*  
 523 *Microbial* **66**: 4605-4614.  
 524 Thompson, J.D., Higgins, D.G., and Gibson, T.J. (1994) CLUSTAL W: improving the sensitivity  
 525 of progressive multiple sequence alignment through sequence weighting, position-specific  
 526 gap penalties and weight matrix choice. *Nucleic Acids Res* **22**: 4673-4680.  
 527 Thompson, J.R., Marcelino, L.A., and Polz, M.F. (2002) Heteroduplexes in mixed-template  
 528 amplifications: formation, consequence and elimination by 'reconditioning PCR'. *Nucleic*  
 529 *Acids Res* **30**: 2083-2088.  
 530 Vaughan, S., Wickstead, B., Gull, K., and Addinall, S.G. (2004) Molecular evolution of FtsZ  
 531 protein sequences encoded within the genomes of archaea, bacteria, and eukaryota. *J Mol*  
 532 *Evol* **58**: 19-29.  
 533 Vaultot, D., and Partensky, F. (1992 ) Cell Cycle Distributions of Prochlorophytes in the North  
 534 Western Mediterranean Sea *Deep Sea Res* **39(5)**: 727-742 511  
 535 Vaultot, D., Marie, D., Olson, R.J., and Chisholm, S.W. (1995) Growth of Prochlorococcus, a  
 536 Photosynthetic Prokaryote, in the Equatorial Pacific-Ocean. *Science* **268**: 1480-1482.  
 537 Weart, R.B., and Levin, P.A. (2003) Growth rate-dependent regulation of medial FtsZ ring  
 538 formation. *J Bacteriol* **185**: 2826-2834.  
 539 Wohlers, J., Engel, A., Zollner, E., Breithaupt, P., Jurgens, K., Hoppe, H.G. et al. (2009)  
 540 Changes in biogenic carbon flow in response to sea surface warming. *Proc Natl Acad Sci*  
 541 **106**: 7067-7072.

542 Zubkov, M.V., Fuchs, B.M., Archer, S.D., Kiene, R.P., Amann, R., and Burkill, P.H. (2001)  
543 Linking the composition of bacterioplankton to rapid turnover of dissolved  
544 dimethylsulphoniopropionate in an algal bloom in the North Sea. *Environ Microbiol* **3**:  
545 304-311.  
546  
547  
548

## Figure legends

Figure 1. Consensus bayesian phylogenetic tree of FtsZ protein sequences from the *Rhodobacterales* group reconstructed using MrBayes v3.1. FtsZ sequences retrieved in this study are marked in boldface. The numbers on nodes represent branch confidence values. The bar represents the mean number of substitutions per residue in the trees used to build the consensus tree.

Figure 2. *FtsZ* gene copy numbers (grey bars) and expression (estimated from the ratio of *ftsZ* cDNA copies to gene copies; white bars) at station RM6 on September 3 and 4, 2005, and tidal heights at Ship Shoal Inlet (37° 13'N, 75° 48'W). Error bars represent the standard deviation of triplicate samples from a single CTD cast. Pre-dawn and post-dusk sampling times are represented by the grayed areas. The tidal height is shown with the line lacking symbols.

Figure 3. Temperature and Salinity plot, with overlaying *in situ* NAC11-7 *ftsZ* expression. Gene normalized *ftsZ* expression levels are proportional to the area of the circles. The area of the circle representing the 3:00 PM sample corresponds to  $2.5 \times 10^{-4}$  mRNA copies per gene copies. Inset: normalized *in situ* *ftsZ* expression in samples with temperatures below and above 25.8 °C.


Figure 4. *ftsZ* gene copy numbers (filled squares) and expression (estimated from the ratio of *ftsZ* cDNA copies to gene copies; open circles) in samples incubated onboard. Error bars represent the standard deviation of triplicate bottles.

566 Supplemental Fig 1. Percentage of different phylotypes identified by ARISA fragment sizes in  
567 the *in situ* samples. The results shown in here indicate that the replicate subsamples were largely  
568 similar in composition, indicative of relatively low within-sample variation. However, in a few  
569 of the samples (i.e. 9:00, 21:00) noticeable variation between replicate subsamples was evident.  
570 These findings highlight the heterogeneity in marine environments and reinforce the value of  
571 replicate subsampling in natural systems.


572 Supplemental Figure 2. A. Alignment of NAC11-7 FtsZ sequences. B alignment of NAC11-7  
573 *ftsZ* sequences

574

575




577 **Figure 2.**



578

579


579 **Figure 3.**



580


581

581 **Figure 4.**



582

583 Supplemental Figure 1



584

585

