



HAL
open science

Diversity of Arctic pelagic Bacteria with an emphasis on photoheterotrophs: a review

Dominique Boeuf, Florian Humily, Christian Jeanthon

► To cite this version:

Dominique Boeuf, Florian Humily, Christian Jeanthon. Diversity of Arctic pelagic Bacteria with an emphasis on photoheterotrophs: a review. *Biogeosciences*, 2014, 11 (12), pp.3309-3322. 10.5194/bg-11-3309-2014 . hal-01097258

HAL Id: hal-01097258

<https://hal.sorbonne-universite.fr/hal-01097258>

Submitted on 19 Dec 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Diversity of Arctic pelagic *Bacteria* with an emphasis on photoheterotrophs: a review

D. Boeuf^{1,2}, F. Humily^{1,2}, and C. Jeanthon^{1,2}

¹CNRS, UMR 7144, Marine Phototrophic Prokaryotes Team, Station Biologique de Roscoff, 29680 Roscoff, France

²Sorbonne Universités, UPMC Univ Paris 06, UMR 7144, Oceanic Plankton Group, Station Biologique de Roscoff, 29680 Roscoff, France

Correspondence to: C. Jeanthon (jeanthon@sb-roscoff.fr)

Received: 21 December 2013 – Published in Biogeosciences Discuss.: 12 February 2014

Revised: 10 April 2014 – Accepted: 2 May 2014 – Published: 20 June 2014

Abstract. The Arctic Ocean is a unique marine environment with respect to seasonality of light, temperature, perennial ice cover, and strong stratification. Other important distinctive features are the influence of extensive continental shelves and its interactions with Atlantic and Pacific water masses and freshwater from sea ice melt and rivers. These characteristics have major influence on the biological and biogeochemical processes occurring in this complex natural system. Heterotrophic bacteria are crucial components of marine food webs and have key roles in controlling carbon fluxes in the oceans. Although it was previously thought that these organisms relied on the organic carbon in seawater for all of their energy needs, several recent discoveries now suggest that pelagic bacteria can depart from a strictly heterotrophic lifestyle by obtaining energy through unconventional mechanisms that are linked to the penetration of sunlight into surface waters. These photoheterotrophic mechanisms may play a significant role in the energy budget in the euphotic zone of marine environments. Modifications of light and carbon availability triggered by climate change may favor the photoheterotrophic lifestyle. Here we review advances in our knowledge of the diversity of marine photoheterotrophic bacteria and discuss their significance in the Arctic Ocean gained in the framework of the Malina cruise.

as the Arctic Mediterranean Sea (Coachman and Aagaard, 1974). However, the Arctic Ocean and its marginal seas (the Chukchi, East Siberian, Laptev, Kara, Barents, White, Greenland, and Beaufort; some oceanographers also include the Bering and Norwegian seas) are the least known basins and bodies of water of the world's oceans owing to their remoteness, hostile weather, and perennial or seasonal ice cover. The Arctic Ocean has the most extensive shelves of any ocean basin, covering about 50 % of its total area. Relative to other ocean basins, rivers play a disproportionately important role in the Arctic Ocean, which contains only about 1 % of the volume of the world's oceans yet receives approximately 10 % of the global terrigenous dissolved organic carbon (DOC) load (Aagaard et al., 1985; Opsahl et al., 1999). The DOC concentrations in coastal Arctic waters are twice higher than those in the Atlantic and Pacific oceans (Cauwet and Sidorov, 1996), highlighting the tight coupling of the Arctic Ocean with terrestrial catchments (Meon and Amon, 2004). In addition to the discharge of large rivers, the Arctic Ocean is also freshened by an inflow of relatively low-salinity Pacific waters through the Bering Strait and net precipitation over the ocean surface (Serreze et al., 2006). Sea ice dynamics also play a pivotal role in the salinity regime, adding salt to the underlying water during ice formation and releasing fresh water during ice melting. As a consequence, estuarine gradients are a defining feature not only nearshore, but throughout this ocean. Due to this large influx of freshwater, the Arctic Ocean is well stratified with a distinctive surface layer of reduced salinity, the polar mixed layer, and density stratification inhibits vertical mixing with warmer, saline Atlantic waters below 200 m, allowing sea ice to form

1 Introduction

The Arctic Ocean is the smallest of the five major oceans, covering 2.8 % of the earth's total surface (Pidwirny, 2006); it is almost completely surrounded by land and can be viewed

(Aagaard and Coachman, 1975). Inorganic nutrient concentrations exhibit strong regional gradients from high nutrient regimes (e.g., the Chukchi Sea shelf) to oligotrophic conditions (e.g., in the Beaufort Gyre and the Beaufort Sea).

The climate of the Arctic marine environment is marked by extreme seasonality in solar radiation, ice cover and atmospheric temperature and, to a lesser extent, water temperature (Carmack et al., 2006). The winter season in the Arctic is characterized by little or no sunlight, of which only a fraction is able to penetrate the thick sea ice and snow layers to the water column below. This mainly affects photosynthetic organisms like phytoplankton that require sunlight for carbon fixation. Most pronounced changes occur during the spring and early summer, when melting sea ice, melt ponds, and rapidly increasing day length allow greater penetration of light to the water column. Despite their low temperatures, Arctic waters support a highly productive ice-free season (Garneau et al., 2008), and bacterial activity has been found to be as high as in lower latitudes (Wheeler et al., 1996). Overall, however, the composition, physiology and function of prokaryotic heterotrophs in the marine Arctic are poorly understood, both in terms of spatial variations as well as temporal dynamics (Amon, 2004).

In this article, we first present an updated overview of the diversity of pelagic heterotrophic bacteria and their influences on biochemistry and upper food webs in the Arctic Ocean. Then, we focus on our current knowledge of the photoheterotrophic bacterial populations. In the light of recent studies that have been conducted during the Malina cruise (<http://malina.obs-vlfr.fr>), we also discuss the possible changes that could occur in the diversity of photoheterotrophic bacteria in coastal environments of the Beaufort Sea.

2 Diversity of pelagic heterotrophic bacteria

Early studies conducted in nearshore areas of the western Beaufort Sea investigated the cultivable fraction of the prokaryotic community (Kaneko et al., 1977; Zobell, 1946). Succession of cultivable bacterial phenotypes was demonstrated according to season and geography and in response to algal blooms. These pioneering studies demonstrated that bacterial isolates tend to differ significantly from those found in other marine environments. This observed potential for high genetic diversity was further confirmed by 16S rRNA gene-based molecular analyses of subsurface Arctic prokaryotic communities (Ferrari and Hollibaugh, 1999). Early molecular studies analyzed compositional variability in prokaryotic assemblages in Arctic waters (Bano and Hollibaugh, 2002). The 16S rRNA gene clone libraries from the central Arctic (Bano and Hollibaugh, 2002), western Arctic (Malmström et al., 2007), Greenland Sea (Pommier et al., 2007; Zaballos et al., 2006), Baffin Bay (Pommier et al., 2007), Laptev Sea (Kellogg and Deming, 2009), and Franklin




Figure 1. Relative abundance of major phyla identified in pyrosequencing studies of the deep Arctic Ocean (Galand et al., 2010); the Arctic Chukchi Beaufort (Kirchman et al., 2010), the central Arctic multiyear ice and the central Arctic seawater (Bowman et al., 2012), the offshore Beaufort Sea and the Mackenzie estuary (Ortega-Retuerta et al., 2013).

Bay (Collins et al., 2010) revealed that bacterial communities were composed of polar and cosmopolitan phylotypes. Alphaproteobacterial and gammaproteobacterial sequences often formed the dominant phylotypes in planktonic bacterial assemblages. However, Malmström et al. (2007) reported high numbers of gammaproteobacterial (53 % of the clones) and Bacteroidetes (29 % of clones) sequences in a Chukchi Sea sample (western Arctic). Gammaproteobacteria and Bacteroidetes were also the dominant groups in surface waters of Baffin Bay and Greenland Sea (Pommier et al., 2007).

Most recently, massively parallel tag sequencing techniques have improved our knowledge of the prokaryotic diversity in the Arctic Ocean (Fig. 1), despite rarefaction analyses still suggested undersampling. Although a greater abundance of Gammaproteobacteria was reported by Kirchman et al. (2010), Alphaproteobacteria typically dominate Arctic surface waters, followed by Gammaproteobacteria and by Bacteroidetes (see references in Fig. 1 and Comeau et al. (2011)). Verrucomicrobia and Actinobacteria are also widely recognized as abundant Arctic seawater clades. Interestingly, the taxonomic composition of the rare phylotypes was similar to that of the most abundant ones (Galand et al., 2009). Low proportions of marine cyanobacterial sequences were generally obtained in marine waters as previously reported (Waleron et al., 2007), although significant contributions were obtained in the attached fraction associated to the Mackenzie plume (Ortega-Retuerta et al., 2013). In the latter study, significant differences between particle-attached and free-living bacterial communities were observed in the open sea, but both fractions showed a similar structure in coastal and river samples. The influence of river inputs on prokaryotic community structure was also demonstrated in previous studies in the Arctic (Galand et al., 2008; Garneau et al., 2006; Kellogg and Deming, 2009) and in a global

comparison of bacterial diversity including Arctic samples (Ghiglione et al., 2012).

Fluorescence in situ hybridization (FISH) has also been used to identify bacterial phylogenetic groups and to assess their actual cell abundance in natural waters (Alonso-Sáez et al., 2008; Elifantz et al., 2007; Garneau et al., 2006; Kirchman et al., 2007; Wells and Deming, 2003). These studies confirmed the dominance of Alphaproteobacteria (Elifantz et al., 2007; Garneau et al., 2006) or Bacteroidetes (Elifantz et al., 2007; Wells and Deming, 2003). With contributions higher than 20 %, SAR11 clade represents the major group (Malmström et al., 2007). Cell abundances of other ubiquitous groups such as SAR86 (Gammaproteobacteria) and *Roseobacter* clade-affiliated (RCA) cluster (8 % and 10 %, respectively) were also significant. Seasonal changes in the relative abundance of major bacterial groups were reported, and some abundant groups showed relatively high activity at the single-cell level (Alonso-Sáez et al., 2008, 2010; Nikrad et al., 2012).

3 Diversity of photoheterotrophic bacteria

Photoheterotrophic bacteria are microorganisms able to harvest light energy and to utilize dissolved organic compounds as carbon and energy source. Three groups of photoheterotrophic bacteria are currently known in the marine environment. The first group includes the abundant autotrophic cyanobacteria like *Prochlorococcus* and *Synechococcus* capable of facultative heterotrophic growth in nutrient-depleted environments. The second group is formed by aerobic anoxygenic phototrophic (AAP) bacteria. These organisms are obligate aerobes using dissolved organic matter as a source of organic carbon for their metabolism and growth. They harvest light using bacteriochlorophyll *a* (BChl *a*)-based pigments and reaction centers in addition to respiration. The third group is composed of diverse bacterial groups that use bacterial rhodopsin as light-driven proton pump.

3.1 Cyanobacteria

Synechococcus spp. and *Prochlorococcus* spp. are prominent constituents of the marine biosphere, which account for a significant percentage of the biomass and oceanic primary production (DuRand et al., 2001; Li, 1994; Veldhuis et al., 1997). Although they depend on light for most of their energy production and CO₂ for carbon acquisition, these cyanobacteria have also been shown to take up diverse organic substrates such as amino acids, nucleosides, oligopeptides, urea, sulfur compounds, or cyanate (Malmström et al., 2005; Palenik et al., 2003; Zubkov and Tarran, 2005), and use them to produce cellular biomass.

3.1.1 Abundance in the Arctic Ocean

Although cyanobacteria predominate in brackish or freshwater polar environments reaching abundances among the highest reported in natural environments (Powell et al., 2005), their abundance are typically low in marine polar waters (Li, 2009). In both polar oceans, picocyanobacteria follow a general trend of decreasing concentrations and relative abundance with increasing latitudes, and strong inverse correlations between cell densities and temperature have been reported (Marchant et al., 1987; Murphy and Haugen, 1985; Rosenbergl, 1993). *Prochlorococcus*, the most abundant photosynthetic organism in the ocean, is widely distributed in subtropical and tropical waters from 40° S to 48° N (Johnson et al., 2006; Partensky et al., 1999b). Due to its temperature sensitivity, its concentrations decline rapidly beyond this band (Marchant et al., 1987). Despite its presence was reported in a sub-Arctic region at latitude 61° N (Buck et al., 1996) and at southern latitudes of the sub-Antarctic (Marchant et al., 1987), *Prochlorococcus* is considered virtually absent in the polar oceans (Baldwin et al., 2005; Cottrell and Kirchman, 2009; Li, 1994; Lin et al., 2012; Zwirgmaier et al., 2008). *Synechococcus* is more ubiquitous and inhabits virtually all marine and freshwater environments (Partensky et al., 1999a). In the Arctic Ocean, *Synechococcus* has been found in many marine cold environments such as the Chukchi Sea (Cottrell and Kirchman, 2009; Huang et al., 2012), the Beaufort Sea (Cottrell and Kirchman, 2009), off Iceland (Michelou et al., 2007) and in an area between the Norwegian, Greenland and Barents seas (Zwirgmaier et al., 2008). Epifluorescence microscopy studies have also reported variable abundances of phycoerythrin-rich cyanobacteria in Resolute passage (Robineau et al., 1999). The highest cell numbers ($> 10^3$ cells mL⁻¹) were measured between the Norwegian, Greenland and Barents seas during the late summer (Zwirgmaier et al., 2008), in marine waters of lower salinities in Resolute passage (Robineau et al., 1999), and in Mackenzie River and estuary, as well as in coastal sites of Franklin Bay and Amundsen Gulf (Waleron et al., 2007). However, *Synechococcus* abundances measured in the Arctic Ocean are generally lower. In two coastal sites from the Chukchi and Beaufort seas, densities never exceed more than 100 cells mL⁻¹ (Cottrell and Kirchman, 2009) and their concentrations decreased by an order of magnitude in offshore sites near Arctic pack ice (Robineau et al., 1999). Low cyanobacterial abundance (less than 24 cells mL⁻¹) was also measured in the East Greenland Current (Gradinger and Lenz, 1989). Interestingly, cyanobacterial abundance did not vary significantly between summer and winter, probably due to heterotrophic activity during the dark period (Cottrell and Kirchman, 2009). During the Malina cruise in summer 2009, *Synechococcus* was completely absent in Arctic waters of the Chukchi and Beaufort seas (Balzano et al., 2012) as observed previously (Li, 1998).




Figure 2. *Synechococcus* community composition calculated from environmental sequences recovered in high-latitude regions. Reprinted by permission from Macmillan Publishers Ltd: The ISME Journal (Huang et al., 2012)

The origin of picocyanobacteria in the Arctic marine environment is controversial. Several authors argue for a strong contribution of allochthonous inputs like rivers or transport by advection from surrounding oceans (Gradinger and Lenz, 1989; Vincent et al.; Waleron et al., 2007) while others hypothesize for brackish communities autochthonous to the Arctic Ocean and adapted to cold environments (Cottrell and Kirchman, 2009; Huang et al., 2012).

3.1.2 Diversity in the Arctic Ocean

Marine *Synechococcus* exhibit a high genetic diversity and are divided into three major subclusters, 5.1, 5.2, and 5.3. Among numerous clades unveiled by different gene markers (more than 20 based on 16S–23S internally transcribed spacer), clades I, II, III and IV are dominant on a global scale (Ahlgren and Rocap, 2012; Zwirgmaier et al., 2008). These lineages show different geographic niche exploitation. Clades I and IV dominate in nutrient-rich coastal waters at high latitudes, while clade II is widely distributed in subtropical and tropical oceanic waters (Huang et al., 2012; Zwirgmaier et al., 2008).

Using 16S rRNA-targeted probes specific for *Synechococcus* clades, Zwirgmaier et al. (2008) showed that clades I and IV dominated *Synechococcus* communities in an area between the Norwegian, Greenland, and Barents seas, both clades yielding approximately equal hybridization signals. The significance of these clades up to 62° N in the north Atlantic was further confirmed by Huang et al. (2012) (Fig. 2). In the southern Bering Sea (56–60° N) where clade IV was not detected, clade I represented a significant portion of the *Synechococcus* community (Huang et al., 2012). The latter study evidenced a clear shift of *Synechococcus* community structure with increasing latitude and decreasing tempera-

ture. Clade I occurred rarely or was not detected in the northern Bering Sea and Chukchi Sea (62–72° N) where the euryhaline clades CB1 and CB5 prevailed. Although these two clades were originally isolated from the Chesapeake Bay, their prevalence in the northern Bering Sea and the Chukchi Sea suggests an autochthonous rather than a riverine or estuarine origin. In the Beaufort Sea that is strongly influenced by the Mackenzie River, only sequences closely related to freshwater and brackish *Synechococcus* were detected (Waleron et al., 2007).

However, regarding their very low abundance in the polar oceans, picocyanobacteria probably play little role on the pelagic carbon and energy flow in this part of the globe (Díez et al., 2012; Gradinger and Lenz, 1989; Koh et al., 2012).

3.2 Aerobic anoxygenic phototrophic bacteria

Cyanobacteria are not the only bacteria that can use light energy in the upper water column. In the late 1970s, novel aerobic photosynthetic bacteria containing bacteriochlorophyll *a* (BChl *a*) were isolated from seaweed, sand, bottom sediments, and seawater in Tokyo Bay (Harashima et al., 1978; Shiba et al., 1979). In contrast to their anaerobic counterparts, these organisms now called AAP bacteria thrive in oxic conditions.

Rediscovered in the early 2000s, AAP bacteria were thought to be abundant and widespread in the euphotic zone of the open ocean and to contribute significantly to photosynthetically driven electron transport (Kolber et al., 2001; Rathgeber et al., 2004). Kolber et al. (2000) suggested that AAP bacteria might be abundant in oligotrophic oceanic regions where the capacity to harvest light energy may provide photoheterotrophs a competitive advantage over chemoheterotrophs. Later studies showed that their abundance and distribution vary greatly among oceanic regimes, suggesting that there is a broad range of potential ecological niches for these microbes (Cottrell et al., 2006; Lehours et al., 2010; Mašín et al., 2006; Sieracki et al., 2006; Yutin et al., 2007). It seems however that AAP bacteria are more abundant in shelf and coastal areas than in the open ocean (Schwalbach and Fuhrman, 2005; Sieracki et al., 2006). The number of AAP bacteria can exceed 10% of total prokaryotes in estuaries (Cottrell et al., 2010; Waidner and Kirchman, 2007) but typically account for a small percentage (2 to 4%) in open ocean waters (Cottrell et al., 2006; Jiao et al., 2007) where high abundances were recorded however (Lami et al., 2007). Despite lower AAP bacterial abundances reported in oligotrophic pelagic environments, these bacteria constitute a very dynamic part of marine microbial communities and may contribute significantly to the upper ocean carbon cycle (Koblížek et al., 2007).




Figure 3. Schematic diagram representing the changes in abundance (a) and diversity (b) of major AAP bacterial clades along an onshore–offshore transect of the Beaufort Sea. AAP proportions were calculated by dividing AAP bacterial numbers (enumerated by infrared epifluorescence microscopy) by the total prokaryotic numbers (4′,6-diamidino-2-phenylindole counts).

3.2.1 Abundance in the Arctic Ocean

The abundance and distribution of AAP bacteria has been poorly studied in high-latitude waters. In the earliest study conducted in Arctic marine waters, AAP bacteria were found to constitute 5 to 8 % of the prokaryotic community in coastal waters of the Beaufort and Chukchi seas (Cottrell and Kirchman, 2009). These authors did not report significant changes in AAP bacterial abundance between summer and winter. During the Malina cruise that crossed the Bering and Chukchi seas and focused on the southern Beaufort Sea in summer 2009, we observed a strong decreasing gradient from the Mackenzie River to the open ocean, with a mean AAP bacterial contribution of 7 % in the Mackenzie mouth, 1 % inshore, and 0.1 % offshore (Boeuf, 2013). The highest AAP abundances were found in surface waters of the Mackenzie plume (14 %) and in the benthic nepheloid layer (BNL) (9 %) (Fig. 3a). These results are in line with those recorded in estuarine systems such as the Long Island Sound, Delaware Bay and Chesapeake Bay, where these bacteria can constitute more than 10 % of the prokaryotic community (Cottrell et al., 2010; Schwalbach and Fuhrman, 2005; Sieracki et al., 2006; Waidner and Kirchman, 2008). The low AAP bacterial percentages in-shelf and offshore Beaufort Sea samples were similar to those reported in Antarctic waters of the Southern Ocean (Schwalbach and Fuhrman, 2005) and in most oligotrophic temperate and tropical regions (Hojerová et al., 2011; Jiao et al., 2007; Lamy et al., 2011).

This is consistent with the trophic status of the Beaufort Sea that is quite oligotrophic in summer (Ortega-Retuerta et al., 2012), with nearly undetectable nitrate levels during the Malina cruise. In contrast to total prokaryote abundance, AAP proportions correlated significantly with ammonium, silicate, total Chl *a*, and the different forms of organic carbon, nitrogen, and phosphorus. This suggests that AAP bacteria may respond to organic supply differently and have higher mineralization capacities than the bacterial community (Boeuf et al., 2013).

3.2.2 Diversity in the Arctic Ocean

Our knowledge on the diversity of AAP bacteria has long been based on cultures (Yurkov and Beatty, 1998). AAP bacteria have been further identified in environmental samples using known photosynthetic reaction center genes (Béjå et al., 2002). Wider polymerase chain reaction (PCR) primers targeting *pufM* gene encoding the M subunit of bacterial photosynthetic reaction centers were then used to expand AAP bacterial diversity in various environments (Yutin et al., 2005). Diverse AAP bacteria were also retrieved in metagenomes from the Atlantic and Pacific oceans (DeLong et al., 2006; Venter et al., 2004; Waidner and Kirchman, 2005; Yutin et al., 2007). Further studies indicated that AAP communities can be structured differently according to geographical location (Jeanthon et al., 2011; Jiao et al., 2007; Lehours et al., 2010). All these efforts have evidenced the genetic diversity of AAP bacteria with members of the Alpha-, Beta-, and Gammaproteobacteria. Both targeted (Hu et al., 2006; Jiao et al., 2007; Lehours et al., 2010) and non-targeted (Yutin et al., 2007) diversity studies have shown that depending on the location and environment, members of either the Alpha- or Gammaproteobacteria typically dominate the marine AAP bacterial community. For example, in the Baltic and Mediterranean seas, most AAP bacteria belonged to Gammaproteobacteria (Jeanthon et al., 2011; Mašín et al., 2006), while in the Global Ocean Sampling (GOS) expedition, the *Roseobacter*-like group of Alphaproteobacteria dominated the oligotrophic AAP bacterial community (Yutin et al., 2007).

Only a limited number of studies have examined the diversity of AAP bacteria in high-latitude waters. First investigations carried out in the permanently frozen freshwater Lake Fryxell (Antarctica) (Karr et al., 2003) revealed that AAP bacteria living in these habitats were distinct from their marine counterparts. The presence of *pufM* genes in Antarctic seawater and sea ice environment was further demonstrated by Koh et al. (2011). Using the same primer set, both these studies found that Alphaproteobacteria dominated the AAP bacterial communities. Seawater clones closely related to *Roseobacter denitrificans* suggest the presence of the ubiquitous RCA cluster in Antarctic waters. In surface waters of the Chukchi and Beaufort seas, although *pufM* genes belong to phylogroups previously identified in the GOS metagenome




Figure 4. Relative abundance of *pufM* genes in clone libraries obtained from Arctic Ocean samples. Clones were retrieved from coastal samples of the Chukchi and Beaufort seas in summer and winter (Cottrell and Kirchman, 2009) and from the Mackenzie plume, coastal and offshore surface waters and other depths (Boeuf et al., 2013). The numbers of clones and libraries analyzed are indicated in bold and between brackets, respectively.

(Yutin et al., 2007), most were distinct from those retrieved at lower latitudes and in Antarctic waters (Boeuf et al., 2013; Cottrell and Kirchman, 2009; Koh et al., 2011). By examining a couple of coastal sites in summer and winter, Cottrell and Kirchman (2009) revealed the presence of three phylogroups (Fig. 4). Two of them (phylogroups E and F) belonged to the Rhodobacterales that is one of the most common alphaproteobacterial order in polar and subpolar oceans (Fu et al., 2010; Ghiglione and Murray, 2012; Niederberger et al., 2010; Prabakaran et al., 2007; Salka et al., 2008; Selje et al., 2004). *Loktanella*-like *pufM* clones (phylogroup F) dominated the libraries at both seasons, whereas proportions of *Sulfitobacter*-related sequences (phylogroup E) increased in winter. Boeuf et al. (2013) investigated more thoroughly the AAP diversity in the transition zone between offshore and Mackenzie-influenced coastal waters (southern Beaufort Sea) in summer. The presence of *pufM* genes affiliated with Rhodobacterales in most of the Beaufort shelf was remarkable (Figs. 3b and 4) and supported by independent isola-

tion of a large number of *Sulfitobacter* and *Loktanella* strains from most of the shelf and offshore stations.

The most striking observation was the widespread distribution of betaproteobacterial clades in the entire Beaufort shelf (Fig. 4). The most abundant betaproteobacterial clade exhibited a strong river-to-ocean gradient (Fig. 3b), suggesting that these bacteria grew in the Mackenzie River and then were mixed with Beaufort coastal waters (Boeuf, 2013). These results were in line with the fact that (i) betaproteobacterial *pufM* genes increased in summer coastal libraries (Cottrell and Kirchman, 2009) at higher river flow, (ii) betaproteobacterial AAP bacteria are found in high proportions in estuarine or freshwater habitats (Salka et al., 2011; Waidner and Kirchman, 2008; Yutin et al., 2007), and (iii) Betaproteobacteria are commonly found in the Mackenzie River where they are dominant (Galand et al., 2008; Garneau et al., 2006). AAP bacterial diversity was highest at depths where Pacific Summer Water mixes with the BNL. At these depths, AAP bacteria affiliated to orders Rhizobiales and Sphingomonadales were common along the Mackenzie shelf but almost absent in surface waters (Figs. 3b and 4).

Although Gammaproteobacteria represent one third of the bacterial community in Arctic marine waters (Kirchman et al., 2010), no gammaproteobacterial AAPs were found at the coastal sites of Chukchi and Beaufort seas (Cottrell and Kirchman, 2009) (Fig. 4). We confirmed this trend during the Malina cruise since only a few *pufM* clones belonging to the gammaproteobacterial phylogroup K were recovered in north Pacific Ocean samples but none were obtained above 62° N. Similarly, no clones of this phylogroup were found in Antarctic sea ice and seawater (Koh et al., 2011). This seems to indicate that polar waters are an exception to its widespread distribution (Jeanthon et al., 2011; Lehours et al., 2010; Yutin et al., 2007).

3.3 Proteorhodopsin-containing bacteria

An additional type of phototrophy has also been recently found in marine surface waters. Proteorhodopsins (PR) are retinal-binding integral membrane proteins discovered via environmental genomic surveys in the early 2000s (Béjà et al., 2000; Béjà et al., 2001; de la Torre et al., 2003; Venter et al., 2004). Proteorhodopsins are homologues of bacteriorhodopsins discovered in the halophilic archeon *Halobacterium salinarum* four decades ago (Oesterhelt and Stoekenius, 1971). Their designation was based on the discovery of metagenomic fragments that linked the first discovered of these PR genes to a small subunit ribosomal RNA (SSU rRNA) gene defining the uncultured SAR86 group II, a gammaproteobacterial group widespread in marine plankton (Béjà et al., 2000; Suzuki et al., 2001).

Subsequent PCR-based gene surveys, screenings of bacterial artificial chromosome and fosmid libraries, and metagenomic and genome analyses have identified a wide variety of PR genes (Béjà et al., 2001; de la Torre et al., 2003; Man




Figure 5. Relative abundance of PR genes in clone libraries obtained from Arctic Ocean samples. Clones were retrieved from coastal samples of the Chukchi and Beaufort seas in summer and winter (Cottrell and Kirchman, 2009) and from the Mackenzie plume, coastal and offshore surface waters and other depths (Boeuf, 2013). The numbers of clones and libraries analyzed are indicated in bold and between brackets, respectively.

et al., 2003). Proteorhodopsins were found in diverse bacterial groups, including the ubiquitous SAR11 groups (Giovannoni et al., 2005; Giovannoni and Stingl, 2005), the abundant coastal clade SAR 92 (Stingl et al., 2007), as well as in strain HTCC 2255 Rhodobacterales, marine Flavobacteria (Gómez-Consarnau et al., 2007; González et al., 2008; Riedel et al., 2010; Yoshizawa et al., 2012) and planktonic Archaea (Frigaard et al., 2006). Phylogenies of PR amino acid sequences are not consistent with those of 16S rRNA, suggesting that PR genes have acquired by lateral gene transfer (Frigaard et al., 2006; Sabehi et al., 2004). Proteorhodopsins have been detected in coastal and open ocean environments, but also in freshwater, estuarine, and brackish ecosystems (Atamna-Ismael et al., 2008; Rusch et al., 2007).

3.3.1 Diversity in the Arctic Ocean

Only very few molecular surveys have reported PR bacteria in marine polar waters. Antarctic PR sequences have been detected in coastal surface waters near Anvers Island, western Antarctic Peninsula (Béjå et al., 2000; Williams et al., 2012) and in sea ice and brine samples of the Ross Sea (Koh et al., 2010). Sequences analyses revealed that Antarctic PR showed similarity to previously reported PR sequences, although most of the sequences were generally distinct. Cottrell and Kirchman (2009) reported the first study on PR bacterial diversity in Arctic waters. Although limited to a couple

of coastal sites, this study revealed a diverse collection of PR genes, only a few of them belonging to previously identified groups. Clone libraries were mainly composed of alpha- and gammaproteobacterial PR types and their composition changed between summer and winter (Fig. 5). During the Malina cruise in the Beaufort Sea, we surveyed the PR diversity in different water masses using the same primer set and compared the composition of DNA and cDNA libraries (Fig. 5). An alphaproteobacterial group entirely composed of Arctic sequences that we called “Arctic” dominated in most surface libraries, confirming earlier results of Cottrell and Kirchman (2009). Contributions of SAR11 sequences were higher in cDNA than in DNA libraries, suggesting that this clade is highly active in Arctic waters (Boeuf, 2013). With the exception of Mackenzie plume samples, high proportions of gammaproteobacterial PR mostly composed of SAR92-like and IMCC2047-like sequences were found in DNA libraries. Only a few clones aligned with the SAR86 clade of PR genes, which was reported to comprise the majority of Antarctic pelagic PR sequences (Béjå et al., 2001). Additionally, coastal surface and Mackenzie plume libraries contained significant amounts of Flavobacteria sequences. In contrast with AAP community composition, only a limited number of few betaproteobacterial PR sequences were obtained along the Mackenzie plume although primer sets targeting freshwater clades were specifically tested. Our study revealed that Beaufort Sea PR sequences did not match closely with Antarctic ones, supporting the fact that PR genes from the Northern and Southern hemispheres are not similar.

3.3.2 Abundance in the Arctic Ocean

Unlike Cyanobacteria and AAP bacteria, abundance of PR prokaryotes cannot be estimated using direct methods such as flow cytometry or epifluorescence microscopy. Quantitative PCR (qPCR) is the technique that has been most used to examine the distribution of PR genes (Campbell et al., 2008; Riedel et al., 2010; Suzuki et al., 2001) and to measure their expression in natural waters (Lami et al., 2009). Alternatively, the abundance of PR genes has been evaluated from the frequency of clones carrying PR genes in environmental DNA libraries (Rusch et al., 2007; Sabehi et al., 2005; Venter et al., 2004).

Proteorhodopsin-containing bacteria have been estimated to contribute for up to 65 % (mean of 50 %) of the bacterial community in the Sargasso Sea, from 7 to 57 % (mean of 23 %) in the north Atlantic Ocean, about 13 % in the Mediterranean Sea and 35 % in the North Sea (Campbell et al., 2008; Riedel et al., 2010; Rusch et al., 2007; Sabehi et al., 2005; Venter et al., 2004). Alphaproteobacterial groups represent generally the highest fraction among PR bacteria. Among them, the ubiquitous SAR11-like cluster is the only one to contribute to 10 % of the overall community and almost 18 % in the Sargasso and Mediterranean seas (Boeuf, 2013; Campbell et al., 2008).




Figure 6. Schematic diagram representing the changes in abundance (a) and diversity (b) of major PR bacterial clades along an onshore–offshore transect of the Beaufort Sea. Total PR abundance (a) was calculated by summing the relative proportions of the eight most abundant clades in the clone libraries (Fig. 5).

In polar marine waters, PR bacteria were detected throughout the annual cycle (Cottrell and Kirchman, 2009; Williams et al., 2012) but their abundance has not been extensively assessed. Cottrell and Kirchman (2009) compared the seasonal abundance of two clades retrieved from two coastal sites of the Chukchi and Beaufort seas. Their abundance was not found to differ significantly between summer and winter. Using the set of PR sequences previously amplified from the Beaufort Sea (Boeuf, 2013), we designed qPCR primers to target the most abundant groups in the libraries. By adding together their abundance estimates, these groups accounted between 0.9 % and 44.7 % (mean of 15.3 %) of the total bacterial community (Fig. 6a) (Boeuf, 2013). The ubiquitous SAR11-like PR clade dominated in the Beaufort Sea and was preferentially distributed offshore and above the deep chlorophyll maximum where mean contributions exceeded 10 % of the total bacterial community (Fig. 6b). This contribution is close to the percentage of SAR11 16S rRNA gene determined from pyrosequencing data obtained from the western Arctic Ocean (Kirchman et al., 2010). Consistent with clone library data, the “Arctic” clade, the second most abundant PR clade in the libraries, was preferentially distributed in surface waters. It represented up to 24 % of the offshore bacterial community. The alphaproteobacterial HOT2C01 cluster was found in high proportions in Sargasso Sea (26 %) while decreasing to 8 % in the north Atlantic (Campbell et al., 2008). In the Beaufort Sea, this cluster was mostly present in coastal areas but its contribution was low (2 %) (Boeuf, 2013). Surprisingly, although proportions of PR Gammaproteobacteria

were high in most clone libraries where they overdominated at depth (Fig. 5), they represented a minor fraction of the total bacterial community in all samples (mean < 1 %). These results indicate that relative frequencies of PR genes in clone libraries do not reflect their real abundance in natural communities. Our quantitative data indicate that PR genes are abundant in the Beaufort Sea, suggesting the high contribution of PR phototrophy in Arctic waters. Alphaproteobacterial PR genes dominate, and their distribution in coastal and offshore waters suggests that factors other than light control their abundance.

4 Potential impact of global change

The Arctic Ocean is a key player in global climate regulation as well as a particularly sensitive system that has already undergone serious perturbations due to climate warming (Camill, 2005; Peterson et al., 2002). The total volume and extent of the sea ice are decreasing (Walsh, 2008) leading to an increase of river runoff (Peterson et al., 2002) and a decrease in the surface layer salinity of the Arctic Ocean (ACIA, 2005). An additional consequence is the accelerating permafrost melting (Nelson, 2003) and coastal erosion (Rachold et al., 2000). The increasing delivery of organic carbon is suspected to enhance the mineralization process carried out by heterotrophic bacteria in the estuarine and coastal Arctic Ocean (Meon and Amon, 2004) and thus release a large amount of CO₂ into the atmosphere (Kling et al., 1991). Excluding the potential shift of surrounding oceans’ connections, currents, and icecap drifting (Hopcroft et al., 2008) that may profoundly affect bacterial communities at both regional and global scales, global warming could impact bacterioplankton by modifying its local environment (physical aspect) and its nutrient sources and interactions in the food web (biotic aspect). The lengthening of the open-water period is likely to stimulate primary production via increased light availability, which in turn would enhance bacteria production (Arrigo et al., 2008; Zhang et al., 2010). These models all assume that more open water will lead to increased mixing and entrainment of nutrients into the upper water column. Alternatively, the freshening of the Arctic could lead to stronger salinity stratification and fewer nutrients being entrained into the euphotic zone.

In the Beaufort Sea, distributions of AAP and PR bacteria and their relationships with environmental parameters show strong differences. The most abundant AAP bacterial clade is strongly influenced by nutrient-rich river inputs. The freshening of the Arctic coastal regions could therefore favor the survival of halotolerant freshwater AAP bacteria and expansion of their habitat (Boeuf, 2013). The impact of the phototrophic lifestyle on energy requirements and carbon metabolism of these bacteria is unknown. Examination of their distribution in other Arctic coastal regions and their contribution to biogeochemical cycling is needed.

The decrease in ice thickness and the shift from perennial ice pack to seasonal ice (Perovich, 2011) could favor massive under-ice phytoplankton blooms in the Arctic (Arrigo et al., 2012). Such structural shifts of Arctic marine ecosystems could in turn be favorable to AAP bacteria of the *Roseobacter* clade (González et al., 2000; Suzuki et al., 2001) and PR groups (Flavobacteriaceae, alphaproteobacterial "Arctic" clade, and gammaproteobacterial SAR92 clade) that exhibit a strong responsiveness to phytoplankton blooms. A long-term consequence of further increasing stratification will be reduction of the vertical flux of nutrients to the euphotic zone and decreasing productivity. Low-nutrient conditions in the upper Beaufort Sea could therefore be particularly favorable to bacterial groups adapted to oligotrophic conditions (SAR11 and other alphaproteobacterial PR groups) whose photoheterotrophic capabilities provide a competitive advantage (Gómez-Consarnau et al., 2010; Steindler et al., 2011).

The current knowledge suggests that microbial processes in the Arctic Ocean are particularly sensitive to environmental changes and have potentially large impacts on carbon flows and other ecosystem functions. The way photoheterotrophic microbes adjust to these changes will probably have impacts on the marine ecosystem of the Arctic Ocean. Understanding how they interact and influence higher food webs as well as biogeochemical cycling in this system is necessary. Future research should examine communities present in less-explored central Arctic regions and include thorough analysis of seasonal influences on microbial processes.

Acknowledgements. This study was conducted as part of the Malina Scientific Program funded by ANR (Agence nationale de la recherche), INSU-CNRS (Institut National des Sciences de l'Univers – Centre National de la Recherche Scientifique), CNES (Centre National d'Etudes Spatiales), and ESA (European Space Agency). This work was supported by the ANR RHOME0 (ANR-11-BSV7-021-02) and the FP7 European Union program MicroB3 (UE-contract-287589). We thank all participants of the Malina cruise for their help, especially M. Babin who coordinated the project, K. Lévesque for the logistics, and all the R/V CCGS *Amundsen* crew members. D. Boeuf and F. Humily were supported by grants from the French Ministry of Higher Education and Research and from ANR, respectively.

Edited by: M. Babin

References

- Aagaard, K. and Coachman, L.: Toward an ice-free Arctic ocean, *Eos*, Transactions American Geophysical Union, 56, 484–486, 1975.
- Aagaard, K., Swift, J. H., and Carmack, E. C.: Thermohaline circulation in the Arctic Mediterranean Seas, *J. Geophys. Res.: Oceans*, 90, 4833–4846, doi:10.1029/JC090iC03p04833, 1985.
- ACIA: Arctic climate impact assessment, Cambridge University Press, 2005.
- Ahlgren, N. A. and Rocap, G.: Diversity and distribution of marine *Synechococcus*: multiple gene phylogenies for consensus classification and development of qPCR assays for sensitive measurement of clades in the ocean, *Frontiers in Microbiology*, 3, 1–24, doi:10.3389/fmicb.2012.00213, 2012.
- Alonso-Sáez, L., Sánchez, O., Gasol, J. M., Balagué, V., and Pedrós-Alió, C.: Winter-to-summer changes in the composition and single-cell activity of near-surface Arctic prokaryotes, *Environ. Microbiol.*, 10, 2444–2454, 2008.
- Alonso-Sáez, L., Galand, P. E., Casamayor, E. O., Pedrós-Alió, C., and Bertilsson, S.: High bicarbonate assimilation in the dark by Arctic bacteria, *ISME J.*, 4, 1581–1590, 2010.
- Amon, R. M. W.: The role of dissolved organic matter for the organic carbon cycle in the Arctic Ocean, in: *The Organic Carbon Cycle in the Arctic Ocean*, edited by: Stein, R. and Macdonald, R. W., Springer, Berlin, 2004.
- Arrigo, K. R., van Dijken, G., and Pabi, S.: Impact of a shrinking Arctic ice cover on marine primary production, *Geophys. Res. Lett.*, 35, 1–6, 2008.
- Arrigo, K. R., Perovich, D. K., Pickart, R. S., Brown, Z. W., van Dijken, G. L., Lowry, K. E., Mills, M. M., Palmer, M. A., Balch, W. M., and Bahr, F.: Massive phytoplankton blooms under Arctic Sea ice, *Science*, 336, 1408–1408, doi:10.1126/science.1215065, 2012.
- Atamna-Ismaeel, N., Sabehi, G., Sharon, I., Witzel, K. P., Labrenz, M., Jürgens, K., Barkay, T., Stomp, M., Huisman, J., and Béjà, O.: Widespread distribution of proteorhodopsins in freshwater and brackish ecosystems, *ISME J.*, 2, 656–662, 2008.
- Baldwin, A. J., Moss, J. A., Pakulski, J. D., Catala, P., Joux, F., and Jeffrey, W. H.: Microbial diversity in a Pacific Ocean transect from the Arctic to Antarctic circles, *Aquat. Microb. Ecol.*, 41, 91–102, 2005.
- Balzano, S., Marie, D., Gourvil, P., and Vaultot, D.: Composition of the summer photosynthetic pico and nanoplankton communities in the Beaufort Sea assessed by T-RFLP and sequences of the 18S rRNA gene from flow cytometry sorted samples, *ISME J.*, 6, 1480–1498, 2012.
- Bano, N. and Hollibaugh, J. T.: Phylogenetic composition of bacterioplankton assemblages from the Arctic Ocean, *Appl. Environ. Microbiol.*, 68, 505–518, 2002.
- Béjà, O., Aravind, L., Koonin, E. V., Suzuki, M. T., Hadd, A., Nguyen, L. P., Jovanovich, S. B., Gates, C. M., Feldman, R. A., Spudich, J. L., Spudich, E. N., and DeLong, E. F.: Bacterial Rhodopsin: evidence for a new type of phototrophy in the sea, *Science*, 289, 1902–1906, 2000.
- Béjà, O., Spudich, E. N., Spudich, J. L., Leclerc, M., and DeLong, E. F.: Proteorhodopsin phototrophy in the ocean, *Nature*, 411, 786–789, 2001.
- Béjà, O., Suzuki, M. T., Heidelberg, J. F., Nelson, W. C., Preston, C. M., Hamada, T., Eisen, J. A., Fraser, C. M., and DeLong, E. F.: Unsuspected diversity among marine aerobic anoxygenic phototrophs, *Nature*, 415, 630–633, 2002.
- Boeuf, D.: Importance Ecologique des Bactéries Photohétérotrophes dans l'Océan Arctique, Ph.D. Ecole Doctorale des Sciences de l'Environnement d'Ile de France, UMR7144 Adaptation et Diversité en Milieu Marin (Station Biologique de Roscoff), Université Pierre et Marie Curie, Paris, 352 pp., 2013.
- Bowman, J. S., Rasmussen, S., Blom, N., Deming, J. W., Rysgaard, S., and Sicheritz-Ponten, T.: Microbial community struc-

- ture of Arctic multiyear sea ice and surface seawater by 454 sequencing of the 16S RNA gene, *ISME J.*, 6, 11–20, 2012.
- Buck, K., Chavez, F., and Campbell, L.: Basin-wide distributions of living carbon components and the inverted trophic pyramid of the central gyre of the North Atlantic Ocean, summer 1993, *Aquat. Microb. Ecol.*, 10, 283–298, 1996.
- Camill, P.: Permafrost thaw accelerates in boreal peatlands during late-20th century climate warming, *Climatic Change*, 68, 135–152, 2005.
- Campbell, B. J., Waidner, L. A., Cottrell, M. T., and Kirchman, D. L.: Abundant proteorhodopsin genes in the North Atlantic Ocean, *Environ. Microbiol.*, 10, 99–109, 2008.
- Carmack, E., Barber, D., Christensen, J., Macdonald, R., Rudels, B., and Sakshaug, E.: Climate variability and physical forcing of the food webs and the carbon budget on panarctic shelves, *Prog. Oceanogr.*, 71, 145–181, 2006.
- Cauwet, G. and Sidorov, I.: The biogeochemistry of Lena River: organic carbon and nutrients distribution, *Mar. Chem.*, 53, 211–227, 1996.
- Coachman, L. K. and Aagaard, K.: Physical Oceanography of Arctic and Subarctic Seas, DTIC Document, 1974.
- Collins, R. E., Rocap, G., and Deming, J. W.: Persistence of bacterial and archaeal communities in sea ice through an Arctic winter, *Environ. Microbiol.*, 12, 1828–1841, 2010.
- Comeau, A. M., Li, W. K. W., Tremblay, J. É., Carmack, E. C., and Lovejoy, C.: Arctic Ocean microbial community structure before and after the 2007 record sea ice minimum, *PLoS ONE*, 6, e27492, doi:10.1371/journal.pone.0027492, 2011.
- Cottrell, M. T. and Kirchman, D. L.: Photoheterotrophic microbes in the Arctic Ocean in summer and winter, *Appl. Environ. Microb.*, 75, 4958–4966, 2009.
- Cottrell, M. T., Mannino, A., and Kirchman, D. L.: Aerobic anoxygenic phototrophic bacteria in the Mid-Atlantic Bight and the North Pacific Gyre, *Appl. Environ. Microb.*, 72, 557–564, 2006.
- Cottrell, M. T., Ras, J., and Kirchman, D. L.: Bacteriochlorophyll and community structure of aerobic anoxygenic phototrophic bacteria in a particle-rich estuary, *ISME J.*, 4, 945–954, 2010.
- de la Torre, J. R., Christianson, L. M., Béjà, O., Suzuki, M. T., Karl, D. M., Heidelberg, J., and DeLong, E. F.: Proteorhodopsin genes are distributed among divergent marine bacterial taxa, *P. Natl. Acad. Sci. USA*, 100, 12830–12835, doi:10.1073/pnas.2133554100, 2003.
- DeLong, E. F., Preston, C. M., Mincer, T., Rich, V., Hallam, S. J., Frigaard, N. U., Martinez, A., Sullivan, M. B., Edwards, R., and Brito, B. R.: Community genomics among stratified microbial assemblages in the ocean's interior, *Science*, 311, 496–503, 2006.
- Díez, B., Bergman, B., Pedrós-Alió, C., Antó, M., and Snoeijs, P.: High cyanobacterial *nifH* gene diversity in Arctic seawater and sea ice brine, *Environ. Microbiol.*, 4, 360–366, 2012.
- DuRand, M. D., Olson, R. J., and Chisholm, S. W.: Phytoplankton population dynamics at the Bermuda Atlantic Time-series station in the Sargasso Sea, *Deep-Sea Res. Pt. II*, 48, 1983–2003, 2001.
- Elifantz, H., Dittel, A. I., Cottrell, M. T., and Kirchman, D. L.: Dissolved organic matter assimilation by heterotrophic bacterial groups in the western Arctic Ocean, *Aquat. Microb. Ecol.*, 50, 39, 2007.
- Ferrari, V. and Hollibaugh, J.: Distribution of microbial assemblages in the Central Arctic Ocean Basin studied by PCR/DGGE: analysis of a large data set, *Hydrobiologia*, 401, 55–68, 1999.
- Frigaard, N. U., Martinez, A., Mincer, T. J., and DeLong, E. F.: Proteorhodopsin lateral gene transfer between marine planktonic Bacteria and Archaea, *Nature*, 439, 847–850, 2006.
- Fu, Y., MacLeod, D. M., Rivkin, R. B., Chen, F., Buchan, A., and Lang, A. S.: High diversity of Rhodobacterales in the subarctic North Atlantic Ocean and gene transfer agent protein expression in isolated strains, *Aquat. Microb. Ecol.*, 59, 283–293, 2010.
- Galand, P. E., Lovejoy, C., Pouliot, J., Garneau, M. E., and Vincent, W. F.: Microbial community diversity and heterotrophic production in a coastal Arctic ecosystem: A stamukhi lake and its source waters, *Limnol. Oceanogr.*, 53, 813–823, 2008.
- Galand, P. E., Casamayor, E. O., Kirchman, D. L., and Lovejoy, C.: Ecology of the rare microbial biosphere of the Arctic Ocean, *Proc. Natl. Acad. Sci.*, 106, 22427–22432, doi:10.1073/pnas.0908284106, 2009.
- Galand, P. E., Potvin, M., Casamayor, E. O., and Lovejoy, C.: Hydrography shapes bacterial biogeography of the deep Arctic Ocean, *ISME J.*, 4, 564–576, 2010.
- Garneau, M. È., Vincent, W. F., Alonso-Sáez, L., Gratton, Y., and Lovejoy, C.: Prokaryotic community structure and heterotrophic production in a river-influenced coastal arctic ecosystem, *Aquat. Microb. Ecol.*, 42, 27–40, 2006.
- Garneau, M. È., Roy, S., Lovejoy, C., Gratton, Y., and Vincent, W. F.: Seasonal dynamics of bacterial biomass and production in a coastal arctic ecosystem: Franklin Bay, western Canadian Arctic, *J. Geophys. Res.*, 113, C07S91, doi:10.1029/2007JC004281, 2008.
- Ghiglione, J. and Murray, A.: Pronounced summer to winter differences and higher wintertime richness in coastal Antarctic marine bacterioplankton, *Environ. Microbiol.*, 14, 617–629, 2012.
- Ghiglione, J. F., Galand, P. E., Pommier, T., Pedrós-Alió, C., Maas, E. W., Bakker, K., Bertilson, S., Kirchman, D. L., Lovejoy, C., and Yager, P. L.: Pole-to-pole biogeography of surface and deep marine bacterial communities, *Proc. Natl. Acad. Sci.*, 109, 17633–17638, 2012.
- Giovannoni, S. J. and Stingl, U.: Molecular diversity and ecology of microbial plankton, *Nature*, 437, 343–348, 2005.
- Giovannoni, S. J., Bibbs, L., Cho, J. C., Stapels, M. D., Desiderio, R., Vergin, K. L., Rappé, M. S., Laney, S., Wilhelm, L. J., and Tripp, H. J.: Proteorhodopsin in the ubiquitous marine bacterium SAR11, *Nature*, 438, 82–85, 2005.
- Gómez-Consarnau, L., González, J. M., Coll-Lladó, M., Gourdon, P., Pascher, T., Neutze, R., Pedrós-Alió, C., and Pinhassi, J.: Light stimulates growth of proteorhodopsin-containing marine Flavobacteria, *Nature*, 445, 210–213, 2007.
- Gómez-Consarnau, L., Akram, N., Lindell, K., Pedersen, A., Neutze, R., Milton, D. L., González, J. M., and Pinhassi, J.: Proteorhodopsin Phototrophy Promotes Survival of Marine Bacteria during Starvation, *PLoS Biol.*, 8, e1000358, doi:10.1371/journal.pbio.1000358, 2010.
- González, J. M., Simó, R., Massana, R., Covert, J. S., Casamayor, E. O., Pedrós-Alió, C., and Moran, M. A.: Bacterial community structure associated with a dimethylsulfoniopropionate-producing North Atlantic algal bloom, *Appl. Environ. Microbiol.*, 66, 4237–4246, 2000.

- González, J. M., Fernández-Gómez, B., Fernández-Guerra, A., Gómez-Consarnau, L., Sánchez, O., Coll-Lladó, M., Del Campo, J., Escudero, L., Rodríguez-Martínez, R., and Alonso-Sáez, L.: Genome analysis of the proteorhodopsin-containing marine bacterium *Polaribacter* sp. MED152 (Flavobacteria), *Proc. Natl. Acad. Sci.*, 105, 8724–8729, doi:10.1073/pnas.0712027105, 2008.
- Gradinger, R. and Lenz, J.: Picocyanobacteria in the high Arctic, *Mar. Ecol. Prog. Ser.*, 52, 99–101, 1989.
- Harashima, K., Shiba, T., Totsuka, T., Simidu, U., and Taga, N.: Occurrence of bacteriochlorophyll *a* in a strain of an aerobic heterotrophic bacterium, *Agric. Biol. Chem.*, 42, 1627–1628, 1978.
- Hojerová, E., Mašín, M., Brunet, C., Ferrera, I., Gasol, J. M., and Koblížek, M.: Distribution and Growth of Aerobic Anoxygenic Phototrophs in the Mediterranean Sea, *Environ. Microbiol.*, 13, 2717–2725, doi:10.1111/j.1462-2920.2011.02540.x, 2011.
- Hopcroft, R., Bluhm, B., Gradinger, R., Whitley, T., Weingartner, T., Norcross, B., and Springer, A.: Arctic ocean synthesis: Analysis of climate change impacts in the Chukchi and Beaufort seas with strategies for future research, Institute of Marine Science, University of Alaska, Fairbanks, 2008.
- Hu, Y., Du, H., Jiao, N., and Zeng, Y.: Abundant presence of the γ -like Proteobacterial *pufM* gene in oxic seawater, *FEMS Microbiol. Lett.*, 263, 200–206, 2006.
- Huang, S., Wilhelm, S. W., Harvey, H. R., Taylor, K., Jiao, N., and Chen, F.: Novel lineages of *Prochlorococcus* and *Synechococcus* in the global oceans, *ISME J.*, 6, 285–297, 2012.
- Jeanthon, C., Boeuf, D., Dahan, O., Le Gall, F., Garczarek, L., Bendif, E. M., and Lehours, A.-C.: Diversity of cultivated and metabolically active aerobic anoxygenic phototrophic bacteria along an oligotrophic gradient in the Mediterranean Sea, *Biogeosciences*, 8, 1955–1970, doi:10.5194/bg-8-1955-2011, 2011.
- Jiao, N., Zhang, Y., Zeng, Y., Hong, N., Liu, R., Chen, F., and Wang, P.: Distinct distribution pattern of abundance and diversity of aerobic anoxygenic phototrophic bacteria in the global ocean, *Environ. Microbiol.*, 9, 3091–3099, 2007.
- Johnson, Z. I., Zinser, E. R., Coe, A., McNulty, N. P., Woodward, E. M. S., and Chisholm, S. W.: Niche partitioning among *Prochlorococcus* ecotypes along ocean-scale environmental gradients, *Science*, 311, 1737–1740, doi:10.1126/science.1118052, 2006.
- Kaneko, T., Atlas, R. M., and Krichevsky, M.: Diversity of bacterial populations in the Beaufort Sea, *Nature*, 270, 596–599, 1977.
- Karr, E. A., Sattley, W. M., Jung, D. O., Madigan, M. T., and Achenbach, L. A.: Remarkable diversity of phototrophic purple bacteria in a permanently frozen Antarctic lake, *Appl. Environ. Microbiol.*, 69, 4910–4914, 2003.
- Kellogg, C. T. and Deming, J. W.: Comparison of free-living, suspended particle, and aggregate-associated bacterial and archaeal communities in the Laptev Sea, *Aquat. Microb. Ecol.*, 57, 1–18, 2009.
- Kirchman, D. L., Elifantz, H., Dittel, A. I., Malmstrom, R. R., and Cottrell, M. T.: Standing stocks and activity of Archaea and Bacteria in the western Arctic Ocean, *Limnol. Oceanogr.*, 52, 495–507, 2007.
- Kirchman, D. L., Cottrell, M. T., and Lovejoy, C.: The structure of bacterial communities in the western Arctic Ocean as revealed by pyrosequencing of 16S rRNA genes, *Environ. Microbiol.*, 12, 1132–1143, 2010.
- Kling, G. W., Kipphut, G. W., and Miller, M. C.: Arctic lakes and streams as gas conduits to the atmosphere: Implications for tundra carbon budgets, *Science*, 251, 298–301, doi:10.1126/science.251.4991.298, 1991.
- Koblížek, M., Mašín, M., Ras, J., Poulton, A. J., and Prášil, O.: Rapid growth rates of aerobic anoxygenic phototrophs in the ocean, *Environ. Microbiol.*, 9, 2401–2406, 2007.
- Koh, E. Y., Atamna-Ismaeel, N., Martin, A., Cowie, R. O. M., Bějá, O., Davy, S. K., Maas, E. W., and Ryan, K. G.: Proteorhodopsin-bearing bacteria in Antarctic sea ice, *Appl. Environ. Microbiol.*, 76, 5918–5925, 2010.
- Koh, E. Y., Phua, W., and Ryan, K. G.: Aerobic anoxygenic phototrophic bacteria in Antarctic sea ice and seawater, *Environ. Microbiol. Rep.*, 3, 710–716, 2011.
- Koh, E. Y., Cowie, R. O. M., Simpson, A. M., O’Toole, R., and Ryan, K. G.: The origin of cyanobacteria in Antarctic sea ice: marine or freshwater?, *Environ. Microbiol. Rep.*, 4, 479–483, 2012.
- Kolber, Z. S., Van Dover, C. L., Niederman, R. A., and Falkowski, P. G.: Bacterial photosynthesis in surface waters of the open ocean, *Nature*, 407, 177–179, 2000.
- Kolber, Z. S., Gerald, F., Plumley, Lang, A. S., Beatty, J. T., Blankenship, R. E., VanDover, C. L., Vetriani, C., Koblizek, M., Rathgeber, C., and Falkowski, P. G.: Contribution of Aerobic Photoheterotrophic Bacteria to the Carbon Cycle in the Ocean, *Science*, 292, 2492–2495, doi:10.1126/science.1059707, 2001.
- Lami, R., Cottrell, M. T., Ras, J., Ulloa, O., Obernosterer, I., Claustre, H., Kirchman, D. L., and Lebaron, P.: High abundances of aerobic anoxygenic photosynthetic bacteria in the South Pacific Ocean, *Appl. Environ. Microbiol.*, 73, 4198–4205, doi:10.1128/AEM.02652-06, 2007.
- Lami, R., Cottrell, M. T., Campbell, B. J., and Kirchman, D. L.: Light-dependent growth and proteorhodopsin expression by Flavobacteria and SAR11 in experiments with Delaware coastal waters, *Environ. Microbiol.*, 11, 3201–3209, doi:10.1111/j.1462-2920.2009.02028.x, 2009.
- Lamy, D., Jeanthon, C., Cottrell, M. T., Kirchman, D. L., Van Wambeke, F., Ras, J., Dahan, O., Pujo-Pay, M., Oriol, L., Bariat, L., Catala, P., Cornet-Barthaux, V., and Lebaron, P.: Ecology of aerobic anoxygenic phototrophic bacteria along an oligotrophic gradient in the Mediterranean Sea, *Biogeosciences*, 8, 973–985, doi:10.5194/bg-8-973-2011, 2011.
- Lehours, A. C., Cottrell, M. T., Dahan, O., Kirchman, D. L., and Jeanthon, C.: Summer distribution and diversity of aerobic anoxygenic phototrophic bacteria in the Mediterranean Sea in relation to environmental variables, *FEMS Microbiol. Ecol.*, 74, 397–409, doi:10.1111/j.1574-6941.2010.00954.x, 2010.
- Li, W. K. W.: Primary production of prochlorophytes, cyanobacteria, and eucaryotic ultraphytoplankton: measurements from flow cytometric sorting, *Limnol. Oceanogr.*, 39, 169–175, 1994.
- Li, W. K. W.: Annual average abundance of heterotrophic bacteria and *Synechococcus* in surface ocean waters, *Limnol. Oceanogr.*, 43, 1746–1753, 1998.
- Li, W. K. W.: From cytometry to macroecology: a quarter century quest in microbial oceanography, *Aquat. Microb. Ecol.*, 57, 239–251, 2009.
- Lin, L., He, J., Zhao, Y., Zhang, F., and Cai, M.: Flow cytometry investigation of picoplankton across latitudes and along the circum Antarctic Ocean, *Acta Oceanol. Sin.*, 31, 134–142, 2012.

- Malmström, R. R., Kiene, R. P., Vila, M., and Kirchman, D. L.: Dimethylsulfoniopropionate (DMSP) assimilation by *Synechococcus* in the Gulf of Mexico and northwest Atlantic Ocean, *Limnol. Oceanogr.*, 50, 1924–1931, 2005.
- Malmström, R. R., Straza, T. R. A., Cottrell, M. T., and Kirchman, D. L.: Diversity, abundance, and biomass production of bacterial groups in the western Arctic Ocean, *Aquat. Microb. Ecol.*, 47, 45–55, 2007.
- Man, D., Wang, W., Sabehi, G., Aravind, L., Post, A. F., Massana, R., Spüdich, E. N., Spüdich, J. L., and Bějí, O.: Diversification and spectral tuning in marine proteorhodopsins, *EMBO J.*, 22, 1725–1731, 2003.
- Marchant, H. J., Davidson, A. T., and Wright, S. W.: The Distribution and Abundance of Chroococoid Cyanobacteria in the Southern Ocean, National Institute of Polar Research, 1987.
- Mašín, M., Zdun, A., Ston-Egiert, J., Nausch, M., Labrenz, M., Moulisova, V., and Koblížek, M.: Seasonal changes and diversity of aerobic anoxygenic phototrophs in the Baltic Sea, *Aquat. Microb. Ecol.*, 45, 247–254, 2006.
- Meon, B. and Amon, R. M. W.: Heterotrophic bacterial activity and fluxes of dissolved free amino acids and glucose in the Arctic rivers Ob, Yenisei and the adjacent Kara Sea, *Aquat. Microb. Ecol.*, 37, 121–135, 2004.
- Michelou, V. K., Cottrell, M. T., and Kirchman, D. L.: Light-stimulated bacterial production and amino acid assimilation by cyanobacteria and other microbes in the North Atlantic Ocean, *Appl. Environ. Microbiol.*, 73, 5539–5546, 2007.
- Murphy, L. and Haugen, E.: The distribution and abundance of phototrophic ultraplankton in the North Atlantic, *Limnol. Oceanogr.*, 30, 47–58, 1985.
- Nelson, F. E.: (Un) frozen in time, *Science*, 299, 1673–1675, doi:10.1126/science.1081111, 2003.
- Niederberger, T. D., Perreault, N. N., Tille, S., Lollar, B. S., Lacrampe-Couloume, G., Andersen, D., Greer, C. W., Pollard, W., and Whyte, L. G.: Microbial characterization of a subzero, hypersaline methane seep in the Canadian High Arctic, *ISME J.*, 4, 1326–1339, 2010.
- Nikrad, M. P., Cottrell, M. T., and Kirchman, D. L.: Abundance and Single-Cell Activity of Heterotrophic Bacterial Groups in the Western Arctic Ocean in Summer and Winter, *Appl. Environ. Microbiol.*, 78, 2402–2409, doi:10.1128/aem.07130-11, 2012.
- Oesterhelt, D. and Stoekenius, W.: Rhodopsin-like protein from the purple membrane of *Halobacterium halobium*, *Nature*, 233, 149–152, 1971.
- Opsahl, S., Benner, R., and Amon, R. M. W.: Major flux of terrigenous dissolved organic matter through the Arctic Ocean, *Limnol. Oceanogr.*, 44, 2017–2023, 1999.
- Ortega-Retuerta, E., Jeffrey, W. H., Babin, M., Bélanger, S., Benner, R., Marie, D., Matsuoka, A., Raimbault, P., and Joux, F.: Carbon fluxes in the Canadian Arctic: patterns and drivers of bacterial abundance, production and respiration on the Beaufort Sea margin, *Biogeosciences*, 9, 3679–3692, doi:10.5194/bg-9-3679-2012, 2012.
- Ortega-Retuerta, E., Joux, F., Jeffrey, W. H., and Ghiglione, J. F.: Spatial variability of particle-attached and free-living bacterial diversity in surface waters from the Mackenzie River to the Beaufort Sea (Canadian Arctic), *Biogeosciences*, 10, 2747–2759, doi:10.5194/bg-10-2747-2013, 2013.
- Palenik, B., Brahamsha, B., Larimer, F., Land, M., Hauser, L., Chain, P., Lamerdin, J., Regala, W., Allen, E., and McCarren, J.: The genome of a motile marine *Synechococcus*, *Nature*, 424, 1037–1042, 2003.
- Partensky, F., Blanchot, J., and Vaulot, D.: Differential distribution and ecology of *Prochlorococcus* and *Synechococcus* in oceanic waters: a review, *Bulletin-Institut Océanographique Monaco*, numéro spécial, 19, 457–476, 1999a.
- Partensky, F., Hess, W., and Vaulot, D.: *Prochlorococcus*, a marine photosynthetic prokaryote of global significance, *Microbiol. Mol. Biol. R.*, 63, 106–127, 1999b.
- Perovich, D. K.: The changing Arctic sea ice cover, *Oceanography*, 24, 162–173, 2011.
- Peterson, B. J., Holmes, R. M., McClelland, J. W., Vörösmarty, C. J., Lammers, R. B., Shiklomanov, A. I., Shiklomanov, I. A., and Rahmstorf, S.: Increasing river discharge to the Arctic Ocean, *Science*, 298, 2171–2173, 2002.
- Pidwirny, M.: Introduction to the Oceans, in: *Fundamentals of Physical Geography*, www.physicalgeography.net, 2nd Edn. Ch. 8, available at: <http://www.physicalgeography.net/fundamentals/8o.html>, 2006.
- Pommier, T., Canbäck, B., Riemann, L., Boström, K. H., Simu, K., Lundberg, P., Tunlid, A., and Hagström, Å.: Global patterns of diversity and community structure in marine bacterioplankton, *Mol. Ecol.*, 16, 867–880, 2007.
- Powell, L., Bowman, J., Skerratt, J., Franzmann, P., and Burton, H.: Ecology of a novel *Synechococcus* clade occurring in dense populations in saline Antarctic lakes, *Mar. Ecol. Prog. Ser.*, 291, 65–80, 2005.
- Prabakaran, S. R., Manorama, R., Delille, D., and Shivaji, S.: Prevalence of *Roseobacter*, *Sulfitobacter*, *Glaciecola* and *Psychrobacter* in seawater collected off Ushuaia, Argentina, Sub-Antarctica, *FEMS Microbiol. Ecol.*, 59, 342–355, 2007.
- Rachold, V., Grigoriev, M. N., Are, F. E., Solomon, S., Reimnitz, E., Kassens, H., and Antonow, M.: Coastal erosion vs riverine sediment discharge in the Arctic Shelf seas, *Int. J. Earth Sci.*, 89, 450–460, 2000.
- Rathgeber, C., Beatty, J. T., and Yurkov, V.: Aerobic phototrophic bacteria: new evidence for the diversity, ecological importance and applied potential of this previously overlooked group, *Photosynth. Res.*, 81, 113–128, 2004.
- Riedel, T., Tomasch, J., Buchholz, I., Jacobs, J., Kollenberg, M., Gerds, G., Wichels, A., Brinkhoff, T., Cypionka, H., and Wagner-Döbler, I.: Constitutive expression of the proteorhodopsin gene by a *Flavobacterium* strain representative of the proteorhodopsin-producing microbial community in the North Sea, *Appl. Environ. Microbiol.*, 76, 3187–3197, 2010.
- Robineau, B., Legendre, L., Michel, C., Budéus, G., Kattner, G., Schneider, W., and Pesant, S.: Ultraphytoplankton abundances and chlorophyll *a* concentrations in ice-covered waters of northern seas, *J. Plankton Res.*, 21, 735–755, 1999.
- Rosenberg, G.: Environmental control and potential fate of size-fractionated phytoplankton production in the Greenland Sea (75° N), *Mar. Ecol. Prog. Ser.*, 98, 297–313, 1993.
- Rusch, D. B., Halpern, A. L., Sutton, G., Heidelberg, K., Williamson, S., and Yooseph, S.: The Sorcerer II Global Ocean Sampling Expedition: Northwest Atlantic through Eastern Tropical Pacific, *PLoS Biol.*, 5, e77, doi:10.1371/journal.pbio.0050077, 2007.

- Sabehi, G., Beja, O., Suzuki, M. T., Preston, C. M., and DeLong, E. F.: Different SAR86 subgroups harbour divergent proteorhodopsins, *Environ. Microbiol.*, 6, 903–910, doi:10.1111/j.1462-2920.2004.00676.x, 2004.
- Sabehi, G., Loy, A., Jung, K. H., Partha, R., Spudich, J. L., Isaacson, T., Hirschberg, J., Wagner, M., and B ej a, O.: New insights into metabolic properties of marine bacteria encoding proteorhodopsins, *PLoS Biol.*, 3, e273, doi:10.1371/journal.pbio.0030273, 2005.
- Salka, I., Moulisova, V., Kobl izek, M., Jost, G., J urgens, K., and Labrenz, M.: Abundance, depth distribution, and composition of aerobic bacteriochlorophyll *a*-producing bacteria in four basins of the central Baltic Sea, *Appl. Environ. Microbiol.*, 74, 4398–4404, 2008.
- Salka, I.,  Cuperov a, Z., Ma s in, M., Kobl izek, M., and Grossart, H. P.: *Rhodospirillum rubrum*-related *pufM* gene cluster dominates the aerobic anoxygenic phototrophic communities in German freshwater lakes, *Environ. Microbiol.*, 13, 2865–2875, 2011.
- Schwalbach, M. S. and Fuhrman, J. A.: Wide-ranging abundances of aerobic anoxygenic phototrophic bacteria in the world ocean revealed by epifluorescence microscopy and quantitative PCR, *Limnol. Oceanogr.*, 50, 620–628, 2005.
- Selje, N., Simon, M., and Brinkhoff, T.: A newly discovered *Roseobacter* cluster in temperate and polar oceans, *Nature*, 427, 445–448, 2004.
- Serreze, M. C., Barrett, A. P., Slater, A. G., Woodgate, R. A., Aagaard, K., Lammers, R. B., Steele, M., Moritz, R., Meredith, M., and Lee, C. M.: The large-scale freshwater cycle of the Arctic, *J. Geophys. Res.: Oceans*, 111, C11010, doi:10.1029/2005JC003424, 2006.
- Shiba, T., Simidu, U., and Taga, N.: Distribution of aerobic bacteria which contain bacteriochlorophyll *a*, *Appl. Environ. Microbiol.*, 38, 43–45, 1979.
- Sieracki, M. E., Gilg, I. C., Thier, E. C., Poulton, N. J., and Goericke, R.: Distribution of planktonic aerobic anoxygenic photoheterotrophic bacteria in the northwest Atlantic, *Limnol. Oceanogr.*, 51, 38–46, 2006.
- Steindler, L., Schwalbach, M. S., Smith, D. P., Chan, F., and Giovannoni, S. J.: Energy Starved Candidatus *Pelagibacter Ubique* Substitutes Light-Mediated ATP Production for Endogenous Carbon Respiration, *PLoS ONE*, 6, e19725, doi:10.1371/journal.pone.0019725, 2011.
- Stingl, U., Desiderio, R. A., Cho, J. C., Vergin, K. L., and Giovannoni, S. J.: The SAR92 clade: an abundant coastal clade of culturable marine bacteria possessing proteorhodopsin, *Appl. Environ. Microbiol.*, 73, 2290–2296, 2007.
- Suzuki, M. T., Preston, C. M., Chavez, F. P., and DeLong, E. F.: Quantitative mapping of bacterioplankton populations in sea-water: field tests across an upwelling plume in Monterey Bay, *Aquat. Microb. Ecol.*, 24, 117–127, 2001.
- Veldhuis, M. J. W., Kraay, G. W., Van Bleijswijk, J. D. L., and Baars, M. A.: Seasonal and spatial variability in phytoplankton biomass, productivity and growth in the northwestern Indian Ocean: The southwest and northeast monsoon, 1992–1993, *Deep-Sea Res. Pt. I*, 44, 425–449, 1997.
- Venter, J. C., Remington, K., Heidelberg, J. F., Halpern, A. L., Rusch, D., Eisen, J. A., Wu, D., Paulsen, I., Nelson, K. E., Nelson, W., Fouts, D. E., Levy, S., Knap, A. H., Lomas, M. W., Nealson, K., White, O., Peterson, J., Hoffman, J., Parsons, R., Baden-Tillson, H., Pfannkoch, C., Rogers, Y.-H., and Smith, H. O.: Environmental Genome Shotgun Sequencing of the Sargasso Sea, *Science*, 304, 66–74, doi:10.1126/science.1093857, 2004.
- Vincent, W., Bowman, J., Rankin, L., and McMeekin, T.: Phylogenetic diversity of picocyanobacteria in Arctic and Antarctic ecosystems, in: Proceedings of the 8th International Symposium on Microbial Ecology, Halifax, Canada, 317–322, 2000.
- Waidner, L. A. and Kirchman, D. L.: Aerobic anoxygenic photosynthesis genes and operons in uncultured bacteria in the Delaware River, *Environ. Microbiol.*, 7, 1896–1908, doi:10.1111/j.1462-2920.2005.00883.x, 2005.
- Waidner, L. A. and Kirchman, D. L.: Aerobic anoxygenic phototrophic bacteria attached to particles in turbid waters of the Delaware and Chesapeake estuaries, *Appl. Environ. Microbiol.*, 73, 3936–3944, 2007.
- Waidner, L. A. and Kirchman, D. L.: Diversity and distribution of ecotypes of the aerobic anoxygenic phototrophy gene, *pufM*, in the Delaware estuary, *Appl. Environ. Microbiol.*, 74, 4012–4021, 2008.
- Waleron, M., Waleron, K., Vincent, W. F., and Wilmotte, A.: Allochthonous inputs of riverine picocyanobacteria to coastal waters in the Arctic Ocean, *FEMS Microbiol. Ecol.*, 59, 356–365, 2007.
- Walsh, J. E.: Climate of the Arctic marine environment, *Ecol. Appl.*, 18, 3–22, 2008.
- Wells, L. E. and Deming, J. W.: Abundance of Bacteria, the Cytophaga-Flavobacterium cluster and Archaea in cold oligotrophic waters and nepheloid layers of the Northwest Passage, Canadian Archipelago, *Aquat. Microb. Ecol.*, 31, 19–31, 2003.
- Wheeler, P. A., Gosselin, M., Sherr, E., Thibault, D., Kirchman, D. L., Benner, R., and Whitley, T. E.: Active cycling of organic carbon in the central Arctic Ocean, *Nature*, 380, 697–699, doi:10.1038/380697a0, 1996.
- Williams, T. J., Long, E., Evans, F., DeMaere, M. Z., Lauro, F. M., Raftery, M. J., Ducklow, H., Grzymiski, J. J., Murray, A. E., and Cavicchioli, R.: A metaproteomic assessment of winter and summer bacterioplankton from Antarctic Peninsula coastal surface waters, *ISME J.*, 6, 1883–1900, 2012.
- Yoshizawa, S., Kawanabe, A., Ito, H., Kandori, H., and Kogure, K.: Diversity and functional analysis of proteorhodopsin in marine *Flavobacteria*, *Environ. Microbiol.*, 14, 1240–1248, 2012.
- Yurkov, V. V. and Beatty, J. T.: Aerobic anoxygenic phototrophic bacteria, *Microbiol. Mol. Biol. Rev.*, 62, 695–724, 1998.
- Yutin, N., Suzuki, M. T., and B ej a, O.: Novel primers reveal wider diversity among marine aerobic anoxygenic phototrophs, *Appl. Environ. Microbiol.*, 71, 8958–8962, 2005.
- Yutin, N., Suzuki, M. T., Teeling, H., Weber, M., Venter, J. C., Rusch, D. B., and B ej a, O.: Assessing diversity and biogeography of aerobic anoxygenic phototrophic bacteria in surface waters of the Atlantic and Pacific Oceans using the Global Ocean Sampling expedition metagenomes, *Environ. Microbiol.*, 9, 1464–1475, 2007.
- Zaballos, M., L opez-L opez, A., Ovreas, L., Bartual, S. G., D’Auria, G., Alba, J. C., Legault, B., Pushker, R., Daae, F. L., and Rodr iguez-Valera, F.: Comparison of prokaryotic diversity at offshore oceanic locations reveals a different microbiota in the Mediterranean Sea, *FEMS Microbiol. Ecol.*, 56, 389–405, 2006.
- Zhang, J., Spitz, Y. H., Steele, M., Ashjian, C., Campbell, R., Berlina, L., and Matrai, P.: Modeling the impact of declining sea

- ice on the Arctic marine planktonic ecosystem, *J. Geophys. Res.: Oceans*, 115, C10015, doi:10.1029/2009JC005387, 2010.
- Zobell, C. E. (Ed.): *Marine microbiology. A monograph on hydrobacteriology*, Chronica Botánica Company, Waltham, USA, 1946.
- Zubkov, M. V. and Tarran, G. A.: Amino acid uptake of *Prochlorococcus* spp. in surface waters across the South Atlantic Subtropical Front, *Aquat. Microb. Ecol.*, 40, 241–249, 2005.
- Zwirgmaier, K., Jardillier, L., Ostrowski, M., Mazard, S., Garczarek, L., Vault, D., Massana, R., Ulloa, O., and Scanlan, D. J.: Global phylogeography of marine *Synechococcus* and *Prochlorococcus* reveals a distinct partitioning of lineages among oceanic biomes, *Environ. Microbiol.*, 10, 147–161, 2008.