



**HAL**  
open science

# Does Encapsulation Protect Embryos from the Effects of Ocean Acidification? The Example of *Crepidula fornicata*

Fanny Noisette, Thierry Comtet, Erwann Legrand, François Bordeyne,  
Dominique Davoult, Sophie Martin

## ► To cite this version:

Fanny Noisette, Thierry Comtet, Erwann Legrand, François Bordeyne, Dominique Davoult, et al.. Does Encapsulation Protect Embryos from the Effects of Ocean Acidification? The Example of *Crepidula fornicata*. PLoS ONE, 2014, 9 (3), pp.e93021. 10.1371/journal.pone.0093021. hal-01097262

**HAL Id: hal-01097262**

**<https://hal.sorbonne-universite.fr/hal-01097262>**

Submitted on 19 Dec 2014

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Does Encapsulation Protect Embryos from the Effects of Ocean Acidification? The Example of *Crepidula fornicata*

Fanny Noisette<sup>1,2\*</sup>, Thierry Comtet<sup>1,2</sup>, Erwann Legrand<sup>1,2</sup>, François Bordeyne<sup>1,2</sup>, Dominique Davoult<sup>1,2</sup>, Sophie Martin<sup>1,2</sup>

<sup>1</sup> Sorbonne Universités, UPMC Univ. Paris 06, UMR 7144, Station Biologique de Roscoff, Roscoff, France, <sup>2</sup> CNRS, UMR 7144, Station Biologique de Roscoff, Roscoff, France

## Abstract

Early life history stages of marine organisms are generally thought to be more sensitive to environmental stress than adults. Although most marine invertebrates are broadcast spawners, some species are brooders and/or protect their embryos in egg or capsules. Brooding and encapsulation strategies are typically assumed to confer greater safety and protection to embryos, although little is known about the physico-chemical conditions within egg capsules. In the context of ocean acidification, the protective role of encapsulation remains to be investigated. To address this issue, we conducted experiments on the gastropod *Crepidula fornicata*. This species broods its embryos within capsules located under the female and veliger larvae are released directly into the water column. *C. fornicata* adults were reared at the current level of CO<sub>2</sub> partial pressure (*p*CO<sub>2</sub>) (390 μatm) and at elevated levels (750 and 1400 μatm) before and after fertilization and until larval release, such that larval development occurred entirely at a given *p*CO<sub>2</sub>. The *p*CO<sub>2</sub> effects on shell morphology, the frequency of abnormalities and mineralization level were investigated on released larvae. Shell length decreased by 6% and shell surface area by 11% at elevated *p*CO<sub>2</sub> (1400 μatm). The percentage of abnormalities was 1.5- to 4-fold higher at 750 μatm and 1400 μatm *p*CO<sub>2</sub>, respectively, than at 390 μatm. The intensity of birefringence, used as a proxy for the mineralization level of the larval shell, also decreased with increasing *p*CO<sub>2</sub>. These negative results are likely explained by increased intracapsular acidosis due to elevated *p*CO<sub>2</sub> in extracapsular seawater. The encapsulation of *C. fornicata* embryos did not protect them against the deleterious effects of a predicted *p*CO<sub>2</sub> increase. Nevertheless, *C. fornicata* larvae seemed less affected than other mollusk species. Further studies are needed to identify the critical points of the life cycle in this species in light of future ocean acidification.

**Citation:** Noisette F, Comtet T, Legrand E, Bordeyne F, Davoult D, et al. (2014) Does Encapsulation Protect Embryos from the Effects of Ocean Acidification? The Example of *Crepidula fornicata*. PLoS ONE 9(3): e93021. doi:10.1371/journal.pone.0093021

**Editor:** Pauline Ross, University of Western Sydney, Australia

**Received:** September 9, 2013; **Accepted:** February 27, 2014; **Published:** March 26, 2014

**Copyright:** © 2014 Noisette et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was supported by the CALCAO project, which received funding from the Region Bretagne, and contributed to the "European Project on Ocean Acidification" (EPOCA) funded by the European Community's Seventh Framework Programme (FP7/2007-2013) under grant agreement n° 211384. It was also supported by the Interreg IV a France (Channel) – England Marinexus project n°4073 funded by the FEDER programme. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: fanny.noisette@sb-roscoff.fr

## Introduction

Early life history stages of marine species, including embryos and larvae, are of crucial importance in population dynamics as they ensure dispersion, colonize new areas and sustain populations [1]. Their success in development and final recruitment are essential for the persistence of viable populations. Early stages of marine invertebrates are in general morphologically and ecologically distinct from the adult stage and are generally thought to be more sensitive to environmental stress [1] although, in some cases, they may be more tolerant than adults, e.g. some Antarctic species exposed to warming [2]. In the context of climate change, early development may be affected by various factors, such as temperature increases, hypoxia zones or ocean acidification. Due to the increase in atmospheric *p*CO<sub>2</sub> predicted for the end of the century (from 475 to 1313 μatm according to the Intergovernmental Panel on Climate Change (IPCC)), pH in surface seawaters is likely to decline by 0.06 – 0.32 units [3], leading to a decrease in carbonate ion concentrations (CO<sub>3</sub><sup>2-</sup>) and a reduction in the calcium carbonate saturation state (Ω) [4]. Due to these changes in seawater carbonate chemistry, ocean acidification is considered a

major threat to calcifying marine species, affecting their physiology and impairing their ability to build calcium carbonate shells and skeletons [8,9,10,11], which can ultimately modify their behavior and distribution [5,6,7]. Early life stages (embryos, larvae and juveniles) of calcifying species are thus expected to be highly affected by ocean acidification [12,13], as opposed to non-calcified larvae which are predicted to be more tolerant [14,15]. This relatively higher vulnerability is likely due to fragile larval skeletons [9] and their high ratio of exposed surface-to-body mass compared to adults [16]. Identifying life history stages that are the most vulnerable to global change is needed to determine bottlenecks for species persistence and addressing their sensitivity to acidification is a major issue in a changing ocean [9].

Responses to near-future (end of century) levels of *p*CO<sub>2</sub> depend on species, populations, habitats and developmental stages [9,17,18,19] and understanding these effects on the early life stages requires taking into account the complete developmental cycle, from egg to juvenile [12]. In particular, the impact of elevated *p*CO<sub>2</sub>/decreased pH on early life stages has been investigated in a broad range of species, including corals [20,21], echinoderms [22,23,24], crustaceans [15,16,25], mollusks [26,27],

and fish [28,29]. In mollusks, which have been studied intensively (see refs. [26,30] for a review), deleterious effects of increased  $p\text{CO}_2$  have been demonstrated on fertilization success [31,32], hatching success [33,34], larval survival [35,36], growth [37,38], shell formation [39,40], development duration [41,42], and settlement [43,44].

Most of the species studied are broadcast spawners, which may be considered particularly vulnerable to ocean acidification because fertilization and complete pelagic larval life occur in the water column [9,45]. Whether alternative reproductive modes are affected in a changing ocean is still poorly documented. Brooding and/or egg laying in egg masses or capsules are typically assumed to confer protection to developing embryos [46,47]. For example, it has been shown that encapsulated embryos of some gastropod species survive better in conditions of salinity stress than embryos removed from their capsule [48,49]. A few studies have explored the effects of decreased pH on embryos brooded and/or laid in benthic gelatinous egg masses or in egg capsules in bivalve [50], gastropod [35,41,51,52] or cephalopod [27,53] mollusks. Depending on the study, reduced pH has different effects that are related to the range of species habitats and the strategy to protect embryos (brooding, egg masses, capsules) studied as well as the source of pH change ( $p\text{CO}_2$  increase, salinity stress). Encapsulation has been suggested to protect embryos against ocean acidification [41,54], whereby the buffer capacity of intracapsular fluids may reduce the potential effect of extracapsular elevated  $p\text{CO}_2$  seawater.

To study this issue in a non-broadcast-spawner species, we chose the slipper limpet *Crepidula fornicata*, Linné 1758 (Gastropoda) as our biological model. Native to the northeast American coast, this species was introduced in Europe at the end of the 19<sup>th</sup> century, primarily via oyster farming [55], and has now become invasive in bays and estuaries where it reaches very high densities of up to several thousands of individuals per  $\text{m}^2$  [56]. It has a benthic-pelagic life cycle, with a number of original features. Benthic adults form stacks with males at the top and females at the bottom. After internal fertilization, females brood their embryos in egg capsules for 3 to 4 weeks [57,58]. Capsules are protected between the neck and the propodium of the female parent and attached to the substratum to which the female is fixed [58]. Each female spawns between 28 and 64 capsules, each containing 300 to 500 embryos [57]. At the end of capsular development, veliger larvae are released at a size of about 400  $\mu\text{m}$  in length into the water column where they spend between 2 and 7 weeks [59,60]. Upon reaching competence (800–1000  $\mu\text{m}$  in length), larvae are able to metamorphose and settle on hard substrata [59,60].

The objective of this work was to investigate the effects of near-future levels of  $p\text{CO}_2$  on the development of *C. fornicata* encapsulated embryos by studying the shell morphology and mineralization level of released larvae. To ensure that the complete development, from the egg to the released larva, occurred under high  $p\text{CO}_2$ , parents were conditioned to the different  $p\text{CO}_2$  levels before mating occurred.

## Methods

### *Crepidula fornicata* adult collection and culture

*C. fornicata* stacks were collected by SCUBA divers on 30 November 2011, after the end of the reproductive period [61] in Morlaix Bay (northwestern Brittany, France), at the “Barre des Flots” site (3°53.015'W; 48°40.015'N). No specific permissions were required for sampling at the selected location, as it is not privately-owned or protected. Field sampling did not involve endangered or protected species.

After being held 6 weeks in natural ambient unfiltered seawater, *C. fornicata* adults were randomly distributed into 18 aquarium tanks of 10 L each (adapted from [62]) and reared for 24 weeks (12 January 2012 to 28 June 2012) in three  $p\text{CO}_2$  treatments selected according to the recommendations of Barry et al. [63]: (1) 390  $\mu\text{atm}$  (pH on the total scale ( $\text{pH}_T$ ) = 8.07) as the current  $p\text{CO}_2$  (control), (2) 750  $\mu\text{atm}$  ( $\text{pH}_T$  = 7.82) and (3) 1400  $\mu\text{atm}$  ( $\text{pH}_T$  = 7.56); the former two  $p\text{CO}_2$  levels are pessimistic scenarios predicted for the end of the century by the IPCC [3]. The  $p\text{CO}_2$  was adjusted by bubbling  $\text{CO}_2$ -free air (current control  $p\text{CO}_2$ ) or pure  $\text{CO}_2$  (two elevated  $p\text{CO}_2$  treatments) in three 100 L header tanks supplied with unfiltered seawater pumped at the foot of the Station Biologique de Roscoff. Each of the three  $p\text{CO}_2$  treatments had six 10 L replicate aquaria. This was an open system, and  $\text{CO}_2$ -treated seawater from the mixing header tanks was continuously supplied to the 18 aquaria (6 per  $p\text{CO}_2$  condition), at a rate of 9  $\text{L h}^{-1}$  (i.e. a renewal rate of 90%  $\text{h}^{-1}$ ). Aquaria were placed in a thermostatic bath where temperature was controlled to within  $\pm 0.2^\circ\text{C}$  using 150 to 250 W submersible heaters. *C. fornicata* adults were grown at four successive temperature levels (10, 13, 16 and  $19^\circ\text{C}$ ) which corresponded to the range of *in situ* temperatures typically encountered in our study area (Service d'Observation de la Mer et du Littoral data). Adults were reared for four weeks at each temperature level. Changes in temperature were implemented slowly, with increases of  $0.2^\circ\text{C day}^{-1}$  over a period of two weeks.

$p\text{CO}_2$  and temperature were monitored and controlled by an off-line feedback system (IKS Aquastar, Karlsbad, Germany) that regulated the addition of gas in the header tanks and the on/off heater switch in the thermostatic bath. The pH values of the system were adjusted from daily measurements of  $\text{pH}_T$  in each of the 18 aquaria using a pH meter (826 pH mobile, Metrohm AG, Herisau, Switzerland) calibrated using Tris/HCl and 2-aminopyridine/HCl buffers [64]. Slipper limpets were fed three times a week with a mix made from a stock solution of *Chaetoceros gracilis* ( $\sim 15 \times 10^6$  cells  $\text{mL}^{-1}$ ) and *Isochrysis affinis galbana* ( $\sim 26 \times 10^6$  cells  $\text{mL}^{-1}$ ). This algal mix (400 mL) was distributed in each aquarium. Seawater flow was stopped for two hours to allow the limpets to feed.

Seawater parameters were monitored throughout the experiment in each of the 18 aquaria.  $\text{pH}_T$  and temperature were recorded daily. Total alkalinity was measured every four weeks, by 0.01 N HCl potentiometric titration on an automatic titrator (Titroline alpha, Schott SI Analytics, Mainz, Germany) in a 20 mL seawater sample taken from each aquarium. Salinity was also measured every four weeks with a conductimeter (LF 330/SET, WTW, Weilheim, Germany) and it varied between  $34.2 \pm 0.1$  and  $35.1 \pm 0.1$  over the course of the experiment. The carbonate chemistry of seawater, i.e. dissolved inorganic carbon (DIC), exact  $p\text{CO}_2$  and saturation state of aragonite ( $\Omega_{Ar}$ ) were calculated for each  $p\text{CO}_2$  and temperature treatment using  $\text{CO}_2\text{SYS}$  software [65] with the constants of Mehrbach et al. [66] refitted by Dickson and Millero [67]. Mean values of these parameters are given in Table 1.


### Larvae collection

In *C. fornicata*, at the end of the embryonic development, the capsule membrane splits and veliger larvae are released in seawater. To prevent released larvae from escaping from their source aquarium, 200  $\mu\text{m}$  mesh size nets covered the overflow outlet of each aquarium. Offspring presence was checked visually every day to collect larvae within 24 h post-hatching. When present, larvae were collected by pouring aquarium seawater on a

**Table 1.** Seawater parameters.

	Temperature		pH <sub>T</sub>		pCO <sub>2</sub>		A <sub>T</sub>		DIC		Ω <sub>Ar</sub>	
	(^°C)		(μEq kg <sup>-1</sup> )		(μEq kg <sup>-1</sup> )		(μEq kg <sup>-1</sup> )		(μmol C kg <sup>-1</sup> )			
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE
<b>10 °C</b>												
390 μatm	23	0.2	8.14	0.01	323	7	2365	2	2138	4	2.47	0.04
750 μatm	23	0.2	7.82	0.01	729	19	2369	2	2270	4	1.33	0.03
1400 μatm	23	0.2	7.55	0.03	1487	75	2377	3	2366	11	0.78	0.08
<b>13 °C</b>												
390 μatm	27	0.2	8.12	0.02	356	25	2418	2	2167	8	2.76	0.07
750 μatm	27	0.1	7.81	0.01	781	20	2416	2	2304	3	1.48	0.03
1400 μatm	27	0.1	7.53	0.01	1557	43	2422	2	2405	4	0.82	0.02
<b>16 °C</b>												
390 μatm	28	0.1	8.08	0.01	376	11	2379	5	2127	5	2.80	0.05
750 μatm	28	0.1	7.82	0.00	748	8	2369	5	2238	2	1.66	0.01
1400 μatm	28	0.1	7.55	0.01	1492	19	2380	5	2345	2	0.94	0.01
<b>19 °C</b>												
390 μatm	23	0.5	8.02	0.01	550	10	2391	2	2152	5	2.70	0.05
750 μatm	23	0.5	7.77	0.01	858	19	2395	3	2266	4	1.68	0.04
1400 μatm	23	0.5	7.51	0.01	1652	41	2394	3	2359	4	0.96	0.03

Legend: Mean parameters of carbonate chemistry in each pCO<sub>2</sub> treatment at each temperature level. The pH on the total scale (pH<sub>T</sub>) was measured daily and total alkalinity (A<sub>T</sub>) was measured every 4 weeks. Other parameters were calculated with the CO<sub>2</sub>SYS software [65]. pCO<sub>2</sub>: CO<sub>2</sub> partial pressure; DIC: dissolved inorganic carbon; Ω<sub>Ar</sub>: saturation state of aragonite.  
doi:10.1371/journal.pone.0093021.t001



**Figure 1. Morphological variables.** All measurements were taken on larvae lying on their right side. A: maximum shell length (L) (in  $\mu\text{m}$ ); B: maximum shell height (h) (in  $\mu\text{m}$ ); C: projected surface area of the left side (in  $\text{mm}^2$ ); D: deformity index, a is the angle of the abnormality (in degrees) and d, the depth of the abnormality (in  $\mu\text{m}$ ). doi:10.1371/journal.pone.0093021.g001

200  $\mu\text{m}$  mesh sieve, rinsed with seawater and preserved in 96% ethanol.

Pools of larvae from adults acclimated to the different  $p\text{CO}_2$  levels since January were collected from the different  $p\text{CO}_2$  conditions at the temperature level of 19°C between 8 and 24 June 2011. Only samples with enough intact larvae were used. Thus two viable samples per  $p\text{CO}_2$  condition were studied.

### Morphological variables


Morphological measurements were performed on a random subsample of 40 larvae when possible or at least 20 larvae from each of the 6 larval pools. Larvae with unbroken shells were isolated in sterile, flat-bottom, 96-well plates and preserved in pure glycerol as described in Auzoux-Bordenave et al. [68]. Each larva was placed on its right side and photographed under light microscopy using an Olympus Camedia C-7070 camera attached to an Olympus SZX 12 dissecting microscope. Pictures were taken without autofocus at  $\times 90$  magnification. Maximum length, height and projected surface area of the left side (Figures 1A, B, and C, respectively) were measured by analyzing images with ImageJ software [69], after calibration with a stage micrometer.

### Abnormalities

In each subsample used for morphological measurements, veliger larvae with abnormal shells were counted and the percentage of abnormal larvae was estimated per  $p\text{CO}_2$  treatment. To be considered as a shell abnormality and not as a broken shell, deformities had to be devoid of fracture lines. A “deformity index” was calculated to quantify the intensity of the shell deformity. It was defined as the ratio between the angle formed by the abnormality and its “depth”, which is the distance between the theoretical curve of the shell and the forest point (extreme point) of the deformity (Figure 1D).

### Shell mineralization

For each  $p\text{CO}_2$  treatment, 5 to 8 larvae were randomly chosen among the previous subsamples and observed under polarized light to determine birefringence patterns with an Olympus dissecting microscope equipped with polarizing filters. All polarized images were acquired with an Olympus camera at  $\times 100$  magnification with 40 ms light exposition. Birefringence under



**Figure 2. Morphological variables.** Mean shell length, height and left surface area ( $\pm$  SE) in the different  $p\text{CO}_2$  treatments. Different letters above bars indicate significant differences between treatments ( $p < 0.05$ , Dunn post-hoc test),  $n = 51$  to 92. doi:10.1371/journal.pone.0093021.g002

**Table 2.** Effect of  $p\text{CO}_2$  on morphological variables, abnormality indices and intensity of birefringence.

	Kruskal-Wallis Test		
	df	H	p
Length	2	37.353	< 0.001
Height	2	16.235	< 0.001
Surface area	2	30.106	< 0.001
Abnormality index	2	6.046	0.049
Intensity of birefringence	2	14.562	< 0.001

Legend: Summary of the non-parametric Kruskal-Wallis tests testing the effect of  $p\text{CO}_2$  on each morphological variable, abnormality index and birefringence intensity. doi:10.1371/journal.pone.0093021.t002

polarized light is due to the mineral phase composing the shell [40,68,70,71]. In the absence of mineralized structures, there is no birefringence and the picture looks totally black. Under identical light conditions, areas appearing more birefringent contain a much larger proportion of crystalline calcium carbonate [70,72]. The intensity of birefringence of each shell was used as a proxy for mineralization level for the three  $p\text{CO}_2$  treatments. It was quantified from pictures by using ImageJ software [69]. Pictures of polarized shells were first transformed into grayscale images. A mean gray value (in pixels) was determined for each birefringent zone. All birefringent zones of the shell were compiled to obtain a global mean gray value, giving the intensity of the birefringence of the whole shell.

### Statistics

All statistical analyses were performed using the free software R 2.15.0 version [73]. Normality and homoscedasticity of the data were first checked using Shapiro and Levene tests, respectively. Due to the non-normality and heterogeneity of variance, the influence of  $p\text{CO}_2$  on morphological variables, deformity indices and birefringence intensity was analyzed using the non-parametric Kruskal-Wallis test followed by the Dunn post-hoc test [74]. A Chi-squared ( $\chi^2$ ) test followed by G-tests (likelihood-ratio test) [75]

were used to compare percentages of anomaly between the three  $p\text{CO}_2$  conditions.


## Results

### Morphological variables

$p\text{CO}_2$  significantly affected length, height and surface area of the hatched larvae (Figure 2, Table 2). These morphological variables are related to each other and were generally influenced in the same way by  $p\text{CO}_2$ . Length and height were the highest at 390  $\mu\text{atm}$  and significantly decreased with increased  $p\text{CO}_2$ . Larvae collected at 750 and 1400  $\mu\text{atm}$   $p\text{CO}_2$  showed a decrease of 5.3% and 5.9% in length, respectively, and 2.6% and 4.5% in height, respectively, compared to control larvae (390  $\mu\text{atm}$ ). Similarly, the greatest shell surface area was observed at 390  $\mu\text{atm}$   $p\text{CO}_2$ , but then significantly decreased by 6.2% and 11.2% at 750 and 1400  $\mu\text{atm}$   $p\text{CO}_2$ , respectively.


### Abnormalities

Abnormalities in larvae were observed as notches located close to the shell aperture (Figure 3). The percentage of abnormal larvae increased with increased  $p\text{CO}_2$  and ranged from 6.7 to 26.5% (Figure 4;  $\chi^2$  test,  $p < 0.05$ ). Abnormalities were 1.5- and 4-fold at  $p\text{CO}_2$  levels of 750 and 1400  $\mu\text{atm}$ , respectively, than at



**Figure 3. Shell abnormalities.** Different intensities of shell abnormalities observed among samples. A, B, C, and D show whole larvae whereas E, F, G, and H show the detail of their respective abnormalities. doi:10.1371/journal.pone.0093021.g003





**Figure 4. Percentage of abnormalities in the different pCO<sub>2</sub> conditions.** Different letters above bars indicate significant differences between treatments (G-test,  $p < 0.05$ ). doi:10.1371/journal.pone.0093021.g004

390 µatm. Furthermore, different intensities of shell abnormalities were observed with variation in notch acuteness. The deformity index varied between 0.03 (390 µatm pCO<sub>2</sub>) and 0.17 (1400 µatm pCO<sub>2</sub>). Although the Kruskal-Wallis test showed a marginally significant pCO<sub>2</sub> effect ( $p = 0.049$ ; Table 2), the pairwise Dunn post-hoc test did not detect significant differences between the three pCO<sub>2</sub> treatments ( $p > 0.05$ ).

Among the abnormal larvae observed under polarized microscopy (see below), some showed abnormalities which appeared less birefringent, and even not mineralized, as revealed by the lack of birefringence in these parts of the shell (Figure 5).

#### Shell mineralization


Pictures taken under polarized light (Figure 6) suggest that the intensity of the birefringence decreased with increased pCO<sub>2</sub>. The measure of birefringence intensity using the mean gray values estimated for each shell clearly confirmed this relationship (Table

2). Mineralization was greatest at 390 µatm pCO<sub>2</sub>, intermediate at 750 µatm pCO<sub>2</sub> and lowest at 1400 µatm pCO<sub>2</sub> (Figure 6).

#### Discussion


In our study, the effects of elevated pCO<sub>2</sub> were integrated from embryo formation and throughout embryonic stages up until the release of veliger larvae. The integration of pCO<sub>2</sub> effects across all developmental stages, from fertilization to settlement and beyond, is particularly instructive because early life stages may respond to environmental stressors in a different way than adults. Acute exposures of early life stages have shown various effects in growth or feeding performances [76,77], but these may not represent field conditions. Results obtained from acclimation to high pCO<sub>2</sub> across different life stages often differ from those arising from acute exposure of a given stage. For example, it has been shown that oyster D-veliger larvae grown from eggs fertilized at elevated pCO<sub>2</sub> were more drastically affected than those first produced at ambient pCO<sub>2</sub> and then reared later (embryo stages) at high pCO<sub>2</sub> [78]. Keeping the parents under different pCO<sub>2</sub> conditions before fertilization and until larval release allowed embryos to develop entirely under a given level of stress. To our knowledge, only Dupont et al. [79] on sea urchins, Parker et al. [80] on mollusks and Vehmaa et al. [81] on copepods acclimated adults to high pCO<sub>2</sub> during reproductive conditioning before studying larvae in the same pCO<sub>2</sub> conditions.

pCO<sub>2</sub> effects were first investigated on the shell morphology of the hatched larvae. The size (length, height and surface area) of the released larvae decreased with increased pCO<sub>2</sub>. Length and height were not significantly different between the pCO<sub>2</sub> levels of 750 and 1400 µatm, whereas the shell surface area progressively decreased with increased pCO<sub>2</sub> from 390 to 1400 µatm. Although in some rare cases, elevated pCO<sub>2</sub> does not affect larval morphology and growth [82,83], a correlation between high pCO<sub>2</sub> and smaller size has been demonstrated in most bivalve and gastropod species studied to date (see review in Table 3), with pCO<sub>2</sub> sometimes affecting the shape of the larval shell [41]. As observed here for length and height (ca. -5% in the two high pCO<sub>2</sub> treatments), shell lengths of *Crassostrea gigas* veliger larvae are smaller under elevated pCO<sub>2</sub>, but are similar in conditions of pH lowered by 0.4 and 0.7 pH units, a range corresponding to our pH values [32]. Smaller size and delayed shell growth can be



**Figure 5. Shell abnormalities under normal (A, C, and E) and polarized (B, D, and F) light.** A-B are pictures of a normal larva presenting the characteristic black cross of birefringence. C-D and E-F show examples of abnormalities observed among the samples, with arrows indicating less calcified zones.

doi:10.1371/journal.pone.0093021.g005



**Figure 6. Birefringence intensity.** Mean gray value (in pixel) of the shells under polarized light in the different  $p\text{CO}_2$  treatments ( $\pm$  SE). Bars with different letters are significantly different (Dunn post-hoc test,  $p < 0.05$ ),  $n = 5$  to 8 larvae per condition. Pictures taken under polarized light (below the graph) show birefringent patterns in the different  $p\text{CO}_2$  treatments. doi:10.1371/journal.pone.0093021.g006

attributed to problems in shell deposition, delayed periostracum formation and/or increased shell dissolution, as hypothesized by Watson et al. [84].

Such processes may lead to developmental abnormalities and to an increase in their frequency under elevated  $p\text{CO}_2$ . Some (7%) *C. formicata* larvae in the control group (390  $\mu\text{atm } p\text{CO}_2$ ) showed mild shell abnormalities in the form of a notch close to the aperture. The frequency of this abnormality increased under high  $p\text{CO}_2$ , being 1.5-fold more frequent at 750  $\mu\text{atm } p\text{CO}_2$  and reaching 26% at 1400  $\mu\text{atm } p\text{CO}_2$ . The intensity of shell abnormality, estimated using the deformity index, did not vary significantly with increasing  $p\text{CO}_2$ , although more pronounced shell deformities were detected at the highest  $p\text{CO}_2$  condition (1400  $\mu\text{atm}$ ). The occurrence of abnormal shells is a common response in mollusk larvae exposed to elevated  $p\text{CO}_2$ . In bivalves for example, abnormalities can occur as shell hinge and edge deformities [85], irregular-shaped shells [40] or protruding mantles [86]. The frequency of abnormalities can reach 40% of shell deformities in *Pecten maximus* larvae reared at 1250  $\mu\text{atm } p\text{CO}_2$  [85], and even 90% of abnormal D-veligers in *Saccostrea glomerata* at 1000  $\mu\text{atm } p\text{CO}_2$  [31]. In gastropods, larval shells are considered abnormal when shells are too small to fully cover the soft body [34] or when dissolution zones are observed at the edge of the aragonitic larval shell [33], with frequencies of abnormality ranging from 20% in *Haliotis discus hannai* at 1650  $\mu\text{atm } p\text{CO}_2$  [33] to 40% in *Haliotis kamschatkana* at 800  $\mu\text{atm } p\text{CO}_2$  [36]. At extremely high  $p\text{CO}_2$  (> 1700  $\mu\text{atm}$ ), some abalone larvae are even unable to precipitate a calcareous shell [36,44].

Such abnormalities may be due to different processes: (i) the production of amorphous  $\text{CaCO}_3$  may be affected by damage to embryonic ectodermic cells and/or (ii) seawater corrosion may induce shell dissolution, affecting the strength and calcification of some parts of the shell [32]. Here, the mineralization level of larval shells was investigated at each  $p\text{CO}_2$  level by observing the veliger aragonitic shell under polarized light [70]. The characteristic dark cross observed in each larval shell indicated a radial arrangement of aragonite crystals [72] and did not have been considered as non-crystalline zones. The intensity of birefringence was used as a proxy for mineralization because increases in birefringence reflect increases in crystalline structure and calcification of the shell. Observed under polarized light, abnormalities appeared less birefringent than the rest of the shell, suggesting that deformities were likely less calcified as proposed by Barros et al. [32]

The birefringence intensity of the larval shells decreased with increased  $p\text{CO}_2$ , and was significantly lower at 1400  $\mu\text{atm } p\text{CO}_2$ . This drop in birefringence revealed a decrease in calcification, which may be related to a less mineralized matrix [87], or more likely to a reduction in shell thickness [72]. Our data did not allow us to discriminate between these two possibilities, but previous studies have already reported a decrease in shell thickness under high  $p\text{CO}_2$  in bivalve larvae. For example, using scanning electron microscopy measurements, Gazeau et al. [39] showed a decrease in thickness of 12% in *Mytilus edulis* larvae at 745  $\mu\text{atm } p\text{CO}_2$ . Talmage and Gobler [43] report a decrease in thickness of *Mercenaria mercenaria* (-43%) and *Argopecten irradians* (-47.5%) larval shells after 17 days of development at 1500  $\mu\text{atm } p\text{CO}_2$ , which was associated with an impact on the integrity and the



**Table 3.** Literature review of reports of high pCO<sub>2</sub> effects on morphological variables in mollusk larvae.

Reproductive mode	Species (veliger stage)	Measured parameter	pH conditions	pCO <sub>2</sub> conditions (µatm)	% decrease due to pCO <sub>2</sub>	Study
Broadcast spawner	<i>Crassostrea gigas</i>	shell length	pH <sub>NBS</sub> 7.4	2268	decrease	[40]
Broadcast spawner	<i>Crassostrea gigas</i>	shell length	pH <sub>NBS</sub> 7.8	1000	16%	[78]
Broadcast spawner	<i>Crassostrea gigas</i>	shell length and height	pH <sub>NBS</sub> 7.76 – 7.37	1386 – 3573	10.6%	[32]
Broadcast spawner	<i>Crassostrea gigas</i>	shell area	pH <sub>NBS</sub> 7.7 – 7.4	1497 – 2386	18.7 – 29%	[101]
Broadcast spawner	<i>Crassostrea virginica</i>	shell length	pH <sub>T</sub> 7.84 – 7.49	≈ 650 – 1500	16.7%	[38]
Broadcast spawner	<i>Crassostrea virginica</i>	shell area	pH <sub>NBS</sub> 7.76	840	16%	[94]
Broadcast spawner	<i>Saccostrea glomerata</i>	shell length	-	1000	22%	[31]
Broadcast spawner	<i>Saccostrea glomerata</i>	shell length	pH <sub>NBS</sub> 7.8	1000	34%	[78]
Broadcast spawner	<i>Saccostrea glomerata</i>	shell length	pH <sub>NBS</sub> 7.9	856	31.6 – 1.3%	[80]
Broadcast spawner	<i>Saccostrea glomerata</i>	shell length	pH <sub>NBS</sub> 7.8 – 7.6	508.8 – 775.6	8.7 – 6.3%	[84]
Broadcast spawner	<i>Saccostrea glomerata</i>	shell height	pH <sub>NBS</sub> 7.8 – 7.6	508.8 – 775.6	7.5 – 5.1%	[84]
Broadcast spawner	<i>Mytilus californianus</i>	shell area	pH <sub>NBS</sub> 7.75	970	5 – 7%	[37]
Broadcast spawner	<i>Mytilus edulis</i>	shell length	pH <sub>NBS</sub> 7.8	1200	4.5 – 6%	[39]
Broadcast spawner	<i>Mytilus edulis</i>	shell thickness	pH <sub>NBS</sub> 7.8	1200	12%	[39]
Broadcast spawner	<i>Mytilus edulis</i>	shell area	pH <sub>NBS</sub> 7.6	1388 – 1493	7 – 8%	[16]
Broadcast spawner	<i>Mytilus galloprovincialis</i>	shell length and height	pH <sub>NBS</sub> 7.4	2000	26 – 20%	[86]
Broadcast spawner	<i>Argopecten irradians</i>	shell length	pH <sub>T</sub> 7.84 – 7.49	≈ 650 – 1500	50%	[38]
Broadcast spawner	<i>Argopecten irradians</i>	shell diameter	pH <sub>T</sub> (?) 7.5	1500	30.7%	[43]
Broadcast spawner	<i>Argopecten irradians</i>	shell thickness	pH <sub>T</sub> (?) 7.5	1500	43%	[43]
Broadcast spawner	<i>Argopecten irradians</i>	shell length	pH <sub>SW</sub> 7.39	1987	18.9 – 7.5%	[102]
Broadcast spawner	<i>Pecten maximus</i>	shell length	pH <sub>NBS</sub> (?) 7.51	1627	10%	[85]
Broadcast spawner	<i>Pecten maximus</i>	shell height	pH <sub>NBS</sub> (?) 7.51	1627	5%	[85]
Broadcast spawner	<i>Macoma balthica</i>	shell length	pH <sub>NBS</sub> 7.7 – 7.2	1700 – 4400	4.3 – 8.5%	[103]
Broadcast spawner	<i>Macoma balthica</i>	shell length	pH <sub>NBS</sub> 7.8 – 7.5	-	16.2 – 16.9%	[42]
Broadcast spawner	<i>Mercenaria mercenaria</i>	shell length	pH <sub>T</sub> 7.84 – 7.49	≈ 650 – 1500	22.7%	[38]
Broadcast spawner	<i>Mercenaria mercenaria</i>	shell diameter	pH <sub>T</sub> (?) 7.5	1500	25.5%	[43]
Broadcast spawner	<i>Mercenaria mercenaria</i>	shell thickness	pH <sub>T</sub> (?) 7.5	1500	43.0%	[43]
Broadcast spawner	<i>Haliotis discus hannai</i>	shell length	pH <sub>T</sub> (?) 7.71	1050	2.50%	[33]
Broadcast spawner	<i>Haliotis discus hannai</i>	shell length	pH <sub>NBS</sub> 7.6 – 7.3	1224.6 – 2543.6	decrease	[34]
Broadcast spawner	<i>Haliotis kamschatkana</i>	shell length	pH <sub>NBS</sub> (?) 8.07	800	5%	[36]
Brooding in pallial cavity	<i>Ostrea chilensis</i>	shell thickness	pH <sub>NBS</sub> (?) 7.7 – 7.0	-	no change	[50]
Brooding in pallial cavity	<i>Ostrea lurida</i>	shell growth	pH <sub>T</sub> 7.76	1000	5 – 14%	[104]
Egg masses	<i>Littorina obtusata</i>	shell length	pH <sub>NBS</sub> (?) 7.6	-	10%	[41]
Egg masses	<i>Stylocheilus striatus</i>	shell area	pH <sub>NBS</sub> (?) 7.6	-	24 – 36%	[52]
Egg	<i>Sepia officinalis</i>	total weight	pH <sub>T</sub> 7.84 – 7.60	750 – 1430	no change	[53]
Encapsulation + brooding	<i>Crepidatella dilatata</i>	shell thickness	pH <sub>NBS</sub> (?) 6	-	30%	[51]
Encapsulation + brooding	<i>Crepidula fornicata</i>	shell length	pH <sub>T</sub> 7.8 – 7.6	750 – 1400	5.3 – 10.7%	Present study
Encapsulation + brooding	<i>Crepidula fornicata</i>	shell height	pH <sub>T</sub> 7.8 – 7.6	750 – 1400	2.6 – 13.1%	Present study
Encapsulation + brooding	<i>Crepidula fornicata</i>	shell area	pH <sub>T</sub> 7.8 – 7.6	750 – 1400	6.2 – 19.8%	Present study

pH<sub>NBS</sub>: pH on the NBS scale; pH<sub>T</sub>: pH on the total scale; pH<sub>SW</sub>: pH on the seawater scale.  
doi:10.1371/journal.pone.0093021.t003

connectedness of the hinge structure. A decrease of 5.7% in shell thickness of brooded larvae of the oyster *Ostrea chilensis* has also been observed following a decrease in pH (down to 6.56) within the mother's pallial cavity due to valve closure under salinity stress [50].

The decrease in larval size and mineralization level of the shell may be due to reduced CaCO<sub>3</sub> saturation or hypercapnic suppression of metabolic pathways involved in the calcification process [9]. Very little is known about the conditions occurring

during intracapsular development and how acidified seawater can affect encapsulated embryos. Previous studies have shown that egg capsules of some gastropods, including *C. fornicata*, are permeable to water and ions (e.g. [48,71,88]) and it can be assumed that the capsule wall in *C. fornicata* is almost impermeable to gas because of its low O<sub>2</sub> conductance [89]. Under "normal" conditions, embryos of *C. fornicata* will progressively be exposed to hypoxia [89] and hypercapnia via their respiration. This may lead to low intracapsular pH, as reported in other gastropod species [90],

without altering development [57]. Under elevated  $p\text{CO}_2$ , diffusion of more protons ( $\text{H}^+$ ) from external seawater to the intracapsular medium may alter intracapsular carbonate chemistry, thus enhancing metabolic acidosis. Similar acidosis can be observed under low salinity stress. For example, a decrease in pH to 6.4 recorded within the pallial cavity of the calyptraeid *Crepidipatella dilatata* [47,51] led to the partial shell decalcification of the brooded encapsulated embryos [51]. This decalcification may cause the release of some carbonate ions ( $\text{CO}_3^{2-}$ ), which could bind to free  $\text{H}^+$  to form bicarbonate ions ( $\text{HCO}_3^-$ ), thus buffering the intracapsular acidosis and limiting the drastic pH effects on larval metabolism. This potential buffering role, in combination with a decrease in the  $\text{CaCO}_3$  saturation state, is likely to affect shell mineralization and calcification. Such processes have been suggested to buffer acidosis resulting from anaerobiosis in *C. formicata* [91], and may also explain our observations. Alternative mechanisms that can decrease the intracapsular acidosis, such as the active excretion of  $\text{H}^+$  out of the capsule through a proton pump, as shown in the cephalopod *Sepia officinalis* [92], need to be investigated.

Altogether, our results show that, despite the potential protective role provided by encapsulation and brooding, elevated seawater  $p\text{CO}_2$  affected the shells of the released larvae in *C. formicata*. Embryos of *C. formicata* were affected by high  $p\text{CO}_2$  during their intracapsular development. However, the overall low abnormality rate and low decrease in size suggested they were likely less affected than other mollusk early life stages. The natural exposure of embryos to low intracapsular pH as demonstrated in cephalopod eggs (pH on the seawater scale of the perivitelline fluid of ca. 7.35 at 16°C [53]) and gastropod capsules (pH of the intracapsular fluid lower than 7 [90]) could confer to *C. formicata* larvae some resilience to elevated  $p\text{CO}_2$  levels. Indeed, it has been shown that bivalves naturally exposed to high  $p\text{CO}_2$  conditions in their habitat (due to high levels of benthic respiration or to seawater naturally enriched in  $\text{CO}_2$ ) are less affected by ocean acidification than other mollusk species [93,94]. Further studies are however needed to determine the pH of the intracapsular fluid in *C. formicata*, and how it will be affected under future scenarios of ocean acidification.

## References

- Pechenik JA (1999) On the advantages and disadvantages of larval stages in benthic marine invertebrate life cycles. *Mar Ecol-Prog Ser* 177: 269–297.
- Peck LS, Souster T, Clark MS (2013) Juveniles are more resistant to warming than adults in 4 species of Antarctic marine invertebrates. *PLoS One* 8: e66033.
- IPCC (2013) Working Group I Contribution to the IPCC Fifth Assessment Report - *Climate Change 2013: The Physical Science Basis - Summary for Policymakers*.
- Feely RA, Doney SC, Cooley SR (2009) Ocean acidification: present conditions and future changes in a high- $\text{CO}_2$  world. *Oceanography* 22: 36–47.
- Pörtner H-O (2008) Ecosystem effects of ocean acidification in times of ocean warming: a physiologist's view. *Mar Ecol-Prog Ser* 373: 203–217.
- Widdicombe S, Spicer JI (2008) Predicting the impact of ocean acidification on benthic biodiversity: what can animal physiology tell us? *J Exp Mar Biol Ecol* 366: 187–197.
- Doney SC, Fabry VJ, Feely RA, Kleypas JA (2009) Ocean acidification: the other  $\text{CO}_2$  problem. *Ann Rev Mar Sc* 1: 169–192.
- Hoegh-Guldberg O (2009) Climate change and coral reefs: Trojan horse or false prophecy? *Coral Reefs* 28: 569–575.
- Byrne M (2011) Impact of ocean warming and ocean acidification on marine invertebrate life history stages: vulnerabilities and potential for persistence in a changing ocean. In: Gibson R, Atkinson R, Gordon J, Smith I, Hughes D, editors. *Oceanography and Marine Biology: An Annual Review*: Taylor & Francis. pp. 1–42.
- Feely RA, Sabine CL, Lee K, Berelson W, Kleypas J, et al. (2004) Impact of anthropogenic  $\text{CO}_2$  on the  $\text{CaCO}_3$  system in the oceans. *Science* 305: 362–366.
- Kroeker KJ, Kordas RL, Crim RN, Singh GG (2010) Meta-analysis reveals negative yet variable effects of ocean acidification on marine organisms. *Ecol Lett* 13: 1419–1434.
- Dupont S, Thorndyke M (2009) Impact of  $\text{CO}_2$ -driven ocean acidification on invertebrates early life-history - What we know, what we need to know and what we can do. *Biogeosciences Discuss.* 6: 3109–3131.
- Kurihara H (2008) Effects of  $\text{CO}_2$ -driven ocean acidification on the early developmental stages of invertebrates. *Mar Ecol-Prog Ser* 373: 275–284.
- Nguyen HD, Doo SS, Soars NA, Byrne M (2012) Noncalcifying larvae in a changing ocean: warming, not acidification/hypercapnia, is the dominant stressor on development of the sea star *Meridiastra calcar*. *Glob Change Biol* 18: 2466–2476.
- Pansch C, Schlegel P, Havenhand J (2013) Larval development of the barnacle *Amphibalanus improvisus* responds variably but robustly to near-future ocean acidification. *ICES J Mar Sci* 70: 805–811.
- Bechmann RK, Taban IC, Westerlund S, Godal BF, Arnborg M, et al. (2011) Effects of ocean acidification on early life stages of shrimp (*Pandalus borealis*) and mussel (*Mytilus edulis*). *J Toxic Envir Health-Part A* 74: 424–438.
- Range P, Pilo D, Ben-Hamadou R, Chicharo M, Matias D, et al. (2012) Seawater acidification by  $\text{CO}_2$  in a coastal lagoon environment: effects on life history traits of juvenile mussels *Mytilus galloprovincialis*. *J Exp Mar Biol Ecol* 424: 89–98.
- Dupont S, Dorey N, Thorndyke M (2010) What meta-analysis can tell us about vulnerability of marine biodiversity to ocean acidification? *Estuar Coast Shelf Sci* 89: 182–185.
- Ries JB, Cohen AL, McCorkle DC (2009) Marine calcifiers exhibit mixed responses to  $\text{CO}_2$ -induced ocean acidification. *Geology* 37: 1131–1134.
- Chua CM, Leggat W, Moya A, Baird AH (2013) Temperature affects the early life history stages of corals more than near future ocean acidification. *Mar Ecol-Prog Ser* 475: 85–92.
- Putnam HM, Mayfield AB, Fan TY, Chen CS, Gates RD (2013) The physiological and molecular responses of larvae from the reef-building coral

## Acknowledgments

The authors thank the “Marine Operations and Services Department” at the Station Biologique de Roscoff for underwater sampling. We also thank the “Multicellular Marine Models” staff for providing microalgae and their help for building the aquarium system. We are grateful to Stéphanie Auzoux-Bordenave and Nathalie Wessel for hosting us at the Station de Biologie Marine de Concarneau and for their help in acquiring the polarized light images. We also thank Frédérique Viard and Thomas Broquet for helpful discussions at various stages of this work. We are grateful to the editor, Pauline Ross, and the two anonymous reviewers for their helpful and constructive comments which greatly improved this manuscript.

## Author Contributions

Conceived and designed the experiments: FN TC DD SM. Performed the experiments: FN EL FB. Analyzed the data: FN EL. Contributed reagents/materials/analysis tools: FN EL TC SM. Wrote the paper: FN EL TC FB DD SM.

- Pocillopora damicornis* exposed to near-future increases in temperature and  $p\text{CO}_2$ . *Mar Biol* 160: 2157–2173.
22. Dupont S, Havenhand J, Thorndyke W, Peck L, Thorndyke M (2008) Near-future level of  $\text{CO}_2$ -driven ocean acidification radically affects larval survival and development in the brittlestar *Ophiothrix fragilis*. *Mar Ecol-Prog Ser* 373: 285–294.
  23. Martin S, Richier S, Pedrotti ML, Dupont S, Castejon C, et al. (2011) Early development and molecular plasticity in the Mediterranean sea urchin *Paracentrotus lividus* exposed to  $\text{CO}_2$ -driven acidification. *J Exp Biol* 214: 1357–1368.
  24. Padilla-Gamino JL, Kelly MW, Evans TG, Hofmann GE (2013) Temperature and  $\text{CO}_2$  additively regulate physiology, morphology and genomic responses of larval sea urchins, *Strongylocentrotus purpuratus*. *P Roy Soc Lond, B Biol* 280: 20130155.
  25. Egilisdottir H, Spicer JI, Rundle SD (2009) The effect of  $\text{CO}_2$  acidified sea water and reduced salinity on aspects of the embryonic development of the amphipod *Echinogammarus marinus* (Leach). *Mar Poll Bull* 58: 1187–1191.
  26. Gazeau F, Parker L, Comeau S, Gattuso J-P, O'Connor W, et al. (2013) Impacts of ocean acidification on marine shelled molluscs. *Mar Biol* 160: 2207–2245.
  27. Gutowska MA, Melzner F (2009) Abiotic conditions in cephalopod (*Sepia officinalis*) eggs: embryonic development at low pH and high  $p\text{CO}_2$ . *Mar Biol* 156: 515–519.
  28. Moran D, Stottrup JG (2011) The effect of carbon dioxide on growth of juvenile Atlantic cod *Gadus morhua* L. *Aquat Toxicol* 102: 24–30.
  29. Munday PL, Gagliano M, Donelson JM, Dixon DL, Thorrold SR (2011) Ocean acidification does not affect the early life history development of a tropical marine fish. *Mar Ecol-Prog Ser* 423: 211–221.
  30. Parker LM, Ross PM, O'Connor WA, Pörtner H-O, Scanes E, et al. (2013) Predicting the response of molluscs to the impact of ocean acidification. *Biology* 2: 651–692.
  31. Parker LM, Ross PM, O'Connor WA (2009) The effect of ocean acidification and temperature on the fertilization and embryonic development of the Sydney rock oyster *Saccostrea glomerata* (Gould 1850). *Glob Change Biol* 15: 2123–2136.
  32. Barros P, Sobral P, Range P, Chicharo L, Matias D (2013) Effects of seawater acidification on fertilization and larval development of the oyster *Crassostrea gigas*. *J Exp Mar Biol Ecol* 440: 200–206.
  33. Kimura R, Takami H, Ono T, Onitsuka T, Nojiri Y (2011) Effects of elevated  $p\text{CO}_2$  on the early development of the commercially important gastropod, Ezo abalone *Haliotis discus hannai*. *Fish Oceanogr* 20: 357–366.
  34. Li J, Jiang Z, Zhang J, Qiu J-W, Du M, et al. (2013) Detrimental effects of reduced seawater pH on the early development of the Pacific abalone. *Mar Poll Bull* 74: 320–324.
  35. Davis AR, Coleman D, Broad A, Byrne M, Dworjanyn SA, et al. (2013) Complex responses of intertidal molluscan embryos to a warming and acidifying ocean in the presence of UV radiation. *PLoS One* 8: e55939.
  36. Crim RN, Sunday JM, Harley CDG (2011) Elevated seawater  $\text{CO}_2$  concentrations impair larval development and reduce larval survival in endangered northern abalone (*Haliotis kamtschatkana*). *J Exp Mar Biol Ecol* 400: 272–277.
  37. Gaylord B, Hill TM, Sanford E, Lenz EA, Jacobs LA, et al. (2011) Functional impacts of ocean acidification in an ecologically critical foundation species. *J Exp Biol* 214: 2586–2594.
  38. Talmage SC, Gobler CJ (2009) The effects of elevated carbon dioxide concentrations on the metamorphosis, size, and survival of larval hard clams (*Mercenaria mercenaria*), bay scallops (*Argopecten irradians*), and Eastern oysters (*Crassostrea virginica*). *Limnol Oceanogr* 54: 2072–2080.
  39. Gazeau F, Gattuso J-P, Dawber C, Pronker AE, Peene F, et al. (2010) Effect of ocean acidification on the early life stages of the blue mussel *Mytilus edulis*. *Biogeosciences* 7: 2051–2060.
  40. Kurihara H, Kato S, Ishimatsu A (2007) Effects of increased seawater  $p\text{CO}_2$  on early development of the oyster *Crassostrea gigas*. *Aquat Biol* 1: 91–98.
  41. Ellis RP, Bersey J, Rundle SD, Hall-Spencer JM, Spicer JI (2009) Subtle but significant effects of  $\text{CO}_2$  acidified seawater on embryos of the intertidal snail, *Littorina obtusata*. *Aquat Biol* 5: 41–48.
  42. Van Colen C, Debusschere E, Braeckman U, Van Gansbeke D, Vincx M (2012) The early life history of the clam *Macoma balthica* in a high  $\text{CO}_2$  world. *PLoS One* 7: e44655.
  43. Talmage SC, Gobler CJ (2010) Effects of past, present, and future ocean carbon dioxide concentrations on the growth and survival of larval shellfish. *P N at Acad Sci* 107: 17246–17251.
  44. Byrne M, Ho M, Wong E, Soars NA, Selvakumaraswamy P, et al. (2010) Unshelled abalone and corrupted urchins: development of marine calcifiers in a changing ocean. *P Roy Soc Lond, B Biol* 278: 2376–2383.
  45. Ross PM, Parker LM, O'Connor WA, Bailey EA (2011) The impact of ocean acidification on reproduction, early development and settlement of marine organisms. *Water* 3: 1005–1030.
  46. Przeslawski R (2004) A review of the effects of environmental stress on embryonic development within intertidal gastropod egg masses. *Molluscan Res* 24: 43–63.
  47. Chaparro OR, Segura CJ, Montroy JA, Navarro JM, Pechenik JA (2009) Brood chamber isolation during salinity stress in two estuarine mollusk species: from a protective nursery to a dangerous prison. *Mar Ecol-Prog Ser* 374: 145–155.
  48. Pechenik JA (1982) Ability of some gastropod egg capsules to protect against low-salinity stress. *J Exp Mar Biol Ecol* 63: 195–208.
  49. Pechenik JA (1983) Egg capsules of *Nuccella lapillus* (L.) protect against low-salinity stress. *J Exp Mar Biol Ecol* 71: 165–179.
  50. Chaparro OR, Montroy JA, Segura CJ, Pechenik JA (2009) Effect of reduced pH on shells of brooded veligers in the estuarine bivalve *Ostrea chilensis* Philippi 1845. *J Exp Mar Biol Ecol* 377: 107–112.
  51. Montroy JA, Chaparro OR, Cubillos VM, Pechenik JA (2009) Isolation of incubation chambers during brooding: effect of reduced pH on protoconch development in the estuarine gastropod *Crepidipelta dilatata* (Calyptreidae). *Mar Ecol-Prog Ser* 374: 157–166.
  52. Allen TR (2012) The effects of ocean acidification and sea surface warming on the embryonic development of the opisthobranch gastropod *Stylocheilus status*. Student Research Papers, Fall 2012, UCB Moorea Class: Biology and Geomorphology of Tropical Islands, Berkeley Natural History Museum, UC Berkeley, 18 p.
  53. Dorey N, Melzner F, Martin S, Oberhaensli F, Teysse JL, et al. (2013) Ocean acidification and temperature rise: effects on calcification during early development of the cuttlefish *Sepia officinalis*. *Mar Biol* 160: 2007–2022.
  54. Fernandes DAO, Podolsky RD (2012) Effects of ocean acidification on growth, development, and calcification of gastropod embryos: does encapsulation matter? *Integrative and comparative biology* 52: e244–e244.
  55. Blanchard M (1995) Origine et état de la population de *Crepidula fornicata* (Gastropoda Prosobranchia) sur le littoral français. *Haliotis* 24: 75–86.
  56. Blanchard M (1997) Spread of the slipper limpet *Crepidula fornicata* (L. 1758) in Europe. Current state and consequences. *Sci Mar* 61: 109–118.
  57. Brante A, Fernandez M, Viard F (2009) Limiting factors to encapsulation: the combined effects of dissolved protein and oxygen availability on embryonic growth and survival of species with contrasting feeding strategies. *J Exp Biol* 212: 2287–2295.
  58. Orton J (1912) An account of the natural history of the slipper-limpet (*Crepidula fornicata*) with some remarks on its occurrence on the oyster grounds on the Essex coast. *J Mar Biol Assoc UK* 9: 437–443.
  59. Rigal F (2009) Dynamique spatio-temporelle du nuage larvaire du gastéropode introduit *Crepidula fornicata* au sein d'une baie mégatidale, la baie de Morlaix (France). PhD thesis. Paris: Université Pierre et Marie Curie. 160p.
  60. Pechenik JA (1984) The relationship between temperature, growth rate, and duration of planktonic life for larvae of the gastropod *Crepidula fornicata* (L.). *J Exp Mar Biol Ecol* 74: 241–257.
  61. Richard J, Huet M, Thouzeau G, Paulet YM (2006) Reproduction of the invasive slipper limpet, *Crepidula fornicata*, in the Bay of Brest, France. *Mar Biol* 149: 789–801.
  62. Noisette F, Duong G, Six C, Davoult D, Martin S (2013) Effects of elevated  $p\text{CO}_2$  on the metabolism of a temperate rhodolith *Lithothamnion corallioides* grown under different temperatures. *J Phycol* 49: 746–757.
  63. Barry JP, Tyrrell T, Hansson L, Plattner GK, Gattuso JP (2010) Atmospheric  $\text{CO}_2$  targets for ocean acidification perturbation experiments. In: Riebesell U, FVJ, Hansson L, & Gattuso J.-P., editor. Guide to best practices for ocean acidification research and data reporting. Luxembourg: Publications Office of the European Union. pp. 260.
  64. Dickson AG, Sabine CL, Christian JR, editors (2007) Guide to best practices for ocean  $\text{CO}_2$  measurements. Sidney, British Columbia: North Pacific Marine Science Organization. 176 p.
  65. Lewis E, Wallace DWR (1998) Program developed for  $\text{CO}_2$  system calculations. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy
  66. Mehrbach C, Culberso C, Hawley JE, Pytkowic RM (1973) Measurement of apparent dissociation-constants of carbonic-acid in seawater at atmospheric-pressure. *Limnol Oceanogr* 18: 897–907.
  67. Dickson AG, Millero FJ (1987) A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep Sea Res* 34: 1733–1743.
  68. Auzoux-Bordenave S, Badou A, Gaume B, Berland S, Helleouet MN, et al. (2010) Ultrastructure, chemistry and mineralogy of the growing shell of the European abalone *Haliotis tuberculata*. *J Struct Biol* 171: 277–290.
  69. Rasband WS (2012) ImageJ version 1.46r. In: Health USNI, editor. imagej.nih.gov/ij/, 1997–2012. Bethesda, Maryland, USA.
  70. Weiss IM, Tuross N, Addadi I, Weiner S (2002) Mollusc larval shell formation: amorphous calcium carbonate is a precursor phase for aragonite. *J Exp Zool* 293: 478–491.
  71. Eyster LS (1986) Shell inorganic composition and onset of shell mineralization during bivalve and gastropod embryogenesis. *Biol Bull* 170: 211–231.
  72. Schöntzner V, Weiss IM (2007) The structure of mollusc larval shells formed in the presence of the chitin synthase inhibitor Nikkomycin Z. *BMC structural biology* 7: 71.
  73. R Core Team (2013) R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing.
  74. Zar JH, editor (1974) Biostatistical analysis: Englewood Cliffs, NJ: Prentice-Hall. 620 p.
  75. Scherrer B, editor (1984) Biostatistiques: Gaetan Morin Editors. 850 p.
  76. Chan KYK, Grunbaum D, O'Donnell MJ (2011) Effects of ocean-acidification-induced morphological changes on larval swimming and feeding. *J Exp Biol* 214: 3857–3867.

77. Kim K-S, Shim J, Kim S (2013) Effects of ocean acidification on the larval growth of olive flounder (*Paralichthys olivaceus*). *Biogeosciences Discuss* 10: 7413–7431.
78. Parker LM, Ross PM, O'Connor WA (2010) Comparing the effect of elevated  $\mu\text{CO}_2$  and temperature on the fertilization and early development of two species of oysters. *Mar Biol* 157: 2435–2452.
79. Dupont S, Dorey N, Stumpp M, Melzner F, Thorndyke M (2013) Long-term and trans-life-cycle effects of exposure to ocean acidification in the green sea urchin *Strongylocentrotus droebachiensis*. *Mar Biol* 160: 1835–1843.
80. Parker LM, Ross PM, O'Connor WA, Borysko L, Raftos DA, et al. (2012) Adult exposure influences offspring response to ocean acidification in oysters. *Glob Change Biol* 18: 82–92.
81. Vehmaa A, Brutemark A, Engström-Öst J (2012) Maternal effects may act as an adaptation mechanism for copepods facing pH and temperature changes. *PLoS One* 7: e48538.
82. Gazeau F, Gattuso J-P, Greaves M, Elderfield H, Peene J, et al. (2011) Effect of carbonate chemistry alteration on the early embryonic development of the pacific oyster (*Crassostrea gigas*). *PLoS One* 6: e23010.
83. Thiagarajan V, Ko GWK (2012) Larval growth response of the Portuguese oyster (*Crassostrea angulata*) to multiple climate change stressors. *Aquaculture* 370–371: 90–95.
84. Watson S-A, Southgate PC, Tyler PA, Peck LS (2009) Early larval development of the Sydney rock oyster *Saccostrea glomerata* under near-future predictions of  $\text{CO}_2$ -driven ocean acidification. *J Shellfish Res* 28: 431–437.
85. Andersen S, Grefsrud ES, Harboe T (2013) Effect of increased  $\mu\text{CO}_2$  level on early shell development in great scallop (*Pecten maximus* Lamarck) larvae. *Biogeosciences* 10: 6161–6184.
86. Kurihara H, Asai T, Kato S, Ishimatsu A (2009) Effects of elevated  $\mu\text{CO}_2$  on early development in the mussel *Mytilus galloprovincialis*. *Aquat Biol* 4: 225–233.
87. Marxen JC, Becker W, Finke D, Hasse B, Epple M (2003) Early mineralization in *Biomphalaria glabrata*: microscopic and structural results. *J Molluscan Stud* 69: 113–121.
88. Maeda-Martinez AN (2008) Osmotic and ionic concentration of the egg capsule fluid of *Crepidula fornicata* in relation to embryonic development. *Mar Biol* 154: 643–648.
89. Brante A, Fernandez M, Viard F (2008) Effect of oxygen conditions on intracapsular development in two calyptraeid species with different modes of larval development. *Mar Ecol-Prog Ser* 368: 197–207.
90. De Mahieu G, Penchaszadeh P, Casal A (1974) Algunos aspectos de las variaciones de proteínas y aminoácidos libres totales del líquido intracapsular en relación al desarrollo embrionario en *Adelomelon brasiliense* (Lamarck, 1811)(Gastropoda, Prosobranchia, Volutidae). *Cah Biol Mar* 15: 215–227.
91. Maeda-Martinez AN (1987) The rates of calcium deposition in shells of molluscan larvae. *Comp Biochem Physiol A-Mol Integr Physiol* 86: 21–28.
92. Melzner F, Gutowska MA, Langenbuch M, Dupont S, Lucassen M, et al. (2009) Physiological basis for high  $\text{CO}_2$  tolerance in marine ectothermic animals: pre-adaptation through lifestyle and ontogeny? *Biogeosciences* 6: 2313–2331.
93. Thomsen J, Gutowska MA, Saphorster J, Heinemann A, Trubenbach K, et al. (2010) Calcifying invertebrates succeed in a naturally  $\text{CO}_2$ -rich coastal habitat but are threatened by high levels of future acidification. *Biogeosciences* 7: 3879–3891.
94. Miller AW, Reynolds AC, Sobrino C, Riedel GF (2009) Shellfish face uncertain future in high  $\text{CO}_2$  world: influence of acidification on oyster larvae calcification and growth in estuaries. *PLoS One* 4: e5661.
95. Hickman CS (2001) Evolution and development of gastropod larval shell morphology: experimental evidence for mechanical defense and repair. *Evolution & Development* 3: 18–23.
96. Hickman CS (1999) Adaptive function of gastropod larval shell features. *Invertebrate biology* 118: 346–356.
97. Pechenik JA (2006) Larval experience and latent effects: metamorphosis is not a new beginning. *Integrative and Comparative Biology* 46: 323–333.
98. Anil AC, Desai D, Khandeparker L (2001) Larval development and metamorphosis in *Balanus amphitrite* Darwin (Cirripedia; Thoracica): significance of food concentration, temperature and nucleic acids. *J Exp Mar Biol Ecol* 263: 125–141.
99. Occhipinti-Ambrogi A (2007) Global change and marine communities: Alien species and climate change. *Mar Poll Bull* 55: 342–352.
100. Lenz M, da Gama BAP, Gerner NV, Gobin J, Groner F, et al. (2011) Non-native marine invertebrates are more tolerant towards environmental stress than taxonomically related native species: Results from a globally replicated study. *Environ Res* 111: 943–952.
101. Ko GWK, Chan VBS, Dineshram R, Choi DKS, Li AJ, et al. (2013) Larval and post-larval stages of Pacific oyster *Crassostrea gigas* are resistant to elevated  $\text{CO}_2$ . *PLoS One* 8: e64147.
102. White MM, McCorkle DC, Mullineaux LS, Cohen AL (2013) Early exposure of bay scallops (*Argopecten irradians*) to high  $\text{CO}_2$  causes a decrease in larval shell growth. *PLoS One* 8: e61065.
103. Jansson A, Norkko J, Norkko A (2013) Effects of reduced pH on *Macoma balthica* larvae from a system with naturally fluctuating pH-dynamics. *PLoS One* 8: e68198.
104. Hettinger A, Sanford E, Hill T, Hosfelt J, Russell A, et al. (2013) The influence of food supply on the response of Olympia oyster larvae to ocean acidification. *Biogeosciences* 10: 6629–6638.