



**HAL**  
open science

# Derivation of the bacterial run-and-tumble kinetic equation from a model with biochemical pathway

Benoît Perthame, Min Tang, Nicolas Vauchelet

► **To cite this version:**

Benoît Perthame, Min Tang, Nicolas Vauchelet. Derivation of the bacterial run-and-tumble kinetic equation from a model with biochemical pathway. *Journal of Mathematical Biology*, 2016. hal-01131101

**HAL Id: hal-01131101**

<https://hal.sorbonne-universite.fr/hal-01131101v1>

Submitted on 12 Mar 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Public Domain

# Derivation of the bacterial run-and-tumble kinetic equation from a model with biochemical pathway

Benoît Perthame<sup>\*†‡</sup>

Min Tang<sup>§</sup>

Nicolas Vauchelet<sup>\*†‡</sup>

March 13, 2015

## Abstract

Kinetic-transport equations are, by now, standard models to describe the dynamics of populations of bacteria moving by run-and-tumble. Experimental observations show that bacteria increase their run duration when encountering an increasing gradient of chemotactic molecules. This led to a first class of models which heuristically include tumbling frequencies depending on the path-wise gradient of chemotactic signal.

More recently, the biochemical pathways regulating the flagellar motors were uncovered. This knowledge gave rise to a second class of kinetic-transport equations, that takes into account an intra-cellular molecular content and which relates the tumbling frequency to this information. It turns out that the tumbling frequency depends on the chemotactic signal, and not on its gradient.

For these two classes of models, macroscopic equations of Keller-Segel type, have been derived using diffusion or hyperbolic rescaling. We complete this program by showing how the first class of equations can be derived from the second class with molecular content after appropriate rescaling. The main difficulty is to explain why the path-wise gradient of chemotactic signal can arise in this asymptotic process.

Randomness of receptor methylation events can be included, and our approach can be used to compute the tumbling frequency in presence of such a noise.

**Key words:** kinetic-transport equations; chemotaxis; asymptotic analysis; run and tumble; biochemical pathway;

**Mathematics Subject Classification (2010):** 35B25; 82C40; 92C17

## 1 Introduction

Two classes of kinetic-transport equations have been proposed to describe, at the cell scale, the movement of bacteria by ‘run and tumble’ in a given external effective signal  $M(x, t)$ , usually related to the extra-cellular chemo-attractant concentration  $S$  by a relation of the type  $M = m_0 + \ln(S)$ .

---

<sup>\*</sup>Sorbonne Universités, UPMC Univ Paris 06, UMR 7598, Laboratoire Jacques-Louis Lions, F-75005, Paris, France

<sup>†</sup>CNRS, UMR 7598, Laboratoire Jacques-Louis Lions, F-75005, Paris, France

<sup>‡</sup>INRIA-Paris-Rocquencourt, EPC MAMBA, Domaine de Voluceau, BP105, 78153 Le Chesnay Cedex, France

<sup>§</sup>Institute of natural sciences and department of mathematics, Shanghai Jiao Tong University, Shanghai, 200240, China. This author is partially supported by NSF of Shanghai under grant 12ZR1445400, NSFC 11301336 and 91330203, and Shanghai Pujiang Program 13PJ140700.

The simplest class is for the probability  $\bar{p}(\mathbf{x}, \mathbf{v}, t)$  to find a bacteria at location  $\mathbf{x} \in \mathbb{R}^d$  and with velocity  $\mathbf{v} \in V$  (a smooth bounded subset of  $\mathbb{R}^d$ , one can choose the unit ball to fix idea ). The evolution of this probability is given by a Boltzmann type equation

$$\partial_t \bar{p} + \mathbf{v} \cdot \nabla_{\mathbf{x}} \bar{p} = \mathcal{T}[D_t M](\bar{p}), \quad \mathbf{x} \in \mathbb{R}^d, \mathbf{v} \in V, t \geq 0, \quad (1)$$

with the path-wise gradient of  $M$  defined as

$$D_t M = \partial_t M + \mathbf{v} \cdot \nabla_{\mathbf{x}} M \quad (2)$$

and  $T$  a tumbling kernel which typically takes the form

$$\mathcal{T}[D_t M](\bar{p}) = \int_V [T(D_t M(\mathbf{x}, \mathbf{v}', t), \mathbf{v}, \mathbf{v}') \bar{p}(\mathbf{x}, \mathbf{v}', t) - T(D_t M(\mathbf{x}, \mathbf{v}, t), \mathbf{v}', \mathbf{v}) \bar{p}(\mathbf{x}, \mathbf{v}, t)] d\mathbf{v}'. \quad (3)$$

Such equations, with  $\mathcal{T}$  depending on  $M$  or  $D_t M$ , were used intensively to model bacterial chemotaxis, possibly with  $M$  connected to the cell density, as a result of chemoattractant release by bacteria. They were first introduced in [14] and the Keller-Segel drift-diffusion system was subsequently derived [15] in the diffusion limit; surprisingly, with a kernel  $T$  depending on  $M$  and not on its gradient, and in opposition to the Keller-Segel system which solutions blow-up for large mass, it was proved that the solutions exist globally [4, 10]. However, experiments show that bacteria as *E.coli* extend their runs when feeling an increasing concentration of chemoattractant and this led to study tumbling kernels  $T$  that depend on  $D_t M$ , see [7, 5]. The nonlinear theory is then more difficult (see [3] and the references therein) and blow-up can occur in finite time [2]. These models with  $T$  depending on  $D_t M$  are able to explain the experimental observation of traveling pulses of bacteria, which cannot be done when  $T$  only depends on  $M$  itself, see [16, 17]. Also, departing from this kinetic-transport equation, it is possible to rescale it and study the diffusion and hyperbolic limit as in [7, 5, 16, 17, 10, 9]. When  $T$  undergoes stiff dependency on  $D_t M$ , the hyperbolic limit is singular and the analysis is particularly delicate [11].

More elaborated kinetic models have been proposed recently that incorporate intracellular chemo-sensory system. In the simplest description of the biochemical pathways, they use a single additional variable  $m \geq 0$ , which represents the intracellular methylation level. Then, the kinetic-transport equation is written for the probability density function  $p(\mathbf{x}, \mathbf{v}, m, t)$  of bacteria at time  $t$ , position  $\mathbf{x} \in \mathbb{R}^d$ , moving at velocity  $\mathbf{v} \in V$  and methylation level  $m > 0$

$$\begin{cases} \partial_t p + \mathbf{v} \cdot \nabla_{\mathbf{x}} p + \partial_m [f(m, M)p] = \mathcal{Q}[m, M](p), \\ p(\mathbf{x}, \mathbf{v}, m = 0, t) = 0. \end{cases} \quad (4)$$

The intracellular adaptation dynamics is described by the reaction rate  $f(\cdot)$  for which we assume  $f(m = 0, M) > 0$ , which allows us to pose the boundary condition at  $m = 0$ . The tumbling term  $\mathcal{Q}[m, M](p)$  is

$$\mathcal{Q}[m, M](p) = \int_V [\lambda(m, M, \mathbf{v}, \mathbf{v}') p(t, \mathbf{x}, \mathbf{v}', m) - \lambda(m, M, \mathbf{v}', \mathbf{v}) p(t, \mathbf{x}, \mathbf{v}, m)] d\mathbf{v}', \quad (5)$$

where  $\lambda(m, M, \mathbf{v}, \mathbf{v}')$  denotes the methylation dependent tumbling frequency from  $\mathbf{v}'$  to  $\mathbf{v}$ , in other words the response of the cell depending on its environment and internal state. We borrow this formalism from [12, 18] even though this type of models, involving more general signal transduction, can

be traced back to [5, 7, 8, 21]. The authors in [5, 7, 8, 18, 21, 22] developed the asymptotic theory which allows to recover, in the diffusion and in the hyperbolic limits, macroscopic equations where the variables are only  $(\mathbf{x}, t)$  as the Keller-Segel system, or  $(\mathbf{x}, m, t)$  for structured Keller-Segel models.

In the program of establishing the relations between these pieces of the model hierarchy for bacterial population motion, a derivation is missing: how are related these two classes of kinetic models (1)–(3) and (4)–(5)?

Our goal is to show how, assuming fast adaptation and stiff response, the methylation level is at equilibrium with the external signal represented by  $M$ , and the equation (1) can be derived from (4). In particular we aim at computing the bulk tumbling kernel  $T(D_t M, \mathbf{v}, \mathbf{v}')$  from the methylation dependent kernel  $\lambda(m, M, \mathbf{v}, \mathbf{v}')$ , a statement we give in the next section. Two difficulties arise here: one is to infer the proper rescaling in the kinetic equations, the second is to carry-out the mathematical analysis for singular limits. Our approach allows us to also include noise resulting from random receptor-methylation and demethylation events. The proof of the formula for  $T$  is given in sections 3 and 4; we show that a direct use of the variable  $m$  is not enough to produce the formula and that a new variable is needed, which zooms on the intra- and extra-cellular methylation equilibrium. We conclude by relating our notations to a more physically based description of the same model where the cell receptors activity is used in the model parameters, see section 5.

To keep simplicity, we assume that the external signal function  $M(\mathbf{x}, t)$  is given and smooth. Therefore questions of existence and blow-up are not considered here.

## 2 Fast adaptation, stiff response

**Assumptions.** For our mathematical derivation, we introduce a small parameter  $\epsilon$  which acts both as a fast time scale for external signal transduction and as a stiffness parameter for the response in terms of tumbling rate. We assume moreover that the reaction rate  $f$  only depends on the difference  $m - M$ , in accordance with the physical models that we recall in Section 5. Therefore, we rescale equation (4)–(5) as

$$\begin{cases} \partial_t p_\epsilon + \mathbf{v} \cdot \nabla_{\mathbf{x}} p_\epsilon + \frac{1}{\epsilon} \partial_m \left( f(m - M) p_\epsilon \right) = \mathcal{Q}_\epsilon[m, M](p_\epsilon), \\ p_\epsilon(\mathbf{x}, \mathbf{v}, m = 0, t) = 0, \end{cases} \quad (6)$$

with the tumbling kernel

$$\mathcal{Q}_\epsilon[m, M](p_\epsilon) = \int_V \left[ \Lambda\left(\frac{m - M}{\epsilon}, \mathbf{v}, \mathbf{v}'\right) p_\epsilon(\mathbf{x}, \mathbf{v}', m, t) - \Lambda\left(\frac{m - M}{\epsilon}, \mathbf{v}', \mathbf{v}\right) p_\epsilon(\mathbf{x}, \mathbf{v}, m, t) \right] d\mathbf{v}'. \quad (7)$$

We complete this equation with an initial data  $p^{\text{ini}} \geq 0$  which satisfies

$$\iint \int_{\mathbb{R}^d \times V \times \mathbb{R}} (1 + m^2) p^{\text{ini}}(\mathbf{x}, \mathbf{v}, m) d\mathbf{x} d\mathbf{v} dm < \infty, \quad (8)$$

$$\bar{p}^{\text{ini}} := \int_{\mathbb{R}} p^{\text{ini}} dm \in L^\infty(\mathbb{R}^d \times V). \quad (9)$$

Also, we are going to use several assumptions for the functions  $M$ ,  $f$  and  $\Lambda$ . We assume they are as smooth as necessary and that for some constants  $m_{\pm}$ ,  $g_{\pm}$ ,  $\lambda_{\pm}$ ,

$$0 < m_- \leq M(\mathbf{x}, t) \leq m_+, \quad M \in C_b^1(\mathbb{R}^d \times [0, \infty)), \quad (10)$$

$$f(y) = -yG(y), \quad \text{with } G \in C_b^1(\mathbb{R}), \quad 0 < g_- \leq G(y) \leq g_+, \quad (11)$$

$$\partial_y \Lambda(y, \mathbf{v}, \mathbf{v}') < 0, \quad 0 < \lambda_- \leq \Lambda(y, \mathbf{v}, \mathbf{v}') \leq \lambda_+. \quad (12)$$

Various scalings have been proposed for kinetic equation and the closest, but still different seems to be the high field limit [1]. In [13], still other scalings or limits are studied.

**The main result.** With these assumptions, we are going to show that as  $\epsilon$  vanishes, we recover the simpler model (1)–(3) as a limit of (6).

**Theorem 2.1 (Derivation of the kinetic equation)** *We make the assumptions (8)–(9) on the initial data, and (10)–(12) on the coefficients. Let  $p_\epsilon$  be the solution to (6). Then, for all  $T > 0$ ,  $\bar{p}_\epsilon$  is bounded in  $L^\infty([0, T] \times \mathbb{R}^d \times V)$  and*

$$\bar{p}_\epsilon := \int_{\mathbb{R}} p_\epsilon dm \xrightarrow[\epsilon \rightarrow 0]{} \bar{p}_0 \quad \text{in } L^\infty([0, T] \times \mathbb{R}^d \times V)\text{-weak-}\star$$

and  $\bar{p}_0$  satisfies equation (1)–(3) with

$$T(u, \mathbf{v}, \mathbf{v}') = \Lambda\left(-\frac{u}{G(0)}, \mathbf{v}, \mathbf{v}'\right).$$

Furthermore, we have  $\bar{p}_0 := \int_{\mathbb{R}} p_0 dm$  with  $p_0$  the weak limit (in measures, see (15)) of  $p_\epsilon$  which is given by

$$p_0(\mathbf{x}, \mathbf{v}, m, t) = \bar{p}_0(\mathbf{x}, \mathbf{v}, t) \delta(m = M(\mathbf{x}, t)).$$

Before we prove this theorem in the next sections, we present a variant of this result.

**Internal noise.** Due to random receptor-methylation and demethylation events, some internal noise can be observed in *E. coli* chemotaxis and we can model it by adding a diffusion term in  $m$  [6]. The model is as follows:

$$\partial_t p_\epsilon + \mathbf{v} \cdot \nabla_{\mathbf{x}} p_\epsilon + \frac{1}{\epsilon} \partial_m \left( f(m - M) p_\epsilon \right) = \epsilon \partial_{mm}^2 p_\epsilon + \mathcal{Q}_\epsilon[m, M](p_\epsilon), \quad (13)$$

with the no-flux boundary condition that now reads

$$f(-M) p_\epsilon(\mathbf{x}, \mathbf{v}, m, t) - \epsilon^2 \partial_m p_\epsilon(\mathbf{x}, \mathbf{v}, m, t) = 0, \quad \text{at } m = 0. \quad (14)$$

**Theorem 2.2 (Limit with noise)** *With the assumptions and notations of Theorem 2.1, the same conclusions hold for the solution  $p_\epsilon$  of (13), with the same expression for  $p_0$  and*

$$T(u, \mathbf{v}, \mathbf{v}') = \sqrt{\frac{G(0)}{2\pi}} \int_{\mathbb{R}} \Lambda(y, \mathbf{v}, \mathbf{v}') e^{-\frac{G(0)}{2} \left(y + \frac{u}{G(0)}\right)^2} dy.$$

**A priori bounds and principle of the proof.** Before we explain the derivation of the formula stated in these theorems, let us make some observations which explain the difficulty. Because we assume that  $M(x, t)$  is given, we handle a linear equation for which existence and uniqueness of weak solutions is well established. The nonlinear case, when the chemoattractant concentration giving rise to  $M$  is coupled to  $p^\epsilon$ , can also be treated, see [13]. In particular we will make use of the uniform estimates (see Section 4)

$$\iiint_{\mathbb{R}^d \times V \times \mathbb{R}} p_\epsilon(\mathbf{x}, \mathbf{v}, m, t) \, d\mathbf{x} \, d\mathbf{v} \, dm = \iiint_{\mathbb{R}^d \times V \times \mathbb{R}} p^{\text{ini}}(\mathbf{x}, \mathbf{v}, m) \, d\mathbf{x} \, d\mathbf{v} \, dm, \quad \forall t \geq 0, \quad (15)$$

$$\bar{p}_\epsilon(\mathbf{x}, \mathbf{v}, t) \leq \|\bar{p}^{\text{ini}}(\mathbf{x}, \mathbf{v})\|_\infty e^{Ct}, \quad \forall t \geq 0, \quad (16)$$

where  $C$  is a nonnegative constant. From these bounds, we conclude that we can extract subsequences (but to simplify the notations we ignore this subsequence) which converge as mentioned in the theorems.

Passing to the limit in the equation on  $p_\epsilon$  (with or without noise) gives us

$$\partial_m \left( f(m - M) p_0 \right) = 0.$$

This tells us that  $f(m - M) p_0 = 0$  (it is constant and  $p_0$  is integrable). Because, with assumption (11),  $f(m - M)$  vanishes only for  $m - M = 0$ , we conclude that  $p_0$  is a Dirac mass at  $m = M$ , hence the expression of  $p_0$  in Theorems 2.1 and 2.2.

However this information is not enough to pass to limit in the equation on  $\bar{p}_\epsilon$  obtained integrating in  $m$  equation (6) or (13), that is

$$\partial_t \bar{p}_\epsilon + \mathbf{v} \cdot \nabla_{\mathbf{x}} \bar{p}_\epsilon = \int_{\mathbb{R}^+} \mathcal{Q}_\epsilon[m, M](p_\epsilon) \, dm.$$

Indeed, in the right hand side, the product  $\Lambda\left(\frac{m-M}{\epsilon}, \mathbf{v}, \mathbf{v}'\right) p_\epsilon(\mathbf{x}, \mathbf{v}', m, t)$  is, in the limit, a discontinuity multiplied by a Dirac mass. For this reason, we have to rescale in  $m$  in order to evaluate this limit, which we do in the next section.

### 3 The change of variable

To get a more accurate view of the convergence of  $p_\epsilon$  to a Dirac mass in  $m$ , and following [7], we introduce a blow-up variable around  $m = M$ . We set

$$y = \frac{m - M}{\epsilon}, \quad q_\epsilon(\mathbf{x}, \mathbf{v}, y, t) = \epsilon p_\epsilon(\mathbf{x}, \mathbf{v}, m, t) \quad (17)$$

so that

$$\bar{q}_\epsilon(\mathbf{x}, \mathbf{v}, t) := \int_{\mathbb{R}} q_\epsilon(\mathbf{x}, \mathbf{v}, y, t) \, dy = \int_{\mathbb{R}} p_\epsilon(\mathbf{x}, \mathbf{v}, m, t) \, dm = \bar{p}_\epsilon(\mathbf{x}, \mathbf{v}, t). \quad (18)$$

Because of these identities, our statements will equivalently be on  $\bar{q}_\epsilon$  and will go through the analysis of  $q_\epsilon$  rather than  $p_\epsilon$  itself.

Also notice that the bounds in (15), (16) also hold true for  $q_\epsilon$  and  $\bar{q}_\epsilon$  and allow us to take weak limits.

(i) **Without noise.** The equation for  $q_\epsilon(t, \mathbf{x}, \mathbf{v}, y)$  is written, using the definition in (2),

$$\begin{aligned} \partial_t q_\epsilon + \mathbf{v} \cdot \nabla_{\mathbf{x}} q_\epsilon & - \frac{1}{\epsilon} D_t M \partial_y q_\epsilon + \frac{1}{\epsilon^2} \partial_y (f(\epsilon y) q_\epsilon) \\ & = \int_V \left[ \Lambda(y, \mathbf{v}, \mathbf{v}') q_\epsilon(\mathbf{x}, \mathbf{v}', y, t) - \Lambda(y, \mathbf{v}', \mathbf{v}) q_\epsilon(\mathbf{x}, \mathbf{v}, y, t) \right] d\mathbf{v}'. \end{aligned}$$

From (11), we can write  $f(\epsilon y) = \epsilon y G(\epsilon y)$  and the above equation becomes

$$\begin{aligned} \partial_t q_\epsilon + \mathbf{v} \cdot \nabla_{\mathbf{x}} q_\epsilon & - \frac{1}{\epsilon} D_t M \partial_y q_\epsilon - \frac{1}{\epsilon} \partial_y (y G(\epsilon y) q_\epsilon) \\ & = \int_V \left[ \Lambda(y, \mathbf{v}, \mathbf{v}') q_\epsilon(\mathbf{x}, \mathbf{v}', y, t) - \Lambda(y, \mathbf{v}', \mathbf{v}) q_\epsilon(\mathbf{x}, \mathbf{v}, y, t) \right] d\mathbf{v}'. \end{aligned} \tag{19}$$

Because  $q_\epsilon$  is a bounded measure on  $\mathbb{R}^d \times V \times \mathbb{R}^+ \times (0, T)$ , for all  $T > 0$ , as  $\epsilon \rightarrow 0$ ,  $q_\epsilon$  has a weak limit  $q_0$  in the sense of measure (again after extraction) and the above equation gives, in the distributional sense,

$$\partial_y (y G(0) q_0 + D_t M(S) q_0) = 0. \tag{20}$$

From this, we infer that

$$q_0(t, \mathbf{x}, \mathbf{v}, y) = \bar{q}_0(t, \mathbf{x}, \mathbf{v}) \delta\left(y = -\frac{D_t M(S)}{G(0)}\right). \tag{21}$$

This information is useful provided we can establish that

$$\bar{q}_0(\mathbf{x}, \mathbf{v}, t) = \int_{\mathbb{R}} q_0(\mathbf{x}, \mathbf{v}, y, t) dy = \text{weak-}\lim_{\epsilon \rightarrow 0} \bar{q}_\epsilon(\mathbf{x}, \mathbf{v}, t). \tag{22}$$

This step is postponed to Section 4 and involves a control of the tail for large values of  $m$ .

We may also integrate equation (19) with respect to  $y$  and find in the limit

$$\partial_t \bar{q}_0 + \mathbf{v} \cdot \nabla_{\mathbf{x}} \bar{q}_0 = \int_V \left[ \Lambda\left(-\frac{D'_t M}{G(0)}, \mathbf{v}, \mathbf{v}'\right) \bar{q}'_0 d\mathbf{v}' - \Lambda\left(-\frac{D_t M}{G(0)}, \mathbf{v}', \mathbf{v}\right) \bar{q}_0 \right], \tag{23}$$

where  $D'_t M(S)$  is the total derivative, as in (2), but in the direction  $\mathbf{v}'$  and where  $\bar{q}'_0$  represents  $\bar{q}_0(\mathbf{x}, \mathbf{v}', t)$ . Finally, for any smooth test function  $\phi$ , we have from the change of variable  $m \mapsto y = (m - M)/\epsilon$ ,

$$\int_{\mathbb{R}} p_\epsilon(\mathbf{x}, \mathbf{v}, m, t) \phi(m) dm = \int_{\mathbb{R}} q_\epsilon(\mathbf{x}, \mathbf{v}, y, t) \phi(M + \epsilon y) dy \rightarrow \bar{q}_0(\mathbf{x}, \mathbf{v}, t) \phi(M),$$

where we use (21). This gives the limiting expression of  $p_0$  in Theorem 2.1.

These are the results stated in Theorem 2.1, if we can establish the relation  $\bar{q}_0 = \bar{p}_0$  as stated in (22), which we do later.

(ii) **With internal noise.** Similarly, after introducing the new variables as in (17), the equation (13) for  $q_\epsilon(t, \mathbf{x}, \mathbf{v}, y)$  writes

$$\begin{aligned} \partial_t q_\epsilon + \mathbf{v} \cdot \nabla_{\mathbf{x}} q_\epsilon & - \frac{1}{\epsilon} D_t M \partial_y q_\epsilon + \frac{1}{\epsilon^2} \partial_y (f(\epsilon y) q_\epsilon) \\ & = \frac{1}{\epsilon} \partial_{yy}^2 q_\epsilon + \int_V \left[ \Lambda(y, \mathbf{v}, \mathbf{v}') q_\epsilon(\mathbf{x}, \mathbf{v}', y, t) - \Lambda(y, \mathbf{v}', \mathbf{v}) q_\epsilon(\mathbf{x}, \mathbf{v}, y, t) \right] d\mathbf{v}'. \end{aligned}$$

From (11), this equation becomes

$$\begin{aligned} \partial_t q_\epsilon + \mathbf{v} \cdot \nabla_{\mathbf{x}} q_\epsilon & - \frac{1}{\epsilon} D_t M \partial_y q_\epsilon - \frac{1}{\epsilon} \partial_y (y G(\epsilon y) q_\epsilon) \\ & = \frac{1}{\epsilon} \partial_{yy}^2 q_\epsilon + \int_V \left[ \Lambda(y, \mathbf{v}, \mathbf{v}') q_\epsilon(\mathbf{x}, \mathbf{v}', y, t) - \Lambda(y, \mathbf{v}', \mathbf{v}) q_\epsilon(\mathbf{x}, \mathbf{v}, y, t) \right] d\mathbf{v}'. \end{aligned} \quad (24)$$

In the limit  $\epsilon \rightarrow 0$ , the above equation converges to, in the sense of distributions,

$$\partial_y (y G(0) q_0 + D_t M q_0) = -\partial_{yy}^2 q_0,$$

which shows that

$$q_0(\mathbf{x}, \mathbf{v}, y, t) = \bar{q}_0(\mathbf{x}, \mathbf{v}, t) \sqrt{\frac{G(0)}{2\pi}} e^{-\frac{G(0)}{2} \left(y + \frac{D_t M}{G(0)}\right)^2}, \quad (25)$$

a useful information, still assuming we have proved the relation (22) for  $\bar{q}_0$ .

We conclude as before. After integration of (24) with respect to  $y$ , passing to the limit  $\epsilon \rightarrow 0$ , we find

$$\begin{cases} \partial_t \bar{q}_0 + \mathbf{v} \cdot \nabla_{\mathbf{x}} \bar{q}_0 = \int_V \left[ T(D_t M, \mathbf{v}', \mathbf{v}) \bar{q}'_0 - T(D_t M, \mathbf{v}, \mathbf{v}') \bar{q}_0 \right] u d\mathbf{v}', \\ T(D_t M, \mathbf{v}, \mathbf{v}') = \sqrt{\frac{G(0)}{2\pi}} \int_{\mathbb{R}} \Lambda(y, \mathbf{v}, \mathbf{v}') e^{-\frac{G(0)}{2} \left(y + \frac{D_t M}{G(0)}\right)^2} dy. \end{cases} \quad (26)$$

The Theorem 2.2 is also proved.  $\square$

## 4 A priori bounds

We now establish the various estimates which justify that we can pass to the limit as indicated in Section 3 and thus we prove the

**Lemma 4.1** *We make the assumptions of Theorem 2.1, then the condition (22) holds and for some constant which depends on  $\iint y^2 q^{\text{ini}} dy dx dv$  and  $\|M\|_{W^{1,\infty}(\mathbb{R}^d \times \mathbb{R}^+)}$ , we have*

$$\iint y^2 q_\epsilon(t) dy dx dv \leq C(q^{\text{ini}}, M).$$

Consequently,  $q_\epsilon$  converges weakly in the sense of measure towards  $q_0$  and

- (i) for  $q_\epsilon$  a solution to (19),  $q_0$  is given by (21) with  $\bar{q}_0$  weak solution of (23),
- (ii) for  $q_\epsilon$  a solution to (24), then,  $q_0$  is given by (25) with  $\bar{q}_0$  weak solution of (26).

**Proof.** We only consider the case (i) without noise, the case (ii) is obtained by the same token. We first prove some estimates which imply weak convergence. Then, we pass to the limit in the equation satisfied by  $\bar{q}_\epsilon$ .

**$L^1$  bound.** For completeness, we recall that equation (19) is positivity preserving and conservative. It follows the uniform, in  $\epsilon$ , bound for  $q_\epsilon$  in  $L^1$ , see (15).

**$L^\infty$  bound on  $\bar{q}_\epsilon$ .** We use the notation (18) for  $\bar{q}_\epsilon$ . Arguing in the spirit of [10, 3, 20]), we first prove the uniform  $L^\infty$  bound on  $\bar{q}_\epsilon$ .



Integrating (19) with respect to  $y$ , from the bound (12) and the nonnegativity of  $q_\epsilon$ , we get

$$\partial_t \bar{q}_\epsilon + \mathbf{v} \cdot \nabla_{\mathbf{x}} \bar{q}_\epsilon \leq \lambda_+ \int \bar{q}_\epsilon d\mathbf{v}'.$$

Then, using the method of characteristics, we have  $\partial_t \bar{q}_\epsilon(t, \mathbf{x} + \mathbf{v}t) \leq \lambda_+ \int \bar{q}_\epsilon(\mathbf{x} + \mathbf{v}t, \mathbf{v}', t) d\mathbf{v}'$ , which implies after integration

$$\bar{q}_\epsilon(t, \mathbf{x}, \mathbf{v}) - \bar{q}^{\text{ini}}(\mathbf{x} - \mathbf{v}t) \leq \lambda_+ \int_0^t \int \bar{q}_\epsilon(\mathbf{x} - \mathbf{v}s, \mathbf{v}', t-s) d\mathbf{v}' ds.$$

Taking the supremum in  $\mathbf{x}, \mathbf{v}$ , we find

$$\|\bar{q}_\epsilon(t)\|_\infty \leq \|\bar{q}^{\text{ini}}\|_\infty + \lambda_+ |V| \int_0^t \|\bar{q}_\epsilon(s)\|_\infty ds,$$

and using Gronwall's inequality, we find the estimate in (16).

**Control on the tail in  $m$ .** In order to prove the condition (22), we need to ensure that there is no mass loss at infinity in  $m$ . To do so, we multiply both sides of (19) by  $y^2$  and integrate by parts with respect to  $x, v$ , and  $y$ . This yields

$$\frac{d}{dt} \iint y^2 q_\epsilon dy dx dv + \frac{2}{\epsilon} \iint y^2 G(\epsilon y) q_\epsilon dy dx dv + \frac{2}{\epsilon} \iint y D_t M q_\epsilon dy dx dv = 0.$$

Using the Cauchy-Schwarz inequality, we deduce

$$\begin{aligned} \frac{d}{dt} \iint y^2 q_\epsilon dy dx dv + \frac{2}{\epsilon} \iint y^2 G(\epsilon y) q_\epsilon dy dx dv &\leq \frac{1}{\epsilon} \iint y^2 G(\epsilon y) q_\epsilon dy dx dv \\ &\quad + \frac{1}{\epsilon} \iint (D_t M)^2 \frac{q_\epsilon}{G(\epsilon y)} dy dx dv. \end{aligned}$$

By assumption (10),  $D_t M$  is bounded in  $L^\infty([0, T] \times \mathbb{R}^d \times V)$ . From assumption (11) and the mass conservation, the last integral of the right hand side is uniformly bounded by a constant denoted by  $C > 0$ . Then from assumption (11), we have,

$$\frac{d}{dt} \iint y^2 q_\epsilon dy dx dv + \frac{g_-}{\epsilon} \iint y^2 q_\epsilon dy dx dv \leq \frac{C}{\epsilon}.$$

From the Gronwall Lemma, we deduce the bound for all  $t > 0$ ,

$$\iint y^2 q_\epsilon(t) dy dx dv \leq e^{-tg_-/\epsilon} \iint y^2 q^{\text{ini}} dy dx dv + \frac{C}{g_-},$$

which implies a uniform bound on  $\iint y^2 q_\epsilon(t) dy dx dv$ .

**Passing to the limit.** From the bound above, we deduce that, we can extract a subsequence which converges weakly in measure  $q_\epsilon \rightharpoonup q_0$  and such that  $\bar{q}_\epsilon \rightharpoonup \bar{q}_0$  in  $L^\infty$ -weak\*. Then we can pass to the limit in the sense of distribution in equation (19) and deduce that the limit  $q_0$  satisfies equation (20) in the sense of distribution. In fact, we notice that from the Lipschitz character of  $G$ , we have

$$\iint y(G(\epsilon y) - G(0)) \bar{q}_\epsilon dx dy \leq C\epsilon \iint y^2 \bar{q}_\epsilon dx dy \rightarrow 0, \quad \text{as } \epsilon \rightarrow 0,$$

thanks to the estimate on the tail above. Finally, (20) implies that  $yG(0)q_0 + D_t M q_0$  is constant a.e., and this constant should be 0 since from the estimate on the tail above, we have that  $yq_0 \in L^1$ . We conclude that  $q_0$  vanishes except when  $yG(0) + D_t M(S) = 0$ . By conservation of the mass, we deduce the expression (21) for  $p_0$ .

## 5 Comments on physical background

The form of the equation (6) corresponds, for *E. coli* chemotaxis, to the formalism in the physical literature. We have simplified the notations for mathematical clarity and we explain now how to relate our notations to known biophysical quantities. Here we have used the same biological parameters as in [12, 19].

- The methylation level  $M(x, t)$  at equilibrium is related to the extra-cellular attractant profile  $S$ , by a logarithmic dependency

$$M = M(S) = m_0 + \frac{f_0(S)}{\alpha}, \quad \text{with} \quad f_0(S) = \ln\left(\frac{1 + S/K_I}{1 + S/K_A}\right).$$

The constants  $m_0$ ,  $K_I$ ,  $K_A$  represent the basic methylation level, and the dissociation constants for inactive, respectively active, receptors. Numerical values are given by  $\alpha = 1.7$ ,  $m_0 = 1$ ,  $K_I = 18.2\mu M$ ,  $K_A = 3mM$ .

- The receptor activity  $a(m, S)$  depends on the intracellular methylation level  $m$  and the extra-cellular chemoattractant concentration  $S$  such that

$$a = (1 + \exp(NE))^{-1}, \quad \text{with} \quad E = -\alpha(m - m_0) + f_0(S) = -\alpha(m - M(S)). \quad (27)$$

The coefficient  $N = 6$  represents the number of tightly coupled receptors.

- The intracellular dynamics and tumbling frequency are given by

$$f(m - M(S)) = F(a) = k_R(1 - a/a_0), \quad \lambda(m - M(S)) = Z(a) = z_0 + \tau^{-1}\left(\frac{a}{a_0}\right)^H,$$

where  $a(m, S)$  is the receptor activity defined in (27). The parameter  $k_R$  is the methylation rate,  $a_0$  is the receptor preferred activity which is such that  $f(a_0) = 0$  and  $f'(a_0) < 0$ . For the tumbling frequency,  $z_0$ ,  $H$ ,  $\tau$  represent the rotational diffusion, the Hill coefficient of flagellar motors response curve and the average run time respectively. All these parameters can be measured biologically, their values are given by  $k_R = 0.01s^{-1} \sim 0.0005s^{-1}$ ,  $a_0 = 0.5$ ,  $z_0 = 0.14s^{-1}$ ,  $\tau = 0.8s$ ,  $H = 10$ .

- Two kinds of noise can be observed in the signaling pathway for *E. coli*, one is from the external fluctuation of the ligand concentration and the other is the internal noise from the random receptor-methylation and demethylation event [6]. For small complexes, the effect on the activity from the external noise is negligible compared to the internal noise.

We refer the readers to [19, 6, 12, 18] and the references therein for the detailed physical meanings of these parameters.

In *E. coli* chemotaxis, the adaptation time is faster than the system time, i.e. the rescaled constant  $k_R$  is large. Let  $k_R = \frac{1}{\epsilon}$ , where  $\epsilon$  is the ratio between the system time and the adaptation time. For example, when the system time is 1000s and the adaptation time is 100s,  $\epsilon = 0.1$ .

Moreover, the Hill coefficient  $H$  is large which indicates that  $\lambda(m - M(S))$  varies fast with respect to  $m - M(S)$ . Therefore, the scaling introduced in (6) is satisfied by *E. coli* chemotaxis. We can use

$$f(r) = 1 - \frac{1}{a_0} (1 + \exp(-N\alpha r))^{-1}.$$

Therefore, the function  $G(\cdot)$  used in (11) is given by

$$G(r) = -\frac{f(r)}{r} = -\frac{1}{r} \left(1 - \frac{1}{a_0} (1 + \exp(-N\alpha r))^{-1}\right), \quad \text{with } G(0) = -f'(0) = \frac{N\alpha}{4a_0}.$$

Besides, from Theorem 2.1 for the case without noise, using  $y = \frac{r}{\epsilon}$  yields

$$\begin{aligned} T(D_t M, \mathbf{v}, \mathbf{v}') &= \Lambda(y, \mathbf{v}, \mathbf{v}') = z_0 + \tau^{-1} \frac{(1 + \exp(-N\alpha \epsilon y))^{-H}}{a_0^H} \\ &= z_0 + \tau^{-1} \frac{\left(1 + \exp(N\alpha \epsilon D_t M(S)/G(0))\right)^{-H}}{a_0^H}. \end{aligned}$$

And from Theorem 2.1, when the noise in the methylation level is considered,

$$T(D_t M, \mathbf{v}, \mathbf{v}') = \sqrt{\frac{G(0)}{2\pi}} \int_{\mathbb{R}} \Lambda(y, \mathbf{v}, \mathbf{v}') e^{-\frac{G(0)}{2} \left(y + \frac{D_t M}{G(0)}\right)^2} dy.$$

Since the run duration last longer when bacteria encounter an increasing gradient of chemotactic molecules, this lead to higher bacteria density at the place where the ligand concentration  $S$  is higher. This phenomena is well explained by the classical Keller-Segel model which can be considered as the parabolic limit of the kinetic-transport models. However, recent experimental observation shows that higher ligand concentration leading to higher bacteria density is only valid in a spatial-temporal slow-varying environment. When the ligand concentration varies fast, there exists a phase shift between the mass center of ligand concentration and of the bacterias [23]. This is due to the memory effect in the slow methylation adaptation rate. In the limiting kinetic model,  $D_t M(S)$  takes into account the memory effect along the trajectory of the moving cell [20]. Then, two interesting questions come: are these two kinds of memory effect the same? Can we see the phase shift between the mass center of ligand concentration and of the bacteria in the limiting kinetic model?

## 6 Numerical illustrations

We performed numerical simulations using the method SPECS [12]. It is a cell based model that takes into account the evolution of each cell intracellular methylation level, which determines the tumbling frequency of each bacteria. As explained in [18, 19], SPECS and the kinetic model that incorporates intracellular chemo-sensory system (4) show a quantitative match. As in [18, 19], we choose the two velocity kinetic model and a periodic 1D traveling wave concentration

$$S(x, t) = S_0 + S_A \sin\left[\frac{2\pi}{\ell}(x - ut)\right],$$

which is spatial-temporal varying and where the wavelength  $\ell$  is equal to the length of the domain. We compare the numerical results of SPECS and the limiting kinetic model in Figure 1. Upwind scheme is used for the transport terms and periodic boundary conditions are considered. The density is scaled at the order of  $10^{-3}$ , it is the ratio between the actual cell number and the total number. In the Figure 1, the  $x$  axis represents the remainder of  $x - ut \bmod \ell$ , i.e. we keep tracking the wave front of the periodic traveling wave.

Two different wave velocities  $u = 0.4\mu\text{m}/\text{s}$  and  $u = 8\mu\text{m}/\text{s}$  are considered. We compare the density profiles  $\rho = \int_V p d\mathbf{v}$  and the cell flux  $J = \int_V \mathbf{v} p d\mathbf{v}$ . When the concentrated wave moves slowly, the




Figure 1: Numerical comparison between the limiting kinetic model and SPECS. The steady state profiles of density  $\rho$  (top) and current  $J$  (bottom) are presented. Left:  $u = 0.4\mu\text{m}/\text{s}$ ; Right:  $u = 8\mu\text{m}/\text{s}$ . In the subfigures, histograms and solid lines are from SPECS, while the dash star and dash dotted lines are calculated using the limiting kinetic equation. Parameters used here are  $[L]_0 = 500\mu\text{M}$ ,  $[L]_A = 100\mu\text{M}$ ,  $\ell = 800\mu\text{m}$ . We have used 20,000 cells for simulation with SPECS.

limiting kinetic model gives good consistency, however in the fast-varying environment, the density and cell flux profiles are different for SPECS and the limiting kinetic model. We refer the readers to [12, 19] for more detailed discussions and physical explanations.

This numerical experiment shows that the memory effect using the model based on  $D_t M(S)$  is different from the memory effect for the complete model (4) when fast external chemoattractant waves are considered. This phenomena can be explained by the slow adaptation rate in the methylation level and the memory along the trajectory compared to the phase shift. In this fast wave regime, our mathematical results do not apply because the scaling assumptions are not satisfied.

## References

- [1] F. Berthelin, N. J. Mauser and F. Poupaud, high-field limit from a kinetic equation to multidimensional scalar conservation laws. *J. Hyp. Diff. Eq.* 4(1) (2007) 123–145.
- [2] N. Bournaveas, V. Calvez, *Critical mass phenomenon for a chemotaxis kinetic model with spherically symmetric initial data*, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 26 (5) (2009), 1871–1895.
- [3] N. Bournaveas, V. Calvez, S. Gutiérrez, and B. Perthame, *Global existence for a kinetic model of chemotaxis via dispersion and Strichartz estimates*, *Comm. P.D.E.* 33 (2008), 79–95.
- [4] F. Chalub, P. A. Markowich, B. Perthame, and C. Schmeiser, *Kinetic models for chemotaxis and their drift-diffusion limits*, *Monatsh. Math.* 142 (2004), 123–141.
- [5] Y. Dolak, C. Schmeiser, *Kinetic models for chemotaxis: Hydrodynamic limits and spatio-temporal mechanisms*, *J. Math. Biol.* 51 (2005), 595–615.
- [6] R. G. Endres, *Physical principles in sensing and signaling, with an introduction to modeling in biology*, Oxford University Press, 2013.

- [7] R. Erban, H. Othmer, *From individual to collective behaviour in bacterial chemotaxis*. SIAM J. Appl. Math. 65(2) (2004), 361–391.
- [8] R. Erban, H. Othmer, *Taxis equations for amoeboid cells*. J. Math. Biol. (2007) 54:847–885.
- [9] T. Hillen and K. Painter, *Transport and anisotropic diffusion models for movement in oriented habitats*, Dispersal, Individual Movement and Spatial Ecology: A mathematical perspective. Eds: M.A. Lewis, P. Maini, S. Petrowskii, Heidelberg, Springer, 2012, 177–222.
- [10] H. J. Hwang, K. Kang, A. Stevens, *Global Solutions of Nonlinear Transport Equations for Chemotaxis*, SIAM. J. Math. Anal. 36 (2005) 1177–1199.
- [11] F. James, N. Vauchelet, *Chemotaxis : from kinetic equations to aggregate dynamics*, Nonlinear Diff. Eq. Appl. 20(1), (2013), 101–127.
- [12] L. Jiang, Q. Ouyang, and Y. Tu, *Quantitative modeling of Escherichia coli chemotactic motion in environments varying in space and time*, PLoS Comput. Biol. 6 (2010), e1000735.
- [13] J. Liao, Global solution for a kinetic chemotaxis model with internal dynamics and its fast adaptation limit. Work in preparation.
- [14] H. Othmer, S. Dunbar, and W. Alt, *Models of dispersal in biological systems*, J. Math. Biol., 26 (1988), 263–298.
- [15] H. G. Othmer, and T. Hillen, *The diffusion limit of transport equations II: Chemotaxis equations*, SIAM J. Appl. Math., 62 (2002), 122–1250.
- [16] J. Saragosti, V. Calvez, N. Bournaveas, A. Buguin, P. Silberzan, and B. Perthame. *Mathematical description of bacterial traveling pulses*. PLoS Comput Biol, 6(8) :e1000890 (2010). doi:10.1371/journal.pcbi.1000890
- [17] J. Saragosti, V. Calvez, N. Bournaveas, B. Perthame, A. Buguin, and P. Silberzan. *Directional persistence of chemotactic bacteria in a traveling concentration wave*. Proceedings of the National Academy of Sciences, 108(39) (2011), 16235–16240.
- [18] G. Si, M. Tang, and X. Yang, *A pathway-based mean-field model for E. coli chemotaxis: mathematical derivation and keller-segel limit*, Multiscale Model Simul. 12(2), (2014), 907–926.
- [19] G. Si, T. Wu, Q. Ouyang, and Y. Tu, *A pathway-based mean-field model for Escherichia coli chemotaxis*, Phys. Rev. Lett. 109 (2012), 048101
- [20] N. Vauchelet, *Numerical simulation of a kinetic model for chemotaxis*, Kin. Rel. Models., Vol 3, no 3 (2010), 501–528.
- [21] C. Xue and H. G. Othmer. *Multiscale models of taxis-driven patterning in bacterial populations*, SIAM J. Appl. Math., Vol. 70, no. 1,(2009), 133–167.
- [22] C. Xue *Macroscopic equations for bacterial chemotaxis: integration of detailed biochemistry of cell signaling*, J. Math. Biol. Vol. 70, (2015), 1–44.
- [23] X. Zhu, G. Si, N. Deng, Q. Ouyang, T. Wu, Z. He, L. Jiang, C. Luo, and Y. Tu, Frequency-dependent Escherichia coli chemotaxis behavior, Phys. Rev. Lett., 108 (2012), 128101.