



HAL
open science

Solid state NMR and diatoms: probing the interfaces

Sylvie Masse, Guillaume Laurent, T. Coradin

► **To cite this version:**

Sylvie Masse, Guillaume Laurent, T. Coradin. Solid state NMR and diatoms: probing the interfaces. Biom mineralization short course, Dec 2014, Paris, France. . hal-01138970

HAL Id: hal-01138970

<https://hal.sorbonne-universite.fr/hal-01138970>

Submitted on 3 Apr 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

INTRODUCTION




Partnership :
V. MARTIN-JEZEQUEL, B. TESSON
Fac. de Sciences et Techniques
ISOMER, NANTES



Gyromagnetic ratio $ \gamma $		$\frac{\gamma}{\text{NA}}$		Relative sensitivity \searrow	
I	γ (10 ⁶)	γ (10 ⁶)	γ (10 ⁶)	γ (10 ⁶)	γ (10 ⁶)
¹ H	100%	100%	100%	100%	100%
³¹ P	100%	100%	100%	100%	100%
¹³ C	11%	11%	11%	11%	11%
²⁹ Si	4.7%	4.7%	4.7%	4.7%	4.7%
¹⁵ N	0.36%	0.36%	0.36%	0.36%	0.36%






CHEMICAL TREATMENT



Whole-cell, SDS-treated and H₂O₂-treated samples were isotopically enriched with ²⁹Si, ¹³C/²⁹Si/¹⁵N and ¹³C/¹⁵N, respectively. While SDS is used to clean the frustule, H₂O₂ treatment seems to be much more aggressive, probably leading to partial dissolution-recrystallization.






SDS-TREATED SAMPLE

MOBILE SPECIES



Liquid-state-inspired NMR experiments mainly highlight the most mobile species: unsaturated lipids. Nevertheless EXSY exp. shows spin diffusion between two broad protons regions.

RIGID SPECIES



Solid-state NMR experiments allow to probe another part of the sample: the rigid one. Playing with the selected nuclei, spatial proximities can be assessed.

CONCLUSION AND REFERENCES

While a lot of work is still needed to fully understand diatoms frustule interface, solid-state NMR appears to be a powerful toolbox. Indeed, this technique is able to selectively probe either mobile or rigid species at a very local scale. Varying methods, species proximity can be checked and connectivity evaluated.

- Tesson, B., Masse, S., Laurent, G., Maquet, J., Livage, J., Martin-Jézéquel, V. and Coradin, T. Contribution of multi-nuclear solid state NMR to the characterization of the Thalassiosira pseudonana diatom cell wall. *Anal Bioanal Chem* 390, 1889–1898 (2008).
- Hatté, C., Hodgins, G., Jull, A. J. T., Bishop, B. & Tesson, B. Marine chronology based on ¹⁴C dating on diatoms proteins. *Marine Chemistry* 109, 143–151 (2008).
- Baccile, N., Laurent, G., Bonhomme, C., Innocenzi, P. & Babonneau, F. Solid-State NMR Characterization of the Surfactant–Silica Interface in Templated Silicas: Acidic versus Basic Conditions. *Chem. Mater.* 19, 1343–1354 (2007).