



HAL
open science

Involvement of BAG3 and HSPB7 loci in various etiologies of systolic heart failure: Results of a European collaboration assembling more than 2000 patients

Sophie Garnier, Christian Hengstenberg, Nicolas Lamblin, Olivier Dubourg, Pascal de Groote, Laurent Fauchier, Jean-Noël Trochu, Eloisa Arbustini, Ulrike Esslinger, Paul Barton, et al.

► **To cite this version:**

Sophie Garnier, Christian Hengstenberg, Nicolas Lamblin, Olivier Dubourg, Pascal de Groote, et al.. Involvement of BAG3 and HSPB7 loci in various etiologies of systolic heart failure: Results of a European collaboration assembling more than 2000 patients. *International Journal of Cardiology*, 2015, 189, pp.105-107. 10.1016/j.ijcard.2015.04.003 . hal-01143379

HAL Id: hal-01143379

<https://hal.sorbonne-universite.fr/hal-01143379>

Submitted on 17 Apr 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Involvement of *BAG3* and *HSPB7* loci in various etiologies of systolic heart failure:**Results of a European collaboration assembling more than 2,000 patients**

Sophie Garnier, PhD^{1,2,3}; Christian Hengstenberg, MD-PhD⁴; Nicolas Lamblin, MD-PhD⁵; Olivier Dubourg, MD-PhD^{6,7}; Pascal De Groote, MD-PhD⁵; Laurent Fauchier, MD⁸; Jean-Noël Trochu, MD-PhD⁹; Eloisa Arbustini, MD-PhD¹⁰; Ulrike Esslinger, PhD^{1,3,4}; Paul Barton, MD-PhD¹¹; Benjamin Meder, MD¹²; Hugo Katus, MD-PhD¹²; Karen Frese, PhD¹²; Michel Komajda, MD-PhD^{1,3,13}; Stuart SA. Cook, PD-PhD^{14,15,16}; Richard Isnard, MD-PhD^{1,13}; Laurence Tiret, PhD^{1,2,3}; Eric Villard, PhD^{1,2,3,13}; Philippe Charron, MD-PhD^{1,3,6,7}

¹INSERM, UMR_S 1166, F-75013, Paris, France; ²Sorbonne Universités, UPMC Univ Paris 06, UMR_S 1166, Team Genomics & Pathophysiology of Cardiovascular Diseases, F-75013, Paris, France; ³ICAN Institute for Cardiometabolism and Nutrition, F-75013, Paris, France; ⁴University of Regensburg, Department of Internal Medicine II, Regensburg, Germany; ⁵Hôpital Régional Universitaire de Lille - Hôpital Cardiologique, Lille, France; ⁶Hôpital Ambroise Paré AP-HP, Paris, France; ⁷Université de Versailles Saint Quentin en Yvelines, Versailles, France; ⁸Hôpital Régional Universitaire de Tours, Tours, France; ⁹INSERM UMR 1087, Nantes, France; ¹⁰IRCCS Foundation Policlinico San Matteo, Center for Genetic Cardiovascular Diseases, Pavia, Italy; ¹¹Royal Brompton & Harefield NHS Foundation Trust's NIHR Cardiovascular biomedical research unit, London, United Kingdom; ¹²University Hospital Heidelberg, Heidelberg, Germany; ¹³Université Paris 6 & AP-HP, Hôpital de la Pitié-Salpêtrière, Paris, France; ¹⁴National Heart and Lung Institute, Imperial College London, London, United Kingdom; ¹⁵National Heart Centre Singapore, Singapore; ¹⁶Duke-NUS, Singapore

Corresponding author: Dr. Sophie Garnier

INSERM UMR S 1166 - Team Genomics & Pathophysiology of Cardiovascular Diseases

Faculté de médecine Pierre et Marie Curie 6ème étage,

91 Boulevard De l'Hôpital, 75634 Paris Cedex 13, France

Tel: +33 1 40 77 81 40

Fax: +33 1 40 77 96 25

e-mail: sophie.garnier@upmc.fr

Word count: 995

Keywords: Genetics, Association studies, Ischemic heart failure, Dilated cardiomyopathy

ACCEPTED MANUSCRIPT

Heart failure (HF), a major public health burden affecting 2% of industrialized populations, is a syndrome resulting from structural or functional myocardial impairment leading to inadequate cardiac output to meet the body's metabolic demands (1). Half of HF patients present systolic dysfunction (systolic-HF), also called reduced ejection fraction (HF-REF), a disease related to various causes including idiopathic Dilated Cardiomyopathy (DCM) and Coronary Artery Diseases (CAD) (ischemic-HF). HF is usually a multifactorial disease but the genetic variants contributing to its susceptibility or its severity may be different according to its underlying causes (2) and their identification is only beginning.

A previous genome wide association study (GWAS) (3) identified two single nucleotide polymorphisms (SNPs) associated with DCM: rs2234962 within *BAG3* gene (B-cell lymphoma 2-associated athanogene 3) and rs10927875 within a locus encompassing five genes including *HSPB7* (heat shock 27 kDa protein family, member 7) (further called *HSPB7* locus). Other studies also reported associations between polymorphisms in the *HSPB7* locus and HF (4-6), two of them assessing the implication of the locus in the subgroup of ischemic-HF patients (4, 5), but none addressed the role of *BAG3* in ischemic-HF nor the potential impact of *HSPB7* and *BAG3* loci on HF severity. We hypothesized that these two loci could be involved in susceptibility to syst-HF due to CAD (ischemic-HF) and/or in severity of syst-HF due to CAD or DCM.

All participants were of European origin, had a diagnosis of systolic-HF (Left Ventricle Ejection Fraction, LVEF, < 45%) and gave written informed consent. The study complies with the principles of the Declaration of Helsinki and was approved by local ethics committees. In total, 1160 ischemic-HF patients and 1612 controls and 1141 DCM patients were studied. Genotyping of rs2234962-*BAG3* and rs10927875-*HSPB7* was performed with TaqMan technology. Disease severity was assessed in patients with Ischemic-HF or DCM as

following: LVEF, Left Ventricle End Diastolic Diameter (LVEDD), NYHA dyspnea at inclusion and Age at diagnosis.

We analyzed the data following this summarized methodology. After quality controls and Hardy-Weinberg equilibrium testing in controls ($p < 0.05$), the association between Ischemic-HF and individual genotypes was tested using a logistic regression model adjusting on age, gender, and study population in the R statistical environment (7). We assumed an additive mode of inheritance and decided to adjust the p-value threshold following Bonferroni correction for two independent tests (p-value < 0.025 considered significant). Severity assessment in patients was performed via linear regression analyses adjusted for age, sex, and origin depending on the SNP genotype exposure.

Considering the allelic odds ratios for rs10927875-*HSPB7* and rs2234962-*BAG3* associations with DCM (0.71 and 0.54 respectively) (3) and the p-value threshold of 0.025, our study had a 95% and 99% power to detect similar associations.

We observed that rs10927875- *HSPB7* is significantly associated with Ischemic-HF (Table 1), with a MAF lower in patients than in controls (0.29 vs. 0.33, $p = 0.0017$, odds ratio 0.96 [0.94-0.98]), which is consistent with a protective effect of the T minor variant against Ischemic-HF. No significant association was detected between rs2234962-*BAG3* and Ischemic-HF (Table 1), even though a trend was detected.

Regarding severity, rs2234962-*BAG3* was significantly associated with LVEDD both in DCM (adjusted-p: 0.0056, Table 2, N=1076 patients) and ischemic-HF patients (adjusted-p: 0.01, N=359 patients). In the ischemic-HF population, the major allele T of rs2234962-*BAG3* was associated with enlarged LVEDD (TT: 64.0 ± 8.7 mm; TC: 62.7 ± 8.0 mm and CC: 61.8 ± 9.8 mm) and thus with a greater severity of the disease. Results are less clear in DCM

patients although the same trend could be observed between major homozygous and heterozygous alleles (TT: 70.9 ± 9.6 mm and TC: 69.1 ± 9.2 mm respectively).

No other association could be revealed between the two SNPs and any of the other severity criteria.

The protective effect of the T minor allele of rs10927875-*HSPB7* against ischemic-HF is consistent with previous findings in DCM patients (3). Matkovich and Cappola also described SNPs in *HSPB7* locus, rs1739840 and rs1739843 respectively, which exhibited a lower MAF in Ischemic-HF cases than in controls (0.4126 vs. 0.4735 and 0.367 vs 0.436 respectively) (5, 6).

Finding an association between ischemic-HF and *HSPB7* locus but not with *BAG3* locus suggests that the genetic background of multifactorial systolic-HF is different according to its underlying causes. This is meaningful since the pathophysiological pathways of these two diseases are different, DCM resulting from a primary myocardial defect whereas ischemic-HF is due to ischemia and necrosis. Since *HSPB7* locus appears involved in both causes, this may reflect a pivotal role of gene(s) in the locus as a poorly adaptive process to various triggers. The restricted association of *BAG3* locus with DCM suggests a more specific role in the context of a primary myocardial defect.

rs10927875-*HSPB7* locates in an intron of *ZBTB17* on chromosome 1p36, but the locus encompasses several other genes—*SPEN*, *HSPB7*, *CLCNKA* and *CLCNKB*—exhibiting strong linkage disequilibrium. *HSPB7* is considered as a strong candidate because it exhibits cardiac-specific expression and some of its variants have been independently associated with advanced HF, systolic-HF (8) or with DCM (4-6). A recent study found that *HSPB7* is early expressed in embryonic heart and have a major role in heart embryogenesis (9).

BAG3 is mainly expressed in striated muscle and is involved in anti-apoptosis and anti-proteotoxicity pathways (10). rs2234962-*BAG3*, a non-synonymous SNP (c.T757C, p.C151R) predicted as probably damaging by Polyphen, is associated in our study with LVEDD in both ischemic-HF and CMD populations. The number of T alleles appears related with enlarged LVEDD in ischemic-HF patients but this was less clear for DCM patients, possibly due to the very low T homozygous DCM patients (9 out of the 1141). A less efficient *BAG3* activity may favour HF severity but the hypothesis is speculative.

Out of the two loci previously associated with DCM, we observed that *HSPB7* locus was also associated with ischemic-HF whereas *BAG3* locus was not. Conversely, *BAG3* locus was associated with LVEDD in both DCM and syst-HF patients. Our results suggest investigating further the role of these two loci in the pathophysiology of HF of various causes.

ACKNOWLEDGMENTS

We thank all the families and physicians for their participation to this study. We also thank all the co-workers who participated to the recruitment of patients, particularly: Dr Aumont M-C (Bichat Claude Bernard Hospital, Paris, France), Dr Gibelin P (Hospital regional University of Nice, Nice, France), Dr Aupetit J-F (Joseph-St Luc Hospital, Lyon, France), Dr. Rachel Buchan (Royal Brompton & Harefield NHS Foundation Trust's NIHR Cardiovascular biomedical research unit, London, United Kingdom) and Dr Leanne Felkin (Imperial College London, Heart Science Centre, London, United Kingdom).

FUNDING

This work was supported by grants from Assistance Publique-Hôpitaux de Paris (PHRC programme hospitalier de recherche Clinique; AOM04041), the 'Fondation LEDUCQ' (11CVD 01 network; Eurogene Heart Failure network), European Union FP7 (INHERITANCE network grant n°241924 and BestAgeing network grant n°306031), Agence Nationale pour la Recherche/Bundesministerium für Bildung und Forschung (INSIGHT-DCM network), the Société Française de Cardiologie/Fédération Française de Cardiologie (2006), the CONNY-MAEVA charitable foundation, The British Heart Foundation, The NMRC Singapore, NIHR cardiovascular BRU and BRC, The MRC UK. The UK Blood Services collection of Common Controls (UKBS collection), part of the Wellcome Trust Case-Control Consortium, was funded by the Wellcome Trust (076113/C104/Z), and the Juvenile Diabetes Research Foundation (WT061858).

CONFLICT OF INTEREST

The authors report no relationships that could be construed as a conflict of interest

ACCEPTED MANUSCRIPT

REFERENCES

1. McMurray JJV, Adamopoulos S, Anker SD, Auricchio A, Böhm M, Dickstein K, Auricchio A, Böhm M, Dickstein K, Falk V, Filippatos G, Fonseca C, Gomez-Sanchez MA, Jaarsma T, Køber L, Lip GY, Maggioni AP, Parkhomenko A, Pieske BM, Popescu BA, Rønnevik PK, Rutten FH, Schwitter J, Seferovic P, Stepinska J, Trindade PT, Voors AA, Zannad F, Zeiher A; ESC Committee for Practice Guidelines. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012. *Eur Heart J*. 2012;33:1787-1847.
2. Friedrichs F, Zugck C, Rauch GJ, Ivandic B, Weichenhan D, Müller-Bardorff M, Meder B, El Mokhtari NE, Regitz-Zagrosek V, Hetzer R, Schäfer A, Schreiber S, Chen J, Neuhaus I, Ji R, Siemers NO, Frey N, Rottbauer W, Katus HA, Stoll M. HBEGF, SRA1, and IK: Three cosegregating genes as determinants of cardiomyopathy. *Genome Res*. 2009;19:395-403.
3. Villard E, Perret C, Gary F, Proust C, Dilanian G, Hengstenberg C, Ruppert V, Arbustini E, Wichter T, Germain M, Dubourg O, Tavazzi L, Aumont MC, DeGroot P, Fauchier L, Trochu JN, Gibelin P, Aupetit JF, Stark K, Erdmann J, Hetzer R, Roberts AM, Barton PJ, Regitz-Zagrosek V; Cardiogenics Consortium, Aslam U, Duboscq-Bidot L, Meyborg M, Maisch B, Madeira H, Waldenström A, Galve E, Cleland JG, Dorent R, Roizes G, Zeller T, Blankenberg S, Goodall AH, Cook S, Tregouet DA, Tiret L, Isnard R, Komajda M, Charron P, Cambien F. A genome-wide association study identifies two loci associated with heart failure due to dilated cardiomyopathy. *European Heart Journal*. 2011;32:1065–1076.
4. Matkovich SJ, Van Booven DJ, Hindes A, Kang MY, Druley TE, Vallania FLM, Mitra RD, Reilly MP, Cappola TP, Dorn GW 2nd. Cardiac signaling genes exhibit unexpected

- sequence diversity in sporadic cardiomyopathy, revealing HSPB7 polymorphisms associated with disease. *J Clin Invest*. 2010;120:280-289.
5. Cappola TP, Li M, He J, Ky B, Gilmore J, Qu L, Keating B, Reilly M, Kim CE, Glessner J, Frackelton E, Hakonarson H, Syed F, Hindes A, Matkovich SJ, Cresci S, Dorn GW 2nd. Common Variants in HSPB7 and FRMD4B Associated With Advanced Heart Failure. *Circ Cardiovasc Genet*. 2010;3:147-154.
 6. Stark K, Esslinger UB, Reinhard W, Petrov G, Winkler T, Komajda M, Isnard R, Charron P, Villard E, Cambien F, Tiret L, Aumont MC, Dubourg O, Trochu JN, Fauchier L, Degroote P, Richter A, Maisch B, Wichter T, Zollbrecht C, Grassl M, Schunkert H, Linsel-Nitschke P, Erdmann J, Baumert J, Illig T, Klopp N, Wichmann HE, Meisinger C, Koenig W, Lichtner P, Meitinger T, Schillert A, König IR, Hetzer R, Heid IM, Regitz-Zagrosek V, Hengstenberg C. Genetic association study identifies HSPB7 as a risk gene for idiopathic dilated cardiomyopathy. *PLoS Genet*. 2010;6:e1001167.
 7. R Development Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing; 2008 (<http://www.R-project.org>).
 8. Zeller T, Wild P, Szymczak S, Rotival M, Schillert A, Castagne R, Maouche S, Germain M, Lackner K, Rossmann H, Eleftheriadis M, Sinning CR, Schnabel RB, Lubos E, Mennerich D, Rust W, Perret C, Proust C, Nicaud V, Loscalzo J, Hübner N, Tregouet D, Münzel T, Ziegler A, Tiret L, Blankenberg S, Cambien F. Genetics and beyond—the transcriptome of human monocytes and disease susceptibility. *PLoS ONE*. 2010;5:e10693.
 9. Rosenfeld GE, Mercer EJ, Mason CE, Evans T. Small heat shock proteins Hspb7 and Hspb12 regulate early steps of cardiomorphogenesis. *Developmental Biology*. 2013;381:389-400.

10. Gamerding M, Carra S, Behl J C. Emerging roles of molecular chaperones and co-chaperones in selective autophagy: focus on BAG proteins. *Mol Med.* 2011; 89:1175–1182.

ACCEPTED MANUSCRIPT

Table 1: Genotypic repartition of cases (ischemic-HF), controls and allelic comparison test assuming an additive allele effect

SNP (genotypes)	Population	Patients Genotype	MAF	Controls Genotypes	MAF	p-adj*	OR [CI 95%]
rs10927875- <i>HSPB7</i> (TT/TC/CC)	Germany	18/133/147	0.28	32/127/115	0.35	0.06	0.95 [0.90-1.00]
	France	28/153/173	0.29	84/310/312	0.34	0.17	0.97 [0.94-1.01]
	UK	33/221/235	0.29	47/271/256	0.32	0.30	0.98 [0.96-1.01]
	All	79/507/555	0.29	163/708/683	0.33	1.7E ⁻³	0.96 [0.94-0.98]
rs2234962- <i>BAG3</i> (CC/TC/TT)	Germany	12/92/192	0.20	12/90/171	0.21	0.75	0.99 [0.93-1.05]
	France	13/103/241	0.18	32/232/433	0.21	0.13	0.96 [0.92-1.00]
	UK	13/151/323	0.18	18/199/346	0.21	0.43	0.99[0.95-1.02]
	All	38/346/756	0.18	62/521/960	0.21	0.07	0.97 [0.94-1.00]

This table gives the genotypic repartition for the three SNPs in the corresponding population.

*The deducted Minor Allele Frequency (MAF) is also given. *p-adj, p-value adjusted for age, gender and population. OR calculation was done on the row genotypes data thus no adjustment was performed.*

Table 2: Association with severity sub-phenotypes for French, German and Italian DCM patients.

Population		France (N=749)	Germany (N=278)	Italy (N=114)	Total (N=1141)
SNP	Severity criteria	p-adj*	p-adj*	p-adj*	p-adj*
rs10927875- HSPB7	LVEF (N=1004)	0.14	0.24	0.05	0.87
	LVEDD (1076)	0.47	0.82	0.79	0.72
	NYHA (738)	0.79	0.02	0.35	0.50
	Age at diagnosis (1141)	0.66	0.07	0.95	0.2
rs2234962- BAG3	LVEF (1004)	0.13	0.40	0.10	0.21
	LVEDD (1076)	4.1E ⁻³	0.37	0.78	5.6E⁻³
	NYHA (738)	0.59	0.09	0.02	0.34
	Age at diagnosis (1141)	0.57	0.74	0.27	0.99

*The phenotypic information was not always available for the 1141 DCM cases. The exact number of tested individuals is given for each sub-phenotype. *p-adj, p-value of the test adjusted on age, gender and population.*