Hydrogen peroxide produced by NADPH oxidases increases proline accumulation during salt or mannitol stress in *Arabidopsis thaliana*

Kilani Ben Rejeb, Delphine Lefebvre-de Vos, Isabel Le Disquet, Anne-Sophie Leprince, Marianne Bordenave, Régis Maldiney, Asma Jdey, Chedly Abdelly, Arnould Savouré

► To cite this version:

Kilani Ben Rejeb, Delphine Lefebvre-de Vos, Isabel Le Disquet, Anne-Sophie Leprince, Marianne Bordenave, et al.. Hydrogen peroxide produced by NADPH oxidases increases proline accumulation during salt or mannitol stress in *Arabidopsis thaliana*. *New Phytologist*, 2015, 208 (4), pp.1138-1148. 10.1111/nph.13550 . hal-01187624

HAL Id: hal-01187624

<https://hal.sorbonne-universite.fr/hal-01187624>

Submitted on 27 Aug 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Hydrogen peroxide produced by NADPH oxidases increases proline accumulation during salt or mannitol stress in *Arabidopsis thaliana*

Kilani Ben Rejeb^{1,2}, Delphine Lefebvre-De Vos¹, Isabel Le Disquet¹, Anne-Sophie Leprince¹, Marianne Bordenave¹, Régis Maldiney¹, Asma Jdey², Chedly Abdelly², Arnould Savouré¹

¹Sorbonne Universités, UPMC Univ Paris 06, Adaptation des Plantes aux Contraintes Environnementales, Case 156, 4 place Jussieu, F-75005, Paris, France ; ²Laboratoire des Plantes Extrêmophiles, Centre de Biotechnologie de Borj-Cedria (CBBC), BP 901, Hammam-Lif, 2050, Tunisia.

*Corresponding author:

Arnould Savouré, email: arnould.savoure@upmc.fr

Phone: +33 (0)1 44 27 26 72, Fax: +33 (0)1 44 27 61 51

Total word count (excluding summary, references and legends):	5307	No. of figures:	8 (Fig 2 in colour)
Summary:	198	No. of Tables:	0
Introduction:	1069	No of Supporting Information files:	5 (Fig. S1-S4; Table S1)
Materials and Methods:	1239		.
Results:	1427		.
Discussion:	1468		
Acknowledgements:	103		

Summary

- Many plants accumulate proline, a compatible osmolyte, in response to various environmental stresses such as water deficit and salinity. In some stress responses, plants generate hydrogen peroxide (H_2O_2) that mediates numerous physiological and biochemical processes. The aim was to study the relationship between stress-induced proline accumulation and H_2O_2 production.

- Using pharmacological and reverse genetic approaches, the role of NADPH oxidases (Rboh) in the induction of proline accumulation was investigated in *Arabidopsis thaliana* in response to stress induced by either 200 mM NaCl or 400 mM mannitol.

- Stress from NaCl or mannitol resulted in a transient increase in H_2O_2 content accompanied by accumulation of proline. Dimethylthiourea, a scavenger of H_2O_2 , and diphenylene iodonium (DPI), an inhibitor of H_2O_2 production by NADPH oxidase, were found to significantly inhibit proline accumulation in these stress conditions. DPI also reduced the expression level of Δ^1 -pyrroline-5-carboxylate synthetase, the key enzyme involved in the biosynthesis of proline. Similarly less proline accumulated in KO-mutants lacking either *AtRbohD* or *AtRbohF* than in wild type in response to the same stresses.

- Our data demonstrate that AtRbohS contribute to H_2O_2 production in response to NaCl or mannitol stress to increase proline accumulation in *A. thaliana*.

Key-words: Abiotic stresses; *Arabidopsis thaliana*; cell signalling; hydrogen peroxide; NADPH-oxidases (Rboh); proline metabolism.

41 Introduction

42 In their natural environments, plants commonly encounter a variety of abiotic
43 constraints like drought and salinity (Nakashima *et al.*, 2009). To overcome these constraints,
44 plants have developed a variety of adaptive mechanisms that allow them to perceive external
45 signals and to optimize adaptive responses. One of these mechanisms is osmotic adjustment
46 through the accumulation of large quantities of osmolytes as it allows plants to avoid water
47 deficit stress by maintaining water uptake. Glycine betaine, polyols, sugars and free amino
48 acids are examples of such osmolytes (Chen & Jiang, 2010; Slama *et al.*, 2015).

49 Proline is the most common free amino acid to accumulate in plants subjected to water
50 deficit stress. Proline has multifunctional roles though which do not necessarily relate to the
51 osmotic balance (for review see Szabados & Savouré, 2010). Proline may stabilize protein
52 complexes, scavenge free radicals and be a source of carbon and nitrogen for growth after
53 stress relief. Proline biosynthesis and degradation are involved in regulating intracellular
54 redox potential and storage as well as the transfer of energy and reducing power (Sharma *et al.*,
55 2011; Szabados & Savouré, 2010). The beneficial effect of proline on plant growth after
56 stress is likely to be the result of changes in proline metabolism rather than the accumulation
57 of the amino acid itself (Sharma *et al.*, 2011; Szabados & Savouré, 2010).

58 The proline content of plant cells depends on tight regulation of its proline
59 biosynthesis and catabolism. Housekeeping levels of proline biosynthesis occur in the cytosol,
60 but stress-induced biosynthesis is thought to be localized in chloroplasts (Székely *et al.*,
61 2008). When under water-deficit stress, proline is mainly synthesized from glutamate. The
62 bifunctional pyrroline-5-carboxylate synthetase (P5CS) reduces glutamate to glutamyl-5-
63 semialdehyde, which is spontaneously converted to pyrroline-5-carboxylate (P5C). P5C is
64 then reduced to proline by P5C reductase (P5CR). Degradation of proline takes place in
65 mitochondria via the sequential action of proline dehydrogenase (ProDH) and P5C
66 dehydrogenase. The rate-limiting steps in proline biosynthesis and degradation are catalyzed
67 by P5CS and ProDH, respectively. Two closely related P5CS-encoding genes were identified
68 in *Arabidopsis thaliana*. *P5CS1* is induced by drought and salt stress (Savouré *et al.*, 1995;
69 Yoshiba *et al.*, 1995), while *P5CS2* is expressed in dividing cells in cell suspension cultures
70 (Strizhov *et al.*, 1997), in meristematic and reproductive tissues (Székely *et al.*, 2008), and in
71 response to biotic stress such as incompatible plant-pathogen interactions (Fabro *et al.*, 2004).
72 Similarly, *A. thaliana* has two genes for ProDH, *ProDH1* and *ProDH2* (Kiyosue *et al.*, 1996;
73 Verbruggen *et al.*, 1996; Funck *et al.*, 2010). *ProDH1* is thought to encode the main isoform

involved in proline degradation (for review see Servet *et al.*, 2012). The expression of *ProDH1* is down-regulated by osmotic stress and upregulated by proline (Kiyosue *et al.*, 1996; Verbruggen *et al.*, 1996).

The regulation of *P5CS* and *ProDH* expression has been studied to identify components of signalling pathways that control proline accumulation. Under non-stress conditions, phospholipase D functions as a negative regulator of proline biosynthesis in *Arabidopsis* (Thiery *et al.*, 2004), whereas calcium signalling and phospholipase C (PLC) trigger *P5CS1* transcription and proline accumulation during salt stress (Parre *et al.*, 2007). *P5CS1* expression has also been found to be stimulated by light (Hayashi *et al.*, 2000) and nitric oxide (Zhao *et al.*, 2009) and it is abscisic acid (ABA) independent under water-deficit stress (Savouré *et al.*, 1997; Sharma & Verslues, 2010). Recently Leprince *et al.* (2015) demonstrated that phosphatidylinositol 3-kinase is involved in the regulation of proline catabolism through transcriptional regulation of *ProDH1*.

Another common plant response to all types of environmental constraints is the accumulation of ROS, which are toxic at high concentrations but at lower concentrations may act as signal molecules in the control of various cellular processes. H₂O₂ is a ROS produced by plant plasma membrane-localized NADPH oxidases, known as respiratory burst oxidase homologues (Rbohs). Rbohs reduce molecular oxygen to superoxide by oxidising NADPH via FAD and two hemes. The superoxide primary product is then converted into H₂O₂ by superoxide dismutase (Sagi & Fluhr, 2001; Sagi & Fluhr, 2006). The *Arabidopsis* genome contains 10 NADPH oxidase-encoding genes, designated *AtRbohA* to *J*, that exhibit different patterns of expression throughout plant development and in response to environmental factors (Fluhr, 2009; Marino *et al.*, 2012). For instance, *AtRbohA*, *B* and *C* are only expressed in roots, especially in the elongation zone. *AtRbohC* was specifically identified as playing a role in root hair development (Foreman *et al.*, 2003). *AtRbohH* and *J* are reported to be expressed only in pollen. Both *AtRbohD* and *AtRbohF* are expressed in all plant organs and are the main isoforms involved in pathogen defence responses (Torres *et al.*, 2002), ABA-induced stomatal closure (Kwak *et al.*, 2003), jasmonic acid signalling regulated by transcription factor MYC2 (Maruta *et al.*, 2011) and ROS-dependent regulation of Na⁺/K⁺ homeostasis under salt stress (Ma *et al.*, 2012). *AtRbohD* has also been demonstrated to mediate rapid systemic signalling triggered by multiple abiotic stresses (Miller *et al.*, 2009) and to be required for salt acclimation signalling mediated by heme oxygenase in *Arabidopsis* (Xie *et al.*, 2011). It was reported that mild salt stress causes a rapid and transient accumulation of ROS in *A. thaliana*

(peak I after 1 h) followed by a second oxidative burst (peak II after 6 h) (Xie *et al.*, 2011). The conclusion is that HY1 heme oxygenase plays an important role in salt acclimation signalling and requires the participation of AtRbohD-derived ROS from peak II. More recently, it has been reported that AtRbohF fulfils a crucial role in protecting shoot cells from transpiration-dependent accumulation of excess Na⁺ (Jiang *et al.*, 2012). Rbohs are thus key regulators of ROS production with pleiotropic functions in plants.

It is possible that there is a link between quantitative changes in ROS and proline. Exogenous H₂O₂ treatment led to a significant accumulation of proline in coleoptiles and radicles of maize seedlings due to the induction of biosynthetic P5CS enzyme activity and a decrease in catabolic ProDH enzyme activity (Yang *et al.*, 2009). Fabro *et al.* (2004) have also demonstrated that ROS can mediate the activation of *AtP5CS2* and proline accumulation during biotic stress. However, reports on the relationship between endogenous ROS increase and proline accumulation under osmotic stress are still limited and the link between NADPH oxidases and proline accumulation has never been addressed. In the present study, the role of ROS-generating NADPH-oxidase enzymes in proline accumulation in response to salt or mannitol stress was investigated in *Arabidopsis thaliana* seedlings.

Materials and Methods

Growth conditions and stress treatments

Arabidopsis thaliana (L.) Heynh ecotype Columbia-0 (Col-0) was used as the wild-type in this study. Homozygous *Arabidopsis thaliana* transposon insertion mutant lines *atrboh3* (European Arabidopsis Stock Centre code N9555) and *atrboh3f* (European Arabidopsis Stock Centre code N9557) and double mutant *atrboh3f* (European Arabidopsis Stock Centre code N9558) (Torres *et al.*, 2002) were ordered from the European Arabidopsis Stock Centre. Homozygous T-DNA insertion lines SALK_070610 (*atrboh3*; seventh exon insertion) and SALK_059888 (*atrboh3f*; third intron insertion) were ordered from the Salk collection (Pogany *et al.*, 2009) and used as controls for the transposon insertion lines.

Surface-sterilized seeds of wild-type (Col-0) and *Arabidopsis* mutant plants were sown onto grids placed on 0.5 × Murashige and Skoog (MS) solid agar medium in Petri dishes according to Parre *et al.* (2007). After 24 h at 4°C to break dormancy, seedlings were allowed to grow at 22°C under continuous light (90 μmol photons m⁻² s⁻¹). Twelve-day-old *Arabidopsis* seedlings were exposed to H₂O₂ (5 to 120 mM), paraquat (PQ, also called methyl viologen, 5 to 120 μM), 200 mM NaCl or 400 mM mannitol for 24 h. To alter ROS levels, the seedlings were

pre-incubated for 4 h without or with H₂O₂ scavenger dimethylthiourea (DMTU, 5 to 40 mM) or flavoenzyme inhibitor diphenylene iodonium (DPI, 5 to 40 µM) in 0.5 × liquid MS medium and then exposed to either NaCl or mannitol for 24 h.

Determination of ion content

Whole plants were harvested, rinsed twice in pure water and quickly blotted. Samples were dried at 60 °C until they reached constant weight then ground. Ions were extracted from samples in 0.5% HNO₃. Na⁺ and K⁺ were assayed by flame emission photometry (Corning, UK).

Proline content measurements

Free proline content was measured according to the Bates method (Bates *et al.*, 1973). Powdered plant samples (50 to 100 mg) were homogenized in 1.5 ml of 3% sulfosalicylic acid and centrifuged at 14,000 × *g* for 10 min at 4 °C. To 1 ml of this extract, 1 ml of acid-ninhydrin and 1ml of glacial acetic acid were added. The reaction mixture was incubated at 100°C for 1 h then placed on ice to stop the reaction. In the presence of proline, the reaction produces a red chromogen. The chromogen was extracted with 2 ml of toluene and the absorbance of the resulting upper phase was read at 520 nm. Proline content of samples was calculated by referring to a standard curve drawn from absorbance readings from samples containing known concentrations of proline.

H₂O₂ content measurements

H₂O₂ content was determined in a horseradish peroxidase-based spectrophotometric assay following the protocol described by Oracz *et al.* (2009). Plant samples (300 mg FW) were ground in a mortar on ice in 1 ml of 0.2 M perchloric acid. After 15 min of centrifugation at 13,000 × *g* at 4°C, the resulting supernatant was neutralized to pH 7.5 with 4 M KOH and then centrifuged at 13,000 × *g* at 4°C. The concentration of H₂O₂ in the supernatant was immediately determined by adding peroxidase with substrates 3-dimethylaminobenzoic acid and 3-methyl-2-benzothiazolidone hydrazone. The increase in absorbance at 590 nm was monitored for 15 min after the addition of peroxidase at 25 °C and compared to a calibration curve obtained with known amounts of fresh H₂O₂.

Histochemical detection of H₂O₂

The production of H_2O_2 was visualized *in vivo* by 2',7'-dichlorofluorescein diacetate (H_2DCFDA) or DAB staining methods. *Arabidopsis* roots were collected after 6-h treatments and immersed in 25 μM H_2DCFDA for 15 min in the dark and then washed with 20 mM potassium phosphate buffer pH 6. Fluorescent signals were visualized using a Zeiss ApoTome microscope (excitation, 488 nm; emission, 525 nm). DAB staining was performed as described by Torres *et al.* (2002). Leaves from PQ-treated seedlings were detached and vacuum infiltrated with DAB solution (1 mg ml^{-1} DAB-HCl, pH 3.8). DAB forms a reddish-brown polymer in the presence of H_2O_2 formation. After staining, leaves were cleared in 96% boiling ethanol and observed using a binocular microscope. For both staining methods, digital images were obtained with an AxioCam camera and AxioVision software (Zeiss).

Cytochemical detection of H_2O_2

Cytochemical detection of H_2O_2 was carried out according to the method described by Bestwick *et al.* (1997). Leaf samples were collected from treated and control seedlings and incubated in freshly prepared 5 mM cerium chloride for 30 min. The leaves were then fixed in a solution containing 4% glutaraldehyde and 1.5% paraformaldehyde for 1 h, post-fixed for 45 min in 1% osmium tetroxide and then embedded in Eponaraldite resin after dehydration in an ethanol series. Sections (60 to 80 nm thick) were cut with a LKB 2128 ultramicrotome, mounted on uncoated copper grids, and stained with 5% uranyl acetate. Sections were observed using a Zeiss912 Omega transmission electron microscope. Digital images were obtained using a Veleta Camera (2kx2k, Olympus) and iTem software (Zeiss).

RT-PCR and qRT-PCR Analysis

Total RNA was extracted from 100 mg of plant tissue ground in liquid nitrogen using a mixer mill (MM301, Retsch, Germany). The powder was suspended in 0.5 mL extraction buffer (0.2 M Tris-HCl, 0.5% (v/v) SDS, 0.25 M NaCl, 25 mM EDTA) and mixed with 0.5 mL of phenol/chloroform/isoamylalcohol (25:24:1, v/v/v). The aqueous phase was extracted 3 times with phenol/chloroform/isoamylalcohol. Total RNA was then precipitated overnight on ice with 2 M LiCl. After centrifugation (15 min, 10,000 $\times g$), the pellet was resuspended in 2 M LiCl and left to precipitate for 6 h on ice. Traces of DNA were removed by DNase treatment. RNA quantified by measuring the absorbance at 260 nm using a Nanovue® spectrophotometer (GE Healthcare Life Science). First-strand cDNA was obtained from 1.5

μg of total RNA using RevertAid™ reverse transcriptase synthesis kit (Fermentas) and oligo(dT)₂₃ as primer.

For RT-PCR, cDNAs were amplified using Taq polymerase and gene-specific primers (Supporting Information Table S1). *APT1* (adenine phosphoribosyltransferase 1; At1g27450) gene transcripts were amplified as a control. Amplified PCR fragments were visualized using ethidium bromide stained 2% (w/v) agarose gels. RT-PCR signals were quantified using the ImageJ 1.48 software (National Institutes of Health, Bethesda, MD).

For quantitative PCR 5 μL of diluted cDNA was used with 10 μL of Maxima SYBR Green/ROX qPCR Master Mix (Fermentas, France) and gene-specific primers (Supporting Information Table S1) in a Mastercycler® ep *realplex* (Eppendorf, France). Critical thresholds (Ct) were calculated using the Realplex 2.0 software (Eppendorf, France). For each gene, a standard curve made with dilutions of cDNA pools was used to calculate the reaction efficiencies, and relative expressions were calculated according to Hellemans *et al.* (2007) with *APT1* (At1g27450) and *AT5G* (At5g13440) as housekeeping genes. A mixture of cDNAs corresponding to each sample was used as reference. All qRT-PCR experiments were carried out with three biological replicates.

Western blots

Total proteins were subjected to SDS-PAGE using 8% acrylamide resolving gels and electroblotted onto a PVDF membrane. Rabbit polyclonal antibodies raised against P5CS or ProDH were used as primary antibodies (Thierry *et al.*, 2004; Parre *et al.*, 2007). Blots were incubated with secondary horseradish peroxidase-conjugated antibodies (GE Healthcare Life Sciences). The highly sensitive ECL Prime detection system (GE Healthcare) was used to quantify proteins using ImageJ 1.48 software (National Institutes of Health, Bethesda, MD) after densitometric scanning of autoradiography films. Alternatively the Storm 840 FluorImager (Molecular Dynamics) was used to visualise and quantify proteins on blots.

Statistical analysis

A one-way analysis of variance (ANOVA) at $P < 0.05$ significance level was performed using the SPSS program for Windows. The Duncan post-hoc test was used to test significant differences between treatments.

Results

Kinetics of ROS and proline accumulation in wild-type *A. thaliana* seedlings

To investigate ROS and proline accumulation in response to stress induced by either salt or mannitol, 12-day-old *A. thaliana* seedlings were exposed to 200 mM NaCl or 400 mM mannitol for up to 24 h. In both cases proline started to accumulate after 6 h. A maximum 16-fold increase in proline content was reached after 18 h of NaCl or mannitol stress (Fig. 1). A transient increase in H₂O₂ content was also observed in stressed seedlings after 6 h. H₂O₂ content had returned to a basal level after 12 h.

Effect of H₂O₂ and paraquat on proline accumulation in *A. thaliana*

The kinetics of H₂O₂ and proline accumulation in response to either NaCl or mannitol stress showed that proline had already started to increase 6 h after stress was applied when the transient increase in H₂O₂ content was also observed (Fig. 1). To determine whether H₂O₂ could be involved in proline accumulation, we first investigated whether adding different concentrations of exogenous H₂O₂ would affect proline accumulation in *A. thaliana*. Results showed that proline accumulation is induced by exogenous H₂O₂ application with 10 mM H₂O₂ having the maximum effect (Fig. 2a).

Paraquat (PQ) is a herbicide which induces superoxide anions and subsequently H₂O₂ generation *in situ*. Like H₂O₂, PQ also stimulated proline accumulation in *Arabidopsis* seedlings in a dose dependent manner with 40 µM PQ having the maximum effect (Fig. 2b). Leaves treated with high concentrations of PQ such as 80 or 100 µM PQ started to bleach after 48 h of treatment due to the high amount of H₂O₂ generated (data not shown) even though proline contents were not higher than at 40 µM PQ. DAB staining revealing the presence of H₂O₂ in leaves demonstrates that exogenously applied PQ gives rise to H₂O₂ *in situ* in a dose dependent manner (Fig. 2c). Taken together, these results indicate a possible causal relationship between H₂O₂ and proline production. However, much less proline accumulated in the presence of 10 mM H₂O₂ (about 5 times less) or 40 µM PQ (about 8 times less) than with NaCl or mannitol stress (Fig. 2a, 2b and Fig.1).

Effect of DMTU, a chemical scavenger of H₂O₂, on proline accumulation in response to stress.

To investigate whether the induction of proline accumulation by salt or mannitol requires H₂O₂, we treated *A. thaliana* seedlings with DMTU, a chemical trap for H₂O₂. As shown in

Fig. 3a, addition of up to 40 mM DMTU inhibited proline accumulation in a dose-dependent manner in seedlings treated with NaCl and to a lesser extent in those treated with mannitol.

Subcellular localisation of H₂O₂ accumulation

Multiple sources of H₂O₂ production inside plant cells have been described (Petrov & Van Breusegem, 2012). To investigate the subcellular localization of H₂O₂ accumulation in leaves of *A. thaliana* seedlings, a cytochemical approach using cerium perhydroxide was taken (Bestwick *et al.*, 1997). No visible cerium perhydroxide deposits were observed at the subcellular level in *A. thaliana* leaves growing under normal growth conditions, indicating the absence of H₂O₂ using this detection method (Fig. 4). However, both salt and mannitol triggered H₂O₂ accumulation in cell walls of mesophyll cells. Importantly no cerium perhydroxide deposits were observed in the cytosol, chloroplasts or mitochondria indicating that no major H₂O₂ accumulation occurred in these organelles.

NADPH oxidases are essential for proline accumulation in response to NaCl and mannitol stresses

Stress-induced H₂O₂ accumulated in the cell wall and apoplast. The main source of cell wall H₂O₂ is the plasma membrane-bound NADPH oxidase (Petrov & Van Breusegem, 2012). We therefore hypothesized that NADPH oxidase activity is required for proline accumulation in response to salt and mannitol stress. We investigated this using a pharmacological approach. Treatments of *A. thaliana* seedlings with DPI, an inhibitor of flavin enzymes such as NADPH oxidases, resulted in a dose-dependent decrease of proline accumulation induced by salt and mannitol stress (Fig. 3b). Addition of various concentrations of DPI did not affect proline accumulation in seedlings growing under normal growth conditions. For further experiments we used 20 µM DPI which diminished proline accumulation by approximately 60%. To aid interpretation of results, the pharmaceutical toxicity of DPI was also tested in seedlings. As shown in Fig. 5, the reduction of proline accumulation caused by the addition of 20 µM DPI to NaCl- and mannitol-treated seedlings for 24 h was totally reversed after washing the seedlings and putting them back in growth medium. In addition no leaf injury or lethality was observed whatever the treatment.

We monitored H₂O₂ production in root tips by adding H₂DCFDA, which reacts with H₂O₂ in living cells to produce fluorescent DCF. As observed in whole seedlings, H₂O₂ is detected in root tips 6 h after exposure to either NaCl or mannitol (Fig. 6a). In contrast H₂O₂ was not

detected in root tips incubated with DPI prior to treatment with either NaCl or mannitol prevented, suggesting H₂O₂ production was inhibited.

The rate limiting steps in proline biosynthesis and degradation in *Arabidopsis* are catalyzed by P5CS and ProDH respectively. To further investigate the role of NADPH oxidase in proline metabolism, steady-state transcript levels of *AtP5CS1* and *AtProDH1* were investigated. Semi-quantitative RT-PCR analysis showed that DPI treatment significantly decreased *P5CS1* transcript levels in response to salt and to a lesser extent to mannitol (Fig. 6c). Steady-state transcript levels of *ProDH1* gene were not altered by DPI. Protein levels of P5CS and ProDH were also investigated by Western blot analyses. In wild-type *A. thaliana* plants, 200 mM NaCl and 400 mM mannitol both caused an increase in P5CS protein content compared with untreated plants (Fig. 6d). However, the increase in P5CS protein levels by salt or mannitol was markedly suppressed by pre-treating seedlings with DPI. In contrast to ProDH, whose protein levels were not affected by the different conditions, a good correlation was observed between *P5CS* transcript and protein levels and proline content.

Proline accumulation in *atrboh* mutants

If ROS produced by NADPH oxidase is involved in regulating proline accumulation due to NaCl and mannitol stress, then KO mutants defective in NADPH oxidase would be expected to accumulate less proline than wild type. Of the ten *Arabidopsis* NADPH oxidase genes identified, *AtRbohD* and *AtRbohF* are the only ones expressed in all plant organs and they are implicated in abiotic stress physiology (Fluhr, 2009; Marino *et al.*, 2012). *AtrbohD-3* and *atrbohF-3* mutants were therefore selected for our study. Semi-quantitative RT-PCR using wild-type and mutant plants revealed that *rbohD-3* and *rbohF-3* homozygous plants contained no detectable levels of *AtRbohD* and *AtRbohF* transcripts in response to salt treatment (data not shown). In both *atrbohD-3* and *atrbohF-3* mutants, the transient increase in H₂O₂ levels at 6 h in response to NaCl and mannitol stresses was not as large as the wild-type increase (Fig. 7). After cerium perhydroxide staining no visible cerium perhydroxide deposits were observed in the leaves of *atrbohD* mutant seedlings grown under mannitol or salt stress, indicating the absence of detectable levels of H₂O₂ (Supporting Information Fig. S1).

The two NADPH oxidase-deficient *atrbohD* and *atrbohF* mutants were examined for the accumulation of proline and P5CS protein in response to either mannitol or salt stress. As expected, application of 200 mM NaCl or 400 mM mannitol induced proline accumulation in wild-type *Arabidopsis* plants (Fig. 8a). Consistent with the results of DPI treatment in wild-

type (see Fig. 6), *atrboh*d-3 and *atrboh*f-3 mutants accumulated less proline and less P5CS protein than wild-type in the presence of 200 mM NaCl (Fig. 8a, b). The results of 400 mM mannitol treatments were similar although less pronounced (Fig. 8a, b). Similar results for proline accumulation were observed with independent alleles of *atrboh*d and *atrboh*f (T-DNA insertion lines SALK_070610 and SALK_059888 respectively) (Supporting Information Fig. S2). In the double *atrboh*d/*f* mutant proline levels reached similar levels to those in the single mutants in response to mannitol stress (Fig. 8). The amount of proline that accumulated in response to NaCl in the double mutant was slightly higher but was not as high as the level induced in wild type. It was noted that the decrease in proline and P5CS accumulation was less pronounced in the *atrboh* mutants than in seedlings treated with DPI (Fig. 5 and 6). Perhaps NADPH oxidases other than AtRbohD and AtRbohF or other sources of H₂O₂ are involved in the regulation of proline biosynthesis. To further demonstrate a role for H₂O₂, we reversed the effect of *atrboh* mutation by applying exogenous H₂O₂. Addition of 10 mM H₂O₂ restored the levels of proline and P5CS protein accumulation in mannitol-treated *atrboh*d-3, *atrboh*f-3 and double *atrboh*d/*f* seedlings (Fig. 8).

Discussion

Plant stress tolerance involves diverse mechanisms such as signal perception and transduction, osmolyte accumulation, ion homeostasis, growth regulation and cellular protection from damage triggered by reactive oxygen species (ROS). It is widely accepted that H₂O₂ and other ROS are also important signalling molecules in the activation of defence genes in response to biotic stress (Foyer & Noctor, 2009, Bartoli *et al.*, 2012). The connection between ROS and proline in response to biotic stress has already been highlighted by Fabro *et al.* (2004). Nevertheless little is known about the involvement of ROS in the regulation of proline metabolism in response to abiotic stresses. The aim of our work was to evaluate the roles of H₂O₂ in the regulation of proline metabolism in response to two different abiotic stresses in *A. thaliana*. Our results showed that both NaCl and mannitol stress induced proline accumulation and a transient increase in H₂O₂ content (Fig. 1). The timing of these changes might suggest that H₂O₂ could act as a secondary messenger involved in triggering proline biosynthesis. This hypothesis is supported by the observation that treatment with H₂O₂ or PQ also promoted proline accumulation (Fig. 2). However these effects are not as strong as those caused by physiological NaCl or mannitol stresses. The fact that much less proline accumulated in response to H₂O₂ and PQ than to the stress conditions is in itself intriguing.

One possibility is that additional signalling pathways are involved in the full stress response. Phospholipase D enzymes are putative signalling components as they have been previously shown to negatively regulate proline accumulation in non-stress or mild stress conditions (Thiery *et al.*, 2004). Following this reasoning, phospholipase D activity may need to be inhibited in order to elicit a full proline stress response by H₂O₂ treatment. As H₂O₂ is able to increase proline accumulation and DMTU is an effective inhibitor of proline accumulation induced by NaCl or mannitol, our data suggest that the stress-induced proline accumulation observed in *A. thaliana* seedlings is mediated at least partly by H₂O₂.

ROS, such as singlet oxygen (¹O₂), H₂O₂, and hydroxyl radical (OH[•]), are produced during normal aerobic metabolism in different cell compartments such as cell walls, plasma membranes, chloroplasts, mitochondria and peroxisomes (Dat *et al.*, 2000). The production of H₂O₂ has been repeatedly demonstrated at the subcellular level by using CeCl₃ techniques, for example, during abscisic acid signalling (Hu *et al.*, 2006), pathogen attack (Bestwick *et al.*, 1997), responses to ozone (Pellinen *et al.*, 1999), drought (Hu *et al.*, 2006), anoxia (Blokhina *et al.*, 2001), and heavy metal excess (Romero-Puertas *et al.*, 2004). Our observation based on CeCl₃ detection revealed that either salt or mannitol stress induced H₂O₂ accumulation in walls of mesophyll cells (Fig. 4). H₂DCFDA fluorescence analysis also revealed an accumulation of H₂O₂ in Arabidopsis root tips in response to either NaCl or mannitol stress (Fig. 6). This is consistent with a report that H₂O₂ accumulation was detected along the plasma membrane of maize leaves challenged with abscisic acid and water-deficit stress (Hu *et al.*, 2006). Using CeCl₃, we did not detect any of H₂O₂ in chloroplasts or in any other organelle of *A. thaliana* leaf cells after 6 h of either salt or mannitol treatment. The apoplastic oxidative burst and resultant H₂O₂ accumulation in the extracellular space is characteristic of plant cells exposed to biotic and abiotic stresses (Bartoli *et al.*, 2012). Studies of different plant species have demonstrated the action of plasma membrane-bound NADPH oxidases in the apoplastic ROS-producing system during early oxidative bursts which is critical in plant signalling and development, including in defence, root hair development, stomatal closure, and early responses to salt stress (Torres *et al.*, 2002; Foreman *et al.*, 2003; Kwak *et al.*, 2003; Leshem *et al.*, 2007). DPI is a commonly used potent inhibitor of flavin enzymes such as NADPH oxidase. We found that DPI efficiently inhibited H₂O₂ production in *A. thaliana* seedling roots exposed to NaCl or mannitol (Fig. 6a), strongly suggesting that at least some of the H₂O₂ production induced by salt or mannitol originates from NADPH oxidase.

In response to either salt or mannitol stress, *P5CS1* proline biosynthesis transcript and protein levels increased at the same time as proline accumulated. Pretreatment of *A. thaliana* seedling with DPI reduced the salt- or mannitol-induced proline accumulation in a dose-dependent manner (Fig. 6). In addition, DPI significantly diminished *P5CS1* transcript and protein accumulation. The reaction catalyzed by NADPH oxidase uses O_2 to generate superoxide ($O_2^{\cdot-}$), which is then converted into H_2O_2 by apoplastic superoxide dismutase. H_2O_2 , as a nonpolar molecule, can easily pass through the plasma membrane probably through aquaporins. H_2O_2 has important roles as a signalling molecule in the regulation of a variety of biological processes. Possibly by redox changes H_2O_2 might directly or indirectly activate unknown signalling components, such as transcription factors, to regulate the transcription of proline biosynthesis genes.

The role of Rbohs in the regulation of proline metabolism was further investigated by reverse genetic approach using *atrbohD* and *atrbohF* KO mutants. Compared to wild type, the strong reduction of H_2O_2 production after 6 h in the *atrbohD* mutant in response to NaCl and mannitol, observed to a lesser extent in the *atrbohF*, could be considered to be consistent with *AtRbohD* being the most highly expressed member of the *AtRboh* gene family in response to salt (Leshem *et al.*, 2007). Furthermore the cytochemical detection of H_2O_2 indicated that H_2O_2 production during salt and mannitol stress is associated with *AtRbohD* expression.

Our study shows that proline accumulation was 20-fold higher in salt-treated wild-type compared to control wild-type plants; however, it was respectively only 5-fold and 9-fold higher in salt-treated *atrbohD* and *atrbohF* seedlings compared to their controls. The double *atrbohD/f* mutant surprisingly produced slightly more proline in response to NaCl than the corresponding single mutants possibly because the double mutant is more sensitive to NaCl. Similarly, a much lower P5CS level was observed in the *atrbohD* mutant, compared to only a marginal reduction in *atrbohF* mutant compared to wild-type plants. Taken together, these results indicate that these NADPH oxidase isoforms are involved in proline accumulation during salt stress with *AtRbohD* having a prominent role. It was noted that the decrease in proline accumulation was more pronounced in wild-type seedlings treated with DPI than in *atrboh* mutants. This result would implicate other NADPH oxidases in the proline accumulation response. Indeed according to qPCR measurements, *AtRbohA*, *AtRbohB* and *AtRbohC* transcripts are more abundant in *atrbohD* and *atrbohF* mutants than in wild-type under stress, which is possibly preliminary evidence of a feed-back mechanism in these mutants (Supporting Information Fig. S3).

Recently, it has been demonstrated that ROS generated by AtrbohF has a specific and predominant role in regulating Na⁺ accumulation and soil-salinity tolerance (Jiang *et al.*, 2012). When grown in saline soil, *atrbohF* mutant accumulated higher levels of Na⁺ than wild-type plants. In addition, Ma *et al.* (2012) have suggested that AtrbohF acts redundantly with AtrbohD in regulating Na⁺/K⁺ homeostasis. These independent observations lead us to postulate that the difference in proline accumulation in the wild-type and *atrboh* mutants plants could be due to impaired Na⁺/K⁺ regulation. No differences in Na⁺ and K⁺ content were found between *atrbohD*, *atrbohF* and wild-type plants subjected to NaCl treatment for 24 h (Supporting Information Fig. S4). However the double *atrbohdf* mutant contained less Na⁺ but had a higher Na⁺/K⁺ ratio. Differences in Na⁺ accumulation in *atrboh* single mutants between our study and the study of Jiang *et al.* (2012) were probably due to large differences in growth and stress conditions in the two experimental systems, such as continuous light versus 16 h light/8 h dark cycles or short versus long durations of stress. However the regulation of proline accumulation by Rboh in Arabidopsis was probably mainly due to the osmotic stress component of salt stress rather than to the ionic component, because Rboh was involved in the response to both NaCl and mannitol stress. Lastly, since the generation of ROS by Rboh causes changes in the cell redox potential, we postulate that redox-sensitive signalling components or transcription factors may be activated and to influence the expression of proline biosynthesis genes.

In conclusion, our results shed new light on the regulation of proline metabolism in response to abiotic stresses showing the involvement of NADPH oxidase and H₂O₂. We show that H₂O₂ is involved in proline accumulation induced by salt and mannitol stresses. First, proline accumulation was preceded by elevated H₂O₂ levels, and scavenging of H₂O₂ by DMTU abolished proline accumulation. Second, we have presented evidence that NADPH oxidases are the potential source of the observed stress-induced H₂O₂ generation. Third, the absence of H₂O₂ production in cell walls and the accumulation of less proline in *atrbohD* and *atrbohF* KO mutants in response to NaCl and mannitol provides convincing genetic evidence that the corresponding NADPH oxidase isoforms contribute to proline accumulation.

Acknowledgements

We thank the IFR 83 Electron Microscopy Service of the CNRS Institute of Integrative Biology (UPMC, Paris, France) for use of transmission electron microscopy facilities. We

thank Christophe Bailly for valuable scientific discussion and advice on ROS measurements and Alban Launay and Pierre Carol for critical reading of the manuscript. K. Ben Rejeb was supported by a PhD scholarship from the Tunisian Ministry of Higher Education and Scientific Research (LR10CBBC02) and the Tunisian-French CMCU (Comité Mixte de Coopération Universitaire) network (29174UF). Part of this study was supported by European Union COST program FA0901 “Putting halophytes to work - From genes to ecosystem”.

References

- Bartoli CG, Casalongue C, Simontacchi M, Márquez-García B, Foyer CH. 2012.** Interactions between hormone and redox signalling pathways in the control of growth and cross tolerance to stress. *Environmental and Experimental Botany* **94**: 73-88.
- Bates LS, Waldren RP, Teare ID. 1973.** Rapid determination of free proline for water stress studies. *Plant and Soil* **39**: 205-207.
- Bestwick CS, Brown IR, Bennett MH, Mansfield JW. 1997.** Localization of hydrogen peroxide accumulation during the hypersensitive reaction of lettuce cells to *Pseudomonas syringae* pv *phaseolicola*. *Plant Cell* **9**: 209-221.
- Blokhina OB, Chirkova TV, Fagerstedt KV. 2001.** Anoxic stress leads to hydrogen peroxide formation in plant cells. *Journal of Experimental Botany* **52**: 1-12.
- Chen H, Jiang JG. 2010.** Osmotic adjustment and plant adaptation to environmental changes related to drought and salinity. *Dossiers environnement* **18**: 309-319.
- Dat J, Vandenabeele S, Vranová E, Van Montagu M, Inzé D, Van Breusegem F. 2000.** Dual action of the active oxygen species during plant stress responses. *Cellular and Molecular Life Sciences* **57**: 779-795.
- Fabro G, Kovács I, Pavet V, Szabados L, Alvarez ME. 2004.** Proline accumulation and *AtP5CS2* gene activation are induced by plant-pathogen incompatible interactions in *Arabidopsis*. *Molecular Plant-Microbe Interactions*. **17**: 343-350.
- Fluhr R. 2009.** Reactive oxygen-generating NADPH oxidases in plants. In: del Rio LA, Puppo A, eds. *Reactive oxygen species in plant signalling*. Berlin, Springer-Verlag, 1-23.
- Foreman J, Demidchik V, Bothwell JHF, Mylona P, Miedema H, Torres MA, Linstead P, Costa S, Brownlee C, Jones JDG, Davies JM, Dolan L. 2003.** Reactive oxygen species produced by NADPH oxidase regulate plant cell growth. *Nature* **422**: 442-446.

502 **Foyer CH, Noctor G. 2009.** Redox regulation in photosynthetic organisms: signaling,
503 acclimation, and practical implications. *Antioxidants and Redox Signaling* **11**: 861-905.

504 **Funck D, Eckard S. Müller G. 2010.** Non-redundant functions of two proline
505 dehydrogenase isoforms in Arabidopsis. *BMC Plant Biology* **10**: 70.

506 **Hayashi F, Ichino T, Osanai R, Wada K. 2000.** Oscillation and regulation of proline
507 content by *P5CS* and *ProDH* gene expressions in the light/dark cycles in *Arabidopsis thaliana*
508 *L. Plant and Cell Physiology* **41**: 1096-1101.

509 **Hellemans J, Mortier G, De Paepe A, Speleman F, Vandesompele J. 2007.** qBase relative
510 quantification framework and software for management and automated analysis of real-time
511 quantitative PCR data. *Genome Biology* **8**: R19.

512 **Hu X, Zhang A, Zheng J, Jiang M. 2006.** Absciscic acid is a key inducer of hydrogen
513 peroxide production in leaves of maize plants exposed to water stress. *Plant and Cell*
514 *Physiology* **47**: 1484-1485.

515 **Jiang C, Belfield EJ, Mithani A, Visscher A, Ragoussis J, Mott R, Smith JAC, Harberd**
516 **NP. (2012).** ROS-mediated vascular homeostatic control of root-to-shoot soil Na delivery in
517 *Arabidopsis*. *EMBO Journal* **31**: 4359.

518 **Kiyosue T, Yoshiba Y, Yamaguchi-Shinozaki K, Shinozaki K. 1996.** A nuclear gene
519 encoding mitochondrial proline dehydrogenase, an enzyme involved in proline metabolism, is
520 upregulated by proline but downregulated by dehydration in *Arabidopsis*. *Plant Cell* **8**: 1323-
521 1335.

522 **Kwak JM, Mori I, Pei Z-M, Leonhardt N, Torres MA, Dangl JL, Bloom R, Bodde S,**
523 **Jones JDG, Schroeder JJ. 2003.** NADPH oxidase *AtrbohD* and *AtrbohF* genes function in
524 ROS-dependent ABA signaling in *Arabidopsis*. *EMBO Journal* **22**: 2623-2633.

525 **Leprince AS, Magalhaes N, De Vos D, Bordenave B, Crilat E, Clément G, Meyer C,**
526 **Munnik T, Saviouré A. 2015.** Involvement of Phosphatidylinositol 3-kinase in the regulation
527 of proline catabolism in *Arabidopsis thaliana*. *Frontiers in Plant Science* **5**: 772-795.

528 **Leshem Y, Seri L, Levine A. 2007.** Induction of phosphatidylinositol 3-kinase-mediated
529 endocytosis by salt stress leads to intracellular production of reactive oxygen species and salt
530 tolerance. *Plant Journal* **51**: 185-197.

531 **Ma L, Zhang H, Sun L, Jiao Y, Zhang G, Miao C, Hao F. 2012.** NADPH oxidase *AtrbohD*
532 and *AtrbohF* function in ROS-dependent regulation of Na⁺/K⁺ homeostasis in *Arabidopsis*
533 under salt stress. *Journal of Experimental Botany* **63**: 305-317.

534 **Marino D, Dunand C, Puppo A, Pauly N. 2012.** A burst of plant NADPH oxidases. *Trends*
535 *in Plant Science* **17**: 9-15.

536 **Maruta T, Inoue T, Tamoi M, Yabuta Y, Yoshimura K, Ishikawa T, Shigeoka S. 2011.**
537 *Arabidopsis* NADPH oxidases, AtrbohD and AtrbohF, are essential for jasmonic acid-induced
538 expression of genes regulated by MYC2 transcription factor. *Plant Science* **180**: 655-660.

539 **Miller G, Suzuki N, Ciftci-Yilmaz S, Mittler R. 2011.** Reactive oxygen species homeostasis
540 and signalling during drought and salinity stresses. *Plant Cell and Environment* **33**: 453-367.

541 **Nakashima K, Ito Y, Yamaguchi-Shinozaki K. 2009.** Transcriptional regulatory networks
542 in response to abiotic stresses in *Arabidopsis* and grasses. *Plant Physiology* **149**: 88-89.

543 **Oracz K, El-Maarouf-Bouteau H, Kranner I, Bogatek R, Corbineau F, Bailly C. 2009.**
544 The mechanisms involved in seed dormancy alleviation by hydrogen cyanide unravel the role
545 of reactive oxygen species as key factors of cellular signaling during germination. *Plant*
546 *Physiology* **150**: 494-505.

547 **Parre E, Ghars MA, Leprince A-S, Thiery L, Lefebvre D, Bordenave M, Luc R, Mazars**
548 **C, Abdelly C, Saviouré A. 2007.** Calcium signaling *via* phospholipase C is essential for
549 proline accumulation upon ionic but not nonionic hyperosmotic stresses in *Arabidopsis*. *Plant*
550 *Physiology* **144**: 503-512.

551 **Pellinen R, Palva T, Kangasjärvi J. 1999.** Subcellular localization of ozone-induced
552 hydrogen peroxide production in birch (*Betula pendula*) leaf cells. *Plant Journal* **20**: 349-356.

553 **Petrov VD, Van Breusegem F. 2012.** Hydrogen peroxide-a central hub for information flow
554 in plant cells. *AoB Plant. pls014* doi: 10.1093/aobpla/pls014.

555 **Pogany M, von Rad U, Grun S, Dongo A, Pintye A, Simoneau P, Bahnweg G, Kiss L,**
556 **Barna B, Durner J. 2009.** Dual roles of reactive oxygen species and NADPH oxidase
557 RBOHD in an *Arabidopsis-Alternaria* pathosystem. *Plant Physiology* **151**: 1459-1475.

558 **Romero-Puertas MC, McCarthy I, Sandalio LM, Palma JM, Corpas FJ, Gomez M, Del**
559 **Rio LA. 1999.** Cadmium toxicity and oxidative metabolism of pea leaves peroxisomes. *Free*
560 *Radical Research* **31**: 25-31.

561 **Sagi M, Fluhr R. 2001.** Superoxide production by plant homologues of the gp91^{phox} NADPH
562 oxidase. Modulation of activity by calcium and by tobacco mosaic virus infection. *Plant*
563 *Physiology* **126**: 1281-1290.

564 **Sagi, M, Fluhr, R. (2006).** *Production of reactive oxygen species by plant NADPH oxidases.*
565 *Plant Physiology* **141**: 336-340.

566 **Savouré A, Hua XJ, Bertauche N, Van Montagu M, Verbruggen N. 1997.** Absciscic acid-
567 independent and absciscic acid-dependent regulation of proline biosynthesis following cold
568 and osmotic stresses in *Arabidopsis thaliana*. *Molecular and General Genetics* **254**, 104-109.

569 **Savouré A, Jaoua S, Hua XJ, Ardiles W, Van Montagu M, Verbruggen N. 1995.**
570 Isolation, characterization, and chromosomal location of a gene encoding the DELTA-1-
571 pyrroline-5-carboxylate synthetase in *Arabidopsis thaliana*. *FEBS Letters* **372**: 13-19.

572 **Servet C, Ghelis T, Richard L, Zilberstein A, Savouré A. 2012.** Proline dehydrogenase: a
573 key enzyme in controlling cellular homeostasis. *Frontiers in Bioscience* **17**: 607-620.

574 **Sharma S, Verslues PE. 2010.** Mechanisms independent of ABA or proline feedback have a
575 predominant role in transcriptional regulation of proline metabolism during low water
576 potential and stress recovery. *Plant Cell and Environment* **33**: 1838-1851.

577 **Sharma S, Villamor JG, Verslues PE. 2011.** Essential role of tissue-specific proline
578 synthesis and catabolism in growth and redox balance at low water potential. *Plant*
579 *Physiology* **157**: 292-304.

580 **Slama I, Abdelly C, Bouchereau A, Flowers T, Savouré A. 2015.** Diversity, distribution
581 and roles of osmoprotective compounds accumulated in halophytes under abiotic stress.
582 *Annals of Botany* **115**: 327-331.

583 **Strizhov N, Abraham E, Okresz L, Blickling S, Zilberstein A, Schell J, Koncz C,**
584 **Szabados L. 1997.** Differential expression of two *P5CS* genes controlling proline
585 accumulation during salt-stress requires ABA and is regulated by ABA1, ABI1 and AXR2 in
586 *Arabidopsis*. *Plant Journal* **12**: 557-569.

587 **Szabados L, Savouré A. 2010.** Proline: a multifunctional amino acid. *Trends Plant Science*
588 **15**: 89-97.

589 **Székely G, Abraham E, Cselo A, Rigo G, Zsigmond L, Csiszar J, Ayaydin F, Strizhov N,**
590 **Jasik J, Schmelzer E, Koncz C, Szabados L. 2008.** Duplicated *P5CS* genes of *Arabidopsis*
591 play distinct roles in stress regulation and developmental control of proline biosynthesis.
592 *Plant Journal* **53**: 11-28.

593 **Thiery L, Leprince A-S, Lefebvre D, Ghars MA, Debarbieux E, Savouré A. 2004.**
594 Phospholipase D is a negative regulator of proline biosynthesis in *Arabidopsis thaliana*.
595 *Journal of Biological Chemistry* **279**: 14812-14818.

596 **Torres MA, Dangl JL, Jones JDG. 2002.** *Arabidopsis* gp91^{phox} homologues AtrbohD and
597 AtrbohF are required for accumulation of reactive oxygen intermediates in the plant defense
598 response. *Proceedings of the National Academy of Sciences USA* **99**: 517-522.

599 **Verbruggen N, Hua XJ, May M, Van Montagu M. 1996.** Environmental and
600 developmental signals modulate proline homeostasis: evidence for a negative transcriptional
601 regulator. *Proceedings of the National Academy of Sciences USA* **93**: 8787–8791.

602 **Xie YJ, Xu S, Han B, Wu MZ, Yuan XX, Han Y, Gu Q, Xu DK, Yang Q, Shen WB.**
603 **(2011).** Evidence of Arabidopsis salt acclimation induced by up-regulation of HY1 and the
604 regulatory role of RbohD-derived reactive oxygen species synthesis. *Plant Journal* **66**: 280-
605 292.

606 **Yang SL, Lan SS, Gong M. 2009.** Hydrogen peroxide-induced proline and metabolic
607 pathway of its accumulation in maize seedlings. *Journal of Plant Physiology* **166**: 1694-1699

608 **Yoshida Y, Kiyosue T, Katagiri T, Ueda H, Mizoguchi T, Yamaguchi-Shinozaki K,**
609 **Wada K, Harada Y, Shinozaki K. 1995.** Correlation between the induction of a gene for
610 delta 1-pyrroline-5-carboxylate synthetase and the accumulation of proline in *Arabidopsis*
611 *thaliana* under osmotic stress. *Plant Journal* **7**: 751-760.

612 **Zhao MG, Chen L, Zhang LL, Zhang WH. 2009.** Nitric reductase-dependent nitric oxide
613 production is involved in cold acclimation and freezing tolerance in *Arabidopsis*. *Plant*
614 *Physiology* **151**: 755-767.

Figure legends

Fig. 1. NaCl and mannitol stresses trigger transient increases in H_2O_2 and proline accumulation.

Twelve-day-old wild-type *A. thaliana* seedlings grown on 0.5 x MS solid medium were transferred to 0.5 x MS liquid medium for treatment. Plants were then exposed to 200 mM NaCl (triangles), 400 mM mannitol (squares) or neither (circles) for 24 h. Open and closed symbols indicate proline and H_2O_2 measurements respectively. The results shown are the means of at least three independent experiments ($\pm$ SE).

Fig. 2. H_2O_2 induces proline accumulation.

Proline accumulation in wild-type *A. thaliana* seedlings treated with H_2O_2 (a) and paraquat (b) for 24 h. Means ($\pm$ SE) of at least three independent experiments with different letters are significantly different at $P < 0.05$. (c) H_2O_2 was visualized by using DAB staining in *A. thaliana* leaves exposed to different concentrations of paraquat for 24 h.

Fig. 3. The NADPH oxidase inhibitor DPI and the H_2O_2 scavenger DMTU affect proline accumulation induced by salt or mannitol stress.

Twelve-day-old *A. thaliana* seedlings grown on 0.5 x MS solid medium were transferred to 0.5 x MS liquid medium for treatment. Plants were preincubated with various concentrations of DMTU (a) or DPI (b) for 4 h and then exposed to 200 mM NaCl (grey bars), 400 mM mannitol (black bars) or neither (white bars) for 24 h. Means ($\pm$ SE) of three independent experiments with different letters are significantly different at $P < 0.05$.

Fig. 4. NaCl and mannitol stresses trigger apoplastic H_2O_2 accumulation in *A. thaliana* leaves.

Twelve-day-old seedlings grown on 0.5 x MS solid medium were transferred to 0.5 x MS liquid medium. Subcellular localization of H_2O_2 was detected by CeCl_3 staining of leaves of wild-type *A. thaliana* grown under normal conditions (control, a, d) or in the presence of either 200 mM NaCl (b, e) or 400 mM mannitol (c, f) for 6 h. Arrows indicate electron-dense deposits of cerium perhydroxides formed in the presence of H_2O_2 and CeCl_3 . Ch, chloroplast; CW, cell wall; M, mitochondria; S, starch; V, vacuole.

Fig. 5. Inhibition of proline accumulation by DPI can be reversed.

A. thaliana seedlings were pre-treated with 20 μ M DPI as described in the legend of Fig. 3 and then treated with either 200 mM NaCl or 400 mM mannitol for 24 h (a). Seedlings were then washed twice and transferred onto NaCl or mannitol medium for another 24 h (b). Means ($\pm$ SE) of three independent experiments with different letters are significantly different at $P < 0.05$.

Fig. 6. DPI inhibition of NADPH oxidase activity affects proline metabolism induced by either salt or osmotic stress.

A. thaliana seedlings were prepared and treated as previously described in the legend of Fig. 3. (a) Sites of H_2O_2 production visualized by using 2',7'-dichlorofluorescein diacetate (H_2DCFDA) in wild-type *A. thaliana* root tips pre-treated with 20 μ M DPI and grown under normal conditions or in the presence of either 200 mM NaCl or 400 mM mannitol for 6 h. (b) Proline accumulation in wild-type seedlings pre-treated with 20 μ M DPI and grown under normal conditions or in the presence of either 200 mM NaCl or 400 mM mannitol for 24 h. Means ($\pm$ SE) of three independent experiments with different letters are significantly different at $P < 0.05$. (c) RT-PCR analysis of *P5CS1* and *ProDH1* expression. RT-PCR was performed using total RNAs extracted from seedlings treated for 24 h as described in (b). *APT1* transcripts were amplified as an internal control. Lower panels, quantification of the expression of *P5CS1* and *ProDH1* normalized to *APT1*. (d) Western blot of P5CS and ProDH proteins from seedlings treated for 24 h as described in (b). Rubisco revealed by Ponceau-S staining of all protein was used as a loading control.

Fig. 7. AtRbohD and AtRbohF are required for transient H_2O_2 accumulation induced by salt or mannitol stress.

A. thaliana seedlings were prepared and treated as described in the legend of Fig. 3. H_2O_2 accumulation was measured in seedlings of wild-type and the transposon insertion mutants *atrbohD-3* and *atrbohF-3* at 6 h after stress treatment with either 200 mM NaCl or 400 mM mannitol. Means ($\pm$ SE) of three independent experiments with different letters are significantly different at $P < 0.05$.

Fig. 8. AtRbohD and AtRbohF are essential for proline accumulation in response to either NaCl or mannitol.

A. thaliana seedlings were prepared and treated as previously described in the legend of Fig. 3. (a) Proline accumulation was determined in transposon insertion mutant lines *atrbohd-3*, *atrbohlf-3* single mutants and in *atrbohd/f* double mutant in comparison to wild-type grown under normal conditions or in the presence of either 200 mM NaCl or 400 mM mannitol for 24 h. Proline content was also measured in wild type and mutant seedlings grown in the same condition after application of 10 mM H₂O₂. Means ($\pm$ SE) of three biological replicates with different letters were significantly different at $P < 0.05$. (b) Western blot of P5CS proteins from seedlings treated as described in (a). Rubisco revealed by Ponceau-S staining of all proteins was used as loading control. Quantification of the western blot normalized to Rubisco is shown as bar graph. C, control; N, NaCl; M, mannitol.

Supporting Information

Fig. S1. Early apoplastic H₂O₂ accumulation is mediated through plasma membrane-bound NADPH oxidases in leaves of *A. thaliana* in response to either NaCl or mannitol.

Fig. S2. Proline accumulation in *atrboh* insertion lines.

Fig. S3. qRT-PCR analysis of differential expression of three *AtRboh* genes in wild-type and *atrbohd* and *atrbohlf* mutants.

Fig. S4. Changes in Na⁺ and K⁺ content and Na⁺/K⁺ ratios in Arabidopsis wild-type (WT) and *atrbohd* and *atrbohlf* mutants upon NaCl stress.

Table S1: List of primers used for RT-PCR and qRT-PCR analysis.

Fig. 1. NaCl and mannitol stresses trigger transient increases in H_2O_2 and proline accumulation. Twelve-day-old wild-type *A. thaliana* seedlings grown on 0.5 x MS solid medium were transferred to 0.5 x MS liquid medium for treatment. Plants were then exposed to 200 mM NaCl (triangles), 400 mM mannitol (squares) or neither (circles) for 24 h. The results shown are the means of at least three independent experiments ($\pm$ SE).

46x27mm (600 x 600 DPI)

Fig.2. H₂O₂ induces proline accumulation. Proline accumulation in wild-type *A. thaliana* seedlings treated with H₂O₂ (a) and paraquat (b) for 24 h. Means (± SE) of at least three independent experiments with different letters are significantly different at *P* < 0.05. (c) H₂O₂ was visualized by using DAB staining in *A. thaliana* leaves exposed to different concentrations of paraquat for 24 h.

102x130mm (300 × 300 DPI)

Fig. 3. The NADPH oxidase inhibitor DPI and the H₂O₂ scavenger DMTU affect proline accumulation induced by salt or mannitol stress.

Twelve-day-old *A. thaliana* seedlings grown on 0.5 × MS solid medium were transferred to 0.5 × MS liquid medium for treatment. Plants were preincubated with various concentrations of DMTU or DPI for 4 h and then exposed to 200 mM NaCl (grey bars), 400 mM mannitol (black bars) or neither (white bars) for 24 h. Means ($\pm$ SE) of three independent experiments with different letters are significantly different at $P < 0.05$.

76x72mm (300 x 300 DPI)

Fig. 4. NaCl and mannitol stresses trigger apoplastic H₂O₂ accumulation in *A. thaliana* leaves. Twelve-day-old seedlings grown on 0.5 × MS solid medium were transferred to 0.5 × MS liquid medium. Subcellular localization of H₂O₂ was detected by CeCl₃ staining of leaves of wild-type *A. thaliana* grown under normal conditions (control, a, d) or in the presence of either 200 mM NaCl (b, e) or 400 mM mannitol (c, f) for 6 h. Arrows indicate electron-dense deposits of cerium perhydroxides formed in the presence of H₂O₂ and CeCl₃. Ch, chloroplast; CW, cell wall; M, mitochondria; S, starch; V, vacuole.

75x43mm (300 x 300 DPI)

Fig. 5. Inhibition of proline accumulation by DPI can be reversed.

A. thaliana seedlings were pre-treated with 20 μM DPI as described in the legend of Fig. 3 and then treated with either 200 mM NaCl or 400 mM mannitol for 24 h (a). Seedlings were then washed twice and transferred onto NaCl or mannitol medium for another 24 h (b). Means (± SE) of three independent experiments with different letters are significantly different at $P < 0.05$.

40x20mm (300 x 300 DPI)

Fig. 6. DPI inhibition of NADPH oxidase activity affects proline metabolism induced by either salt or osmotic stress.

A. thaliana seedlings were prepared and treated as previously described in the legend of Fig. 3. (a) Sites of H₂O₂ production visualized by using 2',7'-dichlorofluorescein diacetate (H₂DCFDA) in wild-type *A. thaliana* root tips pre-treated with 20 μM DPI and grown under normal conditions or in the presence of either 200 mM NaCl or 400 mM mannitol for 6 h. (b) Proline accumulation in wild-type seedlings pre-treated with 20 μM DPI and grown under normal conditions or in the presence of either 200 mM NaCl or 400 mM mannitol for 24 h. Means (± SE) of three independent experiments with different letters are significantly different at *P* < 0.05. (c) RT-PCR analysis of P5CS1 and ProDH1 expression. RT-PCR was performed using total RNAs extracted from seedlings treated for 24 h as described in (b). APT1 transcripts were amplified as an internal control. Lower panels, quantification of the expression of P5CS1 and ProDH1 normalized to APT1. (d) Western blot of P5CS and ProDH proteins from seedlings treated for 24 h as described in (b). Rubisco revealed by Ponceau-S staining of all protein was used as a loading control.

Fig. 7. AtRbohD and AtRbohF are required for transient H₂O₂ accumulation induced by salt or mannitol stress.

A. thaliana seedlings were prepared and treated as described in the legend of Fig. 3. H₂O₂ accumulation was measured in seedlings of wild-type and the transposon insertion mutants *atrboh-3* and *atrboh-3* at 6 h after stress treatment with either 200 mM NaCl or 400 mM mannitol. Means (± SE) of three independent experiments with different letters are significantly different at P < 0.05.

46x26mm (300 x 300 DPI)

Fig. 8. AtRbohD and AtRbohF are essential for proline accumulation in response to either NaCl or mannitol. *A. thaliana* seedlings were prepared and treated as previously described in the legend of Fig. 3. (a) Proline accumulation was determined in transposon insertion mutant lines *atrboh3*, *atrboh3f* single mutants and in *atrboh3f/f* double mutant in comparison to wild-type grown under normal conditions or in the presence of either 200 mM NaCl or 400 mM mannitol for 24 h. Proline content was also measured in wild type and mutant seedlings grown in the same condition after application of 10 mM H₂O₂. Means (± SE) of three biological replicates with different letters were significantly different at $P < 0.05$. (b) Western blot of P5CS proteins from seedlings treated as described in (a). Rubisco revealed by Ponceau-S staining of all proteins was used as loading control. Quantification of the western blot normalized to Rubisco is shown as bar graph. C, control; N, NaCl; M, mannitol.

78x60mm (600 x 600 DPI)