



**HAL**  
open science

## Marine protist diversity in European coastal waters and sediments as revealed by high-throughput sequencing

Ramon Massana, Angélique Gobet, Stéphane Audic, David Bass, Lucie Bittner, Christophe Boutte, Aurélie Chambouvet, Richard Christen, Jean-Michel Claverie, Johan Decelle, et al.

### ► To cite this version:

Ramon Massana, Angélique Gobet, Stéphane Audic, David Bass, Lucie Bittner, et al.. Marine protist diversity in European coastal waters and sediments as revealed by high-throughput sequencing. *Environmental Microbiology*, 2015, 17 (10), pp.4035-4049. 10.1111/1462-2920.12955 . hal-01191731

**HAL Id: hal-01191731**

**<https://hal.sorbonne-universite.fr/hal-01191731v1>**

Submitted on 2 Sep 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Marine protist diversity in European coastal waters and sediments as

5

## revealed by high-throughput sequencing

10

Ramon Massana<sup>1\*</sup>, Angélique Gobet<sup>2,3</sup>, Stéphane Audic<sup>2,3</sup>, David Bass<sup>4</sup>, Lucie Bittner<sup>2,3,5</sup>, Christophe Boutte<sup>2,3</sup>, Aurélie Chambouvet<sup>6</sup>, Richard Christen<sup>7</sup>, Jean-Michel Claverie<sup>8</sup>, Johan Decelle<sup>2,3</sup>, John R. Dolan<sup>9</sup>, Micah Dunthorn<sup>5</sup>, Bente Edvardsen<sup>10</sup>, Irene Forn<sup>1</sup>, Dominik Forster<sup>5</sup>, Laure Guillou<sup>2,3</sup>, Olivier Jaillon<sup>11</sup>, Wiebe H.C.F. Kooistra<sup>12</sup>, Ramiro Logares<sup>1</sup>, Frédéric Mahé<sup>5</sup>, Fabrice Not<sup>2,3</sup>, Hiroyuki Ogata<sup>13</sup>, Jan Pawlowski<sup>14</sup>, Massimo C. Pernice<sup>1</sup>, Ian Probert<sup>2,3</sup>, Sarah Romac<sup>2,3</sup>, Thomas Richards<sup>6</sup>, Sébastien Santini<sup>8</sup>, Kamran Shalchian-Tabrizi<sup>10</sup>, Raffaele Siano<sup>15</sup>, Nathalie Simon<sup>2,3</sup>, Thorsten Stoeck<sup>5</sup>, Daniel Vaultot<sup>2,3</sup>, Adriana Zingone<sup>12</sup>, and Colomán de Vargas<sup>2,3</sup>

15

<sup>1</sup> Institut de Ciències del Mar (CSIC), Passeig Marítim de la Barceloneta, 37-49, ES-08003, Barcelona, Catalonia, Spain

<sup>2</sup> CNRS, UMR 7144, Station Biologique de Roscoff, FR-29682 Roscoff, France

<sup>3</sup> UPMC Université Paris 06, UMR 7144, Station Biologique de Roscoff, France

<sup>4</sup> The Natural History Museum, London SW7 5BD & Cefas, Weymouth, Dorset DT4 8UB, United Kingdom

<sup>5</sup> University of Kaiserslautern, D-67663 Kaiserslautern, Germany

20

<sup>6</sup> University of Exeter, Biosciences, Exeter, EX4 4QD, United Kingdom

<sup>7</sup> CNRS, UMR 7138, Université Nice Sophia Antipolis, FR-06108 Nice, France

<sup>8</sup> CNRS, UMR 7256, Aix-Marseille Université, FR-13288 Marseille, France

<sup>9</sup> CNRS UMR 7093, UPMC Université Paris 06, Laboratoire d'Océanographie de Villefranche, FR-06230 Villefranche-sur-Mer, France

<sup>10</sup> University of Oslo, Department Biosciences, N-0316 Oslo, Norway

25

<sup>11</sup> CEA, Genoscope, 2 rue Gaston Crémieux, FR-91000 Evry, France

<sup>12</sup> Stazione Zoologica Anton Dohrn, Villa Comunale, I-80121 Naples, Italy

<sup>13</sup> Institute for Chemical Research, Kyoto University, Uji, Kyoto 611-0011, Japan

<sup>14</sup> University of Geneva, CH-1211 Geneva, Switzerland

<sup>15</sup> Ifremer DYNECO/Pelagos, BP 7029280 Plouzané, France

30

\* Corresponding author: Ramon Massana. Department of Marine Biology and Oceanography, Institut de Ciències del Mar (CSIC), Passeig Marítim de la Barceloneta 37-49, ES-08003 Barcelona, Catalonia, Spain.

E-mail: ramonm@icm.csic.es. Phone: 34-93-2309500. Fax: 34-93-2309555.

Running title: Protist diversity in European coastal areas

35

## Summary

Although protists are critical components of marine ecosystems they are still poorly characterized. Here we analyzed the taxonomic diversity of planktonic and benthic protist communities collected in six distant European coastal sites. Environmental DNA and RNA from three size-fractions (pico-, nano-, and micro/meso-plankton) as well as from dissolved DNA and surface sediments were used as templates for tag pyrosequencing of the V4 region of the 18S rDNA.  $\beta$ -diversity analyses split the protist community structure into three main clusters: picoplankton-nanoplankton-dissolved DNA, micro/mesoplankton, and sediments. Within each cluster, protist communities from the same site and time clustered together, while communities from the same site but different seasons were unrelated. Both DNA and RNA-based surveys provided similar relative abundances for most class-level taxonomic groups. Yet, particular groups were overrepresented in one of the two templates, such as MALV-I and MALV-II that were much more abundant in DNA surveys. Overall, the groups displaying the highest relative contribution were Dinophyceae, Diatomea, Ciliophora and Acantharia. Also well represented were Mamiellophyceae, Cryptomonadales, MALV and MAST in the picoplankton, and Monadofilosa and basal Fungi in sediments. Our extensive and systematic sequencing of geographically separated sites provides the most comprehensive molecular description of coastal marine protist diversity to date.

55

Key words: 18S rDNA / dissolved DNA / environmental DNA and RNA extracts / marine protist diversity / pyrosequencing / taxonomy

## Introduction

Protists or unicellular eukaryotes cover a wide spectrum of cell sizes, shapes, and taxonomic affiliations (Schaechter, 2012). They represent the majority of eukaryotic lineages, so studying their diversity is of primary interest for understanding the eukaryotic tree of life (Keeling *et al.*, 2005; Burki, 2014). Moreover, protists play a variety of crucial roles in marine ecosystems from primary producers, predators, decomposers to parasites (Sherr *et al.*, 2007), leading to much effort in quantifying particular species and inferring their ecological functions. A vast literature exists in which species of dinoflagellates (e.g. Graham *et al.*, 2004), diatoms (e.g. Olguín *et al.*, 2006) and ciliates (e.g. Dolan *et al.*, 2013) have been studied based on morphological features observable in light microscopy (LM), a task that requires considerable expertise and time to key out species accurately. Even for these relatively visible groups, examples are known of morphologically similar individuals belonging to different cryptic species (Amato *et al.*, 2007) or morphologically distinct types from the same species (Pizay *et al.*, 2009). Accurate identification may thus not always be discerned from cell morphology alone, and this is more critical for protists below 20 µm in size that often lack conspicuous shapes (Massana, 2011). Over the last decades, DNA sequencing of environmental phylogenetic markers has changed our perception of microbial diversity in most ecosystems. These molecular surveys have been instrumental in decoding the large protist diversity and in unveiling new lineages, such as Picozoa (Not *et al.*, 2007; Seenivasan *et al.*, 2013), MALV (Marine Alveolates) clades (Guillou *et al.*, 2008) and MAST (Marine Stramenopiles) clades (Massana *et al.*, 2004; 2014).

Earlier molecular surveys were based on clone libraries of near full-length 18S rDNA genes followed by Sanger sequencing of a subset of the clones (Díez *et al.*, 2001; López-García *et al.*, 2001; Moon-van der Staay *et al.*, 2001). The resulting high-quality, often manually

checked environmental sequences have been crucial for the phylogenetic placement of novel clades and, together with sequences from monoclonal cultures, are the basis of reference rDNA databases (Guillou *et al.*, 2013; Pernice *et al.*, 2013). However, traditional clone  
85 libraries only capture the most dominant species in the community (Pedrós-Alió, 2006), a limitation bypassed by high-throughput sequencing (HTS) methods. By providing the deep inventories needed both for taxonomic descriptions and sample comparisons, HTS has enabled microbial ecology to advance greatly. HTS has been applied to study protist diversity in a wide variety of systems, including surface and deep marine waters (Amaral-Zettler *et al.*,  
90 2009; Cheung *et al.*, 2010; Edgcomb *et al.*, 2011; de Vargas *et al.*, 2015), marine sediments (Bik *et al.*, 2012), lakes (Mangot *et al.*, 2013), soils (Bates *et al.*, 2013), and metazoan hosts (He *et al.*, 2014). In the case of marine protists, most studies have targeted a specific size-fraction or a particular location. In addition, these surveys generally used environmental DNA as template for PCR amplification, and it has been shown that using RNA extracts instead can  
95 provide a different picture of biodiversity (Stoeck *et al.*, 2007; Not *et al.*, 2009; Lejzerowicz *et al.*, 2013) and useful complementary information (Blazewicz *et al.*, 2013).

The present study is an investigation of benthic-pelagic protists in marine habitats along the European coastline, sampled between 2009 and 2010 during the research program *BioMarKs*. The 95 different pyrosequenced samples analyzed herein address total protist diversity from  
100 benthic and planktonic (size-fractionated) communities using an eukaryotic "universal" primer set to PCR amplify the V4 rDNA pre-barcode (Pawlowski *et al.*, 2012) from both DNA and RNA extracts. Previous studies using this sequencing dataset focused on particular taxonomic groups, such as uncultured MAST (Logares *et al.*, 2012), cercozoan amoebae (Berney *et al.*, 2013) or diatoms (Nanjappa *et al.*, 2014). More recently, we used a subset of  
105 the samples (23 planktonic RNA samples) and newly collected HTS reads (Illumina sequencing of the V9 18S rDNA region) to investigate the patterns of a particular community

property, the rare biosphere (Logares *et al.*, 2014). Here we analyze the complete 454 dataset from a taxonomic community perspective to address the following questions: How different are the protist communities found in the pico-, nano-, micro/mesoplankton and sediments?

110 Does the dissolved DNA fraction originate from particular taxonomic groups and/or organismal size-fractions? Do DNA and RNA surveys provide similar protist diversity profiles? Which taxonomic groups are differentially represented in either survey? Which groups dominate in each plankton organismal size fraction and associated sediments? Overall, our study highlights fundamental questions on the diversity of protists, an important but less  
115 known component of marine microbial ecosystems (Caron *et al.*, 2009).

## Results

As a product of the joint effort within the *BioMarkS* project we sampled six European coastal sites in the water column and sediments (Table 1). The thirteen planktonic communities were  
120 size fractionated (pico-, nano-, micro/mesoplankton and dissolved DNA) and, together with 7 benthic communities, used to obtain environmental DNA and RNA for pyrosequencing the V4 region of the 18S rDNA. Most assayed templates were successful, and we obtained 26 pyrosequenced samples for picoplankton, 26 for nanoplankton, 21 for micro/mesoplankton, 8 for dissolved DNA and 14 for sediments (Table 1). The pyrotag number for each 454-sample  
125 is detailed in Table S1 (~9000 pyrotags per sample on average). Pyrotags from these 95 samples clustered in 15,295 OTU<sub>97</sub> (Table S2), which were classified into 99 eukaryotic taxonomic groups (excluding unknown, metazoan and nucleomorphs). The broad coverage of this dataset across ecological and taxonomic scales provided a unique opportunity to study the diversity of both planktonic and benthic marine coastal protists.

We explored the reproducibility of sequencing outputs using a subset of nucleic acid extracts (11 DNA and 14 RNA) derived from three cruises (2 from Blanes, 11 from Naples-2009 and 12 from Oslo-2009) and including the four compartments (8 picoplankton, 7 nanoplankton, 5 micro/mesoplankton and 5 sediments). Each duplicated pair (same nucleic acid template and separate PCR and 454 reactions) was selected from the OTU table of 120 samples (Table S2), and linear regressions were performed by plotting OTU abundances in each pair. The pyrotag ratio between duplicates ranged from ~1 to 0.05 and was used to display the statistics of all regressions at once (Fig. 1). The  $R^2$  coefficients (Fig. 1A) were always high (0.89 on average) and independent of pyrotag ratios. This indicated that the same abundant OTUs were found at similar relative abundances in duplicates. The regression slopes were explained by pyrotag ratios ( $R^2$  of 0.92; slope of 0.84;  $p < 0.001$ ), indicating that OTU abundances increased proportionally with the number of pyrotags in the sample. Next we explored the pyrotags and OTUs shared between duplicates and calculated the shared percentages of the pair with fewer pyrotags (Fig. 1B). The amount of shared OTUs ranged from 32 to 89% and was higher at low pyrotag ratios, displaying a significant relationship ( $R^2 = 0.52$ ,  $p < 0.001$ ). Indeed, it was easier to find OTUs from the sample with less pyrotags in its duplicate with higher sequencing effort. The percentage of pyrotags included in the set of shared OTUs was always high (average of 91%) and its variation was poorly related to pyrotag ratios ( $p = 0.048$ ), indicating that OTUs appearing in only one duplicate contained few pyrotags. Thus, as expected, most differences between duplicates are found among the rare OTUs.

#### *Comparisons between communities*

$\beta$ -diversity patterns of protist assemblages were explored in a dendrogram with 92 samples. Structuring patterns revealed three main groups of samples: (i) picoplankton, nanoplankton

and dissolved DNA, (ii) micro/mesoplankton, and (iii) sediments (Fig. 2). Within the  
155 planktonic size fractions, samples from each sampling event grouped together (large grey dots  
in Fig. 2), regardless of water depth or nucleic acid template (only Naples-2010 did not form a  
cluster due to the atypical sample "DCM\_picoplankton\_DNA"). Planktonic samples from the  
same site collected at different seasons and years did not cluster together, indicating the  
significant impact of seasonal community changes. By contrast, sediment samples collected in  
160 different periods in Naples or Oslo clustered together, suggesting a more stable community in  
sediments than in plankton.

Protist community structure derived from DNA or RNA-based exhibited only minor  
differences. In total, 24 of the 41 DNA/RNA pairs appeared closest in the dendrogram (small  
black dots in Fig. 2). Most of the other pairs were still very close, but their coupling was less  
165 tight than that between pico- and nanoplankton from the same site, micro/mesoplankton from  
surface and DCM, or sediment cores from the same site. Only in two cases (the DCM  
picoplankton in Naples-2010 and Oslo-2010) the DNA and RNA samples were rather  
different, although still belonging to the same geographic cluster.

The dendrogram showed a close association of dissolved DNA samples with the pico- and  
170 nanoplankton from the same water body (Fig. 2). We then analyzed the overlap of individual  
OTUs amongst size fractions in the planktonic samples that had a complete sequence report of  
the four fractions in the DNA survey (Fig. 3). Many of the OTUs from dissolved samples  
(59% on average) were shared with the pico- and nanoplankton (alone, combined, or together  
with the micro/mesoplankton). Still, a substantial number of OTUs (35%) remained unique to  
175 dissolved samples, whereas only a few (6%) were shared with the micro/mesoplankton. We  
then computed the number of pyrotags from dissolved samples within each shared subarea  
(Fig. 3). OTUs shared with the small size fractions explained 89% of pyrotags while unique



OTUs accounted for 8% of pyrotags. Thus, the dissolved DNA composition largely derived from the pico- and nanoplankton fractions.

180 *Taxonomic representations derived from DNA and RNA surveys*

Relative abundances of taxonomic groups seen in DNA and RNA-based surveys were analyzed separately for picoplankton (13 DNA/RNA pairs), nanoplankton (13 pairs), micro/mesoplankton (10 pairs), and sediment (7 pairs) samples (Fig. 4). Within the picoplankton we focused in the 26 taxonomic groups more abundant (96.4% pyrotags on average; range of 92.1% to 99.2% in the 26 samples). Most were equally represented in DNA and RNA surveys, as shown by their placement near the 1:1 line (Fig. 4A). Two groups, MALV-I and MALV-II, were notably overrepresented in DNA surveys, averaging 15.4% and 19.7% of DNA pyrotags but only 1.8% and 2.0% of RNA pyrotags. Conversely, seven groups were overrepresented in RNA surveys. These RNA-prevalent groups were Prymnesiophyceae (averaged RNA/DNA pyrotag ratio of 7.4), MOCH-2 (6.0), Pelagophyceae (4.6), Telonema (2.6), Choanomonada (2.1), Ciliophora (2.0) and Chrysophyceae (1.6). In the nanoplankton, we identified 23 taxonomic groups that accounted for 97.1% of pyrotags (between 94.7% and 99.6% amongst samples), and many were equally represented in both surveys (Fig. 4B). As for the picoplankton fraction, MALV-I and -II were prevalent in nanoplankton DNA surveys (9.8% of DNA signal versus 2.3% in RNA), while virtually the same groups were prevalent in RNA surveys: Prymnesiophyceae (RNA/DNA pyrotag ratio of 8.6), MOCH-2 (5.4), Pelagophyceae (5.3), Choanomonada (3.6), Ciliophora (2.9), and Chrysophyceae (3.7). In the micro/mesoplankton, 9 taxonomic groups explained 98.6% of pyrotags (96.6% to 99.9%). Within this compartment MALV-I was overrepresented in DNA surveys and Acantharia showed the opposite trend (Fig. 4C). In sediments, 28 groups accounted for 95.8% of pyrotags (91.4 to 98.4%), and the DNA versus RNA-based taxonomic abundances were more biased than in pelagic samples (Fig. 4D). Seven groups were more abundant in DNA surveys

185  
190  
195  
200

(Apicomplexa, basal Fungi, Dinophyceae, MALV-I, MALV-II, Peronosporomycetes and RAD-B) while 4 were more abundant in RNA surveys (Bicosoecida, Chrysophyceae, Ciliophora and Tubulinea). In general, taxonomic groups exhibited a consistent RNA/DNA trend across the four major compartments analyzed.

*An overview of marine coastal protist diversity based on total RNA extracts*

We averaged the contribution of each taxonomic group for all samples within the 9 different combinations of planktonic size fractions, sediments, and nucleic acid templates (Fig. S1). Here we used the RNA survey to provide an overview of protist diversity within the different ecological compartments (Fig. 5), based on the 40 most prevalent taxonomic groups (which account for 97.7% pyrotags in picoplankton, 98.5% in nanoplankton, 99.4% in micro/mesoplankton and 95.2% in sediment samples). Within the picoplankton, Dinophyceae (25.3%), Ciliophora (11.9%) and Mamiellophyceae (9.3%) were the most abundant, followed by 19 additional groups with 1 to 8% relative abundance. The taxonomic composition of the nanoplankton was similar to that of the picoplankton, being dominated by Dinophyceae (39.3%), Diatomea (24.9%), and 13 additional groups at 1-8% abundance. Some contributed similarly in both size fractions, but others were clearly overrepresented in the picoplankton (Mamiellophyceae, Chrysophyceae, Pelagophyceae, Picozoa, Chlorarachniophyta, MAST-7, and MAST-4) or in the nanoplankton (Diatomea, Telonema, Katablepharidae and Trebouxiophyceae). The micro/mesoplankton was dominated by Acantharia (35.1%), Dinophyceae (28.9%) and Diatomea (22.4%), and only Ciliophora and MALV-IV displayed 1-8% abundance. Finally, sediments showed remarkable differences with the plankton. Diatomea (40.9%), Ciliophora (19.4%) and Monadofilosa (11.8%) were dominant in these samples, whereas 8 additional groups appeared at 1-8% abundances, seven of them being very rare in the plankton: basal Fungi, Labyrinthulomycetes, MAST-6, Granofilosea, Apusomonadida, Apicomplexa and Centrohelida.

### *Dominant OTUs in RNA surveys*

All previous analyses were done with OTUs clustered at 97% similarity, which sometimes  
230 enclose more than one species or genera. In our dataset, this was obvious within Dinophyceae  
(the most abundant OTU<sub>97</sub> retrieved 44 genera above 97% similarity), Diatomea and  
Cryptomonadales. Consequently, we used OTUs clustered at 99% similarity (a total of 63,365  
OTU<sub>99</sub>) to report the dominant taxa (>0.8% of pyrotags) from the RNA survey (Table 2).

Dominant OTU<sub>99</sub> were often 100% similar to described species. Picoplankton samples were  
235 the most distinct, containing small-sized genera like *Micromonas*, *Pelagomonas*, *Florenciella*,  
and *Minorisa* and several MAST and MALV phytotypes. They also included species within  
the Acantharia, Ciliophora and Dinophyceae known to belong to the microplankton, so their  
detection could be due to the presence of small life-cycle stages or to filtration artifacts.

Dominant OTU<sub>99</sub> within the other ecological compartments belonged mostly to Diatomea,  
240 Dinophyceae, Ciliophora and Acantharia. Of particular interest were the MALV-IV OTUs in  
the micro/mesoplankton (MALV-IV are crustaceans' parasites), and OTUs of uncultured basal  
Fungi, MAST-6 and Monadofilosa in sediments.

Picoplanktonic dominant OTU<sub>99</sub> tended to exhibit a high frequency across the investigated  
sites (Table 2). On average they occurred in 74% of the sites, while the occurrence of  
245 dominant OTUs in the other compartments was lower, with on average 65% of sites in the  
nanoplankton, 56% in the micro/mesoplankton and only 55% in sediments. On note, while the  
dominant OTU<sub>99</sub> of Ciliophora, Acantharia and Diatomea generally appeared in half of the  
samples, those of Dinophyceae were generally more widespread (86% of sites).

## 250 **Discussion**

Molecular surveys of marine protist diversity initiated in 2001 (Díez *et al.*, 2001; López-García *et al.*, 2001; Moon-van der Staay *et al.*, 2001), and developed together with the availability of new molecular tools including HTS (Amaral-Zettler *et al.*, 2009; Cheung *et al.*, 2010; Edgcomb *et al.*, 2011; Logares *et al.*, 2014). Previous surveys were partial, restricted to  
255 a cell-size fraction, a given geographic area, or a single nucleic acid template. We study here planktonic and benthic protist diversity inhabiting distant coastal places, from the smallest picoeukaryotes to larger colonial cells up to 2 mm in size (including also dissolved DNA), and derived from DNA and RNA templates. Given that the sampling plan was designed to maximize the ecological space covered, the emerging patterns are likely generalizable to other  
260 similar coastal locations (in terms of latitude and/or bathymetry). Seasonal studies at each site, required for a complete diversity description (Nolte *et al.*, 2010), were not addressed in our survey, but time-series HTS datasets are underway in most sites. Moreover, HTS studies using group-specific primers are being conducted to obtain a finer phylogenetic resolution of particular taxonomic groups (Egge *et al.*, 2015). Altogether, our study gives an improved  
265 view of marine protists diversity and clearly illustrates the differences between the taxonomic composition derived in DNA and RNA surveys.

### *Technical considerations*

There has been concern about the lack of replication in molecular surveys (Prosser, 2010; Robasky *et al.*, 2014), and our strategy was to test the reproducibility of OTU profiles in a set  
270 of 25 samples processed twice for the PCR and pyrosequencing steps. Duplicated pairs were usually very similar (range of  $R^2$  from 0.64 to 1.00) and systematically retrieved the same dominant OTUs at comparable relative abundances. Thus, our molecular surveys were well suited for obtaining robust  $\beta$ -diversity and taxonomic descriptions. Nevertheless, many low

abundant OTUs were found in only one pair, implying that undersampling was affecting the  
275 richness observed and/or the finding of different errors in each pair (Decelle *et al.*, 2014).

During the preparation of the OTU table for downstream analysis we removed unknown  
OTUs that contributed substantially to the OTU number (~13%) but little to the pyrotag  
number (~1%). This removed sequencing artifacts and chimeras while did not compromise  
our goal of a comprehensive protist description since they represented few pyrotags. Novel  
280 diversity was likely removed, however investigating novel branches on the tree of life was not  
the aim of this study and deserves a separate analysis. We also removed OTUs affiliating to  
metazoans, which contributed 1-10% of pyrotags in picoplankton and nanoplankton and about  
50% in micro/mesoplankton and sediment samples. The presence of metazoans in smaller  
fractions could be due to minute life-cycle stages but also to the breakdown of animals during  
285 filtration. The later certainly occurs in some cases, such as picoplankton OTUs affiliating to  
copepods (these have internal fecundation and never release gametes). So, although  
metazoans are worthy targets for molecular studies (Fonseca *et al.*, 2010), we preferred to  
exclude them from our analysis. Finally, our survey aimed to cover all eukaryotic lineages but  
it is important to keep in mind that universal primers may miss some relevant taxonomic  
290 groups. In our case, this negative selection was certain for Foraminifera, Prymnesiophyceae,  
and several excavate lineages.

#### *DNA versus RNA-based surveys of total marine protist diversity*

Comparing diversity surveys using environmental DNA or RNA is a common practice in  
microbial ecology (Weinbauer *et al.*, 2002). Initially, the rDNA/rRNA ratios measured from a  
295 given microbial population were proposed as proxies of *in situ* growth rates (Poulsen *et al.*,  
1993), but soon this moved to a more qualitative scenario where OTUs found in DNA surveys  
indicated species present, while OTUs found in RNA represented active species (Stoeck *et al.*,

2007; Jones and Lennon, 2010). Even this simplified view might not be universal, and differences in life histories, life strategies, and non-growth activities among species might  
300 confound this interpretation (Blazewicz *et al.*, 2013). Nevertheless, protistologists have generally accepted this scenario, and RNA surveys were recently used to identify the active protist community (Stoeck *et al.*, 2007; Not *et al.*, 2009; Lejzerowicz *et al.*, 2013). However, there is an additional factor that may severely affect rDNA/rRNA comparisons in eukaryotes, which is the large variation (up to orders of magnitude) in the rDNA copy number among  
305 species (Zhu *et al.*, 2005; Weber and Pawlowski, 2013).

In our dataset, community structure derived from DNA or RNA templates was similar, as paired samples clustered very close in the dendrogram and many taxonomic groups exhibited similar relative abundances. Among the groups with differential abundance, the most extreme were MALV-I and MALV-II that dominated picoplankton DNA surveys as in other studies  
310 (Massana, 2011) and contributed little to RNA surveys. One explanation for this discrepancy is that the MALV I-II parasites are abundant in the picoplankton as dispersal, relatively inactive stages with few ribosomes. However, FISH (Fluorescence In situ Hybridization) direct cell counts showed relative abundances much lower than in DNA surveys (Siano *et al.*, 2010), so a most plausible explanation is that MALV I-II cells have a higher genomic rDNA  
315 copy number than other picoeukaryotes. Similarly, groups more abundant in RNA surveys may have lower rDNA copy numbers. Given the consistent trends found across size-fractions, our DNA/RNA ratios could reflect different genomic architectures among taxonomic groups, and not necessarily different relative activities. At any rate, RNA surveys should provide a better representation of *in situ* protist biomass and diversity, since labile RNA likely derives  
320 from living cells, in contrast with DNA that can be preserved in dead cells or in the dissolved extracellular pool (Karl and Bailiff, 2012). Our analysis of the eukaryotic diversity derived from dissolved DNA samples showed that, in contrast to what was found in anoxic deep-sea

sediments (Corinaldesi *et al.*, 2011), these samples did not contain a particular phylogenetic signature. Dissolved DNA samples from these coastal sites resembled the pico- and  
325 nanoplankton from the same seawater, suggesting cell breakage during filtration.

*Marine protist diversity across organismal size-fractions and habitats*

Our analysis confirms a well-known observation that planktonic and benthic protists are different (Lee and Patterson, 1998; Bik *et al.*, 2012). This was evident in the  $\beta$ -diversity analysis as well as in the taxonomic list, where 13 out of 40 groups were much more abundant  
330 in sediments than in the water column. Sediments have also been considered seed banks of planktonic biodiversity, out of which taxa can re-emerge into the plankton at appropriate seasons (Satta *et al.*, 2010). Our data is consistent with this view, since dominant OTUs in the plankton generally were also found, at lower abundances, in sediments.

Regarding planktonic assemblages, our data show a broad differentiation among larger  
335 (micro/mesoplankton) and smaller (pico- and nanoplankton) protists. This could be due partly to the different collection protocols (net tows and pressure filtration, respectively), but also from the known different cell size of given species. Within the micro/mesoplankton, only four taxonomic groups dominated, Acantharia, Dinophyceae, Diatomea and Ciliophora, and we did not find other relevant groups, consistent with classical inverted microscopy inspections  
340 of the plankton. At the other end of the size spectrum, many taxonomic groups were identified, highlighting the high-rank diversity of small marine protists (Massana, 2011).

Picoplankton and nanoplankton assemblages from the same seawater were always related, but a closer inspection revealed marked differences in the relative abundances of key groups. The diversity of both large and small protists was very different in each coastal site, so a standard  
345 protist community could not be delineated. Samples from the same site but different seasons

were also unrelated, indicating the impact of seasonal successions and highlighting that one or few samples cannot represent the complete diversity at one site.

Molecular surveys of protist diversity have often used size-fractionated samples to discern among different size classes. This step may introduce artifacts, derived from cell breakage and retention of smaller fragments in the smaller size fraction, or from the retention of smaller cells in filters whose larger pores have been obstructed after processing large sample volumes. Our study has ample evidences of the first case, since about 40% of picoplankton pyrotags comes from the four dominating micro/mesoplankton groups. Although part of this signal might derive from gametes or spores (Amato *et al.*, 2007; Kimoto *et al.*, 2011), the most plausible explanation is that larger cells from these groups (typically >10  $\mu\text{m}$ ) are broken during the filtration (Sørensen *et al.*, 2013). This bias could have been even more dramatic had we used a smaller pore-size filter (such as 0.2  $\mu\text{m}$ ) for picoplankton collection (Sørensen *et al.*, 2013). Picoeukaryote diversity excluding this microplanktonic signal would be composed by a set of photosynthetic groups (58% of pyrotags), heterotrophic groups (16%), MALV clades (15%) and MAST clades (11%). Following this reasoning, it is expected that nanoplanktonic cells are also represented in the picoplankton (such as Cryptomonadales). The second case of filtration artifact, smaller cells retained in larger size fractions, is not obvious from our data, and groups known to be picoeukaryotes (like Mamiellophyceae, MAST-4 or MAST-7) are only minority in the nanoplankton. So, our data show clear evidence of larger cells collected in smaller size fractions but little support of the other way around. Filtration artifacts need to be considered when interpreting molecular surveys, and cell sizes should be established by direct cell observations by FISH (Siano *et al.*, 2010, Massana, 2011).



### *Concluding remarks*

This study presents a description of marine coastal protist diversity obtained from 95  
370 independent HTS samples. Our data show that larger protists affiliate to a few taxonomic  
groups well studied by microscopy for decades, whereas smaller protists, known as "small  
flagellates", include a wide variety of less-known taxonomic groups. Our results provide  
answers to the initial questions, with a special emphasis on (i) the general similarity between  
DNA and RNA surveys despite clear biases in few groups, (ii) the taxonomic discontinuity  
375 between micro/mesoplankton and the two smaller size fractions, (iii) the phylogenetic  
signature of dissolved DNA deriving from the smallest cells in the sample, and (iv) the  
distinct composition of planktonic and benthic communities. Our data also suggests a higher  
seasonal variability in the plankton compared to the benthos. These patterns emerging from  
the large spatial sampling strategy adopted here would likely apply to other coastal sites,  
380 while open ocean assemblages require similar approaches (de Vargas *et al.*, 2015). In the  
future, diversity studies based on sequencing surveys, complementary FISH counts and SAGs  
(Single Amplified Genomes) analyses will contribute to infer the genetic potential and  
ecological roles of the key protist players.

## 385 **Experimental Procedures**

### *Sampling*

Marine samples were collected through the *BioMarKs* project (<http://biomarks.eu/>) in coastal  
sites near Blanes (Balearic Sea, Spain), Gijon (Gulf of Biscay, Spain), Naples (Tyrrhenian  
Sea, Italy), Oslo (Skagerrak, Oslofjorden, Norway), Roscoff (Western English Channel,  
390 France) and Varna (Black Sea, Bulgaria) (Table 1). Some sites are long-term observatories,

the Blanes Bay Microbial Observatory (BBMO), the Long Term Ecological Research station MareChiara (Naples), and the SOMLIT-Astan site (Roscoff). Seawater temperature, salinity and chlorophyll *a* concentration were recorded with CTD (conductivity-temperature-depth) and fluorometer sensors. Water samples were collected with Niskin bottles at the sea surface  
395 in all sites (and at the DCM [deep chlorophyll maximum] in a few sites), and passed through a 20  $\mu\text{m}$  metallic sieve. The nanoplankton (3-20  $\mu\text{m}$ ) and picoplankton (0.8-3  $\mu\text{m}$ ) fractions were sampled from ~20 liters of seawater filtered sequentially with a peristaltic pump through 3  $\mu\text{m}$  and 0.8  $\mu\text{m}$  polycarbonate (PC) membranes (142 mm diameter) for less than 40 min to minimize RNA degradation. For dissolved DNA, 20 liters of 0.2  $\mu\text{m}$ -filtered seawater were  
400 mixed with 400 mL of 0.5% CTAB (pH=8) for 5 h and filtered onto 0.2  $\mu\text{m}$  PC membranes (142 mm). A plankton net of 20- $\mu\text{m}$  mesh-size was towed for 5-15 min and the large protists collected were rinsed with 0.2  $\mu\text{m}$  filtered seawater, passed through a 2000  $\mu\text{m}$  metallic sieve and filtered on 12  $\mu\text{m}$  PC membranes (47 mm), to collect the micro- (20-200  $\mu\text{m}$ ) and meso- (200-2000  $\mu\text{m}$ ) planktonic fractions (micro/mesoplankton). Filters were flash-frozen and  
405 stored at -80° C until processed. Finally, sediment cores were taken with a Multicorer sampler or by Scuba divers and kept at -80°C until processed.

#### *Nucleic acid extraction and pyrosequencing*

Total DNA and RNA were extracted simultaneously from a complete filter using the NucleoSpin® RNA kit (Macherey-Nagel) or from 2.5 g of surface sediment (~1 cm upper  
410 layer) using the Power Soil RNA kit (MoBio). Extracts were quantified using a Nanodrop ND-1000 Spectrophotometer and checked on a 1.5% agarose gel. Contaminating DNA was removed from RNA extracts using the Turbo DNA-free kit (Ambion). Complete DNA removal was verified by PCR using eukaryotic primers, and in the few cases of positive PCR we did a second DNase treatment. One hundred ng of extracted RNA were immediately

415 reverse transcribed to cDNA using the RT Superscript III random primers kit (Invitrogen).  
Both DNA extracts and cDNA products were kept at -80°C until processed.

Environmental DNA or cDNA were used as templates for PCR amplification of the V4 region of the 18S rDNA (~380 bp) using primers TAREuk454FWD1 and TAREukREV3 (Stoeck *et al.*, 2010) that amplify most eukaryotic groups. The forward primer had a barcode and both  
420 primers were adapted for 454 sequencing. PCR reactions (25 µl) contained 1x Master Mix Phusion® High-Fidelity DNA Polymerase (Finnzymes), 0.35 µM of each primer, 3% DMSO, and 5 ng of DNA or cDNA. The PCR program had an initial denaturation step at 98°C during 30 sec, 10 cycles of 10 sec at 98°C, 30 sec at 53°C and 30 sec at 72°C, then 15 similar cycles but with 48°C annealing temperature, and a final step at 72°C for 10 min. PCR triplicates  
425 were purified and eluted (30 µl) with NucleoSpin® Gel and PCR Clean-Up kit (Macherey-Nagel), and quantified with the Quant-It™ PicoGreen® dsDNA Assay kit (Invitrogen). About 1 µg of pooled amplicons were sent to Genoscope (<http://www.genoscope.cns.fr>, France) for pyrosequencing on a 454 GS FLX Titanium system (454 Life Sciences, USA). The complete sequencing dataset is available at the European Nucleotide Archive (ENA) under the study  
430 accession number PRJEB9133 (<http://www.ebi.ac.uk/ena/data/view/PRJEB9133>).

#### *Pyrotag processing and OTU tables*

Sequences obtained through 454 sequencing (pyrotags) were demultiplexed using the barcode identifier in the forward primer. Pyrotags 150-600 bp long, with exact primer sequences, and with homopolymers no longer than 8 bases, were retained. Identical pyrotags within each  
435 sample were dereplicated to keep a single representative sequence. For quality check, errors were computed in sliding windows of 50 bp and pyrotags containing a window with an error >1% and appearing only once in the dataset were removed. Chimera check was run with UCHIME (Edgar *et al.*, 2011), using *de novo* and reference-based chimera searches against

the SILVA108 release (Quast *et al.*, 2013). Additional chimera searches were done using  
440 ChimeraSlayer (Haas *et al.*, 2011) and SILVA108. A local BLAST search (Altschul *et al.*,  
1990) against SILVA108 was used to exclude 16S rDNA prokaryotic or plastidial sequences.

This initial procedure yielded ~1.3 million curated pyrotags derived from 120 samples (95  
distinct samples, 25 of them in duplicate [same nucleic acid extract but separate PCR and  
sequencing reactions]). These were clustered into OTUs (Operational Taxonomic Units) at  
445 97% similarity (OTU<sub>97</sub>) with USEARCH (Edgar, 2010). A local BLAST was done to  
compare the representative sequence of each OTU (the most abundant one) against the  
GenBank Release 183.0. Distant OTUs having an e-value  $>10^{-100}$  (below ~85% similarity)  
were considered as "unknown" and removed. Taxonomic OTU assignment was performed by  
best hit BLAST against two reference databases, PR<sup>2</sup> (Guillou *et al.*, 2013) and a smaller and  
450 phylogenetically fully-validated database (Pernice *et al.*, 2013). Metazoans and nucleomorphs  
were removed, leaving OTUs classified into 99 eukaryotic groups, including 64 described  
groups generally at class level (Adl *et al.*, 2012), 27 environmental ribogroups (Guillou *et al.*,  
2008; Massana *et al.*, 2014), and 8 unidentified categories within each supergroup. The initial  
OTU table (120 samples) was used to extract duplicated samples. A second table with the  
455 distinct 95 samples was prepared for the general diversity analyses. A third table (only the 44  
RNA samples) was used for the taxonomic description (see Table S2 for the number of OTUs  
and pyrotags within these tables).

For beta diversity analyses, three samples with few 300-500 pyrotags (Table S1) were  
removed from the OTU table and the remaining 92 samples were subsampled to 1402  
460 pyrotags using the tool "rrarefy" of the Vegan R package (Oksanen *et al.*, 2008). Pyrotag  
counts were log transformed to diminish the effect of the most abundant OTUs, and the table  
subsampled again with rrarefy (after multiplying the log-transformed values times 1000) to  
obtain the same signal per sample. A distance matrix was computed with the Bray-Curtis

index and a dendrogram was constructed using the UPGMA function in Vegan. Venn  
465 diagrams were generated with the R package VennDiagram.

## Acknowledgments

Financial support has been provided by the European project *BioMarkS* (2008-6530, ERA-net  
Biodiversa, EU), and state-based projects FLAME (CGL2010-16304, MICINN, Spain) to  
470 RM, OCEANOMICS (ANR-11-BTBR-0008, France) to CdV, grant 31003A-125372 (Swiss  
National Foundation) to JP, and RCN grant 190307 (Norway) to BE.

## Conflict of Interest

The authors declare no conflict of interest.

475

## References

- Adl, S.M., Simpson, A.G.B, Lane, C.E., Lukes, J., Bass, D., Bowser, S.S., *et al.* (2012) The revised  
classification of eukaryotes. *J Eukaryot Microbiol* **59**: 429-493.
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990) Basic local alignment search tool. *J*  
480 *Mol Biol* **215**: 403-410.
- Amaral-Zettler, L.A., McCliment, E.A., Ducklow, H.W., and Huse, S.M. (2009) A method for studying protistan  
diversity using massively parallel sequencing of V9 hypervariable regions of small-subunit ribosomal RNA  
genes. *PLoS ONE* **4**: e6372.

- Amato, A., Kooistra, W.H.C.F., Ghiron, J.H.L., Mann, D.G., Pröschold, T., and Montresor, M. (2007)  
485 Reproductive isolation among sympatric cryptic species in marine diatoms. *Protist* **158**: 193-207.
- Bates, S.T., Clemente, J.C., Flores, G.E., Walters, W.A., Parfrey, L.W., Knight, R., and Fierer, N. (2013) Global  
biogeography of highly diverse protistan communities in soil. *ISME J* **7**: 652-659.
- Berney, C., Romac, S., Mahé, F., Santini, S., Siano, R., and Bass, D. (2013) Vampires in the oceans: predatory  
cercozoan amoebae in marine habitats. *ISME J* **7**: 2387-2399.
- 490 Bik, H.M., Sung, W., De Ley, P., Baldwin, J.G., Sharma, J., Rocha-Olivares, A., and Thomas, W.K. (2012)  
Metagenetic community analysis of microbial eukaryotes illuminates biogeographic patterns in deep-sea and  
shallow water sediments. *Mol Ecol* **21**: 1048-1059.
- Blazewicz, S.J., Barnard, R.L., Daly, R.A., and Firestone, M.K. (2013) Evaluating rRNA as an indicator of  
microbial activity in environmental communities: limitations and uses. *ISME J* **7**: 2061- 2068.
- 495 Burki, F. (2014) The eukaryotic tree of life from a global phylogenomic perspective. *Cold Spring Harb Perspect  
Biol* **6**: a016147.
- Caron, D.A., Worden, A.Z., Countway, P.D., Demir, E., and Heidelberg, K.B. (2009) Protists are microbes too:  
a perspective. *ISME J* **3**: 4-12.
- Cheung, M.K., Au, C.H., Chu, K.H., Kwan, H.S., and Wong, C.K. (2010) Composition and genetic diversity of  
500 picoeukaryotes in subtropical coastal waters as revealed by 454 pyrosequencing. *ISME J* **4**: 1053-59.
- Corinaldesi, C., Barucca, M., Luna, G.M., and Dell'Anno, A. (2011) Preservation, origin and genetic imprint of  
extracellular DNA in permanently anoxic deep-sea sediments. *Mol Ecol* **20**: 642-654.
- Decelle, J., Romac, S., Sasaki, E., Not, F., and Mahé, F. (2014) Intracellular diversity of the V4 and V9 regions  
of the 18S rRNA in marine protists (Radiolarians) assessed by high-throughput sequencing. *PLoS ONE* **9**:  
505 e104297.
- de Vargas, C., Audic, S., Henry, N., Decelle, J., Mahé, F., Logares, R., *et al.* (2015) Eukaryotic plankton diversity  
in the sunlit ocean. *Science* **348**: 1261605-1-11.

- Díez, B., Pedrós-Alió, C., and Massana, R. (2001) Study of genetic diversity of eukaryotic picoplankton in different oceanic regions by small-subunit rRNA gene cloning and sequencing. *Appl Environ Microbiol* **67**: 2932-2941.
- 510
- Dolan, J.R., Landry, M.R., and Ritchie, M.E. (2013) The species-rich assemblages of tintinnids (marine planktonic protists) are structured by mouth size. *ISME J* **7**: 1237-1243.
- Edgar, R.C. (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**: 2460-2461.
- 515
- Edgar, R.C., Haas, B.J., Clemente, J.C., Quince, C., and Knight, R. (2011) UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* **27**: 2194–2200.
- Edgcomb, V., Orsi, W., Bunge, J., Jeon, S., Christen, R., Leslin, C., *et al.* (2011) Protistan microbial observatory in the Cariaco Basin, Caribbean. I. Pyrosequencing vs. Sanger insights into species richness. *ISME J* **5**: 1344-1356.
- 520
- Egge, E.S., Johannessen, T.V., Andersen, T., Eikrem, W., Bittner, L., Larsen, A., *et al.* (2015) Seasonal diversity and dynamics of haptophytes in the Skagerrak, Norway, explored by high-throughput sequencing. *Mol Ecol* doi: 10.1111/mec.13160.
- Fonseca, V.G., Carvalho, G.R., Sung, W., Johnson, H.F., Power, D.M., Neill, S.P., *et al.* (2010) Second-generation environmental sequencing unmasks marine metazoan biodiversity. *Nat Commun* **1**: 98.
- 525
- Graham, J.M., Kent, A.D., Lauster, G.H., Yannarell, A.C., Graham, L.E., and Triplett, E.W. (2004) Seasonal dynamics of phytoplankton and planktonic protozoan communities in a Northern temperate humic lake: Diversity in a dinoflagellate dominated system. *Microb Ecol* **48**: 528-540.
- Guillou, L., Viprey, M., Chambouvet, A., Welsh, R.M., Kirkham, A.R., Massana, R., *et al.* (2008) Widespread occurrence and genetic diversity of marine parasitoids belonging to *Syndiniales* (*Alveolata*). *Environ Microbiol*
- 530
- 10**: 397-408.
- Guillou, L., Bachar, D., Audic, S., Bass, D., Berney, C., Bittner, L., *et al.* (2013) The Protist Ribosomal Reference database (PR<sup>2</sup>): a catalog of unicellular eukaryote Small SubUnit rRNA sequences with curated taxonomy. *Nucleic Acids Res* **41**: D597-D604.

- Haas, B.J., Gevers, D., Earl, A.M., Feldgarden, M., Ward, D.V., Giannoukos, G., *et al.* (2011) Chimeric 16S  
535 rRNA sequence formation and detection in Sanger and 454-pyrosequenced PCR amplicons. *Genome Res* **21**:  
494-504.
- He, L., Liu, F., Kauppih, V., Ren, Y., and Li, Z. (2014) Comparisons of the fungal and protistan communities  
among different marine sponge holobionts by pyrosequencing. *Microb Ecol* **67**: 951-961.
- Jones, S.E., and Lennon, J.T. (2010) Dormancy contributes to the maintenance of microbial diversity. *Proc Natl*  
540 *Acad Sci USA* **107**: 5881- 5886.
- Karl, D.M., and Bailiff, M.D. (2012) The measurement and distribution of dissolved nucleic acids in aquatic  
environments. *Limnol Oceanogr* **34**: 543–558.
- Keeling, P.J., Burger, G., Durnford, D.G., Lang, B.F., Lee, R.W., Pearlmans, R.E., *et al.* (2005) The tree of  
eukaryotes. *Trends Ecol Evol* **20**: 670-676.
- 545 Kimoto, K., Yuasa, T., and Takahashi, O. (2011) Molecular identification of reproductive cells released from  
*Cyprina irregularis* Nigrini (Radiolaria). *Environ Microbiol Rep* **3**: 86–90.
- Lee, W.J., and Patterson, D.J. (1998) Diversity and geographic distribution of free-living heterotrophic  
flagellates - Analysis by PRIMER. *Protist* **149**: 229-244.
- Lejzerowicz, F., Voltsky, I., and Pawlowski, J. (2013) Identifying active foraminifera in the Sea of Japan using  
550 metatranscriptomic approach. *Deep Sea Res II* **86-87**: 214-220.
- Logares, R., Audic, S., Santini, S., Pernice, M.C., de Vargas, C., and Massana, R. (2012) Diversity patterns and  
activity of uncultured marine heterotrophic flagellates unveiled with pyrosequencing. *ISME J* **6**: 1823-1833.
- Logares, R., Audic, S., Bass, D., Bittner, L., Boutte, C., Christen, R., *et al.* (2014) Patterns of rare and abundant  
marine microbial eukaryotes. *Curr Biol* **24**: 813-821.
- 555 López-García, P., Rodríguez-Valera, F., Pedrós-Alió, C., and Moreira, D. (2001) Unexpected diversity of small  
eukaryotes in deep-sea Antarctic plankton. *Nature* **409**: 603-607.



- Mangot, J.-F., Domaizon, I., Talb, N., Marouni, N., Duffaud, E., Bronner, G., and Debroas, D. (2013) Short-term dynamics of diversity patterns: evidence of continual reassembly within lacustrine small eukaryotes. *Environ Microbiol* **15**: 1745-1758.
- 560 Massana, R., Castresana, J., Balagué, V., Guillou, L., Romari, K., Groisillier, A., *et al.* (2004) Phylogenetic and ecological analysis of novel marine stramenopiles. *Appl Environ Microbiol* **70**: 3528-3534.
- Massana, R. (2011) Eukaryotic picoplankton in surface oceans. *Annu Rev Microbiol* **65**: 91-110.
- Massana, R., del Campo, J., Sieracki, M.E., Audic, S., and Logares, R. (2014) Exploring the uncultured microeukaryotic majority in the oceans: reevaluation of ribogroups within stramenopiles. *ISME J.* **8**: 854-866.
- 565 Moon-van der Staay, S.Y., De Wachter, R., and Vaultot, D. (2001) Oceanic 18S rDNA sequences from picoplankton reveal unsuspected eukaryotic diversity. *Nature* **409**: 607-610.
- Nanjappa, D., Audic, S., Romac, S., Kooistra, W.H.C.F., and Zingone, A. (2014) Assessment of species diversity and distribution of an ancient diatom lineage using a DNA metabarcoding approach. *PLoS ONE* **9**: e103810.
- 570 Nolte, V., Pandey, R.V., Jost, S., Medinger, R., Ottenwälder, B., Boenigk, J., and Schlötterer, C. (2010) Contrasting seasonal niche separation between rare and abundant taxa conceals the extent of protist diversity. *Mol Ecol* **19**: 2908-2915.
- Not, F., Valentin, K., Romari, K., Lovejoy, C., Massana, R., Töbe, K., *et al.* (2007) Picobiliphytes: A marine picoplanktonic algal group with unknown affinities to other eukaryotes. *Science* **315**: 252-254.
- 575 Not, F., del Campo, J., Balagué, V., de Vargas, C., and Massana, R. (2009) New insights into the diversity of marine picoeukaryotes. *PLoS ONE* **4**: e7143.
- Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Simpson, G.L., Solymos, P., *et al.* (2008) Vegan: community ecology package. R package version 1.15-0 (Finland: University of Oulu).
- Olguín, H.F., Boltovskoy, D., Lange, C.B., and Brandini, F. (2006) Distribution of spring phytoplankton (mainly diatoms) in the upper 50 m of the Southwestern Atlantic Ocean (30-61°S). *J Plankton Res* **28**: 1107-1128.
- 580

- Pawlowski, J., Audic, S., Adl, S., Bass, D., Belbahri, L., Berney, C., *et al.* (2012) CBOL Protist Working Group: Barcoding eukaryotic richness beyond animal, plant, and fungal kingdoms. *PLoS Biol* **10**: e1001419.
- Pedrós-Alió, C. (2006) Marine microbial diversity: can it be determined? *Trends Microbiol* **14**: 257–263.
- 585 Pernice, M.C., Logares, R., Guillou, L., and Massana, R. (2013) General patterns of diversity in major marine microeukaryote lineages. *PLoS ONE* **8**: e57170.
- Pizay, M.-D., Lemée, R., Simon, N., Cras, A.-L., Laugier, J.-P., and Dolan, J.R. (2009) Night and day morphologies in a planktonic dinoflagellate. *Protist* **160**: 565-575.
- Poulsen, L.K., Ballard, G., and Stahl, D.A. (1993) Use of rRNA fluorescence in situ hybridization for measuring the activity of single cells in young and established biofilms. *Appl Environ Microbiol* **59**: 1354–1360.
- 590 Prosser, J.I. (2010) Replicate or lie. *Environ Microbiol* **12**: 1806-1810.
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., *et al.* (2013) The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res* **41**: D590-D596.
- Robasky, K., Lewis, N.E., and Church, G.M. (2014) The role of replicates for error mitigation in next-generation sequencing. *Nature Rev Genet* **15**: 56–62.
- 595 Satta, C.T., Anglès, S., Garcés, E., Lugliè, A., Padedda, B.M., and Sechi, N. (2010) Dinoflagellate cysts in recent sediments from two semi-enclosed areas of the Western Mediterranean Sea subject to high human impact. *Deep-Sea Res II* **57**: 256-267.
- Schaechter, M. (2012) *Eukaryotic Microbes* (editor). Elsevier, Oxford, UK.
- 600 Seenivasan, R., Sausen, N., Medlin, L.K., and Melkonian, M. (2013) *Picomonas judraskeda* Gen. et Sp. Nov.: The first identified member of the Picozoa phylum Nov., a widespread group of piceoekaryotes, formerly known as 'Picobiliphytes'. *PLoS ONE* **8**: e59565.
- Sherr, B.F., Sherr, E.B., Caron, D.A., Vaulot, D., and Worden, A.Z. (2007) Oceanic protists. *Oceanography* **20**: 130-134.

- 605 Siano, R., Alves-de-Souza, A., Foulon, E., Bendif, E.M., Simon, N., Guillou, L., and Not, F. (2010) Distribution and host diversity of Amoeboophryidae parasites across oligotrophic waters of the Mediterranean Sea. *Biogeosci Discuss* **7**: 7391-7419.
- Sørensen, N., Daugbjerg, N., and Richardson K. (2013) Choice of pore size can introduce artefacts when filtering picoeukaryotes for molecular biodiversity surveys. *Microb Ecol* **65**: 964-968.
- 610 Stoeck, T., Zuendorf, A., Breiner, H.-W., and Behnke, A. (2007) A molecular approach to identify active microbes in environmental eukaryote clone libraries. *Microb Ecol* **53**: 328-339.
- Stoeck, T., Bass, D., Nebel, M., Christen, R., Jones, M.D., Breiner, H.W., and Richards, T.A. (2010) Multiple marker parallel tag environmental DNA sequencing reveals a highly complex eukaryotic community in marine anoxic water. *Mol Ecol* **19 Suppl 1**: 21-31.
- 615 Weber, A., and Pawlowski, J. (2013) Can abundance of protists be inferred from sequence data? A case study of cultured Foraminifera. *PLoS ONE* **8**: e56739.
- Weinbauer, M.G., Fritz, I., Wenderoth, D.F., and Höfle, M.G. (2002) Simultaneous extraction from bacterioplankton of total RNA and DNA suitable for quantitative structure and function analyses. *Appl Environ Microbiol* **68**: 1082–1087.
- 620 Zhu, F., Massana, R., Not, F., Marie, D., and Vaultot, D. (2005) Mapping of picoeukaryotes in marine ecosystems with quantitative PCR of the 18S rRNA gene. *FEMS Microbiol Ecol* **52**: 79-92.

## Table legends

625 **Table 1.** Coastal marine sites visited and water column (surface and DCM) and sediment samples taken for investigating global protist diversity by pyrosequencing. Each planktonic community is analyzed in up to seven 454-samples (pico, nano and micro/mesoplankton by RNA/DNA plus dissolved DNA) and each sediment sample in two (by RNA/DNA). Cases with no sequencing results are noted. See Table S1 for details on the number of pyrotags for each 454-sample.

630 **Table 2.** Averaged relative abundance (% tags), occurrence (number of times detected), and taxonomic affiliation of the most abundant OTU<sub>99</sub> derived from the RNA survey in the four ecological compartments. Relative abundances in the other compartments (Pic= Picoplankton; Nan= Nanoplankton; Mic= Micro/mesoplankton; Sed= Sediments) are also shown (in grey when >0.8%).

635

## Figure legends

**Fig. 1.** Summary of the comparison of duplicated 454-samples (n=25) shown in relation to the pyrotag ratio between duplicates. A. Statistics of the linear regressions between the OTU abundance in each duplicate:  $R^2$  coefficients (white dots) and slope values (black dots; regression line shown). B. Percentage of OTUs shared in duplicate pairs (black dots; regression line shown) and percentage of pyrotags within these shared OTUs (white dots). These later percentages apply to the pair with less pyrotags.

**Fig. 2.** UPGMA dendrogram based on the Bray-Curtis dissimilarity matrix from an OTU table with 92 samples (subsamped to 1402 pyrotags and log-transformed). Sample names have four components: sampling cruise, depth, size fraction, and nucleic acid template. The three main clusters are highlighted. Large grey dots indicate grouping of samples from the same sampling event, while small black dots indicate a closest relation of the same assemblage analyzed by DNA and RNA.

**Fig. 3.** Number of shared OTUs among plankton size-fractions derived from the DNA survey in five cases. Areas covering the dissolved samples are shaded. Below the number of shared OTUs, the percentages of pyrotags they represent in dissolved samples are shown.

**Fig. 4.** Relative pyrotag abundance in RNA and DNA surveys of taxonomic groups in picoplankton (A), nanoplankton (B), micro/mesoplankton (C) and sediment (D) samples. Each point represents paired percentages (in logarithmic scales) of a given taxonomic group in a given sample. Groups selected are those with highest pyrotag abundance (together account for 96-98% of pyrotags). Groups overrepresented in DNA surveys appear above the 1:1 line as colored triangles, those overrepresented in RNA surveys appear below the line as colored circles, and those equally represented appear as small black circles.

**Fig. 5.** Percentages of pyrotags (average and standard error) of taxonomic groups in RNA  
660 surveys in the picoplankton (13 samples), nanoplankton (13 samples), micro/mesoplankton  
(11 samples) and sediments (7 samples). Note the different vertical scale in the four plots.  
Groups are ordered by their overall pyrotag abundance in the whole dataset.

665

For Peer Review Only

## Supporting Information

**Table S1.** Number of pyrotags per 454-sample in the global survey of protist diversity in the plankton (several organismal size-fractions) and sediments analyzed in DNA and RNA surveys (total of 95 samples). See Table 1 for additional information on sampling cruises.

670 **Table S2.** Number of OTUs and pyrotags included in OTU97 tables, showing the initial numbers and the final numbers after excluding unknown, metazoan, and nucleomorph OTUs. Three OTU tables are used in this study: 120 samples (all datasets including duplicates), 95 samples (duplicates removed), and 44 samples (only the RNA survey).

**Fig. S1.** Percentages of pyrotags (average and standard error) of taxonomic groups in RNA  
675 and DNA surveys in the picoplankton, nanoplankton, micro/mesoplankton, sediments and the dissolved fraction (the latter only DNA survey). Note the different vertical scale in the four plots. Groups are ordered by their overall pyrotag abundance in the whole dataset.




Fig. 1. Summary of the comparison of duplicated 454-samples (n=25) shown in relation to the pyrotag ratio between duplicates. A. Statistics of the linear regressions between the OTU abundance in each duplicate: R2 coefficients (white dots) and slope values (black dots; regression line shown). B. Percentage of OTUs shared in duplicate pairs (black dots; regression line shown) and percentage of pyrotags within these shared OTUs (white dots). These later percentages apply to the pair with less pyrotags.

133x141mm (300 x 300 DPI)






Fig. 2. UPGMA dendrogram based on the Bray-Curtis dissimilarity matrix from an OTU table with 92 samples (subsamped to 1402 pyrotags and log-transformed). Sample names have four components: sampling cruise, depth, size fraction, and nucleic acid template. The three main clusters are highlighted. Large grey dots indicate grouping of samples from the same sampling event, while small black dots indicate a closest relation of the same assemblage analyzed by DNA and RNA.  
252x571mm (300 x 300 DPI)






Fig. 3. Number of shared OTUs among plankton size-fractions derived from the DNA survey in five cases. Areas covering the dissolved samples are shaded. Below the number of shared OTUs, the percentages of pyrotags they represent in dissolved samples are shown.  
159x188mm (300 x 300 DPI)



161x134mm (300 x 300 DPI)

www.Only



302x425mm (300 x 300 DPI)

**Table 1.** Coastal marine sites visited and water column (surface and DCM) and sediment samples taken for investigating global protist diversity by pyrosequencing. Each planktonic community is analyzed in up to seven 454-samples (pico, nano and micro/mesoplankton by RNA/DNA plus dissolved DNA) and each sediment sample in two (by RNA/DNA). Cases with no sequencing results are noted. See Table S1 for details on the number of pyrotags for each 454-sample.

Site	Coordinates	Date	Depth (m)	Temperature (°C)	Salinity	Chl a ( $\mu\text{g L}^{-1}$ )	454-samples	Pyrotags
<b>Blanes</b>	41°40'N, 2°48'E	9/02/2010	Surface (1)	12.5	37.6	0.7	7	28569
			Sediment (20)	12.6	37.8	-	2	3899
<b>Gijon</b>	43°40'N; 5°35'W	14/09/2010	Surface (1)	20.2	35.7	0.6	4 <sup>a,b,c</sup>	49747
<b>Naples</b>	40°48'N, 14°15'E	13/10/2009	Surface (1)	22.8	37.7	1.7	6 <sup>b</sup>	100567
			DCM (26)	19.2	37.9	1.5	7	82327
			Sediment (78)	14.6	37.9	-	2	20545
		14/05/2010	Surface (1)	19.2	37.2	1.1	6 <sup>c</sup>	31185
			DCM (34)	15.5	37.7	1.0	6 <sup>c</sup>	49402
			Sediment (78)	14.0	37.9	-	2	9680
<b>Oslo</b>	59°16'N, 10°43'E	22/09/2009	Surface (1)	15.5	25.2	2.5	6 <sup>c</sup>	61963
			DCM (20)	16.1	29.2	1.1	6 <sup>c</sup>	70006
			Sediment-1 (103)	8.2	35.0	-	2	10974
			Sediment-2 (103)	8.2	35.0	-	2	12661
		22/06/2010	Surface (1)	15.0	21.5	1.1	7	56190
			DCM (10)	11.9	29.5	1.9	5 <sup>a,b</sup>	61067
			Sediment (103)	6.0	35.0	-	2	10480
<b>Roscoff</b>	48°46'N, 3°57'W	20/04/2010	Surface (1)	9.9	34.9	0.2	7	33142
			Sediment (60)	9.9	34.9	-	2	3122
<b>Varna</b>	43°10'N, 28°50'E	27/05/2010	Surface (3)	18.0	16.5	5.2	7	60352
			DCM (40)	8.7	17.9	6.1	7	84620

<sup>a</sup> No results for Micro/mesoplankton-RNA; <sup>b</sup> No results for Micro/mesoplankton-DNA; <sup>c</sup> No results for Dissolved-DNA

**Table 2.** Averaged relative abundance (% tags), occurrence (number of times detected), and taxonomic affiliation of the most abundant OTU<sub>99</sub> derived from the RNA survey in the four ecological compartments. Relative abundances in the other compartments (Pic= Picoplankton; Nan= Nanoplankton; Mic= Micro/mesoplankton; Sed= Sediments) are also shown (in grey when >0.8%).

<b>A. Picoplankton samples (n=13)</b>									
OTU#	% tags	Occurrence	Closest match to described species	%	Taxonomic group	Nan	Mic	Sed	
13	4.5	8	<i>Micromonas pusilla</i> CCMP1195	100	Mamiellophyceae	0.2	0.0	0.0	
1	3.2	7	<i>Acanthostaurus purpurascens</i>	100	Acantharia	1.6	25.3	0.0	
10	2.2	11	<i>Teleaulax amphioxeia</i>	100	Cryptomonadales	1.8	0.0	0.0	
36	1.6	6	<i>Pelagomonas calceolata</i>	100	Pelagophyceae	0.3	0.0	-	
51	1.6	10	<i>Micromonas pusilla</i> RCC829	99.5	Mamiellophyceae	0.1	-	0.0	
5	1.5	4	<i>Lynnella semiglobulosa</i>	100	Ciliophora	0.1	0.0	0.0	
14	1.4	11	<i>Gyrodinium fusiforme</i>	100	Dinophyceae	1.3	0.9	0.0	
28	1.2	4	<i>Polykrikos kofoidii</i>	100	Dinophyceae	0.2	0.2	0.2	
46	1.2	10	<i>Pseudotontonia simplicidens</i>	99.7	Ciliophora	0.4	0.0	-	
22	1.2	13	<i>Katodinium rotundatum</i>	100	Dinophyceae	1.5	0.0	0.0	
44	1.1	12	<i>Brachidinium capitatum</i>	93.2	MALV-III	0.7	0.0	-	
26	1.1	13	<i>Azadinium concinnum</i>	99.7	Dinophyceae	3.2	0.9	0.0	
73	1.1	12	<i>Florenciella parvula</i>	100	Dictyochophyceae	0.4	0.0	-	
9	1.0	10	<i>Lepidodinium viride</i>	100	Dinophyceae	2.4	0.2	-	
50	0.9	12	<i>Teleaulax gracilis</i>	100	Cryptomonadales	0.6	-	-	
107	0.9	11	<i>Pirsonia verrucosa</i>	86.3	MAST-7B	0.1	-	-	
32	0.9	10	<i>Strombidium</i> sp. SNB99-2	97.3	Ciliophora	1.2	0.0	0.0	
87	0.9	13	<i>Prorocentrum triestinum</i>	91.6	MALV-III	0.3	-	-	
145	0.8	7	<i>Minorisa minuta</i>	100	Chlorarachniophyta	0.1	0.0	-	
<b>B. Nanoplankton samples (n=13)</b>									
OTU#	% tags	Occurrence	Closest match to described species	%	Taxonomic group	Pic	Mic	Sed	
0	4.8	6	<i>Leptocylindrus aporus</i>	100	Diatomea	0.4	0.4	0.1	
26	3.2	13	<i>Azadinium concinnum</i>	99.7	Dinophyceae	1.1	0.9	0.0	
9	2.4	11	<i>Lepidodinium viride</i>	100	Dinophyceae	1.0	0.2	-	
3	2.4	3	<i>Bacterosira bathyomphala</i>	100	Diatomea	0.0	0.5	0.9	
10	1.8	10	<i>Teleaulax amphioxeia</i>	100	Cryptomonadales	2.2	0.0	0.0	
1	1.6	8	<i>Acanthostaurus purpurascens</i>	100	Acantharia	3.2	25.3	0.0	
22	1.5	13	<i>Katodinium rotundatum</i>	100	Dinophyceae	1.2	0.0	0.0	
14	1.3	12	<i>Gyrodinium fusiforme</i>	100	Dinophyceae	1.4	0.9	0.0	
32	1.2	9	<i>Strombidium</i> sp. SNB99-2	97.3	Ciliophora	0.9	0.0	0.0	
12	1.2	6	<i>Skeletonema marinoi</i>	100	Diatomea	0.0	0.1	3.6	
174	1.1	3	<i>Thalassiosira profunda</i>	99.7	Diatomea	0.0	0.1	1.1	
15	1.0	8	<i>Chaetoceros setoense</i>	100	Diatomea	0.2	0.4	1.2	
<b>C. Micro/mesoplankton samples (n=11)</b>									
OTU#	% tags	Occurrence	Closest match to described species	%	Taxonomic group	Pic	Nan	Sed	
1	25.3	7	<i>Acanthostaurus purpurascens</i>	100	Acantharia	3.2	1.6	0.0	
17	6.0	5	<i>Noctiluca scintillans</i>	100	Dinophyceae	0.0	0.0	-	
53	3.2	9	<i>Neoceratium fusus</i>	100	Dinophyceae	0.0	0.0	-	
24	3.1	5	<i>Skeletonema pseudocostatum</i>	100	Diatomea	0.1	0.4	0.9	
63	2.2	10	<i>Neoceratium azoricum</i>	99.5	Dinophyceae	0.0	0.0	-	
18	2.0	2	<i>Favella markusovszkyi</i>	100	Ciliophora	-	-	-	

77	1.7	4	<i>Hexaconus serratus</i>	98.9	Acantharia	0.1	0.0	-
23	1.5	6	<i>Biddulphia alternans</i>	86.1	Diatomea	0.1	0.0	0.8
271	1.5	4	<i>Hematodinium</i> sp. ex Nephrops	95.3	MALV-IV	0.1	0.0	0.0
52	1.4	11	<i>Neoceratium furca</i>	100	Dinophyceae	0.0	0.0	-
54	1.4	5	<i>Xiphacantha alata</i>	100	Acantharia	0.0	0.1	0.0
191	1.2	2	<i>Stenosemella pacifica</i>	100	Ciliophora	0.0	-	-
284	1.1	1	<i>Amphorides quadrilineata</i>	98.9	Ciliophora	0.0	-	-
283	1.1	6	<i>Thalassiosira rotula</i>	100	Diatomea	0.0	0.1	0.2
173	1.0	11	<i>Dinophysis acuminata</i>	100	Dinophyceae	0.0	0.0	-
26	0.9	11	<i>Azadinium concinnum</i>	99.7	Dinophyceae	1.1	3.2	0.0
14	0.9	9	<i>Gyrodinium fusiforme</i>	100	Dinophyceae	1.4	1.3	0.0
385	0.8	2	<i>Thalassiosira anguste-lineata</i>	99.7	Diatomea	-	0.0	0.1
161	0.8	8	<i>Syndinium</i> sp. ex Corycaeus	100	MALV-IV	0.1	0.0	-

#### D. Sediment samples (n=7)

OTU#	% tags	Occurrence	Closest match to described species	%	Taxonomic group	Pic	Nan	Mic
12	3.6	3	<i>Skeletonema marinoi</i>	100	Diatomea	0.0	1.2	0.1
30	2.4	3	<i>Chaetoceros</i> cf. <i>neogracile</i>	100	Diatomea	0.0	0.0	0.0
57	2.2	3	<i>Powellomycetaceae</i> sp.	85.7	Basal fungi	0.0	0.1	0.1
275	1.3	1	<i>Pirsonia formosa</i>	89.4	MAST-6	-	-	-
33	1.3	3	<i>Monodinium</i> sp.	98.7	Ciliophora	0.4	0.0	0.0
249	1.3	3	<i>Protaspis obliqua</i>	93.6	Monadofilosa	-	-	-
15	1.2	5	<i>Chaetoceros setoense</i>	100	Diatomea	0.2	1.0	0.4
623	1.1	6	<i>Psammodictyon</i> sp.	99.7	Diatomea	-	0.0	0.0
174	1.1	6	<i>Thalassiosira profunda</i>	99.7	Diatomea	0.0	1.1	0.1
24	0.9	4	<i>Skeletonema pseudocostatum</i>	100	Diatomea	0.1	0.4	3.1
3	0.9	5	<i>Bacterosira bathyomphala</i>	100	Diatomea	0.0	2.4	0.5