



**HAL**  
open science

## Interplay between the genetic clades of *Micromonas* and their viruses in the Western English Channel

Anne-Claire Baudoux, H Lebretonchel, H Dehmer, M Latimier, R Edern, Fabienne Rigaut-Jalabert, P Ge, Laure Guillou, Elodie Foulon, Yves Bozec, et al.

### ► To cite this version:

Anne-Claire Baudoux, H Lebretonchel, H Dehmer, M Latimier, R Edern, et al.. Interplay between the genetic clades of *Micromonas* and their viruses in the Western English Channel. *Environmental Microbiology Reports*, 2015, 7 (5), pp.765-773. 10.1111/1758-2229.12309 . hal-01196410

**HAL Id: hal-01196410**

<https://hal.sorbonne-universite.fr/hal-01196410v1>

Submitted on 9 Sep 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Interplay between the genetic clades of *Micromonas* and their viruses in the Western**  
2 **English Channel**

3

4 A-C. Baudoux<sup>1,2\*</sup>, H. Lebreton<sup>3,4</sup>, H. Dehmer<sup>1,2</sup>, M. Latimier<sup>1,2</sup>, R. Edern<sup>1,2</sup>, F. Rigaut-  
5 Jalabert<sup>1,2</sup>, P. Ge<sup>1,2</sup>, L. Guillou<sup>1,2</sup>, E. Foulon<sup>1,2</sup>, Y. Bozec<sup>1,2</sup>, T. Cariou<sup>1,2</sup>, Y. Desdevises<sup>3,4</sup>, E.  
6 Derelle<sup>3,4</sup>, N. Grimsley<sup>3,4</sup>, H. Moreau<sup>3,4</sup>, and N. Simon<sup>1,2</sup>

7

8 <sup>1</sup> Sorbonne University, UPMC Paris 06, Station Biologique de Roscoff, UMR7144, Roscoff,  
9 France

10 <sup>2</sup> CNRS, Station Biologique de Roscoff, UMR7144 and FR2424, Roscoff, France

11 <sup>3</sup> Sorbonne University, UPMC Paris 06, Observatoire Océanologique de Banyuls-sur-mer,  
12 UMR7232, Banyuls-sur-mer, France

13 <sup>4</sup> CNRS, Observatoire Océanologique de Banyuls-sur-mer, UMR7232, Banyuls-sur-mer,  
14 France

15

16 Running title: *Micromonas* – virus interactions

17

18 For correspondence. Email [acbaudoux@sb-roscoff.fr](mailto:acbaudoux@sb-roscoff.fr); Tel (+33)2 98 29 25 37; Fax (+33)2 98

19 29 23 24

20

21

22 **Abstract**

23 The genus *Micromonas* comprises distinct genetic clades that commonly dominate eukaryotic  
24 phytoplankton community from polar to tropical waters. This phytoplankter is also recurrently  
25 infected by abundant and genetically diverse prasinoviruses. Here we report on the interplay  
26 between prasinoviruses and *Micromonas* with regards to the genetic diversity of this host.  
27 During one year, we monitored the abundance of 3 clades of *Micromonas* and their viruses in  
28 the Western English Channel both in the environment, using clade-specific probes and flow  
29 cytometry, and in the laboratory, using clonal isolates of *Micromonas* clades to assay for their  
30 viruses by plaque-forming units. We showed that the seasonal fluctuations of *Micromonas*  
31 clades were closely mirrored by the abundances of their corresponding viruses, indicating that  
32 the members of *Micromonas* genus are susceptible to viral infection, regardless of their  
33 genetic affiliation. The characterization of 45 viral strains revealed that *Micromonas* clades  
34 are attacked by specific virus populations , which exhibit distinctive host clade specificity,  
35 life history traits, and genetic diversity. However, some viruses can also cross-infect different  
36 host clades suggesting a mechanism of horizontal gene transfer within the genus *Micromonas*  
37 . This study provides novel insights into the impact of viral infection for the ecology and  
38 evolution of the prominent phytoplankter *Micromonas*.

39

40 Key words: Mamiellophyceae, Phycodnaviridae, Prasinovirus, host specificities, life history  
41 traits, DNA polymerase, English Channel

42

## 43 **Introduction**

44 Viruses are undoubtedly the most abundant biological entities in the ocean and they  
45 intimately interact with every facet of the marine biosphere. Through these interactions,  
46 viruses profoundly influence the global biogeochemical cycles by altering the structure and  
47 the function of marine communities and by contributing to the cycling of major elements (for  
48 review see Suttle, 2005, 2007; Sime-Ngando, 2014). In spite of their global-scale  
49 implications, the nature and the dynamics of virus-host interactions in marine environments  
50 are far from understood and many basic, yet fundamental, questions remain unsolved: Who  
51 are the hosts infected by marine viruses? What are the infection strategies evolved by viruses?  
52 How do virus-host interactions vary in time and space? In this respect, characterizing the  
53 interplay of ecologically relevant virus-host model systems is a prerequisite for advancing our  
54 understanding of virus impacts in nature.

55 Viruses that infect the cosmopolitan green picoalga *Micromonas* were first reported by  
56 (Pienaar, 1976) and they were found to co-occur with their hosts in a wide range of marine  
57 environments, from temperate to cold waters (e.g., Cottrell and Suttle, 1991, 1995a; Sahlsten,  
58 1998; Zingone *et al.*, 1999). The majority of known *Micromonas* viruses belong to the genus  
59 *Prasinovirus* in the Phycodnaviridae family. Virions are tailless, they exhibit icosahedral  
60 capsids of 100 -125 nm in diameter and dsDNA genome of approx. 200 kb (Mayer and Taylor  
61 1979; Cottrell and Suttle 1991; Zingone *et al.*, 1999). Yet, these prasinoviruses have variable  
62 tolerance to chloroform (Martinez *et al.* 2015) suggesting ultrastructural divergence and they  
63 exhibit significant level of genetic variation (Cottrell and Suttle, 1995b). These viruses  
64 display a marked seasonal dynamics (Sahlsten, 1998; Zingone *et al.*, 1999) and they were  
65 shown to cause considerable amount of mortality in their host population (Cottrell and Suttle,  
66 1995a; Evans *et al.*, 2003). *Micromonas* strains show a great variability with respect to their  
67 susceptibility to infection, indicating that viruses do not only quantitatively but also

68 qualitatively regulate their host populations (Sahlsten, 1998; Zingone *et al.*, 1999, 2006). So  
69 far, the infection patterns and processes underlying these complex interactions are poorly  
70 understood. Of particular interest, the relation between the susceptibility to infection and the  
71 genetic diversity of *Micromonas* has seldom been investigated (Zingone *et al.*, 2006).

72 The ubiquitous genus *Micromonas* is genetically diverse and comprises an assemblage  
73 of 3 (Guillou *et al.*, 2004) to 5 (Šlapeta *et al.*, 2006; Worden, 2006) discrete phylogenetic  
74 clades (or lineages) of flagellated cells that correspond, at least, to 3 distinct species (Simon,  
75 unpublished). These clades are often sympatric in marine ecosystems (Foulon *et al.*, 2008;  
76 Šlapeta *et al.*, 2006). Yet, their relative contribution to total *Micromonas* abundance varies in  
77 time and space, suggesting that they occupy specific niches (Foulon *et al.*, 2008). The factors  
78 that regulate this clade dynamics are not clearly understood. In all likelihood, *Micromonas*  
79 clades exhibit differential responses to abiotic factors but they might also respond differently  
80 to predation risks, including those imposed by viruses.

81 To test whether *Micromonas* clades display differential susceptibility to virus  
82 infection, we combined field and laboratory experiments on 45 novel viral isolates in order to  
83 characterize the interactions between the genetic clades of *Micromonas* and their viruses in  
84 the Western English Channel (WEC) throughout the year 2009. The WEC constitutes an ideal  
85 study site where *Micromonas* is known to dominate the picophytoplankton community (Not *et*  
86 *al.*, 2004). The three main *Micromonas* genetic types, designated as clades [A.ABC.12],  
87 [B.E.3], and [C.D.5] (Worden, 2006) and here referred to as clades A, B, and C, respectively,  
88 are recorded year round in this ecosystem (Foulon *et al.*, 2008). Our study revealed that  
89 *Micromonas* clades interact with specific viral populations that display distinctive dynamics  
90 and life history traits.

91

## 92 **Results and Discussion**

93

#### 94 **Physico-chemistry at the sampling station**

95 Strong tidal mixing produces a permanently mixed water column at the long term  
96 monitoring coastal station SOMLIT-Astan (48°46'N, 3°58'W, Marrec *et al.*, 2013). During the  
97 sampling period (February to December 2009), the water temperature progressively increased  
98 from 8.8°C in February to a maximum of 16.9°C in September while salinity varied between  
99 34.8 (March) and 35.3 (September). Nutrient dynamics varied according to the classical  
100 pattern observed in this area (Not *et al.*, 2004). Phosphate and nitrate concentrations showed  
101 comparable dynamics with minima recorded in summer (0.16 and 0.7 µM, respectively) and  
102 maxima during the winter period (0.53 and 11 µM, respectively). Detailed dynamics of  
103 biogeochemical variables at SOMLIT-Astan are publicly available on the observatory website  
104 (<http://somlit-db.epoc.u-bordeaux1.fr/bdd.php>).

105

#### 106 **Dynamics of *Micromonas* clades in the Western English Channel**

107 The phytoplankton community at SOMLIT-Astan during this period was numerically  
108 dominated by picophytoplanktonic cells, which comprised, on average, 90% of the total  
109 community in 2009 (data not shown). Among these, the abundance of the picoeukaryotes  
110 varied between 2.1 and  $16.5 \times 10^3$  cells mL<sup>-1</sup> with *Micromonas* spp. accounting for 20% to  
111 80% of the total counts as shown by a combination of flow cytometry (FCM) and TSA-FISH  
112 analyses (Fig. 1A). As reported previously, this prominent genus displayed marked seasonal  
113 dynamics with major peaks of abundance recorded in late spring ( $6.6 \times 10^3$  cells mL<sup>-1</sup>), early  
114 summer ( $6.3 \times 10^3$  cells mL<sup>-1</sup>), and mid-autumn ( $4.4 \times 10^3$  cells mL<sup>-1</sup>) (Foulon *et al.*, 2008;  
115 Not *et al.*, 2004).

116 The 3 main *Micromonas* genetic clades (hereafter referred to as clades A, B, and C)  
117 exhibit recurring seasonal dynamics in the WEC according to previous time series (Foulon *et*

118 *al.*, 2008). We confirmed this pattern of succession in 2009 (Fig. 1B). *Micromonas* clade A  
119 numerically dominated the bloom from February to mid-May comprising 63 to 77% of the  
120 total counts. From June to December, both genetic types A and B equally contributed to  
121 *Micromonas* counts, although clade B cells showed a sudden, yet unexplained, drop in  
122 abundance on June 16. Cells that belonged to clade C were the least abundant accounting, on  
123 average, for  $7 \pm 6\%$  of the total counts with somewhat higher contribution during the winter  
124 period (14%) than the remainder of the year (5%).

125         The ecological processes that influence the shifts in the relative abundance of  
126 *Micromonas* clades are unclear. It is traditionally believed that global-scale distribution and  
127 diversity of species are mostly driven by ability to withstand abiotic controls (Wiens et al.  
128 2011). Previous study on the global distribution of *Micromonas* clades suggested that clade B  
129 tend to thrive in warmer coastal waters whereas clade C mostly occur during low-light  
130 conditions (Foulon *et al.*, 2008). At local spatial extents, it is however well accepted that  
131 species distribution is also influenced by biotic controls (Wiens et al. 2011). It is thus likely  
132 that besides eco-physiological adaptations, biotic interactions also regulate the observed  
133 *Micromonas* clade dynamics in the Western English Channel.

#### 134 **Dynamics of *Micromonas* viruses in the Western English Channel**

135         Viruses that infect *Micromonas* (hereafter referred to as MicV), mainly prasinoviruses,  
136 have been observed and isolated from different oceanic regions (Cottrell and Suttle, 1991,  
137 1995a; Sahlsten, 1998; Zingone *et al.*, 1999; Short and Short, 2008). FCM analyses of  
138 cultured isolates showed that these prasinoviruses consistently cluster within a well-defined  
139 population based on their nucleic acid fluorescence (upon Sybr green I staining) and side  
140 scatter properties (Fig. S1). In the WEC, this specific population accounted for a small but  
141 significant proportion (on average 10%) of the total virus counts and displayed a marked  
142 seasonal dynamics with concentration ranging from  $3.4 \times 10^5 \text{ mL}^{-1}$  in February to  $8.8 \times 10^5$

143 mL<sup>-1</sup> in July 2009 (Fig. 2A). This FCM population probably does not exclusively comprise  
144 *Micromonas* viruses, but counts of this viral population covary strikingly well with  
145 *Micromonas* abundances ( $r=0.675$ ,  $n=20$ ,  $p<0.005$ ), suggesting that viruses might perhaps  
146 control abundance of this genus during the sampling period.

147 To further investigate this hypothesis, we quantified infectious MicV in the WEC by  
148 plaque assay using cultures of each 3 *Micromonas* clades. Significant differences in the  
149 number of PFU were detected depending on the phylotype of the host culture (Fig. 2B). As  
150 also recorded for their hosts, PFU obtained on *Micromonas* clade A and B (hereafter, PFU-A  
151 and PFU-B, respectively) were substantially more abundant than PFU formed on clade C-  
152 hosts (hereafter PFU-C). During the sampling period, PFU-A abundance increased  
153 concomitantly with the development of *Micromonas* clade A. Their maximum abundance ( $8$   
154  $\times 10^2$  virus mL<sup>-1</sup>) was observed 2 weeks after the peak of host abundance (Figs. 1B and 2B).  
155 The monitoring of PFU-B showed low concentration during the winter-spring period but their  
156 dynamics was tightly coupled to their host abundance from the summer period until the end of  
157 the year, reaching a peak of  $11 \times 10^2$  virus mL<sup>-1</sup> in autumn. Regarding PFU-C, infectious  
158 particles were detected year round but their abundance was 10 to 100-fold lower than PFU-A  
159 and -B with no clear relation to their specific host dynamics. Because of variable strain  
160 susceptibilities to viral infection, it is difficult to compare our results to former PFU  
161 monitoring on *Micromonas* host cell clades although maximal abundance were within the  
162 same range ( $1.0 - 4.6 \times 10^3$  PFU mL<sup>-1</sup>, Cottrell and Suttle, 1991; Sahlsten, 1998; Zingone *et*  
163 *al.*, 1999). For the same reason, this assay most likely underestimates the global abundance of  
164 viruses that infect each *Micromonas* phylotype. Nonetheless, it provides unequivocal  
165 evidence that *Micromonas*, regardless of their genetic clade, were subject to viral attack year  
166 round in the WEC.

167



168 ***Micromonas* virus - host interactions**

169 The above results led us to question whether specific virus populations are associated to  
170 *Micromonas* clades and, if so, how these virus populations interact with their hosts. To  
171 tentatively address this question, 45 viruses (14-16 per host clade) were isolated throughout  
172 the sampling period and they were characterized. These viruses were selected randomly from  
173 a collection of 176 viral isolates obtained from plaques on *Micromonas* clade A, B, and C  
174 along the year 2009. Regardless of the host on which they were isolated, MicVs exhibited  
175 sizeable icosahedral capsids (110 to 130 nm in diameter) as determined by negative staining  
176 using transmission electron microscopy, and large dsDNA genome, as suggested by their flow  
177 cytometry signature upon Sybr green I staining (Fig. S1). We therefore assigned our isolates  
178 to the virus family Phycodnaviridae and the genus *Prasinovirus*.

179

180 *Host specificities* - The host range of these 45 prasinoviruses was then determined by pair-  
181 wise infection using 14 strains belonging to the 3 *Micromonas* clades (Table 1, Table S1). The  
182 large majority of our isolates (35 MicVs) were strictly clade-specific. They infected host  
183 strains that belonged only to the clade of their initial host. The remaining 10 viruses could  
184 propagate at comparable yield on hosts belonging to clades A and B. Six viruses (out of 15)  
185 isolated on a clade A strain could also infect clade B strains, and 4 viruses (out of 14) isolated  
186 on a clade B strain could also infect clade A strains. Within each main pattern of specificity  
187 (A, B, AB, and C), a remarkable clonal diversity was observed with 26 unique specificity  
188 profiles that included variable virus infection types ranging from generalist to specialist.

189 In order to test the structure of this infection network, we applied a network-based  
190 analysis according to Flores *et al.* (2011) and Weitz *et al.* (2013). As observed previously for  
191 phage-bacteria infection networks (Flores *et al.*, 2011), the structure of *Micromonas* - virus  
192 interactions was statistically modular. This statistical analysis applied at the whole matrix

193 scale and intra-module scale discriminated 3 modules, which comprised the viruses isolated  
194 on hosts from clade A, clade B, and clade C, respectively (Fig. S2). Altogether, these results  
195 suggest that *Micromonas* clade A, B, and C interact with distinct, yet diversified, viral  
196 populations. The observed variability in strain-specificity indicates that viruses influence the  
197 intraspecific diversity within each *Micromonas* clades. We also showed that host switches can  
198 occur between the more closely related host clades (those that belong to clades A and B),  
199 suggesting that viruses could also promote *Micromonas* diversity through horizontal gene  
200 transfers within this genus.

201

202 *Virus infection strategies* - One-step growth experiments conducted on 4 to 5 representative  
203 viruses per host clade indicated a similar clade-specific grouping of our isolates. This assay  
204 provided estimates of the virus burst size and latent period, which can be related to the virus  
205 life strategy. By analogy to the theory of the r- and K-selection (MacArthur, 1967),  
206 opportunistic (the most r-selected) viruses are those that exhibit short generation times and  
207 high burst sizes whereas less virulent (the most K-selected) viruses are those that induce low  
208 mortality in their host population (Suttle, 2007) and they propagate through latent or chronic  
209 infection as observed for specific *Ostreococcus* viruses (Thomas *et al.*, 2011). Based on these  
210 parameters, we detected divergent life strategies in *Micromonas* viruses depending on the host  
211 clades that they attack (Fig. 3). The viruses isolated on clade B hosts tended to be the most  
212 virulent with short latent period and relatively high progeny production, regardless of their  
213 infection range types (i.e. specialist or generalist) whereas the viruses isolated on clade C  
214 strains tended to be the least virulent with generally extended latent period (up to 30 h) and  
215 moderate to high burst size (Fig. 3). Viruses isolated on clade A hosts exhibited an  
216 intermediate phenotype with moderate latent period and low burst sizes.

217 In contrast to past theories (trade-offs hypothesis, (Poulin, 1998)), generalism (i.e.,  
218 viruses that propagate on several host clades) was not associated with any compensatory  
219 effect on the burst size or on the latent period. The infection kinetics of the selected viral  
220 isolates does not correlate to the growth rates of their host (Fig. S3). , The observed  
221 differences in infection strategies could be explained by the dynamics of their respective host  
222 clade in the studied area. For example, *Micromonas* clade C is present year round at low  
223 abundance. It is thus tempting to speculate that viruses specific to this host clade had to  
224 evolve long latent periods to be maintained within the host population and fairly high progeny  
225 production to increase the probability of encounter with their host (Abedon *et al.*, 2001). By  
226 contrast, *Micromonas* clade B displays a marked seasonal dynamics in the WEC with high  
227 abundance recorded between mid-June and end-September. We could thus expect that  
228 corresponding viruses had to evolve an opportunist lifestyle to efficiently exploit their  
229 transient resources.

230

### 231 **Phylogenetic relationships between *Micromonas* viruses**

232 To investigate the phylogenetic relationships between these viral isolates, we  
233 amplified a fragment of the gene encoding their DNA polymerase (*polB*, Chen and Suttle,  
234 1996, Table S2). This sequence is well conserved among Phycodnaviridae, yet there is  
235 sufficient sequence variability to build phylogenetic relationships between viruses from the  
236 same genus (Bellec *et al.*, 2009). In spite of their affiliation to the Phycodnaviridae family, the  
237 amplification of the *polB* gene using the AVS1/AVS2 primers failed for 8 of the 45 selected  
238 isolates including B-, C- and AB-specific viruses. The reason for this failure is unclear. We  
239 cannot exclude that the sequence targeted by the degenerated primers might be too divergent.  
240 It is also possible that these polymerases carry an intein that makes the PCR product too long  
241 to amplify easily under the tested conditions (Clerissi *et al.*, 2013).

242           The partial *polB* sequences of the remaining 37 isolates fell into 29 haplotypes (Fig.  
243 4). The large majority of these haplotypes (24 out of 29) were new, i.e. different from *polB*  
244 sequences in public databases (BLAST analyses, <http://www.ncbi.nlm.nih.gov/>). In most  
245 cases, viral isolates that belonged to the same haplotype exhibited different specificity  
246 patterns, suggesting that *polB* sequencing underestimates the functional diversity of  
247 phycodnaviruses as reported previously (Baudoux and Brussaard, 2005; Bellec *et al.*, 2009;  
248 Clerissi *et al.*, 2012). In the phylogenetic trees, *polB* sequences of viruses infecting clade C  
249 hosts formed a well-supported cluster while viruses that infect clade A and clade B hosts  
250 distributed into multiple branches suggesting a higher level of genetic diversity. Interestingly,  
251 regardless of the host strain they infect, viruses that were highly specific (that infect one  
252 single strain among all strains tested) fell apart from other viruses (i.e. RCC4232 and  
253 RCC4240, 4242, 4243, Fig. 4 and Table 1). The prasinoviruses that infect *Ostreococcus tauri*  
254 showed a similar phylogenetic pattern and tended to cluster with counterparts that exhibit  
255 similar infection range types (Clerissi *et al.*, 2012).

256           This recurrent correlation between *polB* phylogeny and the number of hosts infected  
257 by a given prasinovirus is somewhat puzzling. The *polB* gene is indeed used as a neutral  
258 marker of evolution in Phycodnaviridae (Chen and Suttle, 1996; Dunigan *et al.*, 2006), that is,  
259 this gene is well-conserved among this virus family whereas specificity markers are expected  
260 to be more variable genes (Clerissi *et al.*, 2012). Furthermore, virus-host specificity is  
261 typically determined during events upstream the virus DNA replication. Several studies  
262 reported that the virus attachment on host cell surface determines the susceptibility of marine  
263 plankton to infection (e.g., Tarutani *et al.*, 2006; Mizumoto *et al.*, 2007; Stoddard *et al.*,  
264 2007). In other words, the evolution of resistance to viral infection is mostly due to changes in  
265 host surface properties that limit viral attachment. Recently, Thomas *et al.* (2011) evidenced  
266 an intriguing mode of resistance in the prasinophyte *Ostreococcus tauri*. This study

267 demonstrated that prasinoviruses could still attach on resistant strains but they did not produce  
268 viral progeny. It is thus likely that the resistance mechanism was not due to a change in the  
269 host receptor but rather occurred at the DNA entry or DNA replication stage, in which the  
270 *polB* gene product would be involved. Alternatively, viral attachment and internalisation  
271 might be a two-stage process as in reoviruses (Reiter *et al.*, 2011). The notion that DNA  
272 replication machinery might be involved in host specificity is of course highly speculative but  
273 the repeatedly observed link between *polB* clustering and prasinovirus infectivity range  
274 certainly deserves appropriate investigation.

275

## 276 **Concluding remarks**

277 In the Western English Channel, a complex assemblage of viruses is associated to the  
278 dominating picophytoplankton species *Micromonas spp.* Our study revealed, for the first time,  
279 that the main *Micromonas* genetic clades (clade A, clade B, and clade C) are attacked by  
280 specific virus populations which displayed distinctive dynamics and life history traits. The  
281 apparently high variability in *Micromonas* virus specificities suggests that viruses maintain a  
282 high genetic diversity within each of these clades. However, host switching can occur  
283 (particularly for hosts that belong to the clade A and B) suggesting that virus could also  
284 influence *Micromonas* diversity through horizontal gene transfers. The complete genome  
285 sequencing of representative viruses and hosts should provide insightful information on the  
286 extent of these putative genetic exchanges. It is of course critical to keep in mind that our  
287 model-based approach only reflects a partial picture of the actual diversity of virus-host  
288 interactions. Still, these approaches provide invaluable information to improve our  
289 understanding of the mechanisms of virus-host interactions as well as the evolution and  
290 outcomes of these interactions in natural environments.

291

292 **Acknowledgments**

293 The Captain and the crew of R.V. Neomysis are acknowledged for their excellent assistance  
294 onboard. We thank the “Service d'Observation en Milieu Littoral (SOMLIT), INSU-CNRS”  
295 from Station Biologique de Roscoff for the data on the biogeochemistry and biology at  
296 ASTAN. We are grateful to Sophie Lepanse from the Microscopy core facilities, Morgan  
297 Perennou from the GENOMER core facilities at the Station Biologique de Roscoff and David  
298 Demory for stimulating discussions. We thank the two anonymous reviewers for their  
299 constructive comments on a previous version of this manuscript. This research was funded by  
300 the ANR funding agency (PICOVIR; grant no. BLAN07-1\_200218 and REVIREC; grant no.  
301 12-BSV7-0006-01).

302

303 **References**

- 304 Abedon, S. T., Herschler, T. D., and Stopar, D. (2001). Bacteriophage latent-period evolution  
305 as a response to resource availability. *Appl Environ Microbiol* 67: 4233-4241.
- 306 Baudoux, A. C., and Brussaard, C. P. D. (2005). Characterization of different viruses  
307 infecting the marine harmful algal bloom species *Phaeocystis globosa*. *Virology* 341: 80-  
308 90.
- 309 Bellec, L., Grimsley, N., Moreau, H., and Desdevises, Y. (2009). Phylogenetic analysis of  
310 new Prasinoviruses (Phycodnaviridae) that infect the green unicellular algae  
311 *Ostreococcus*, *Bathycoccus* and *Micromonas*. *Environ Microbiol Reports* 1: 114-123
- 312 Brussaard, C. P. D. (2004). Optimization of procedures for counting viruses by flow  
313 cytometry. *Appl Environ Microbiol* 70: 1506-1513.
- 314 Chen, F., and Suttle, C. A. (1995). Amplification of DNA polymerase gene fragments from  
315 viruses infecting microalgae. *Appl Environ Microbiol* 61: 1274–1278.

316 Chen, F., and Suttle, C. A. (1996). Evolutionary relationships among large double-stranded  
317 DNA viruses that infect microalgae and other organisms as inferred from DNA  
318 polymerase genes. *Virology* 219: 170-178.

319 Clerissi, C., Desdevises, Y., and Grimsley, N. (2012). Prasinoviruses of the marine green alga  
320 *Ostreococcus tauri* are mainly species specific. *J Virol* 86: 4611-4619.

321 Clerissi, C., Grimsley, N., and Desdevises, Y. (2013). Genetic exchanges of inteins between  
322 prasinoviruses (phycodnaviridae). *Evolution* 67: 18-33.

323 Cottrell, M. T., and Suttle, C. A. (1991). Wide-spread occurrence and clonal variation in  
324 viruses which cause lysis of a cosmopolitan, eukaryotic marine phytoplankter,  
325 *Micromonas pusilla*. *Mar Ecol Prog Ser* 78: 1-9.

326 Cottrell, M. T., and Suttle, C. A. (1995a). Dynamics of a lytic virus infecting the  
327 photosynthetic marine picoflagellate *Micromonas pusilla*. *Limnol and Oceanogr* 40:  
328 730-739.

329 Cottrell, M. T., and Suttle, C. A. (1995b). Genetic diversity of algal viruses which lyse the  
330 photosynthetic picoflagellate *Micromonas pusilla* (Prasinophyceae). *Appl Environ*  
331 *Microbiol* 61: 3088-3091.

332 Dunigan, D. D., Fitzgerald, L. A., and Van Etten, J. L. (2006). Phycodnaviruses: A peek at  
333 genetic diversity. *Virus Res* 117: 119-132.

334 Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high  
335 throughput. *Nucleic Acids Res* 32: 1792-1797.

336 Evans, C., Archer, S. D., Jacquet, S., and Wilson, W. H. (2003). Direct estimates of the  
337 contribution of viral lysis and microzooplankton grazing to the decline of a  
338 *Micromonas* spp. population. *Aquat Microbial Ecol* 30: 207-219.

339 Flores, C. O., Meyer, J. R., Valverde, S., Farr, L., and Weitz, J. S. (2011). Statistical structure  
340 of host–phage interactions. *Proc Natl Acad Sci USA* 108: E288-E297.

341 Foulon, E., Not, F., Jalabert, F., Cariou, T., Massana, R., and Simon, N. (2008). Ecological  
342 niche partitioning in the picoplanktonic green alga *Micromonas pusilla*: evidence from  
343 environmental surveys using phylogenetic probes. *Environ Microbiol* 10 : 2433-2443.

344 Guillou, L., Eikrem, W., Chrétiennot-Dinet, M.-J., Le Gall, F., Massana, R., Romari, K., et al.  
345 (2004). Diversity of picoplanktonic prasinophytes assessed by direct nuclear SSU  
346 rDNA sequencing of environmental samples and novel isolates retrieved from oceanic  
347 and coastal marine ecosystems. *Protist* 155: 193-214.

348 MacArthur, R. H. (1967). *The theory of island biogeography* (Vol. 1): Princeton University  
349 Press.

350 Marrec, P., Cariou, T., Collin, E., Durand, A., Latimier, M., Macé, E., et al. (2013). Seasonal  
351 and latitudinal variability of the CO<sub>2</sub> system in the western English Channel based on  
352 Voluntary Observing Ship (VOS) measurements. *Mar Chem* 155: 29-41.

353 Martinez Martinez, J., Boere, A., Gilg, I., van Lent, J. W. M., Witte, H. J., van Bleijswijk J. D.  
354 L., Brussaard, C.P.D. (2015) New lipid envelope-containing dsDNA virus isolates  
355 infecting *Micromonas pusilla* reveal a separate phylogenetic group. *Aquat Microb*  
356 *Ecol* 74: 17-28

357

358 Mizumoto, H., Tomaru, Y., Takao, Y.,  
359 Shirai, Y., and Nagasaki, K. (2007). Intraspecies host specificity of a single-stranded  
360 RNA virus infecting a marine photosynthetic protist is determined at the early steps of  
361 infection. *J Virol* 81: 1372-1378.

362 Not, F., Latasa, M., Marie, D., Cariou, T., Vaultot, D., and Simon, N. (2004). A single species,  
363 *Micromonas pusilla* (Prasinophyceae), dominates the eukaryotic picoplankton in the  
364 western English channel. *Appl Environ Microbiol* 70: 4064-4072.



365 Posada, D., and Crandall, K.A. (1998). MODELTEST: testing the model of DNA  
366 substitution. *Bioinformatics* 14: 817-818.

367 Pienaar, R. N. (1976). Virus-like particles in three species of phytoplankton from San Juan  
368 Island, Washington. *Phycologia* 15: 185-190.

369 Poulin, R. (1998). Large-scale patterns of host use by parasites of freshwater fishes. *Ecol*  
370 *Letters* 1: 118-128.

371 Reiter, D. M., Frierson, J. M., Halvorson, E. E., Kobayashi, T., Dermody, T. S., and Stehle, T.  
372 (2011). Crystal structure of reovirus attachment protein  $\sigma 1$  in complex with sialylated  
373 oligosaccharides. *PLoS pathogens* 7: e1002166.

374 Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D. L., Darling, A., Höhna, S., et al.  
375 (2012). MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice  
376 across a large model space. *Syst Biol* 61: 539-542.

377 Sahlsten, E. (1998). Seasonal abundance in Skagerrak-Kattegat coastal waters and host  
378 specificity of viruses infecting the marine photosynthetic flagellate *Micromonas*  
379 *pusilla*. *Aquat Microbial Ecol* 16: 103-108.

380 Short, S. M., and Short, C. M. (2008). Diversity of algal viruses in various North American  
381 freshwater environments *Aquat Microbial Ecol* 51: 13–21.

382 Sime-Ngando, T. (2014). Environmental bacteriophages: viruses of microbes in aquatic  
383 ecosystems. *Front Microbiol* 5: 355

384 Šlapeta, J., López-García, P., and Moreira, D. (2006). Global dispersal and ancient cryptic  
385 species in the smallest marine eukaryotes. *Mol Biol Evol* 23: 23-29.

386 Stoddard, L. I., Martiny, J. B., and Marston, M. F. (2007). Selection and characterization of  
387 cyanophage resistance in marine *Synechococcus* strains. *Appl Environ Microbiol* 73:  
388 5516-5522.

389 Suttle, C. A. (2005). Viruses in the sea. *Nature* 437: 356-361.

- 390 Suttle, C. A. (2007). Marine viruses - major players in the global ecosystem. *Nature Rev*  
391 *Microbiol* 5: 801-812.
- 392 Swofford, D. L. (2003). PAUP: Phylogenetic analysis using parsimony. Version 4.  
393 Sunderland, Massachusetts: Sinauer Associates.
- 394 Tarutani, K., Nagasaki, K., and Yamaguchi, M. (2006). Virus adsorption process determines  
395 virus susceptibility in *Heterosigma akashiwo* (Raphidophyceae). *Aquat Microbial*  
396 *Ecol* 42: 209-213.
- 397 Thomas, R., Grimsley, N., Escande, M. I., Subirana, L., Derelle, E., and Moreau, H. (2011).  
398 Acquisition and maintenance of resistance to viruses in eukaryotic phytoplankton  
399 populations. *Environ Microbiol* 13: 1412-1420.
- 400 Weitz, J. S., Poisot, T., Meyer, J. R., Flores, C. O., Valverde, S., Sullivan, M. B., et al. (2013).  
401 Phage–bacteria infection networks. *Trends Microbiol* 21: 82-91.
- 402 Worden, A. Z. (2006). Picoeukaryote diversity in coastal waters of the Pacific Ocean. *Aquat*  
403 *Microbial Ecol* 43: 165-175.
- 404 Zingone, A., Sarno, D., and Forlani, G. (1999). Seasonal dynamics in the abundance of  
405 *Micromonas pusilla* (Prasinophyceae) and its viruses in the Gulf of Naples  
406 (Mediterranean Sea). *J Plank Res* 21: 2143-2159.
- 407 Zingone, A., Natale, F., Biffali, E., Borra, M., Forlani, G., and Sarno, D. (2006) Diversity in  
408 morphology, infectivity, molecular characteristics and induced host resistance between  
409 two viruses infecting *Micromonas pusilla*. *Aquat Microbial Ecol* 45: 1–14.