



HAL
open science

High connectivity in a long-lived high-Arctic seabird, the ivory gull *Pagophila eburnea*

Glenn Yannic, Jonathan M Yearsley, Roberto Sermier, Christophe Dufresnes,
Olivier Gilg, Adrian Aebischer, Maria V. Gavrilov, Hallvard Strøm, Mark L.
Mallory, R I Guy Morrison, et al.

► **To cite this version:**

Glenn Yannic, Jonathan M Yearsley, Roberto Sermier, Christophe Dufresnes, Olivier Gilg, et al.. High connectivity in a long-lived high-Arctic seabird, the ivory gull *Pagophila eburnea*. *Polar Biology*, 2016, 39 (2), pp.221-236. 10.1007/s00300-015-1775-z . hal-01196647

HAL Id: hal-01196647

<https://hal.sorbonne-universite.fr/hal-01196647>

Submitted on 10 Sep 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

2 **High connectivity in a long-lived High-Arctic seabird, the ivory gull *Pagophila eburnea***

4 Glenn Yannic^{1,2,3*}, Jonathan M. Yearsley⁴, Roberto Sermier⁵, Christophe Dufresnes⁵, Olivier
Gilg^{3,6}, Adrian Aebischer^{3,7}, Maria V. Gavrilov^{8,9}, Hallvard Strøm¹⁰, Mark L. Mallory¹¹, R. I.
6 Guy Morrison¹², H. Grant Gilchrist¹³, Thomas Broquet^{14,15}

8 1. Département de Biologie, Université Laval, Québec, Québec, G1V 0A6, Canada

2. Canada Research Chair in Polar and Boreal Ecology, Department of Biology,
10 Université de Moncton, Moncton, New-Brunswick, E1A 3E9, Canada

3. Groupe de Recherche en Ecologie Arctique (GREA), 21440 Francheville, France

12 4. School of Biology and Environmental Science, University College Dublin, Belfield,
Dublin, Ireland

14 5. Department of Ecology and Evolution, University of Lausanne, CH-1015 Lausanne,
Switzerland

16 6. Laboratoire Biogéosciences, UMR CNRS 5561, Equipe Ecologie Evolutive,
Université de Bourgogne, 6 Boulevard Gabriel, 21000 Dijon, France

18 7. Museum of Natural History, Chemin du Musée 6, 1700 Fribourg, Switzerland

20 8. National Park Russian Arctic, 57 Sovetskikh Kosmonavtov pr., 168000, Archangelsk,
Russia

22 9. Joint Directorate of Taimyr Nature Reserves, 22 Talnakhsкая, 663300, Norilsk,
Russia

10. Norwegian Polar Institute, Fram Centre, 9296 Tromsø, Norway

24 11. Department of Biology, Acadia University, 33 Westwood Avenue, Wolfville, Nova
Scotia B4P 2R6, Canada

26 12. Environment Canada, National Wildlife Research Centre, Ottawa, Ontario, K1A0H3,
Canada

28 13. Environment Canada, National Wildlife Research Centre and Department of Biology,
Carleton University, Ottawa, Ontario, K1S5B6, Canada

30 14. CNRS, team diversity and connectivity of coastal marine landscapes, Station
Biologique de Roscoff, 29680 Roscoff, France

32 15. Sorbonne Universités, UPMC Univ Paris 06, UMR 7144, Station Biologique de
Roscoff, 29680 Roscoff, France

34 Correspondence and present address:

Glenn Yannic, glenn.yannic@gmail.com

36 LECA - Laboratoire d'Ecologie Alpine - CNRS UMR 5553

Université Savoie Mont Blanc

38 73376 Le Bourget-Du-Lac cedex, France

+(33) 04 79 75 88 65

40

Keywords: natal dispersal, breeding dispersal, effective number of breeders, population

42 genetic structure, overlapping generation model

44 **Abstract**

Species may cope with rapid habitat changes by distribution shifts or adaptation to new
46 conditions. A common feature of these responses is that they depend on how the process of
dispersal connects populations, both demographically and genetically. We analyzed the
48 genetic structure of a near threatened High Arctic seabird, the ivory gull (*Pagophila eburnea*)
in order to infer the connectivity among gull colonies. We analyzed 343 individuals sampled
50 from 16 localities across the circumpolar breeding range of ivory gulls, from northern Russia
to the Canadian Arctic. To explore the roles of natal and breeding dispersal we developed a
52 population genetic model to relate dispersal behavior to the observed genetic structure of
worldwide ivory gull populations. Our key finding is the striking genetic homogeneity of
54 ivory gulls across their entire distribution range. The lack of population genetic structure
found among colonies, in tandem with independent evidence of movement among colonies,
56 suggests that on-going effective dispersal is occurring across the Arctic Region. Our results
contradict the dispersal patterns generally observed in seabirds where species movement
58 capabilities are often not indicative of dispersal patterns. Model predictions show how natal
and breeding dispersal may combine to shape the genetic homogeneity among ivory gull
60 colonies separated by up to 2800 km. Although field data will be key to determine the role of
dispersal for the demography of local colonies and refine the respective impacts of natal
62 versus breeding dispersal, conservation planning needs to consider ivory gulls as a genetically
homogeneous, Arctic-wide metapopulation effectively connected through dispersal.

64

Introduction

66 The distribution of natural habitats worldwide is currently changing as a direct consequence
of global climate trends, and this is happening particularly fast in the Arctic, where climate
68 warming is maximal (ACIA 2004; IPCC 2007). Species that live in the Arctic or in other
rapidly changing environments might cope with this rapid change by shifting their
70 distributions, by adjusting through phenotypic plasticity or by evolving adaptations to the new
local climatic conditions (reviewed by Chen et al. 2011; Gienapp et al. 2008; Gilg et al. 2012;
72 Hoffmann and Sgro 2011; Parmesan 2006). These responses partly depends on the process of
dispersal; that is, the movement of individuals between birth and reproduction (natal
74 dispersal), and possibly between successive reproduction events (breeding dispersal). Besides
its role in the spatial structure and demographic dynamics of populations, dispersal is
76 important in the context of habitat change because it is one key driver of the potential rate of
spread of a population and, as the process by which genes are moved among populations, it
78 influences the rate of adaptation to changing conditions and the potential for evolutionary
rescue (Bell and Gonzalez 2011; Travis et al. 2013). Thus, understanding, predicting and
80 managing biodiversity responses to rapid climate change demands a full consideration of a
species' dispersal characteristics and their demographic and genetic consequences.

82 We focus here on the ivory gull *Pagophila eburnea*, a bird that completes its life-cycle
entirely in the Arctic. Over its entire breeding range (Canadian Arctic, Greenland, Svalbard
84 and Russian Arctic islands) it breeds either on inland cliffs and 'nunataks', *i.e.* rocky outcrops
emerging from icecaps, or on high-Arctic barren islands or flatlands (Gilg et al. 2009;
86 Mallory et al. 2008). In Canada, where the status of the species has been designed
'Endangered' (COSEWIC 2006), studies indicated that 80% of the breeding population was
88 lost during the past 20 years (Gilchrist and Mallory 2005). The species is listed as Near
Threatened by the IUCN (BirdLife International 2012) and an international circumpolar

90 'Conservation Strategy and Action Plan' has been presented by leading seabird experts from
Arctic countries to gain more insight into how this bird responds to increasing threats from
92 disappearance of sea ice habitat, natural resource exploration and increased contaminant loads
(Gilchrist et al. 2008).

94 Ivory gulls are capable of travelling thousands of kilometers either on single foraging
trips or to reach wintering grounds in the north Pacific (Bering Sea and Sea of Okhotsk) and
96 in the northwest Atlantic (Davis Strait and Labrador Sea) where most of the world population
is thought to spend the winter (Gilg et al. 2010; Mallory et al. 2008). However, most seabirds
98 have an extraordinary ability to travel long distances and yet show evidence of restricted gene
flow and exhibit high levels of philopatry, sometimes returning to breed within a few meters
100 of their natal nest (Friesen et al. 2007). A species' movement capabilities thus do not
automatically inform us about demographic and genetic connectivity among colonies. This is
102 the "seabird paradox", *i.e.*, the apparent paradox between high vagility and low effective
dispersal (Milot et al. 2008).

104 Dispersal may take place at different stages of an individual's life. For ivory gulls, natal
dispersal may happen during the two first years of life before the individual becomes sexually
106 mature and joins a breeding colony. However, the behavior of ivory gulls during that time is
almost completely unknown. In addition, adult ivory gulls may disperse among colonies from
108 one breeding season to the next. Such breeding dispersal could effectively contribute to
demographic and genetic exchanges among colonies, but our knowledge of these aspects for
110 ivory gulls currently relies only on incidental observations (O. Gilg, A. Aebischer and M.L.
Mallory, unpubl. data).

112 Here we take a genetic approach to investigating dispersal in order to complement
ongoing mark-recapture and satellite tracking efforts (Gilg et al. 2010; Spencer et al. 2014).
114 Genetic data can complement other approaches to measure dispersal either by providing direct

information on individual movements (e.g., through parentage or population assignment) or
116 indirect signatures of dispersal patterns (e.g., through analyses of genetic structure).

Disentangling the effects of natal dispersal and breeding dispersal on realized gene flow is,
118 however, challenging, and has rarely been addressed in the molecular ecology literature
(Broquet and Petit 2009), although Rousset (2001) and Laporte and Charlesworth (2002)
120 present general class-structured models that lay the foundations to such an endeavor.

The aim of this study was to explore population structure and spatial dispersal pattern in
122 the ivory gull and to infer natal versus breeding dispersal among colonies. For that purpose,
we analyzed a genetic data set representative of the entire species range and developed a
124 population genetic model to infer lower bounds on natal and breeding dispersal consistent
with the observed genetic structure of ivory gull populations worldwide.

126

MATERIAL AND METHODS

128 Study species

The ivory gull is a long-lived High Arctic seabird (annual survival estimated to 0.86;
130 Stenhouse et al. 2004; and maximum record 28 years; Mallory et al. 2012), which is
associated with sea ice all year round (Gilg et al. 2010; Spencer et al. 2014). Breeding
132 colonies are scattered in Arctic Canada, Greenland, Svalbard, and the northern islands of
Russia in the Barents and Kara seas (Table 1). The current total global population of the ivory
134 gull was estimated to be approximately 19,000-27,000 breeding pairs (BirdLife International
2012). The Russian population is estimated to number in the range of 14,500-22,000
136 individuals (Gavrilo 2011). The population in Canada has declined since the 1980s (Mallory
et al. 2008). In Norway (Svalbard) the population probably declined in the first part of last
138 century, but after 1970 the trend is uncertain (Mallory et al. 2008). Population trends in
Greenland are unclear due to sparse historical information (Gilg et al. 2009). Ivory gulls are

140 thought to first breed after their second year, based on the fact that they acquire adult plumage
in their second winter, and that individuals in less than full adult plumage are rarely seen at
142 breeding colonies (Mallory et al. 2008). Unlike most gulls, which regularly lay 3 eggs, the
ivory gull usually lays 1–2 eggs, more rarely 3 eggs. Most of the world population is thought
144 to spend the winter in two main wintering grounds (Mallory et al. 2008): the north Pacific
(Bering Sea and Sea of Okhotsk) and the northwest Atlantic (Davis Strait and Labrador Sea,
146 Figure 1).

148 **Sample collection**

Field works took place in summers 2006 to 2012, during the breeding season (late June to
150 August). Sample locations were distributed across the entire breeding range of the species,
including the Canadian Arctic Archipelago, north-eastern Greenland, Svalbard Archipelago,
152 Franz Josef Land Archipelago, Severnaya Zemlya Archipelago and Kara Sea islands (16
sampling locations overall, listed in Table 1 and Figure 1). We collected samples either in
154 breeding colonies or opportunistically near two military stations where ivory gulls are
attracted by food remains (namely Alert, Canada and Station Nord, Greenland). Three
156 nondestructive DNA sampling methods (mouth swabs, plucked feathers and blood) and a
noninvasive sampling method (shed feathers) were used. Pieces of tissue were also
158 opportunistically collected on dead birds.

Juveniles (chicks of the year) were sampled in two sites from Greenland in 2009: Amdrup
160 Land and Station Nord (Table 1 and Figure 1) in order to perform parentage analyses. In these
cases, buccal swabs and tissue samples were used as DNA sources. All other samples were
162 taken from adult birds, where we considered two classes of individuals according to their
breeding status. Field observations suggest that non-breeding adults visit or stay in colonies
164 during the breeding season. Moreover, satellite transmitters indicated that breeding birds

visited colonies as far as 200 km from their own breeding colony (O. Gilg & A. Aebischer,
166 unpublished data). Hence in any one site adult birds were classified as "breeding" only if they
were seen hatching eggs or raising chicks, and "unknown status" otherwise. Thus, "unknown"
168 birds included: i) the non-breeding component of the population (the so-called "floaters";
Penteriani et al. 2011), but also ii) birds found in colonies but that were not reproducing
170 locally and that may reproduce elsewhere in an unknown colony; and ii) birds that they were
identified from shed feathers collected on the ground. This distinction is relevant for
172 analyzing the genetic structure of colonies since non-breeding birds or individuals lacking
information on their breeding location (all called here "unknown") could be transient visitors.
174 Due to field constraints we generally have information on only one of these two classes of
adults within each sampling site (reported in Table 1), either because samples were taken only
176 from breeding individuals or because the breeding status was ignored altogether (e.g., shed
feathers or transient birds). However in one repeatedly visited site from Greenland (called
178 Station Nord), we could collect precise mark-resight data on both breeding and unknown (see
above) individuals, and obtain sizable samples from these two classes of birds (referred to as
180 "breeding" and "unknown" in Table 1).

All samples from Greenland, Norway and Russia (Table 1) were obtained using non-
182 destructive (collection of mouth swabs and plucked feathers) and non-invasive DNA
sampling methods (collection of shed feathers) as described in Yannic *et al.* (2011). In
184 addition, birds from Alert (Canada) were caught with rocket nets near a military base.
Immediately following capture, a blood sample (about 0.3 ml) was collected from the brachial
186 vein in heparinized micro-hematocrit capillary tubes, before release. Blood samples were
centrifuged on site at 13,200 g for 15 min. Red blood cells and plasma were separated and
188 stored frozen at -20°C until laboratory analyses).

190 **DNA extraction and genotyping**

Genomic DNA from all individuals was extracted from shed and plucked feathers, tissue,
192 blood or buccal swab following protocols described in Yannic *et al.* (2011) (see also
Supplementary material 1). Previously optimized microsatellite markers were used in four
194 polymerase chain reaction (PCR) multiplexes, totaling 22 markers (Yannic *et al.* 2011). For
samples obtained from shed feathers we performed three independent PCR replicates of each
196 locus to obtain reliable genotypes (see Yannic *et al.* 2011). The microsatellite amplicons were
loaded on an ABI PRISM 3100 (Applied Biosystems Foster City, CA, USA) automated DNA
198 sequencer. Microsatellite alleles were detected, scored, and manually verified using
GENEMAPPER 3.7 (Applied Biosystems).

200

Genetic structure

202 All loci were found to be independent of one another (linkage disequilibrium test performed
in FSTAT 2.9.4 (Goudet 2005), using 10 000 permutations and *p-values* adjusted for multiple
204 comparisons using the Benjamini and Yekutieli false discovery rate procedure with initial $\alpha =$
0.05). We used two sets of loci depending on downstream analyses. All 22 loci were used for
206 the parentage analyses because: i) all genetic data from the juveniles came from good quality
samples (tissue and buccal swab); and ii) genotyping errors or null alleles can be identified
208 and taken into account (see below). For the analysis of spatial genetic structure some data
come from "low quality" samples (shed feathers, Table 1). Hence for these analyses we used a
210 subset of 13 loci (listed in Table 2) chosen for their polymorphism and reliability as reported
in Yannic *et al.* (2011).

212 We investigated the differentiation among ivory gulls sampling sites by estimating F_{ST} (Weir
and Cockerham 1984). We ran some of the analyses using only the samples with >10 adults.

214 Global F_{ST} was computed with FSTAT for different combinations of samples: overall adults (n

= 15 localities), over sites with >10 adults ($n = 9$ localities), and among breeders only ($n = 6$
216 localities). The significance of the differentiation was tested using two approaches. First we
used the log-likelihood G statistic calculated for observed data and compared to that of 10 000
218 randomized datasets obtained through permutation of individuals among samples (as
implemented in Fstat; Goudet 2005; Goudet et al. 1996). Second, for a strict comparison with
220 results from our evaluation of power (see below), we also used Fishers' Exact Test as
implemented in GENEPOP. In that case the distribution of alleles within individuals is ignored
222 and thus genic rather than genetic differentiation among samples is tested. Furthermore, the
null distribution is obtained using a Markov Chain algorithm rather than permutations,
224 performed here with defaults GENEPOP parameters. Pairwise F_{ST} among all samples were also
calculated with FSTAT.

226 The statistical power to detect a significant genetic heterogeneity at various true levels of
differentiation for the present set of samples, number of loci and allele frequencies was
228 evaluated using POWSIM 4.1 (Ryman and Palm 2006). POWSIM simulates samples of genes
from a specified number of populations that have drifted to an expected predefined level of
230 differentiation (measured as F_{ST}). These samples are then used for testing genetic
homogeneity using Fisher's Exact Test. With this procedure we estimated the power that we
232 had when looking for genetic differentiation using all adults and breeders only (see Table 1).
Estimates of power were given by the proportion of significant outcomes when repeating the
234 simulations 1000 times for each level of simulated F_{ST} . The use of post hoc power analyses
should however be used with caution as stressed by Hoenig and Heisey (2001). But, here our
236 goal is not to modify a hypothesis test a posteriori (the problematic situation identified by
Hoenig and Heisey (2001)) but rather to give an idea of the degree to which our data are
238 informative.

240 **Reproductive success and effective number of breeders**

To interpret our observations of genetic structure across colonies we needed an estimate of
242 effective colony sizes. This can be approximated using the effective number of breeders (N_b)
(N_b ; Waples and Teel, 1990), a parameter that depends on the census number of adults in a
244 colony (here noted N_c) and the distribution of reproductive success among individuals within
colonies following (Kimura and Crow 1963): $N_b = (N_c k - 1) / [k - 1 + (V_k / k)]$, where k is the
246 mean and V_k the variance in reproductive success among individuals. As a first approximation
we estimated these figures from field observations of the number of juveniles per nest in
248 colonies Amdrup Land and Station Nord, considering that there are two and only two adults
associated with a given nest.

250 This approach assumes that juveniles within a nest descend from the adult pair providing
parental care to these offspring. This is a weak assumption since extra-pair paternity is
252 frequent in socially monogamous birds (Westneat and Stewart 2003), meaning that some
males may not have sired the juveniles they are taking care of, whilst other males may have
254 parented offspring with more than one female. Hence males may have a slightly higher
variance in reproductive success than those calculated from field observations. To check
256 whether social monogamy reflects the actual breeding system we performed genetic parentage
assignments in colony Station Nord, where we had DNA samples from a number of juveniles
258 (n=20) and presumed parents (n=24), that is, adults seen hatching eggs or raising chicks.

Details of the parentage analysis, performed with the method implemented in COLONY 2.0.4.5
260 (Jones and Wang 2009; Wang 2012) are described fully in Electronic Supplementary Material
(Supplementary material 1). We repeated these analyses in the colony of Amdrup Land,
262 where 65 juveniles (but no parents) were sampled.

264 **Model of genetic structure: overlapping generations, natal and breeding dispersal**

Interpreting genetic differentiation in terms of connectivity and dispersal behavior is not
266 trivial given that it requires some knowledge of the effective number of breeders within
colonies (Nb , which we investigated in this study) and the effect of life-history traits such as
268 longevity and the potential movement behavior of juveniles (natal dispersal) and adults
between breeding seasons (breeding dispersal). We therefore used a 2-sample coalescent
270 approach to describe an island model, with overlapping generations and both natal and
breeding dispersal. The model is used to explore the dispersal scenarios that are consistent
272 with the observed level of population differentiation (global F_{ST}) among arctic-wide
populations of ivory gulls.

274 The model builds upon Yearsley et al. (2013) to introduce overlapping generations following
the general approach laid out by Laporte and Charlesworth (2002). Each deme contains N
276 diploid non-selfing individuals, of which $N_a = \nu N$ are adults who have survived at least one
breeding cycle and $N_j = (1 - \nu) N$ are first-year juveniles (ν is the adult survival probability per
278 breeding cycle). One breeding-cycle going forward in time represents a unit time step and is
composed of: reproduction, mutation, dispersal, adult mortality, juveniles either mature into
280 adults or die, population regulation (*i.e.*, the population size remains constant at every
breeding cycle). The model simplifies certain aspects of the ivory gull's life-history.

282 Maturation for ivory gulls is known to be longer than one year whereas our model, to enable
an analytical solution, assumes that a maximum juvenile period of one year. From numerical
284 simulations (results not shown) the effect of a prolonged juvenile stage on F_{ST} is small when
adult mortality is low (as for the ivory gull). All adults in the model are assumed to have
286 equal reproductive success. Individual variation in reproductive success and non-breeding
adults must be accounted for by the effective population size, N . The model also does not
288 describe sex-linked differences in life-history, such as dispersal or survival. At present we do

not have sufficient sex-specific data for the ivory gull to know whether such differences exist
290 for this species.

292 The model estimate expected coalescence times, genetic diversities, and F -statistics for a
DNA sequence under the infinite-sites model (Kimura 1969) with a mutation rate μ /
294 generation/sequence. For our model parameterizations the force of mutation upon genetic
diversities is weak compared to the forces of genetic drift and gene flow. Our model considers
296 the coalescent for a sample of two DNA sequences that are randomly sampled just prior to
population regulation. We define three states for a pair of sampled sequences: two sequences
298 in the same diploid individual, two sequences in different individuals in the same deme, and
two sequences in different individuals in different demes (states 1, 2 and 3 respectively).

300 The ancestral history of a pair of sequences can be defined by a transition matrix, \mathbf{G} , where an
element, $G_{i,j}$, gives the probability that a pair of sequences in state i had ancestors from the
302 previous generation in state j (the rate of coalescence per generation for a pair of sequences in
state i is then given by $G_{i0} = 1 - \sum_j G_{i,j}$). Using first-step analysis (Wakeley 2009) the expected
304 times to coalescence of two lineages in state i , T_i , can be calculated by solving

$$306 \quad T_i = 1 + \sum_{j=1}^3 G_{i,j} T_j \quad (1)$$

308 This equation is analogous to equation 8 in Laporte and Charlesworth (2002), and details of
the approach used to derive equation 1 are given in Yearsley et al. (2013). Using Slatkin's
310 approximation (Slatkin 1991), these coalescence times can be used to approximate F -statistics
in the small mutation limit as

$$312 \quad F_{IS} = \frac{T_2 - T_1}{T_2}$$

$$F_{ST} = \frac{T_3 - T_2}{T_3} \quad (2)$$

314 Alternatively, mutations can be included in the matrix \mathbf{G} and F -statistics calculated from
recurrence relationships for identity by descent.

316 We specified the transition matrix, \mathbf{G} , by identifying three types of sequence pair: sequences
from two juveniles (*i.e.* newly born in the current breeding cycle), sequences from two adults
318 (*i.e.* individuals surviving from the previous breeding cycle), and sequences from one juvenile
and one adult (these types are labelled -, +, \pm respectively). The transition matrix can be

320 written as $G = G^- + G^\pm + G^+$ where

$$G^- = (1-\nu)^2 \begin{pmatrix} 0 & 1/(1-\nu) & 0 \\ \alpha^-/2N & \alpha^-(N-1)/N & 1-\alpha^- \\ \beta^-/2N & \beta^-(N-1)/N & 1-\beta^- \end{pmatrix} \quad (3a)$$

322 is the transition matrix when the both sequences in the pair are from juveniles (possibly the
same juvenile),

$$324 \quad G^\pm = 2\nu(1-\nu) \begin{pmatrix} 0 & 0 & 0 \\ \alpha^\pm/2N & \alpha^\pm(N-1)/N & 1-\alpha^\pm \\ \beta^\pm/2N & \beta^\pm(N-1)/N & 1-\beta^\pm \end{pmatrix} \quad (3b)$$

is the transition matrix when one sequence is from a juvenile and one from an adult and

$$326 \quad G^+ = \nu^2 \begin{pmatrix} 1/\nu & 0 & 0 \\ 0 & \alpha^+ & 1-\alpha^+ \\ 0 & \beta^+ & 1-\beta^+ \end{pmatrix} \quad (3c)$$

is the transition matrix when the both sequences in the pair are from adults (possibly the same
328 adult). The other parameters in the many-deme limit are $\alpha^- = (1-m_j)^2$, $\alpha^\pm = (1-m_j)(1-m_a)$, $\alpha^+ =$
 $(1-m_a)^2$, with m_j and m_a the juvenile and adult migration rates, respectively (*i.e.*, m_j and m_a
330 represent natal and breeding dispersal). The parameters β^x make a negligible contribution to
 F_{ST} in the many-deme limits because they are inversely proportional to the number of demes.

332

Substituting equation 3a-c into equation 1 and solving, and taking the many-deme limit gives

$$T_1 = T_2 = 2 N_a D / (1 - \nu^2) \quad (4a)$$

$$T_3 = T_2 + D (1 - m_j) (1 + p) M / [(1 - M^2) (1 + \nu)] \quad (4b)$$

where $M = (1 - \nu) (1 - m_j) + \nu (1 - m_a)$, $p = \nu (1 - m_a) / M$ and time units are in breeding cycles. To express these coalescence times in numbers of generations they should be divided by generation time (equal to $1 / (1 - \nu)$).

Substituting equations 4 into equations 2 gives the F -statistics, $F_{IS} = 0$ and an expression for F_{ST} in the small mutation limit of

$$\frac{1 - F_{ST}}{F_{ST}} = 2 N_a \frac{1 - M^2}{M^2} \frac{1}{1 - p^2} \quad (5)$$

For non-overlapping generations ($\nu = 0$) and small migration rates equation 5 gives the classic result $(1 - F_{ST}) / F_{ST} = 4 N_a m_j$ (Wright 1931). The model also correctly predicts the inbreeding effective population size $N_e = N_a / (1 + \nu)$ for a single isolated population with overlapping generations (Felsenstein 1971; Hill 1972), equivalent to the case when $m_j = m_a = 0$.

Equation 5 shows how the genetic differentiation among ivory gull colonies depends upon adult survival (ν), effective colony size (N_a), natal dispersal (m_j) and breeding dispersal (m_a).

Adult annual survival rate was estimated to $\nu = 0.86 \pm 0.04$ (95% CI: 0.75;0.91) (Stenhouse et al. 2004). We used our model with the mean annual survival rate, $\nu = 0.86$ and the upper limit of the confidence interval $\nu = 0.91$. Using a higher survival value will tend to underestimate migration rates, making our interpretation more conservative. Effective colony size N_a cannot be precisely parameterized because contrary to our model's assumptions the number of breeding adults is variable across colonies. Known colony sizes (reviewed in Table 3) show a skewed distribution, with a few large colonies (in the order of 100 – 2000 breeding pairs) and many smaller ones (below 100 pairs). Furthermore we did not know the prevalence and year-to-year behavior of adults that are apparently non-breeding at some observation time point.

Such individuals can inflate N_a if they have or will enter reproduction at some other breeding
358 season. Based upon i) field observations of colony sizes (Table 3), ii) the fact that low
variance in reproductive success should inflate local effective numbers of breeders (see results
360 for N_b/N_c in colonies Amdrup Land and Station Nord), and iii) remaining uncertainties about
the resulting parameter N_a , we explored the model behavior for N_a ranging 50-1000. Using
362 equation 5 we then worked out the conditions of juvenile and adult migration that would
result in a F_{ST} value equal to the observed global $F_{ST} = 0.001$. This allows us to estimate and
364 discuss the lower bound on migration rates for ivory gulls.

366 RESULTS

Genetic structure

368 Number of alleles, observed and expected heterozygosity in each sample and for each of the
13-microsatellite loci are shown in Table 1 and in Electronic Supplementary Material (Table
370 S1 in Supplementary material 1), respectively. With 13 loci examined in 15 samples, nine
locus/site combinations showed a significant deficit in heterozygotes. There was no consistent
372 pattern across samples or loci, and only one locus in one sample (B125, Schmidt Island,
Russia) remains significant if one corrects for multiple testing. Yet it is plausible that a small
374 number of allelic dropouts remained undetected in genotypes obtained from shed feathers
despite marker selection and genotyping repetitions. The number of genotyping repetitions
376 that we used is based upon average error rates reported in Yannic et al. (2011) but individual
shed feathers may happen to be unusually poor sometimes (Yannic et al. 2011). For this
378 reason, we reported differentiation statistics with and without data from shed feathers. The
mean observed heterozygosity (0.63–0.85) and mean expected heterozygosity (0.73–0.82)
380 across loci are shown in Table 1.

No genetic differentiation was observed among breeding samples ($n = 6$, $F_{ST} = 0.000$, 95%CI: -0.006;0.005; G statistic permutation test $p = 0.61$, Fisher's Exact Test $p = 0.40$; Table 2) or among samples containing more than 10 individuals ($n = 9$, $F_{ST} = 0.000$, 95%CI: -0.002;0.003; G statistic permutation test $p = 0.15$, Fisher's Exact Test $p = 0.14$; Table S2 in Supplementary material 1), while very low and non-significant differentiation was found overall adult samples ($n = 15$, $F_{ST} = 0.001$, 95%CI: -0.002;0.005; G statistic permutation test $p = 0.09$, Fisher's Exact Test $p = 0.09$; Table 2). This figure was unaffected when removing all shed feather samples ($n = 8$, $F_{ST} = 0.000$). Pairwise F_{ST} values were also very low, ranging from -0.032 to 0.043 and none of these pairwise values was significant after correction for multiple testing (Benjamini–Yekutieli correction; Table S3 in Supplementary material 1). There was no significant difference in relatedness among breeders vs among unknown birds sampled the same year in the same colony (*i.e.*, Station Nord in 2009: “Effect of transient individuals on genetic structure” section in Supplementary material 1), suggesting that breeders and unknown birds belong to a homogeneous pool. These results were further confirmed by model-based clustering that suggests that our ivory gulls most likely form one worldwide population (“Model-based clustering” section in Supplementary material 1) and by the absence of isolation-by-distance over long distance (“Isolation by distance” section in Supplementary material 1).

Simulations demonstrated that our sample sizes and genetic markers provided sufficient power to detect weak population structure. Population structure was found significant for all simulated populations (*i.e.* power = 100%) with an F_{ST} of 0.006 when using all adult sampling sites ($n = 17$; Figure 2). Even when F_{ST} was reduced to 0.0035, structure was correctly detected in 90% of the simulations. For F_{ST} values as low as, or lower than the observed value (*i.e.*, global F_{ST} among all adults = 0.001), power drops to 25%. When using only breeders sampling sites ($n = 6$ sites), the sample sizes and the genetic markers contain

406 sufficient power to detect population structure with 90% accuracy for simulated populations
with F_{ST} values ≥ 0.007 (Figure 2).

408

Reproductive success and effective number of breeders

410 In the colony Amdrup Land, we counted 98 adults (49 nests) with one offspring, 82 adults (41
nests) with two offspring, and 12 adults (6 nests) with an unknown number of offspring
412 (Yannic et al. 2014a). Assuming that the latter show the same distribution of reproductive
success than all other adults, this gives $k \approx 1.46$, $V_k \approx 0.25$, and $N_b \approx 445$. In Station Nord we
414 observed 24 adults with one offspring and 48 adults with two, which gives $k \approx 1.67$, $V_k \approx 0.23$,
and $N_b \approx 148$.

416 Genetic parentage assignment at Station Nord identified the two parents (from our sample of
adults) for 6 juveniles out of 20. Twelve additional juveniles had one of their parents
418 identified from the candidate adults. The "second parents" of these juveniles and the two
parents of the remaining juveniles ($n = 2$) were not identified from the adult samples but their
420 genotype was reconstructed by the software COLONY, meaning that these adults could still be
used to check for extra-pair paternity (*e.g.*, if one unsampled male had sired three of our
422 offspring with different unsampled females, this would be visible in the data). As it turned
out, the parent-offspring relationships observed in the field were all confirmed by the genetic
424 assignment (that is, for all the individuals with a DNA sample available), with one exception:
one adult that was observed caring for a juvenile did not appear to be its genetic parent.
426 Moreover, this true parent was identified from our sample of adults and it was found to have a
second offspring with a different mating partner (field observation, independently confirmed
428 by the genetic data). This suggests one plausible event of extra-pair paternity.

We repeated these analyses in the colony of Amdrup Land, where 65 juveniles (but no
430 parents) were sampled. But with such small clutch size (one or two offspring in general) and

without any actual parent genotyped, we did not succeed to recover reliable sibship
432 information in this colony (data not shown).

In summary, observable parental behavior seems a reliable indicator of parentage, and field
434 observations suggest that the effective breeding size N_b is approximately twice the census
colony size N_c . This figure results from the near-zero variance in breeding success among
436 birds seen in colonies. This variance could be slightly inflated by extra-pair paternity, but
with very little consequences for the N_b/N_c ratio (*e.g.*, N_b decreases from 148 to 142 in colony
438 Station Nord if one considers one event of extra-pair paternity where one bird has no success
and another one has fathered three offspring).

440

Model of genetic structure: natal versus breeding dispersal

442 We explored the conditions of natal dispersal (dispersal of juveniles) and breeding dispersal
(movement of adults among colonies across breeding seasons) that would be consistent with
444 the low level of observed genetic structure.

A general result obtained with the model is that breeding dispersal is very effective at
446 homogenizing the distribution of the genetic variation across populations in long-lived species
with overlapping generations. For instance with the ivory gulls, with $v = 0.91$ (Figure 3B) and
448 $N_a = 1000$ and no natal dispersal (that is, perfect philopatry) then a breeding dispersal of only
4.6% is required to yield an F_{ST} as low as 0.001. By contrast, above 30% natal dispersal
450 would be required in the absence of breeding dispersal.

The above scenario is conservative, providing lower bounds on dispersal rates because we
452 used our highest observation of global F_{ST} ($F_{ST} = 0.001$; see Table 2), large colony size, and
high survival. A slightly less conservative scenario ($F_{ST} = 0.001$, $N_a = 500$, $v = 0.86$, visible in
454 Figure 3A) gives: 14% breeding dispersal or 48% natal dispersal (or any combination along
the $F_{ST} = 0.001$ contour line in Figure 3A). Any smaller (*i.e.*, less conservative) value for N_a

456 or F_{ST} will increase the minimum level of dispersal. As expected, predictions of genetic
structure were highly sensitive to effective colony size (as shown by the different contour
458 lines within Figures 2A and 2B) and survival (compare Figure 3A against 3B).

460 **Discussion**

The key finding from this research is the striking genetic homogeneity of the ivory gull across
462 its entire distribution range. Even with conservative assumptions for local effective breeding
numbers and survival rate this suggests that gene flow regularly occurs among distant regions
464 in order for populations to become, and remain, genetically homogenous. We develop below
the interpretation of these results indicating genetic homogeneity among populations
466 separated by up to 2800 km.

468 **A single Arctic-wide population**

Information retrieved from microsatellites suggests that the ivory gull represents a single,
470 Arctic-wide metapopulation. We found no significant genetic differentiation among breeding
colonies of ivory gull ($F_{ST} = 0.000$, $CI_{95\%}$: -0.006; 0.005) or among overall adult samples (F_{ST}
472 $= 0.001$, $CI_{95\%}$: -0.002; 0.005). We did not observe significant isolation-by-distance among
breeding colonies and among overall adult samples across the range of the species (“Isolation
474 by distance” section in Supplementary material 1). These results agree with the weak
differentiation found using mitochondrial data (Royston and Carr 2014 and this study;
476 Supplementary material 1).

This absence of genetic structure is *a priori* not surprising for a species capable of travelling
478 thousands of kilometers either on single foraging trips or to reach its wintering grounds (Gilg
et al. 2010). Genetic homogeneity is, however, not the rule in seabird species with similar
480 flying capability. Out of forty-seven seabird species reviewed by (Friesen et al. 2007), only

few were reported to have as little genetic structure as the ivory gull. The grey-faced petrel
482 *Pterodroma macroptera gouldi* (Lawrence et al. 2014), the little auk *Alle alle* (Wojczulanis-
Jakubas et al. 2014) and the wandering albatross *Diomedea exulans* (Milot et al. 2008) are
484 examples of seabird that present weak genetic structure throughout their distribution. But the
vast majority of seabird species rather seem to show a stronger level of genetic divergence,
486 even among geographically proximate colonies (*e.g.*, the Hawaiian petrel *Pterodroma*
sandwichensis (Welch et al. 2012) or Cory's shearwater *Calonectris diomedea* (Genovart et
488 al. 2013). Genetic divergence among seabird populations inhabiting the Polar Regions seems
then to be generally lower in comparison with those breeding at lower latitudes.

490 Patterns of genetic structuring in species capable of long-distance dispersal may be driven by
multiple mechanisms, including restricted gene flow as a result of high natal philopatry,
492 cryptic barriers to dispersal, or behavioral mechanisms (Friesen et al. 2007). In addition, local
adaptation to differing ecological conditions and strong selective pressures may promote
494 geographic patterns of differentiation. Our results show that such gene flow limiting processes
are not at work in the ivory gull population and high intercolony dispersal genetically
496 homogenizes the populations. It is however worth noting that our results are based on neutral
genetic loci (*i.e.*, microsatellite loci) and adaptive differences could exist among colonies.

498

Our interpretation of the data assumes that the F_{ST} is at migration-drift equilibrium. With
500 small deme size and large migration rates, F_{ST} reaches equilibrium very rapidly [*i.e.* in the
order of a few dozen of generations, Rousset (2004)], contrary to gene diversity which may
502 take a much longer time to reach equilibrium (Crow and Aoki 1984). The hypothesis that we
believe to be most parsimonious in the case of ivory gulls is that F_{ST} has long been
504 equilibrated and there is large-scale genetic exchange between colonies, most likely due to a
combination of natal and breeding dispersal. An alternative hypothesis may be that the

506 worldwide population is sub-structured into poorly connected demes and the genetic
homogeneity observed in ivory gull today is a consequence of the evolutionary history of the
508 species, *i.e.*, a northward expansion of population from a single homogeneous refugia after
the deglaciation of the Arctic region (e.g., Wojczulanis-Jakubas et al. 2014). However, while
510 it is temperate species were restricted to refugial area during glacial stages, taxa found in
more northern latitudes today are known to have had greater distributions during the glacial
512 phases (e.g., Lorenzen et al. 2011; Yannic et al. 2014b). This suggests that colder adapted
species were in more restricted areas during interglacial and not during glacial stages (Stewart
514 and Dalen 2008; Stewart and Lister 2001). From this perspective, ivory gulls could be said to
be in “refugia” today and not necessarily in the Late Pleistocene.

516

Natal versus breeding dispersal

518 To disentangle the respective role of natal dispersal, *i.e.* the movement from the natal site to
the site of first reproduction (Greenwood and Harvey 1982), and breeding dispersal, *i.e.*
520 movement between successive breeding attempts in the ivory gull, we developed an infinite
island model with overlapping generations that we used to calculate the expected global F_{ST} at
522 equilibrium for a range of adult and juvenile migration rate scenarios. Our results show that
breeding dispersal is very effective at reducing genetic differentiation across populations in
524 long-lived seabird with overlapping generations. We used this model here in an attempt to
better understand the demo-genetics of a featured high-artic seabird species, but the modeling
526 framework that we presented here is very general. Our model could be used further to look at
the effect of overlapping generations and variations in natal vs breeding dispersal, two aspects
528 that have largely been ignored from empirical molecular ecology research so far.

530 Long-term field data are lacking for the ivory gull (see next section below), but breeding
dispersal is thought to be less than natal dispersal for seabirds in general (e.g., Gauthier et al.
532 2010). In many long-lived seabird species with low reproductive rate, breeding philopatry is
believed to be very high, although actual dispersal rates have been rigorously quantified for a
534 few species only: roseate tern *Sterna dougallii* ($m_a=0.00-0.09$ yr⁻¹; Lebreton et al. 2003),
common tern *Sterna hirundo* ($m_a=0.04-0.08$ yr⁻¹; Nisbet and Cam 2002), wandering
536 albatross ($m_a=0.00-0.30$ yr⁻¹; Gauthier et al. 2010) or Adélie penguin *Pygoscelis adeliae*
($m_a < 0.01$ yr⁻¹; Dugger et al. 2010). In these species, breeding dispersal rates appear to be
538 very low and strongly limited by the distance among colonies, although dispersal could vary
with ice conditions (e.g., Dugger et al. 2010). These observations suggest that there are
540 behavioral constraints on adult movement amongst breeding colonies (Friesen et al. 2007).
Many seabirds have an extraordinary ability to travel long distances and yet show evidence of
542 restricted gene flow and exhibit high levels of philopatry, sometimes returning to breed within
a few meters of their natal nest (Friesen et al. 2007). The ultimate causes for such philopatric
544 behavior are not known, although familiarity with natal and/or previous breeding habitats
(Friesen et al. 2007) and fitness costs incurred by dispersal itself (Clobert et al. 2001) seem
546 likely to be involved.

Our results contradict in some ways the general pattern found in the literature (Friesen et al.
548 2007). According to our models (and recalling that we are considering lower bounds on
migration rates), it seems unlikely that the low breeding dispersal rates reported above for
550 seabirds are compatible with the genetic pattern observed here for the ivory gull, even if natal
dispersal is strong. To be compatible with our observations, a level of breeding dispersal
552 below 0.1 would have to be associated with extremely frequent natal dispersal (that is, a
complete mixture of young adults, see Figure 3 with m_a in 0-0.1). Demographic data from the

554 field will be very important to test this suggestion. Information on the movement behavior of
juvenile birds and additional estimates of adult survival would be particularly valuable.

556

Movement of adult ivory gulls inferred from ecological data

558 Ring recoveries are in line with large-scale movement in ivory gulls and suggest long distance
travel events (> 3400 km; Gaston et al. 2008; Lyngs 2003). However, it is often not known
560 whether recovered birds were actually breeding in the areas where they were found, making
inferences about the frequency of effective dispersal at large spatial scales difficult. Recent
562 advances in movement ecology using satellite transmitters indicated similar post-breeding
flyways over long distance for ivory gulls breeding in the north east Atlantic, *i.e.*, for birds
564 breeding in north Greenland, Svalbard and Franz Josef Land, Russia (Gilg et al. 2010).

Wintering grounds were reached in December, in southeast Greenland and along the Labrador
566 Sea ice-edge, where Canadian birds also overwinter or in the Bering Strait region (Gilg et al.
2010; Mallory et al. 2008). Data also indicate that birds from different colonies, however,
568 migrate eastwards towards wintering area in the Bering Strait region, hence demonstrating a
bi-directional migration pattern (Figure 1).

570 Similar flyways and wintering area for birds from different colonies over the entire species
range may result in the recruitment of birds to distant colonies after the overwinter period
572 (*i.e.*, birds never return to the natal colony). Such movement events may be accidental (*i.e.*,
birds are unable to return to the natal area) or may reflect behavioral variation in philopatry
574 among individuals (Weatherhead and Forbes 1994). The tendency for birds to disperse may
also be linked to the conditions in the natal colony the year they were born and to the local
576 dynamics of the colonies that they recruit to. Such long-distance dispersal events or
reshuffling of individuals on the pre-breeding flyways may be sufficient to eliminate the
578 traces of regional structure among populations. The fidelity of ivory gulls to the breeding site

is unknown but at least some marked individuals return to the same breeding colony from one
580 year to the next (MacDonald 1976), and an example of extreme breeding site fidelity has been
reported (Mallory et al. 2012). Populations that breed on flat land of Russia, where the highest
582 census population size are observed (Table 3), are often prone to move from site to site (de
Korte and Volkov 1993).

584

Dispersal and connectivity under climate change

586 Climate change is geographically shifting the climatic envelope of many species and this is
predicted to occur rapidly in the Arctic (up to ~ 0.40 km/yr; Loarie et al. 2009). The capacity
588 of populations to respond to climate change will depend of evolutionary and demographic
processes (*i.e.*, plasticity, adaptation or migration) (Bourne et al. 2014). Specifically, level of
590 additive genetic variance within population can directly influence evolutionary outcomes in
response to environmental change by providing the necessary genetic variation upon which
592 selection can act (Bourne et al. 2014; Lande and Shannon 1996). Now, our genetic results
suggest high connectivity and gene flow among populations that furthermore still maintain
594 high level of genetic diversity and higher evolutionary potential within each population,
despite recent declines in population census size in some regions (*e.g.*, Canada).

596 Following these results, two important points call however for further investigations. First, the
very high level of genetic connectivity revealed by this study remains difficult to translate
598 into an estimate of demographic connectivity. We have made some efforts to disentangle the
effects of dispersal from local population size, but there remains too much uncertainty in our
600 estimates to determine whether the extent of dispersal that ensures genetic homogeneity is
enough to have an effect on local demography (a relevant issue in high gene flow species;
602 Waples 1998; Waples and Gaggiotti 2006). Information on the behavior of first- and second-
year ivory gulls and adult survival estimates will be key to reduce the space of dispersal

604 parameters that are compatible with our genetic findings. We need to know more about the
movement of birds between their natal site and first breeding attempt. Second, while our
606 findings show that the genetic diversity within colonies is currently high, further studies will
have to determine whether this state is stable or show signs of disequilibrium (*e.g.* in line with
608 findings from demographic surveys that show a strong decline in colony numbers and size).
The effects of overlapping generations and metapopulation functioning will have to be taken
610 into account when looking for genetic signatures of demographic stability or decline (Broquet
et al. 2010; Chikhi et al. 2010; Leblois et al. 2006).

612

Conservation implications

614 Resources for conservation management of endangered species are always limited, and
therefore an understanding of population differentiation and connectivity can help identify
616 conservation priorities and inform management decisions. Here our results indicate that the
ivory gull should be considered a wide-range, genetically homogeneous metapopulation. The
618 lack of population genetic structure found among colonies, in tandem with independent
evidence of movement among colonies, suggests ongoing effective dispersal is occurring
620 across ocean basins. This intercolony movement over large spatial scales can potentially
enhance the persistence of highly fragmented seabird colonies. The generally large
622 nonbreeding component of populations may also play an important role on the structure,
dynamics and persistence of populations in buffering the effects of mortality with
624 compensatory recruitment (although it may also hide a recent population decline, Penteriani et
al. 2011; Votier et al. 2008). Our study suggests immigrant recruitment from distant
626 populations could have similar effects. Understanding patterns of connectivity among disjunct
populations of highly vagile colonial seabirds is vital to appropriately manage their
628 populations and help predict the effect of future environmental change.

630

Acknowledgments

632 This work was supported by grants from foundation Ellis Elliot (Switzerland), Société
vaudoise des Sciences naturelles (Switzerland), and Nos Oiseaux (Switzerland) to GY, by a
634 foundation Agassiz (Switzerland) grant to TB and by the Nicolas Perrin's research group,
Department of Ecology and Evolution at University of Lausanne, Switzerland. Field work in
636 Greenland was supported by the National Geographic Society, Prix Gore-Tex initiative,
Fondation Avenir Finance, the Arctic Ocean Diversity Census of Marine Life Project, CNES,
638 CLS and F. Paulsen. The ivory gull project was part of the work plan of the Joint Norwegian-
Russian Commission on Environmental Protection and fieldwork in Norway and Russia was
640 funded by the Norwegian Ministry of Environment, the Norwegian Polar Institute, Arctic and
Antarctic Research Institute and Russian IPY 2007/08 program. We are grateful to
642 Emmanuelle Pouivé, Brigitte Sabard, Vladimir Sokolov and Oleg Prodan, Vorkuta helicopter
crew commanded by Sergey Kiryushkin for logistical assistance, Russian colleagues Mikhail
644 Ivanov, Andrey and Elena Volkov for participating in fieldwork. Constructive comments by
Nigel G. Yoccoz and two anonymous referees improved the manuscript.

646

References

- 648 ACIA (2004) Impacts of a Warming Arctic: Arctic Climate Impact Assessment. Cambridge
University Press, Cambridge, UK.
- 650 Bell G, Gonzalez A (2011) Adaptation and evolutionary rescue in metapopulations
experiencing environmental deterioration. *Science* 332:1327-1330
- 652 BirdLife International (2012) *Pagophila eburnea*. In: IUCN 2013. IUCN Red List of
Threatened Species. Version 2013.2, <http://www.iucnredlist.org> Accessed 23 January 2014
- 654 Bourne EC, Bocedi G, Travis MJJ, Pakeman RJ, Brooker RW, Schiffrers K (2014) Between
migration load and evolutionary rescue: dispersal, adaptation and the response of
656 spatially structured populations to environmental change. *Proc R Soc Lond, Ser B:
Biol Sci* 281:20132795
- 658 Broquet T, Angelone S, Jaquière J, Joly P, Léna JP, Lengagne T, Plénet S, Luquet E, Perrin N
(2010) Genetic bottlenecks driven by population disconnection. *Conserv Biol*
660 24:1596-1605
- Broquet T, Petit EJ (2009) Molecular Estimation of Dispersal for Ecology and Population
662 Genetics. *Annu Rev Ecol Evol Syst* 40:193-216
- Chen IC, Hill JK, Ohlemüller R, Roy DB, Thomas CD (2011) Rapid range shifts of species
664 associated with high levels of climate warming. *Science* 333:1024-1026
- Chikhi L, Sousa VC, Luisi P, Goossens B, Beaumont MA (2010) The confounding effects of
666 population structure, genetic diversity and the sampling scheme on the detection and
quantification of population size changes. *Genetics* 186:983-995
- 668 Clobert J, Danchin E, Dhondt A, Nichols J (2001) Dispersal. Oxford University Press, New
York
- 670 COSEWIC (2006) COSEWIC Assessment on the Ivory Gull *Pagophila eburnea* in Canada
Committee on the Status of Endangered Wildlife in Canada, Ottawa, ON, Canada
- 672 Crow JF, Aoki K (1984) Group selection for a polygenic behavioral trait: estimating the
degree of population subdivision. *Proc Natl Acad Sci USA* 81:6073-6077
- 674 de Korte J, Volkov AE (1993) Large colony of ivory gulls *Pagophila eburnea* at Domashniy
Island, Severnaya Zemlya. *Sula* 7:107-110
- 676 Dugger KM, Ainley DG, Lyver POB, Barton K, Ballard G (2010) Survival differences and
the effect of environmental instability on breeding dispersal in an Adelie penguin
678 meta-population. *Proc Natl Acad Sci USA* 107:12375-12380
- Environment Canada (2013) Recovery Strategy for the ivory gull (*Pagophila eburnea*) in
680 Canada. Recovery Planning, Environment Canada, Gatineau, Quebec, Canada
- Felsenstein J (1971) Inbreeding and variance effective numbers in populations with
682 overlapping generations. *Genetics* 168:581-597
- Friesen VL, Burg TM, McCoy KD (2007) Mechanisms of population differentiation in
684 seabirds. *Mol Ecol* 16:1765-1785
- Gaston AJ, Brewer D, Diamond AW, Woodsworth EJ, Collins BT (2008) Canadian Atlas of
686 Bird Banding. Volume 2, Seabirds, 1921-1995. Canadian Wildlife Service,
Environment Canada, Canada
- 688 Gauthier G, Milot E, Weimerskirch H (2010) Small-scale dispersal and survival in a long-
lived seabird, the wandering albatross. *J Anim Ecol* 79:879-887
- 690 Gavriilo MV (2011) Ivory gull *Pagophila eburnea* (Phipps, 1774) in the Russian Arctic:
breeding patterns of species within the current species range optimum. PhD, Saint-
692 Petersburg State University, Saint-Petersburg, Russia
- Genovart M, Thibault JC, Igual JM, Bauza-Ribot MD, Rabouam C, Bretagnolle V (2013)
694 Population structure and dispersal patterns within and between Atlantic and
Mediterranean populations of a large-range pelagic seabird. *Plos One* 8

- 696 Gienapp P, Teplitsky C, Alho JS, Mills JA, Merila J (2008) Climate change and evolution:
disentangling environmental and genetic responses. *Mol Ecol* 17:167-178
- 698 Gilchrist HG, Mallory ML (2005) Declines in abundance and distribution of the ivory gull
(*Pagophila eburnea*) in Arctic Canada. *Biol Conserv* 121:303-309
- 700 Gilchrist HG, Strøm H, Gavrilov MV, Mosbech A (2008) International ivory gull conservation
strategy and action plan vol No. 18. Conservation of Arctic Flora and Fauna (CAFF)
702 International Secretariat, Circumpolar Seabird Group (CBird), Akureyri
- Gilg O, Aars J, Fort J, Gauthier G, Gremillet D, Ims RA, Kovacs KM, Meltofte H, Moreau J,
704 Post E, Schmidt NM, Yannic G, Bollache L (2012) Climate change and the ecology
and evolution of Arctic vertebrates. *Ann N Y Acad Sci* 1249:166-190
- 706 Gilg O, Boertmann D, Merkel F, Aebischer A, Sabard B (2009) Status of the endangered
ivory gull, *Pagophila eburnea*, in Greenland. *Polar Biol* 32:1275-1286
- 708 Gilg O, Strøm H, Aebischer A, Gavrilov MV, Volkov A, Miljeteig C, Sabard B (2010) Post-
breeding movements of the northeast Atlantic ivory gull *Pagophila eburnea*
710 populations. *J Avian Biol* 41:532-542
- Goudet J (2005) FSTAT: a program to estimate and test gene diversities and fixation indices
712 (version 2.9.3). Available from http://www.unil.ch/dee/page6767_en.html. University of
Lausanne, Lausanne
- 714 Goudet J, Raymond M, deMeeus T, Rousset F (1996) Testing differentiation in diploid
populations. *Genetics* 144:1933-1940
- 716 Greenwood PJ, Harvey PH (1982) The natal and breeding dispersal of birds. *Annu Rev Ecol
Syst* 13:1-21
- 718 Hill WG (1972) Effective size of populations with overlapping generations. *Theor Popul Biol*
3:278-289
- 720 Hoenig JM, Heisey DM (2001) The abuse of power: The pervasive fallacy of power
calculations for data analysis. *The American Statistician* 55:19-24
- 722 Hoffmann AA, Sgro CM (2011) Climate change and evolutionary adaptation. *Nature*
470:479-485
- 724 IPCC (2007) Climate change 2007: impacts, adaptation and vulnerability. Contribution of
working group II to the Fourth Assessment Report of the intergovernmental panel on
726 climate change. Cambridge University Press, New York, USA.
- Jones O, Wang J (2009) COLONY: a program for parentage and sibship inference from
728 multilocus genotype data. *Mol Ecol Resour* 10:551-555
- Kimura M (1969) Number of heterozygous nucleotide sites maintained in a finite population
730 due to steady flux of mutations. *Genetics* 61:893-903
- Kimura M, Crow J (1963) The measurement of effective population number. *Evolution*
732 17:279-288
- Lande R, Shannon S (1996) The role of genetic variation in adaptation and population
734 persistence in a changing environment. *Evolution* 50:434-437
- Laporte V, Charlesworth B (2002) Effective population size and population subdivision in
736 demographically structured populations. *Genetics* 162:501-519
- Lawrence H, Lyver P, Gleeson D (2014) Genetic panmixia in New Zealand Grey-faced
738 Petrel: Implications for conservation and restoration. *Emu* 114:249-258
- Leblois R, Estoup A, Streiff R (2006) Genetics of recent habitat contraction and reduction in
740 population size: does isolation by distance matter? *Mol Ecol* 15:3601-3615
- Lebreton JD, Hines JE, Pradel R, Nichols JD, Spendelov JA (2003) Estimation by capture-
742 recapture of recruitment and dispersal over several sites. *Oikos* 101:253-264
- Loarie SR, Duffy PB, Hamilton H, Asner GP, Field CB, Ackerly DD (2009) The velocity of
744 climate change. *Nature* 462:1052-1055

- 746 Lorenzen ED, Nogues-Bravo D, Orlando L, Weinstock J, Binladen J, Marske KA, Ugan A,
748 Borregaard MK, Gilbert MTP, Nielsen R, Ho SYW, Goebel T, Graf KE, Byers D,
750 Stenderup JT, Rasmussen M, Campos PF, Leonard JA, Koepfli KP, Froese D, Zazula
752 G, Stafford TW, Aaris-Sorensen K, Batra P, Haywood AM, Singarayer JS, Valdes PJ,
754 Boeskorov G, Burns JA, Davydov SP, Haile J, Jenkins DL, Kosintsev P, Kuznetsova
756 T, Lai XL, Martin LD, McDonald HG, Mol D, Meldgaard M, Munch K, Stephan E,
758 Sablin M, Sommer RS, Sipko T, Scott E, Suchard MA, Tikhonov A, Willerslev R,
760 Wayne RK, Cooper A, Hofreiter M, Sher A, Shapiro B, Rahbek C, Willerslev E
(2011) Species-specific responses of Late Quaternary megafauna to climate and
762 humans. *Nature* 479:359-364
- 764 Lyngs P (2003) Migration and winter ranges of birds in Greenland: An analysis of ringing
766 recoveries. *Dansk Ornitologisk Forenings Tidsskrift* 97:1-167
- 768 MacDonald SD (1976) Phantoms of the polar pack-ice. *Audubon* 78:2-19
- 770 Mallory ML, Allard KA, Braune BM, Gilchrist HG, Thomas VG (2012) New longevity
772 record for ivory gulls (*Pagophila eburnea*) and evidence of natal philopatry. *Arctic*
774 65:98-101
- 776 Mallory ML, Stenhouse IJ, Gilchrist HG, Robertson G, Haney JC, Macdonald SD (2008)
778 Ivory gull (*Pagophila eburnea*). *The Birds of North America Online* (A. Poole, Ed.).
780 Ithaca: Cornell Laboratory of Ornithology; Retrieved from *The Birds of North*
782 *America Online*: <http://bna.birds.cornell.edu/BNA/species/175/>. DOI: 10.2173/bna.175.
- 784 Milot E, Weimerskirch H, Bernatchez L (2008) The seabird paradox: dispersal, genetic
786 structure and population dynamics in a highly mobile, but philopatric albatross
788 species. *Mol Ecol* 17:1658-1673
- 790 Nisbet ICT, Cam E (2002) Test for age-specificity in survival of the common tern. *Journal of*
792 *Applied Statistics* 29:65-83
- 794 Parmesan C (2006) Ecological and evolutionary responses to recent climate change. *Annual*
796 *Review of Ecology Evolution and Systematics* 37:637-669
- 798 Penteriani V, Ferrer M, Delgado MM (2011) Floater strategies and dynamics in birds, and
800 their importance in conservation biology: towards an understanding of nonbreeders in
802 avian populations. *Anim Conserv* 14:233-241
- 804 Rousset F (2001) Genetic approaches to the estimation of dispersal rates. In: Clobert J,
806 Danchin E, Dhondt AA, Nichols JD (eds) *Dispersal*. Oxford University Press, New
808 York,
- 810 Rousset F (2004) *Genetic Structure and Selection in Subdivided Populations*. Monographs in
812 *Population Biology*. Princeton University Press, Princeton and Oxford
- 814 Royston S, Carr S (2014) Conservation genetics of high-arctic Gull species at risk: I.
816 Diversity in the mtDNA control region of circumpolar populations of the Endangered
818 Ivory Gull (*Pagophila eburnea*). *Mitochondrial DNA* 26:1-4
- 820 Ryman N, Palm S (2006) POWSIM: a computer program for assessing statistical power when
822 testing for genetic differentiation. *Mol Ecol Notes* 6:600-602
- 824 Slatkin M (1991) Inbreeding coefficients and coalescence times. *Genetics Research* 58:167-
826 175
- 828 Spencer NC, Gilchrist HG, Mallory ML (2014) Annual movement patterns of endangered
830 ivory gulls: the importance of sea ice. *Plos One* 9:e115231
- 832 Stenhouse IJ, Robertson GJ, Gilchrist HG (2004) Recoveries and survival rates of ivory gulls
834 (*Pagophila eburnea*) banded in Nunavut, Canada, 1971–1999. *Waterbirds* 27:486-492
- 836 Stewart JR, Dalen L (2008) Is the glacial refugium concept relevant for northern species? A
838 comment on Pruett and Winker 2005. *Clim Change* 86:19-22
- 840 Stewart JR, Lister A (2001) Cryptic northern refugia and the origins of the modern biota.
842 *Trends in Ecology and Evolution* 16:608-613

- 796 Travis JMJ, Delgado M, Bocedi G, Baguette M, Barton K, Bonte D, Boulangeat I, Hodgson
JA, Kubisch A, Penteriani V, Saastamoinen M, Stevens VM, Bullock JM (2013)
Dispersal and species' responses to climate change. *Oikos* 122:1532-1540
- 798 Votier SC, Birkhead TR, Oro D, Trinder M, Grantham MJ, Clark JA, McCleery RH,
Hatchwell BJ (2008) Recruitment and survival of immature seabirds in relation to oil
800 spills and climate variability. *J Anim Ecol* 77:974-983
- 802 Wakeley J (2009) Coalescent theory. Roberts and Company Publishers, Greenwood Village,
Colorado
- 804 Wang J (2012) Computationally efficient sibship and parentage assignment from multilocus
marker data. *Genetics* 191:183-194
- 806 Waples RS (1998) Separating the wheat from the chaff: patterns of genetic differentiation in
high gene flow species. *J Hered* 89:438-450
- 808 Waples RS, Gaggiotti O (2006) What is a population? An empirical evaluation of some
genetic methods for identifying the number of gene pools and their degree of
connectivity. *Mol Ecol* 15:1419-1439
- 810 Waples RS, Teel DJ (1990) Conservation genetics of Pacific salmon. I. Temporal changes in
allele frequency. *Conserv Biol* 4:144-156
- 812 Weatherhead P, Forbes MRL (1994) Natal philopatry in birds: genetic or ecological
influences? *Behavioural Ecology* 5:426-433
- 814 Weir B, Cockerham C (1984) Estimating F-statistics for the analysis of population structure.
Evolution 38:1358-1370
- 816 Welch AJ, Fleischer RC, James HF, Wiley AE, Ostrom PH, Adams J, Duvall F, Holmes N,
Hu D, Penniman J, Swindle KA (2012) Population divergence and gene flow in an
818 endangered and highly mobile seabird. *Heredity* 109:19-28
- 820 Westneat DF, Stewart IRK (2003) Extra-pair paternity in birds: causes, correlates, and
conflict. *Annual Review of Ecology, Evolution, and Systematics* 34:365-396
- 822 Wojczulanis-Jakubas K, Kilikowska A, Harding AMA, Jakubas D, Karnovsky NJ, Steen H,
Strom H, Welcker J, Gavriilo M, Lifjeld JT, Johnsen A (2014) Weak population
genetic differentiation in the most numerous Arctic seabird, the little auk. *Polar Biol*
824 37:621-630
- 826 Wright S (1931) Evolution in mendelian populations. *Genetics* 16:97-159
- 828 Yannic G, Aebischer A, Sabard B, Gilg O (2014a) Complete breeding failures in ivory gull
following unusual rainy storms in North Greenland. *Polar Res* 33:22749
- 830 Yannic G, Pellissier L, Ortego J, Lecomte N, Couturier S, Cuyler C, Dussault C, Hundermark
KJ, Irvine RJ, Jenkins DA, Kolpashikov L, Mager K, Musiani M, Parker KL, Røed
KH, Sipko T, Þórisson SG, Weckworth BV, Guisan A, Bernatchez L, Côté SD
832 (2014b) Genetic diversity in caribou linked to past and future climate change. *Nature
Climate Change* 4:132-137
- 834 Yannic G, Sermier R, Aebischer A, Gavriilo MV, Gilg O, Miljeteig C, Sabard B, Strom H,
Pouive E, Broquet T (2011) Description of microsatellite markers and genotyping
performances using feathers and buccal swabs for the ivory gull (*Pagophila eburnea*).
836 *Mol Ecol Resour* 11:877-889
- 838 Yearsley J, Viard F, Broquet T (2013) The effect of collective dispersal on the genetic
structure of a subdivided population. *Evolution* 67:1649-1659

840 **Table 1.** Estimates of genetic variability for sampled sites of ivory gull (*Pagophila eburnea*). *N* gives the number of samples genotyped at 13
 microsatellite markers. Statistics include number of alleles (*nA*), allelic richness (*Ar*; estimated for $n \geq 10$ individuals and based on min. sample
 842 size of 8 diploid individuals successfully genotyped at 13 loci), observed heterozygosity (H_O) and expected heterozygosity (H_E). Sampled areas
 are identified by their abbreviation (Abbr.) throughout the study

ID	Country	Estimated regional population	Region	Site	Abbr.	Latitude	Longitude	Status	DNA source	N	<i>nA</i>	<i>Ar</i>	H_O	H_E		
#1	Greenland	> 2000 pairs ^a	National Park	Station Nord	StNo	81.60	-16.66	adult unknown	swab	81	9.9	6.01	0.76	0.78		
#2							StBr	81.61	-16.49	adult breeding	swab	25	8.3	6.08	0.77	0.78
#3							StJv	81.61	-16.49	juvenile	swab/tissue	19/1	7.6	5.95	0.82	0.79
#4						Amdrup Land	AmLa	80.85	-14.63	juvenile	swab/tissue	33/12	9.2	6.04	0.79	0.79
#5	Norway	350-500 pairs ^b	Svalbard	Svenskoya	Sven	78.72	26.63	adult breeding	blood	9	6.8	-	0.84	0.82		
#6						Auga	Auga	78.50	21.74	adult breeding	swab/blood	1/17	7.5	6.05	0.76	0.78
#7						Hübnerbreen	HübN	78.41	21.69	adult breeding	swab	7	5.7	-	0.85	0.76
#8						Freemanbreen	Free	78.38	21.43	adult breeding	swab/plucked feathers	34/2	9.7	6.50	0.76	0.80
#9	Russia	14,500 – 22,000 pairs ^c	Franz Josef Land	Nagurskoje	Nagu	80.72	48.22	adult unknown	shed feathers	5	3.9	-	0.63	0.73		
#10						Rudolf Island	Rudo	81.75	58.39	adult unknown	shed feathers	17	7.4	6.12	0.67	0.79
#11						Eva-Liv Island	EvLi	81.64	63.22	adult unknown	shed feathers	5	4.8	-	0.70	0.80
#12						Severnaya Zemlya	SchI	81.04	90.76	adult unknown	shed feathers	12	6.6	5.94	0.73	0.76
#13						Domashny Island	Doma	79.51	94.84	adult unknown	shed feathers/swab	17/6	8.5	6.19	0.80	0.77
#14						Komsomalets Island	Koms	80.77	91.05	adult unknown	shed feathers	6	5.7	-	0.83	0.80
#15						Sukhaya River	Sukh	80.77	96.75	juvenile	shed feathers	7	5.8	-	0.76	0.80
#16					Kara Sea Islands	Heiberg Islands	HeiI	77.61	101.51	adult unknown	shed feathers	4	4.2	-	0.73	0.77
#17	Canada	900 pairs ^d	Nunavut	Seymour Island	SeyI	76.80	-101.27	adult breeding	Swab/plucked feathers	11	6.5	5.84	0.75	0.78		
#18						Ellesmere Island (Alert)	AlEI	82.50	-62.33	adult unknown	blood	12	6.7	5.87	0.80	0.77
		19,000 – 27,000 pairs											343			

844 ^a Gilg *et al.* (2009); ^b Gilchrist *et al.* (2008); ^c Gavrilov (2011); ^d Environment Canada (2013)

Table 2. F_{ST} and exact G -test probability values obtained for each autosomal microsatellite846 and over all loci for two different datasets of ivory gull (*Pagophila eburnea*)

Loci	All adults sites ($n = 15$)		Breeding colonies ($n = 6$)	
	F_{ST}	P -value	F_{ST}	P -value
A111	-0.005	0.78	-0.014	0.94
B125	0.002	0.32	0.004	0.61
C7	0.009	0.83	0.003	0.45
D126	0.004	0.27	0.002	0.62
D5	0.004	0.17	0.013	0.16
D9	-0.008	0.76	0.001	0.23
A112	-0.003	0.65	-0.012	0.99
A132	0.010	0.37	-0.011	0.85
B114	-0.009	0.76	-0.010	0.92
D103	0.006	0.11	-0.004	0.62
C6	0.002	0.20	0.018	0.54
B103	0.008	0.62	-0.006	0.08
D1	0.007	0.04	0.013	0.04
Over all loci	0.001	0.09	-0.000	0.61
Jackknifing over loci	0.001		0.003	
Bootstrapping 95% CI	-0.002;0.005		-0.006;0.005	

848

Table 3. Census colony size across the breeding distribution of ivory gull (*Pagophila*

850 *eburnea*)

Country	Number of ivory gulls	Number of colonies
Greenland ^a	Records between 1854 and 2009	
	<5	13
	5-24	6
	25-99	11
	100-300	5
Norway ^b	Maximum records	
	<5	7
	4-10	5
	11-30	19
	31-60	7
	61-100	3
Russia ^c	Historically maximum records	
	2 – 20	>10
	22 – 100	19
	200 – 700	13
	800 – 1600	7
	2000 +	5
	1990s - 2000s	
	2 – 20	0
	22 – 100	17
	200 – 700	11
800 – 1600	6	
2000 +	3	
Canada ⁴	Historically records	
	Between 1976 and 1992	
	<5	0
	5 – 24	3
	25 – 50	6
	50 – 99	6
	100 – 340	2
	Recent time records between	
	2001 and 2003	
	<5	10
5 – 24	9	
25 – 50	2	
50 – 99	1	
100 – 300	0	

¹ Gilg et al. (2009); ² Norwegian Polar Institute; ³ Maria Gavrilo, unpublished data; ⁴ Gilchrist and Mallory

852 (2005)

854 **Titles and legends to figures**

856 **Figure 1.** Map of the study area illustrating the Holarctic distribution of ivory gull (*Pagophila*
eburnea) breeding colonies. Sampling localities are indicated by the ID corresponding with
858 Table 1; orange dots depict known breeding sites (Gilchrist et al. 2008). Dashed lines:
wintering grounds (variable during the winter and among years according to the extension of
860 the sea-ice; modified from Gilg *et al.* (2010)). Background map represent the maximum sea-
ice extent in July between 1979-2013 (light blue) and the sea-ice extent in July 2013 (dark
862 blue) (data from the National Snow and Ice Data Centre, Boulder, Colorado;
<http://nsidc.org/>).

864

Figure 2. Statistical power for obtaining significant outcomes in tests of genetic
866 differentiation involving the specific marker characteristics and sample sizes of ivory gull for
i) all adults localities and ii) breeding colonies only. Simulations were performed using
868 POWSIM version 4.1 (Ryman and Palm 2006). The dotted lines indicate the level of genetic
differentiation that can be detected with 90% statistical power for the two data sets.

870

Figure 3. The parameter space (natal dispersal, m_j , breeding dispersal, m_a , effective colony
872 size, N_a (contour lines) and adult survival probability, v) of an overlapping generation model
that predicts a global equilibrium F_{ST} (equation 5) equal to the observed value for ivory gull
874 (using all samples $F_{ST} = 0.001$). Given N_a , the plot shows the combinations of natal and
breeding dispersal that are required to yield the observed genetic structure in ivory gulls
876 across its distribution range. The dashed lines in panel A show an example: with $v = 0.86$ and
 $N_a = 250$, a combination of 25% natal dispersal and 16.5% breeding dispersal would predict
878 $F_{ST} = 0.001$ in the simplified conditions of our model





