Unravelling the impact of harvesting pressure on canopy-forming macroalgae

Doriane Stagnol, Renaud Michel, Dominique Davoult

► To cite this version:

Doriane Stagnol, Renaud Michel, Dominique Davoult. Unravelling the impact of harvesting pressure on canopy-forming macroalgae. *Marine and Freshwater Research*, 2015, 67 (1), pp.153-161. 10.1071/MF14198 . hal-01203192

HAL Id: hal-01203192

<https://hal.sorbonne-universite.fr/hal-01203192>

Submitted on 22 Sep 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Unravelling the impact of harvesting pressure on canopy-forming macroalgae

Doriane Stagnol ^{a,b*}, Renaud Michel ^{a,b}, Dominique Davoult ^{a,b}

^a Sorbonne Universités, UPMC Univ Paris 6, Station Biologique, Place Georges Teissier, F-29682 Roscoff, France

^b CNRS, UMR 7144 AD2M, Place Georges Teissier, F-29682, Roscoff, France

* Corresponding author: doriane.stagnol@gmail.com

Abstract

Canopy-forming macroalgae create a specific surrounding habitat (the matrix) with their own ecological properties. Previous studies have shown a wide range of responses to canopy removal. Magnitude and strength of the effects of harvesting are thought to be context-dependent, with the macroalgal matrix that can either soften or exacerbate the impact of harvesting. We experimentally examined *in situ* the effect of harvesting on targeted commercial species, and how these potential impacts might vary in relation to its associated matrix. We found that patterns of recovery following the harvesting disturbance were variable and matrix specific, suggesting that local factors and surrounding habitats characteristics mediated the influence of harvesting. The greatest and longest effects of harvesting were observed for the targeted species that created a dominant and monospecific canopy on their site prior to the disturbance. Another relevant finding was the important natural spatiotemporal variability of macrobenthic assemblages associated with canopy-forming species, which raises concern about the ability to discriminate the natural variability from the disturbance impact. Finally, our results support the need to implement ecosystem-based management, assessing both the habitats conditions and ecological roles of targeted commercial species, in order to insure the sustainability of the resource.

Keywords: benthic, biodiversity, ecology, ecosystem-based management (EBM)

Running head: Unravelling the impact of harvesting pressure

Introduction

Intertidal macroalgae are commonly recognized as foundation species that have an important effect on community structure by creating habitats and modifying environmental conditions, species interactions, and resource availability (Bruno and Bertness 2001). Loss of habitat-forming species, for example through harvesting, is a key threat to ecosystems. Previous studies have shown a wide range of responses to canopy removal, including reduction or loss of diversity, decrease in invertebrate abundance and richness (Benedetti-Cecchi *et al.* 2001), shifts in community composition (Schiel and Lilley 2007), replacement by grazers or turfs (Perkol-Finkel and Airoidi 2010), or reduction in algal biomass and primary productivity (Gollty *et al.* 2008; Tait and Schiel 2011). Recovery of algal and invertebrate communities may take up to 4, 6 or even 12 years (Foster *et al.* 2003; Jenkins *et al.* 2004; Araujo *et al.* 2012).

Although the short and long-term consequences of losing habitat-forming species on community structure and composition are progressively understood (Benedetti-Cecchi *et al.* 2001), very little research has been conducted *in situ* to examine the effect of macroalgal manual harvesting depending on the identity of the targeted species, which have different life cycles and population structures. Each canopy-forming macroalgal species creates a specific surrounding habitat (the matrix) with its own ecological properties. The magnitude and the strength of the effects of harvesting are thought to be context-dependent, with the macroalgal matrix that can either soften or exacerbate the impact of harvesting (Mayer-Pinto *et al.*, 2015). The existence of a current global decline in rocky shore habitat-forming macroalgae from a wide range of stressors has been broadly documented (e.g. Airoidi *et al.* 2008; Halpern *et al.* 2007). Intertidal macroalgae are harvested worldwide for human consumption, alginate production, cosmetic and pharmaceutical industries, agricultural supply, or potentially biofuel. As the demand has intensified in recent decades and will most certainly continue to expand, assessing the harvesting impact is of major importance to define the potential consequences of disturbances to critical ecosystem functions (e.g. primary productivity).

In this context, we ask (1) whether harvesting has significant impacts on the targeted species, and (2) how these potential impacts might vary in relation to the targeted species and its associated macroalgal matrix. Further we discuss the suitability of management procedure in place (in Brittany) from an ecosystem-based management (EBM) perspective. EBM is widely accepted as the new paradigm to manage marine ecosystems (McLeod and Leslie, 2009), but

is still relatively new in marine habitats and in particular for rocky shores. Besides the scientific interest, this study is also of political and commercial importance since it addresses a practical management issue - the effect of resource harvesting - using the approach of fundamental and applied research. Given the global trend of declining biodiversity (Sala and Knowlton 2006; Butchart *et al.* 2010), the present challenge is to understand the interactive effects of this stressor along with our fundamental knowledge of seaweed ecosystem structure and function, and use the information to facilitate EBM of these important and productive coastal marine systems.

Materials and methods

Study area and studied species

This study was performed in 2011 in Brittany, the French region in which commercial seaweed harvesting is the most important. Seven study locations, where macroalgae are commonly harvested were selected on the coast between Bloscon (48°N43'30.59", 3°W58'05.75") and Landunvez (48°N31'51.10", 4°W45'56.92"). These locations are dominated by hard substratum, they differ, however, in terms of geomorphology, habitat structure (heterogeneity), and exposition to wave action. They were mainly chosen for large seaweed standing crop and accessibility.

In each location, the targeted species were making up the canopy on the study site and were considerably larger than the other members of the algal assemblage, forming an extensive layer above them: *Chondrus crispus* at location 1, *Fucus serratus* at location 2, *Palmaria palmata* at location 3 (epiphytic on a *F. serratus* canopy), *Porphyra linearis* at location 4, *Porphyra umbilicalis* at location 5, *P. palmata* at location 6 (epilithic), and *Himanthalia elongata* at location 7. It is important to note that at location 1, the dominant canopy was a mixture of small canopy species *C. crispus* and *Mastocarpus stellatus* that could not be harvested separately.

Matrices

Here we defined 3 types of matrices (derived from T0 sampling) with their representatives locations: matrix A, associated with a monospecific canopy accounting for a high percentage of the seaweed cover in the community and with an intermediate to high associated fauna

diversity (locations 2 and 7; with *F. serratus* (A1) and *H. elongata* (A2) as respective dominant canopy-forming species); matrix B, associated with a mixed canopy accounting for an intermediate percentage of the seaweed cover in the community and with an intermediate associated fauna diversity (locations 3, 6, and 1; with *P. palmata*, either epiphytic (B1) or epilithic (B2), and *C. crispus* (B3) as respective dominant canopy-forming species); and matrix C, associated with a monospecific canopy accounting for an intermediate to high percentage of the seaweed cover in the community and with a low associated fauna diversity (locations 4 and 5, with *P. linearis* (C1) and *P. umbilicalis* (C2) as respective dominant canopy-forming species). These seaweed species were chosen because they have been significantly exploited for many decades and because they are mostly found in the midlittoral zone.

Experimental design

Traditional before–after control–impact (BACI) designs consider one impact and one control location without replicating sites (Underwood 1994). Using only one control site, however, prevents from disentangling the effect of the impact from that of different natural variability between the impact and control sites.. Here, the impact of a harvest disturbance event on the ecosystem was investigated experimentally using a multiple before–after control–impact (MBACI) design to address concerns of BACI related to the confounding effects of spatial and temporal variation (Underwood 1994; Stewart-Oaten and Bence 2001). Sampling multiple control sites and multiple impact sites increases the probability that observed differences are due to the impact. Each studied matrix has replicated locations: locations 2 and 7 for matrix A, locations 1, 3 and 6 for matrix B; and locations 4 and 5 for matrix C. On each location, one site was experimentally impacted and one site served as control. Harvest was simulated by removing the species biomass. The harvest was carried out according to the legal French recommendations into force in 2011: all individuals of *F. serratus* and *C. crispus* were removed, while only *H. elongata* individuals longer than 80 cm and *P. palmata* and *Porphyra* sp. individuals longer than 25 cm were removed. To really assess the impact of traditional harvesting in Brittany, however, we choose to reproduce the harvest practices as they are really applied by the harvesters. Therefore, the harvesting of *P. palmata*, *P. linearis*, and *P. umbilicalis* was made by tearing off the individuals (and not by cutting above the holdfast as stated by the legal French recommendations), which appears to be the most suitable way to harvest, if not the only technically feasible for these species.

Community structure

On each site, five 0.1 m² (32 cm x 32 cm) random quadrats were sampled. Sampling was undertaken prior to manipulation and at intervals thereafter for 12 months. For each sample, the cover index of the target species was estimated from the percentage cover of the sampling surface, using the Braun-Blanquet cover scale: 0 (absent), 1]0-25%], 2]25-50%], 3]50-75%] and 4]75-100%]. The taxonomic richness was recorded *in situ* by identifying every taxon (fauna and flora) visible to the eye to the lowest possible taxonomic level. Diversity is thus referred as «taxonomic diversity» hereafter. Animal abundance (as number of individuals) was also recorded *in situ* for each animal taxa, excluding colonial organisms and two polychaetes *Pomatoceros triqueter* and Spirorbidae, whose densities are very difficult to estimate. We also calculated evenness for each quadrat. The removal of the canopy may have different impacts on associated species, depending whether canopy provides them with support, shelter or food. It is therefore interesting to study the effect of harvesting on macrofauna by distinguishing species according to their motility: mobile or sessile, and according to their trophic regime: suspension feeders, grazers or carnivores in the broadest sense, i.e. predators, omnivores, scavengers, etc.

Statistical Analyses

For each matrix type analyses were made to test the effect of the experimental harvest disturbance using a permutational multivariate analysis of variance (PERMANOVA) design that included four factors: (1) before vs. after the disturbance event (BA: 2 levels, fixed and crossed), (2) controls vs. impacted (CsI: 2 levels, fixed and crossed), (3) time (T: 4 levels, fixed and nested within BA), and (4) locations (L: 2 or 3 levels, random and nested in CsI). In the MBACI design, the impact of harvesting is identified by a significant Before-vs-After x Controls-vs-Impacted (BA x CsI) interaction indicating an overall difference between the impacted sites compared to the controls from before to after the harvest disturbance. A significant CsI x T(BA) could also indicate a delayed impact of the harvesting disturbance. We examined biological responses to the harvesting disturbance for four compound measures of community structure (i.e., canopy cover index, algal taxonomic diversity, animal taxonomic diversity, and total animal abundance). We also examined multivariate differences in assemblage structure of the animal community associated with the matrices, using the same design as above. Analyses were made with the PERMANOVA+ add-on package for PRIMER v6 (Anderson *et al.*, 2008). To test for any animal taxonomic composition differences

between the assemblages of control and impact sites, separate analysis of similarity (ANOSIM) tests were done for each matrix at each sampling period on animal presence/absence data. If overall significant differences in animal assemblages between the two sites were obtained, we used an analysis of similarity percentages (SIMPER) in PRIMER to determine the species mainly responsible for the dissimilarity between treatments. A Principal Component Analysis (PCA) was made, using Statbox software ©, for the ordination of quadrats based on standardized data of community structure (i.e., canopy cover index, algal taxonomic diversity, taxonomic diversity of both sessile and mobile taxa, total animal abundance, and evenness).

Results

Effects of loss of canopy on harvested species

Harvesting of canopy significantly affected the cover index of the targeted species associated with the matrix A (PERMANOVA, BA x CsI, $p < 0.05$, Table 1). Differences in the cover index of the targeted species between the control and the impact sites were maintained up to 9 months after the disturbance event (Fig. 1a). The cover index of the targeted species associated with the matrix B varied substantially at the spatial scale (PERMANOVA, L(CsI), $p < 0.001$, Table 1), but no significant interaction was observed. Finally, the cover index of the targeted species associated with the matrix C varied significantly before and after the disturbance but with different effects between the two locations (PERMANOVA, BA x L(CsI), $p < 0.001$, Table 1). Note that one month after the initial harvest of *P. linearis*, sand started to silt the study site, reaching a thickness of 1.5 m seven months after the start of the study. After the sand was removed, the initial patchy distribution of *P. linearis* was replaced by a *P. linearis* bloom with a cover of 100% (see Stagnol *et al.* 2013).

Effects of loss of canopy on community structure

Harvesting of canopy affected algal communities associated with the dominant-canopy forming species of matrix A, but only at one location (PERMANOVA, BA x L(CsI), $p < 0.01$, Table 1). Indeed, we observed a significant increase in algal diversity associated with *F. serratus* on the impact site three months after the disturbance event. In contrast, no differences were observed between the control and the impacts sites of *H. elongata* over the whole study. PERMANOVA revealed a significant CsI x T(BA) interaction (Table 1) for the animal taxonomic diversity associated with the targeted species of matrix A, indicating that

effects of harvesting were significant but variable over time (Fig. 1, Table 1). Regarding the animal abundance, we observed significant effects of disturbance, variable over the two locations (PERMANOVA, BA x L(CsI), $p < 0.05$, Table 1).

The algal and animal diversities and the fauna abundance of communities within matrix B displayed a significant spatial variability (PERMANOVA, L(CsI), $p < 0.05$, Table 1, Fig. 1). This spatial variability between and within locations made it difficult to determine whether the harvesting of the canopy was having an impact that was separate and distinct from natural spatial and temporal variation.

PERMANOVA revealed a significant T(BA) x L(CsI) interaction for the algal and animal taxonomic diversity associated with the targeted species within matrix C (Table 1, $p > 0.001$). This indicates that these parameters varied substantially over spatial and temporal scales. It is interesting to note that similarly to matrix A, we observed a significant increase in algal diversity associated with *P. umbilicalis* on the impact sites three months after the disturbance event. No significant effects or differences were observed for the animal abundance associated with the dominant-canopy forming species of the matrix C.

Harvesting of canopy has significantly affected the benthic macrofauna assemblages associated with the species of matrix A three months after the disturbance event (Table 2). For matrix A, these differences were mostly the results of a lesser occurrence of sessile invertebrates on the impact sites, most of them being suspension-feeders, such as the polychaete *Pomatoceros triqueter*, the tunicates *Botryllus schlosseri* and Polyclinidae, the bryozoans *Alcyonidium* sp. and *Schizoporella unicornis*. In addition, we observed a great decrease in the abundance of the dominant species, the gastropod *Gibbula pennanti*, on the impact sites increasing the evenness (and then the diversity) of the associated benthic macrofauna assemblages (SIMPER routine, Primer). We also observed a significant difference between the control and impact sites in the benthic macrofauna assemblages associated with the species of matrix C three months after the disturbance event (Table 2). These differences were mostly the results of a lesser occurrence of the gastropods *Nucella lapillus* and *G. pennanti* on the impact sites (SIMPER routine, Primer).

Variation in relation to the targeted species and its associated matrix

The means of canopy cover index ranged from 0.8 to 4, number of sessile taxa from 0 to 5.2 species, number of mobile taxa from 0 to 7.4 species, animal abundance from 0 to 676

individuals, evenness from 0 to 1. PCA revealed the dynamics of impacted and control communities of the six species through the 1-year study (Fig. 2). The first two axes explained 62% of the variance in the benthic macrofauna assemblages, with 40% explained by the first axis and 22% by the second one. Pearson correlations of input variables suggested a significant correlation of both the number of sessile and mobile taxa with the algal diversity ($p < 0.05$), while the canopy cover index is significantly correlated with the animal abundance ($p < 0.05$).

The first principal component axis (PC1) differentiated plots along a gradient from high number of mobile taxa ($r = 0.93$) and high animal abundance ($r = 0.84$) at positive PC1 scores, to low number of mobile taxa and low animal abundance at negative PC1 scores. To a lesser extent, the number of sessile taxa ($r = 0.62$) and algal diversity ($r = 0.58$) also contributed to the first axis. The second principal component axis (PC2) differentiated plots along a gradient from high evenness ($r = 0.7$) and low canopy cover index at positive PC2 scores, to low evenness and high canopy cover index ($r = -0.48$) at negative PC2 scores.

The impact of harvesting was not clearly witnessed for the communities associated with *P. palmata* (B1), i.e. epiphytic *P. palmata*, *P. linearis* (C1), and *P. umbilicalis* (C2). The eight samples of *P. palmata* (B1), i.e. epiphytic *P. palmata*, were noticeably adjacent regardless of the sampling time, and visibly separated from the other communities by the second axis. The benthic community associated with *P. palmata* had generally a great evenness, which is the major determinant factor for this community. Benthic communities associated with *P. linearis* (C1) and *P. umbilicalis* (C2) were structurally similar, occupying a similar PCA space. The close association of the *P. linearis* and *P. umbilicalis* samples in the PCA (Fig. 2) reflects the relative similarity of the benthic communities associated with these two species. They are characterized by low number of mobile taxa, low animal abundance, high cover index and low evenness (except 4 plots noticeably separated from the others by the second axis). Besides, the natural variability that occurred on the *P. umbilicalis* site at T9, in the form of a significant sand burial of the site, affected both control and impact plots (C2C9 and C2I9).

On the other hand, communities associated with *F. serratus* (A1), *H. elongata* (A2), *P. palmata* (B2), i.e. epilithic *P. palmata*, and *C. crispus* (B3) displayed an initial impact of harvesting along with a progressive recovery. The benthic community associated with *F. serratus* (A1) changed little over time in the control plots, showing a high number of mobile taxa and a great animal abundance (described by the first axis) as well as a high cover index

and a low evenness (described by the second axis). The impacted plots followed a trajectory away from the control plots, with 2 impacted plots (A1I3 and A1I9) noticeably separated from the others. The impact plot at T3 was separated from the others by both axes, displaying a low number of mobile taxa, a low animal abundance, a low cover index and a relatively high evenness ($E = 0.70$). On the other hand, the impact plot at T9 was separated from the others by the second axis only, also displaying a low cover index and a relatively high evenness ($E = 0.70$) but a quite high number of mobile taxa and high animal abundance. A similar pattern was observed for the community associated with *H. elongata* (A2), although the eight plots were more dispatched on the PCA space. Still, two impact plots (A2I3 and A2I9) were remarkably detached from the others by the first axis. Indeed, they both showed a low animal abundance and a low algal diversity, coupled with a low cover index (although this last variable is not discriminated by the first axis).

Although communities associated with *P. palmata* (B2), were not visibly separated from the other communities, the trajectories were different between the control and impact sites in spite of the very little temporal variation. Finally, the benthic community associated with *C. crispus* (B3) also showed different time trajectories between the control plots and the impact plots, in spite all the plots being rather adjacent in the PCA space.

Discussion

1. Different patterns of recovery in relation to the matrix identity (surrounding habitat)

There has been some evidence over the study that ecosystem recovery (after harvest disturbance) has occurred. Twelve months after the initial harvesting, the effects of harvesting on the ecosystem were not detectable at any location: all community structures and assemblages on the impact sites were similar to the ones observed on the control sites. The associated ecosystems impacted by the canopy removal have all returned to a state comparable to the ones seen on an undisturbed zone. Patterns of recovery, however, were quite variable and matrix specific, suggesting that local factors and surrounding habitat characteristics mediated the influence of harvesting on the ecosystem (Foster *et al.* 2003).

The three matrices in our study demonstrated strikingly different patterns of recovery following experimental harvesting. The greatest and longest effects of seaweed harvesting were observed for the targeted species associated with the matrix A, *F. serratus* and *H. elongata*. These brown algae clearly created a dominant (in terms of cover and biomass)

and monospecific canopy on their site prior to the disturbance. While the canopy cover index was strongly affected up to 9 months after the disturbance event for both species, the cascading effects triggered by the canopy loss strongly diverged between the two species. Indeed, our study showed a significant decrease of the animal diversity and abundance associated with *F. serratus* up to 9 months after the harvesting, followed by a significant increase in algal diversity. On the other hand, the only cascading effect observed after the harvesting of *H. elongata* was a decrease in the animal diversity 3 months after the initial disturbance. The main explanation is that *F. serratus* forms a perennial canopy while *H. elongata* is a seasonal canopy-forming species. Therefore, while the removal of *F. serratus* canopy left a barren area, the *H. elongata* canopy loss on the impact sites was mediated by the natural and seasonal reduction in *H. elongata* canopy on the control sites. Recent monospecific studies on perennial brown algae have shown that harvesting affects the size structure of the population (Vega *et al.* 2014) and the abundance of invertebrates (Phillippi *et al.* 2014). Besides, Migné *et al.* (2015) concluded that *F. serratus* canopy was acting on the community not by reducing the abiotic stress, but by providing food, habitat, or both. We also observed significant short-term effects of the harvesting on the communities associated with species of the matrix C but the natural variability of this matrix was over and above the disturbance effects. The removal of targeted species within matrix B (*P. palmata* either epiphytic or epilithic, and *C. crispus*) did not clear the impact sites of its algal biomass, leaving patches of non-targeted species (e.g., *F. serratus*). It explains partly why we did not observe significant effects on either the community structure or the assemblage following the canopy removal. Moreover, recovery is assumed to be higher in the presence of herbivores (Aquilino and Stachowicz 2012), whose densities were maintained in impact plots with non-targeted canopy species present. Herbivores promoted succession by grazing on early successional fast-growing ephemeral species that would otherwise prevent establishment of perennial algae (Aquilino and Stachowicz 2012).

2. Important natural spatiotemporal variability in benthic assemblages

Another relevant finding of this study was the important natural spatiotemporal variability of communities associated with the canopy-forming species within the matrices B and C. For example, *Porphyra* spp. live on upper rocky shore where abiotic conditions are highly variable, creating a stressful environment (high light, long exposure to ultraviolet light, desiccation, extremes of temperature and salinity, and nutrient unavailability). Eulittoral species are expected to be better adapted to tolerate such environmental challenges than

sublittoral species (Lüning *et al.* 1990). Moreover, we observed that both *Porphyra* spp. were restricted to spatially patchy populations, which is thought to be the results of grazing patterns (Branch *et al.* 1990; Griffin *et al.* 1998). Grazers may act to clear young sporelings from areas shortly after their recruitment (Griffin *et al.* 1998). This hypothesis is supported by our observation that, after the sand burial on the *P. linearis* site had killed most grazers present, the initial patchy distribution of *P. linearis* was replaced by a *P. linearis* bloom with a cover of 100%. Branch *et al.* (1990) observed the same pattern, after freshwater floods had killed most grazers present on their study site. Therefore, at a broad scale, abiotic factors define distribution patterns on the upper rocky shore, and within these patterns, other processes may occur at smaller scale and modify the species distribution and abundance (Gimenez *et al.* 2005; Smale *et al.* 2010). As a result, any potential impact of *Porphyra* sp. harvesting is unclear.

Such complex heterogeneity may have obstructed the detection of canopy harvesting effects. Marine benthic assemblages have been found to be highly variable across different scales of time and space (Fraschetti *et al.* 2005), which raises concern about the ability to discriminate this natural variability. The scale and relative magnitude of variability has important implications for the design of the experiment because it can give rise to ambiguity about the effects of putative impacts or stressors (Underwood, 1994; Heino *et al.*, 2004). Although our data clearly cannot provide mechanistic explanations for the patterns observed, it is quite conceivable that most of the variation among assemblages could be explained at the scale of quadrats. Unfortunately, as small-scale variability in macrobenthic assemblage properties is often remarkable (Fraschetti *et al.* 2005), a few small-scale samples may not be enough to detect the overall effect of canopy harvesting. Such small-scale spatial and temporal variance should not be considered simply as a statistical nuisance (Coleman 2002). Indeed, quantifying the range of the natural heterogeneity of marine benthic assemblages may help to identify the physical and biological factors that are the most relevant to be explored first under an experimental approach. There is thus a need for monitoring methods that allow clear discrimination between natural spatiotemporal variability and human pressures.

3. Conclusions and implications for the management of seaweeds harvest

Canopy-forming macroalgae provide a wide range of ecosystem services, notably through their contribution to the primary production and carbon cycling (Kaldy and Dunton 2000; Gollty *et al.* 2008) and the formation of biogenic habitat that create the foundation for entire

communities, enhancing the species diversity and the abundance of many species (Wikstrom and Kautsy 2006; Smale 2010). Intuitively, these foundational species should be the focus of conservation efforts but our study also highlights that seaweed management plans need to move away from a single species focus and account for the whole ecological role of targeted commercial species.

In the light of our results, the recovery of the canopies and of their associated communities seems realistic within the current frame of management plans in France. The current methods used coupled to the French laws regulating seaweeds exploitation result generally in continuous vegetative growth of remaining individuals and greater recruitment of new individuals. Seaweed harvest conservation plans, however, are currently based on single-species management in order to maximize seaweed population biomass while the underlying effects of harvesting on the whole ecosystem are mostly ignored. In Chile, the management of *Lessonia nigrescens*, a brown alga, protects its associated biodiversity by focusing on the harvest methodology, which is based on ecological and biological recommendations, rather than on the harvest quantities (Vasquez *et al.* 2012). On the Atlantic coast of Canada, the management program of the perennial brown alga *Ascophyllum nodosum* focused on a precautionary approach to habitat use and natural bed sustainability, for almost two decades (Ugarte and Sharp 2011). There is also a need to complement the management plans with monitoring and research programs centered on the biology, habitat, and associated fauna of the harvested seaweeds.

There are environmental dynamics that occur and also need to be approached. These are storm frequency and severity, sand flooding events and continued warming of waters. This appears to be crucial given the large body of literature that suggests that the cumulative effects of local anthropogenic and other stressors (e.g. rising sea surface temperatures and increasing wave exposure) have negative effects on the growth and survival of canopy-forming algae (Halpern *et al.* 2007). Therefore, these environmental factors should be monitored on a regular basis and if necessary, the legislation should be amended accordingly.

EBM is an integrated approach to manage marine resources that considers the entire ecosystem, not simply individual components of the ecosystem (McLeod and Leslie 2009), seeking to sustain resources and promote the conservation of the biodiversity as a keystone to maintain overall ecosystem health, diversity, and resilience (Curtin and Prellezo 2010). The application of EBM, however, is still largely based on single-species assessments and

disregards the broader ecosystem context and impact. Here, our results strongly support the need to implement ecosystem-based management, assessing both the habitat conditions and ecological roles of targeted commercial species, in order to insure the sustainability of the resource. Understanding the mechanisms driving the patterns of spatio-temporal variability should also be an important step to devise sensible and efficient policies for conservation and management.

Acknowledgement

The authors thank all the people who helped carry out the fieldwork and anonymous reviewers for reviewing and greatly improving the manuscript. Our study was funded by the ALGMARBIO project, InitiativeBioBretagne (IBB), FranceAgriMer (National Institute of Agricultural and Marine Products), the Brittany Regional Council, the Côte d'Armor Departmental Council, the European Fisheries Fund (EFF), and benefited from the support of the French Government run by the National Research Agency and with regards to the investment expenditure programme IDEALG ANR-10-BTBR-04.

References

- Airoidi, L., Balata D., and Beck, M.W. (2008). The Gray Zone: relationships between habitat loss and marine diversity and their applications in conservation. *Journal of Experimental Marine Biology and Ecology* **366**, 8-15.
- Anderson, M. J., Gorley R. N., and Clarke, K. R. (2008). PERMANOVA+ for PRIMER: guide to software and statistical methods. PRIMER-E, Plymouth, UK.
- Aquilino, K.M., and Stachowicz, J.J (2012). Seaweed richness and herbivory increase rate of community recovery from disturbance. *Ecology* **93**, 879-890.
- Araujo, R., Isabel, S.P., Serrao, E.A., and Per, A. (2012). Recovery after trampling disturbance in a canopy-forming seaweed population. *Marine Biology* **159**, 697-707.
- Benedetti-Cecchi, L., Pannacciulli, F., Bulleri, F., Moschella, P.S., Airoidi, L., Relini, G., and Cinelli, F. (2001). Predicting the consequences of anthropogenic disturbance: large-scale effects of loss of canopy algae on rocky shores. *Marine Ecology Progress Series* **214**, 137-150.

409 Branch, G. M., Eekhout, S., and Bosman, A. L. (1990). Short-term effects of the 1988 Orange
 410 River floods on the intertidal rocky- shore communities of the open coast. *Transactions of the*
 411 *Royal Society of South Africa* **47**, 331-354.

412 Bruno, J.F., and Bertness, M.D. (2001). Habitat modification and facilitation in benthic
 413 marine communities. In: *Marine Community Ecology* (ed. Bertness, M.D.). Sinauer
 414 Associates Inc., Sunderland, MA, pp. 201–218.

415 Butchart, S. H. M., Walpole, M. , Collen, B. , van Strien, A., Scharlemann, J. P. W., Almond,
 416 R. E. A., et al. (2010). Global biodiversity: indicators of recent declines. *Science* **328**, 1164-
 417 1168.

418 Coleman, M.A. (2002). Small-scale spatial variability in intertidal and subtidal turfing algal
 419 assemblages and the temporal generality of these patterns. *Journal of Experimental Marine*
 420 *Biology and Ecology* **267**, 53-74.

421 Curtin, R., and Pallezo, R. (2010). Understanding ecosystem based management: a literature
 422 review. *Marine Policy* **34**, 821-830.

423 Foster, M.S., Nigg, E.W., Kiguchi, L.M., Hardin, D.D., and Pearse, J.S. (2003). Temporal
 424 variation and succession in an algal-dominated high intertidal assemblage. *Journal of*
 425 *Experimental Marine Biology and Ecology* **289**, 15-39.

426 Fraschetti, S., Terlizzi, A., and Benedetti-Cecchi, L. (2005). Patterns of distribution of marine
 427 assemblages from rocky shores: evidence of relevant scales of variation. *Marine Ecological*
 428 *Progress Series* **296**, 13–29.

429 Gimenez, L., Borthagaray, A.I., Rodríguez, M., Brazeiro, A., and Dimitriadis, C. (2005).
 430 Scale-dependent patterns of macrofaunal distribution in soft-sediment intertidal habitats along
 431 a large-scale estuarine gradient. *Helgoland Marine Research* **59**, 224-236.

432 Griffin, N. J., Bolton, J. J., and Anderson, R. J. (1998). Potential for harvest of *Porphyra*
 433 species in the south western Cape. Unpublished Sea Fisheries Research Institute Report, Sea
 434 Fisheries Research Institute, Roggebaai, 48 pp.

435 Gollety, C., Migne, A., and Davoult, D. (2008). Benthic metabolism on a sheltered rocky
 436 shore: role of the canopy in the carbon budget. *Journal of Phycology* **44**, 1146-1153.

437 Halpern, B.S., Selkoe, K.A., Micheli, F., and Kappel, C.V. (2007). Evaluating and ranking the
 438 vulnerability of global marine ecosystems to anthropogenic threats. *Conservation Biology* **21**,
 439 1301-1315.

440 Heino, J., Louhi, P., and Muotka, T. (2004). Identifying the scales of variability in stream
 441 macroinvertebrate abundance, functional composition and assemblage structure. *Freshwater*
 442 *Biology* **49**, 1230–1239.

443 Jenkins, S.R., Norton, T.A., and Hawkins, S.J. (2004). Long term effects of *Ascophyllum*
 444 *nodosum* canopy removal on mid shore community structure. *Journal of the Marine*
 445 *Biological Association of the United Kingdom* **84**, 327-329.

446 Kaldy, J.E., and Dunton, K.H. (2000). Above- and below-ground production, biomass and
 447 reproductive ecology of *Thalassia testudinum* (turtle grass) in a subtropical coastal lagoon.
 448 *Marine Ecology Progress Series* **193**, 271-283.

449 Lüning, K., Yarish, C., and Kirkman, H. (1990). Seaweeds. Their Environment,
 450 Biogeography and Ecophysiology. Wiley, New York.

451 Mayer-Pinto, M., Underwood, A.J., and Marzinelli, E.M (2015). The matrix influences direct
 452 and indirect effects of an anthropogenic disturbance on marine organisms. *Environmental*
 453 *Research* **136**, 15-20.

454 McLeod, K., and Leslie H. (2009). Ecosystem-Based Management for the Oceans.
 455 Washington, DC: Island. 392 pp.

456 Migné, A., Golléty, C., and Davoult, D. (2014). Effect of canopy removal on a rocky shore
 457 community metabolism and structure. *Marine Biology* **162**, 449-457.

458 Morrissey, D.J., Howitt, L., Underwood, A.J., and Stark, J.S. (1992). Spatial variation in soft
 459 sediment benthos. *Marine Ecology Progress Series* **81**, 197-204.

460 Perkol-Finkel, S., and Airoidi, L. (2010). Loss and recovery potential of marine habitats: an
 461 experimental study of factors maintaining resilience in subtidal algal forests at the Adriatic
 462 Sea. *PloS One* **5**, e10791.

463 Phillippi, A., Tran, K., and Perna, A. (2014). Does intertidal canopy removal of *Ascophyllum*
 464 *nodosum* alter the community structure beneath? *Journal of Experimental Marine Biology and*
 465 *Ecology* **461**, 53-60.

466 Sala, E., and Knowlton, N. (2006). Global marine biodiversity trends. *Annual Review of*
467 *Environmental Resources* **31**, 93-122.

468 Schiel, D.R., and Lilley, S.A. (2007). Gradients of disturbance to an algal canopy and the
469 modification of an intertidal community. *Marine Ecology Progress Series* **339**, 1-11.

470 Smale, D.A. (2010). Monitoring marine macroalgae: the influence of spatial scale on the
471 usefulness of biodiversity surrogates. *Diversity and Distribution* **16**, 985-995.

472 Stagnol, D., Renaud, M., and Davoult, D. (2013). Effects of commercial harvesting of
473 intertidal macroalgae on ecosystem biodiversity and functioning. *Estuarine, Coastal and Shelf*
474 *Science* **130**, 99-110.

475 Tait, L.W., and Schiel, D.R. (2011). Dynamics of productivity in naturally structured
476 macroalgal assemblages: importance of canopy structure on light-use efficiency. *Marine*
477 *Ecology Progress Series* **421**, 97-107.

478 Ugarte, R., and Sharp G. (2011). Management and production of the brown algae
479 *Ascophyllum nodosum* in the Canadian maritimes. *Journal of Applied Phycology* **24**, 409-416.

480 Underwood, A. J. (1994). On Beyond BACI: Sampling designs that might reliably detect
481 environmental disturbances. *Ecological Society of America* **4**, 3-15.

482 Vasquez, J. A., Piaget, N., and Vega, J. M. A. (2012). The *Lessonia nigrescens* fishery in
483 northern Chile: “how you harvest is more important than how much you harvest”. *Journal of*
484 *Applied Phycology* **24**, 417-426.

485 Vega, J. M. A., Broitman, B. R., and Vasquez, J.A. (2014). Monitoring the sustainability of
486 *Lessonia nigrescens* (Laminariales, Phaeophyceae) in northern Chile under strong harvest
487 pressure. *Journal of Applied Phycology* **26**, 791-801.

488 Wikstrom, S. L., and Kautsky, L. (2006). Structure and diversity of invertebrate communities
489 in the presence and absence of canopy-forming *Fucus vesiculosus* in the Baltic Sea.
490 *Estuarine, Coastal and Shelf Science* **72**, 168-176.

491

Fig. 1 Temporal variation in the mean ($\pm$ SE) canopy cover index (a), taxonomic algal diversity (b), taxonomic animal diversity (c), and animal abundance (d) in both control and impact plots (grey and black lines respectively) for each studied species and site. Significance codes: * $P < 0.05$.

Table 1. Summary of Permutational Multivariate Analysis of Variance of the MBACI model for canopy cover index, algal taxonomic diversity, animal taxonomic diversity and total animal abundance.

Matrix A									
Source	df	Canopy cover Index		Algal taxonomic diversity		Animal taxonomic diversity		Total animal abundance	
		MS	F	MS	F	MS	F	MS	F
BA	1	4680.30	20.547	721.94	0.727	45.27	0.099	1134.30	1.449
Csl	1	4231.50	3.727	63.66	0.008	506.88	0.293	586.63	0.109
T(BA)	2	469.50	0.752	1450.00	2.359	829.94	9.791**	7219.20	5.501*
L(Csl)	2	1135.40	2.700*	8099.70	40.166***	1731.40	5.363**	5387.80	7.470***
BAXCsl	1	3693.50	16.215*	535.81	0.539	2048.10	4.473	93.68	0.120
BAXL(Csl)	2	227.78	0.542	993.20	4.925**	457.85	1.418	782.99	1.086
CslxT(BA)	2	2661.40	4.261	216.47	0.352	1056.30	12.462**	1122.10	0.856
T(BA)xL(Csl)	4	624.66	1.486	614.65	3.048**	84.76	0.26	1310.30	1.817*
Res	64	420.45		201.66		322.84		721.26	

Matrix B									
Source	df	Canopy cover Index		Algal taxonomic diversity		Animal taxonomic diversity		Total animal abundance	
		MS	F	MS	F	MS	F	MS	F
BA	1	199.66	0.674	432.17	2.990	622.68	2.800	295.79	0.294
Csl	1	163.18	0.037	817.75	1.144	649.48	1.036	1483.20	0.163
T(BA)	2	968.97	1.515	280.50	1.818	1083.30	2.519	1189.80	1.274
L(Csl)	4	4411.10	10.306***	714.94	6.915***	626.79	2.614*	9119.50	10.822***
BAXCsl	1	546.01	1.842	716.23	4.955	696.06	3.130	837.60	0.833
BAXL(Csl)	4	296.40	0.692	144.54	1.398	222.36	0.927	1005.40	1.193
CslxT(BA)	2	677.49	1.059	163.10	1.057	381.88	0.888	482.05	0.516
T(BA)xL(Csl)	8	639.51	1.494	154.27	1.49	430.04	1.79	933.63	1.11
Res	96	428.02		103.39		239.82		842.70	

Matrix C									
Source	df	Canopy cover Index		Algal taxonomic diversity		Animal taxonomic diversity		Total animal abundance	
		MS	F	MS	F	MS	F	MS	F
BA	1	4362.60	2.070	99.53	0.721	18812.00	6.076	19274.00	4.648
Csl	1	1004.40	1.997	58.72	0.054	574.95	0.192	1192.60	0.339
T(BA)	2	14686.00	26.613**	1469.80	0.425	3145.90	0.537	4520.10	0.881
L(Csl)	2	502.85	1.261	1082.00	3.8761*	2991.60	1.301	3518.80	1.299
BAXCsl	1	284.36	0.135	250.41	1.814	617.99	0.200	791.83	0.191
BAXL(Csl)	2	2107.90	5.284***	138.01	0.494	3096.10	1.346	4146.80	1.531
CslxT(BA)	2	348.06	0.631	369.80	0.107	2863.40	0.489	3687.90	0.719
T(BA)xL(Csl)	4	551.83	1.383	3455.80	12.380***	5859.00	2.547***	5132.50	1.894
Res	64	398.93		279.14		2300.10		2709.30	

Significance codes: * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

493

494

495

Table 2 ANOSIM results: global R and significance level of differences between the assemblages of the control and the impact sites for each matrix at 3 months, 9 months and 12 months after the simulated harvest, based on presence/absence data matrices. Bold labels indicate a significant difference.

	Before harvest		3 months		9 months		12 months	
	R-value	Signif. Level	R-value	Signif. Level	R-value	Signif. Level	R-value	Signif. Level
Matrix A	-0.042	0.771	0.163	0.025	0.081	0.123	-0.041	0.644
Matrix B	0.005	0.393	-0.027	0.717	-0.014	0.570	-0.030	0.749
Matrix C	-0.041	0.651	0.112	0.039	-0.044	0.838	-0.040	0.822

Fig. 2 Plot of Principal components 1 and 2, showing variables vectors. Species coded as A1 for *F. serratus*, A2 for *H. elongata*, B1 for *P. palmata* (epiphytic), B2 for *P. palmata* (epilithic), B3 for *C. crispus*, C1 for *P. linearis*, and C2 for *P. umbilicalis*, for each zone (circle = control; square = impact) and each matrix (black filled = matrix A; grey filled = matrix B; white filled = matrix C). Labels code for species, treatment zone (C or I) and the number of months after the harvest. Bold labels have a significant contribution to the first or second axis (or both): underlined labels significantly contribute to the first axis while framed labels significantly contribute to the second axis.