

No evidence of pre-copulatory mate choice by gynes in the facultatively parthenogenetic ant *Cataglyphis cursor*

Florence Helft, C. Doums, T. Monnin

► **To cite this version:**

Florence Helft, C. Doums, T. Monnin. No evidence of pre-copulatory mate choice by gynes in the facultatively parthenogenetic ant *Cataglyphis cursor*. *Insectes Sociaux*, Springer Verlag, 2016, 63, pp.199-201. 10.1007/s00040-015-0444-0 . hal-01233413

HAL Id: hal-01233413

<https://hal.sorbonne-universite.fr/hal-01233413>

Submitted on 25 Nov 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **No evidence of pre-copulatory mate choice by gynes in the facultatively parthenogenetic**
2 **ant *Cataglyphis cursor***

3

4 Florence Helft^{1,2}, Claudie Doums², Thibaud Monnin¹

5

6 ¹ Sorbonne Universités, UPMC Univ Paris 06, UMR 7618 Institute of Ecology and
7 Environmental Sciences of Paris, 75005, Paris, France

8 ² Institut de Systématique, Évolution, Biodiversité, ISYEB, UMR 7205, CNRS, UPMC, École
9 Pratique des Hautes Études, MNHN Département Systématique et Évolution, 45 rue Buffon,
10 CP 39, 75005 Paris, France

11

12 Corresponding author: Florence Helft

13 Address: Sorbonne Universités, UPMC Univ Paris 06, UMR 7618 Institute of Ecology and
14 Environmental Sciences of Paris, 75005, Paris, France

15 E-mail: Florence.helft@gmail.com

16 Tel: +33 1 44 27 58 61

17

18 **ABSTRACT**

19 Virgin queens (gynes) of the ant *Cataglyphis cursor* mate on the ground at the nest entrance,
20 where they benefit from the protection of nestmate workers. By carrying out mating tests in
21 the laboratory, we show that gynes mate with the first males they encounter, suggesting the
22 absence of pre-copulatory mate choice by gynes. This is in agreement with the recent finding
23 that workers behave aggressively towards foreign males and could thereby exercise a form
24 of inclusive mate choice, which could replace pre-copulatory mate choice by gynes.

25 **KEYWORDS**

26 Eusocial insects, Sexual selection, Inclusive mate choice, Multiple mating.

27

28

29 Sexual selection favours phenotypic traits affecting reproductive success, for instance
30 through better access to a sexual partner resulting from increased competitiveness among
31 individuals of the same sex (intra-sexual selection) or from increased attractiveness towards
32 individuals of the opposite sex (inter-sexual selection). While sexual selection has received
33 considerable attention in solitary organisms it has been the subject of less attention in social
34 species, especially in social insects (Davidson 1982; Boomsma *et al.* 2005; Boomsma 2007;
35 den Boer *et al.* 2010; Boomsma 2013). This might be linked to the fact that in the latter
36 sexual partners are committed for life, which minimizes the potential for sexual selection
37 (Boomsma 2013). Furthermore, the widespread occurrence of nuptial flights associated with
38 a high mortality of sexuals favours the evolution of traits that optimize and access to
39 partners (e.g. flight abilities, partner detection) but not those linked with pre-copulatory
40 sexual selection (e.g. weaponry, ornaments, courtship behaviours) (Boomsma *et al.* 2005;
41 Baer 2011).

42 However, some eusocial insects mate multiply on the ground near the nest entrance.
43 This lowers the mortality of gynes (young virgin queens) since they are in the presence of
44 protective workers and can retreat into the nest. Moreover, workers may directly participate
45 in gyne mate choice, i.e. exercise a form of inclusive mate choice (Cronin *et al.* 2011).
46 Indeed, gynes and workers have the same interests in choosing the “best” males. This is
47 because workers typically do not reproduce and obtain indirect fitness by helping related
48 queens. Workers would therefore benefit from helping related gynes to choose good mates,

49 and could achieve this by physically preventing males of poor quality from accessing gynes.
50 Gynes would benefit from deferring mate choice to workers if doing so minimises their
51 exposure to desiccation and predation. In addition, workers may collectively be in a better
52 position to make an optimal choice than hard-pressed gynes.

53 The involvement of workers in mate choice has been suggested in army ants (Franks
54 & Hölldobler 1987), Argentine ants (in the invasive range, Sunamura *et al.* 2011) and the
55 facultatively parthenogenetic ant *Cataglyphis cursor* (Cronin *et al.* 2011; Helft *et al.* 2015). In
56 the latter two species, workers are aggressive towards all foreign males who enter or come
57 near the nest searching for a receptive gyne. Worker aggression may be a form of inclusive
58 sexual selection by workers if they differentially affect some male phenotypes. This is indeed
59 what occurs in *C. cursor*. Gynes are produced by the multiply mated queen (Lenoir *et al.*
60 1988; Percy *et al.* 2004) and workers are highly related to both parthenogenetically
61 produced gynes (clones of their mother, $r \sim 0.5$) and sexually produced gynes ($r = 0.44$ on
62 average) (Doums *et al.* 2013), and a recent study showed that worker aggression toward
63 males affected male relative mating potential, increasing it for lighter males and decreasing
64 it for heavier males (Helft *et al.* 2015). The present study therefore aims at determining
65 whether *C. cursor* gynes exercise pre-copulatory mate choice or not. We observed mating
66 behaviour of gynes in the presence of two males under laboratory conditions. The males
67 differed in the level of harassment they had experienced from workers, with one of the two
68 males previously exposed to worker aggression for 10 minutes while the other was not
69 exposed to workers.

70 We used 115 sexuals (46 males and 69 females) from 22 colonies which had been
71 produced in nature but which emerged in the laboratory, allowing us to be certain of their
72 virginity (see Supplementary material). We carried out 23 mating tests, each using one focal

73 gyne and two males. In addition, two other gynes were used during the exposure phase of
74 males to worker aggression immediately prior to the mating test (below). The focal gyne was
75 placed alone in a mating arena to which it could habituate. Simultaneously, one male was
76 placed in another box containing a second gyne in a mesh cage, so that he could perceive
77 her presence but could not mate (treatment without workers or W-). A second male was
78 placed in a third box containing the third gyne, also encaged, but with the addition of 50 of
79 the gyne's nestmate workers that could aggress the male as shown in Helft *et al.* (2015)
80 (male W+). After 10 minutes, the two males were moved to the mating arena containing the
81 focal gyne and matings were observed for 10 minutes. All individuals were used only once.
82 The three gynes used in each test were sisters and at least 19 days old, since previous
83 observations suggest that most gynes are sexually active at this age. The two males were
84 brothers and unrelated to the gynes. We selected males of the same age and size (see
85 Supplementary material), hence they differed only in their previous exposure to worker
86 harassment.

87 Mating attempts were unambiguous. The male mounted above the gyne with his
88 front and middle legs embracing her thorax and only his hind legs remaining on the ground,
89 made frenzied antennae contacts and attempted to copulate by rubbing the tip of his
90 abdomen around that of the gyne. A successful attempt ended by the intromission of the
91 male's genitalia into the female gaster for several seconds and was thus recorded as a
92 mating even though we do not know whether sperm transfer occurred. Using BioLogic
93 software (Dimitri Missoh, <http://swarmy.free.fr/wordpress/2009>) we recorded the latency
94 before the first mating attempt, the number of mating attempts, and the number and
95 duration of mating (i.e. duration of copulation). All statistical analyses were conducted using
96 R-Gui software (version 3.0.2, 2008).

97 The gyne refused to mate with either male in nine out of the 23 tests. This may be
98 because she was not receptive or stressed by the experimental setup, e.g. by the absence of
99 nestmate workers or colony odour. Twenty matings occurred in the remaining 14 tests. They
100 lasted 1 min 10 s \pm 30 s (mean \pm SD). *C. cursor* queens mate multiply (Lenoir *et al.* 1988,
101 Pearcy *et al.* 2004) and multiple mating by gynes was indeed observed during our 10
102 minutes tests. Nine gynes mated once, four mated twice (two with the same male and two
103 with both males) and one mated thrice (with both males). Similarly, males can mate multiply
104 in this species (Cronin *et al.* 2011) and multiple mating by males was also recorded, with 14
105 males mating once and three mating twice (Fig. 1).

106 Although all W+ males were aggressed by workers during the exposure phase that
107 immediately preceded the mating test (from 22 to 201 aggressions in 10 minutes, median =
108 94), this had no effect on the outcome of mating tests. W+ and W- males did not differ in
109 latency before the first mating attempt (Wilcoxon test: $p = 0.9$), number of mating attempts
110 (Wilcoxon test: $p = 0.7$), number of matings (Fisher exact test: $p = 0.54$) or duration of
111 matings (Wilcoxon test: $p = 0.7$) (Fig. 1).

112 Both males attempted mating in 11 out of the 14 tests where mating occurred. The
113 gyne always mated with the first male who attempted mating, usually on his second or third
114 attempt, and she always mated before the second male attempted to mate (Fig. 1). This
115 suggests that gynes mated indiscriminately, at least for their first mating, and supports the
116 hypothesis that gynes largely defer pre-copulatory mate choice to workers (Helft *et al.*
117 2015). Note, however, that this does not preclude post-copulatory sperm selection (Baer
118 2011). For instance, cryptic mate choice by gynes could occur by rejecting the sperm of less
119 fit partners and/or by controlling the duration of mating (Oppelt and Heinze 2007). In
120 several instances we observed that gynes in copula turned around and bit the mounting

121 male and that this ended the copulation. This suggests that gynes have at least some degree
122 of control over mating duration, and it is interesting in this context to note that mating
123 duration was indeed very variable, lasting from 10 s to 1 min 57 s (Fig. 1).

124 The indiscriminate mating by gynes observed in our experiment is in apparent
125 contradiction with previous field observations that gynes can refuse to mate and indeed
126 actively avoid most mating attempts (Lenoir *et al.* 1988, Cronin *et al.* 2011). However, in
127 these field observations gynes were likely already mated since most had lost their wings and
128 wing loss is a good predictor of being mated (Doums *et al.* 2013), hence they were likely not
129 seeking to mate but were seeking to be transported by workers to a new colony (Lenoir *et*
130 *al.* 1988, Chéron *et al.* 2011). Further studies should investigate whether all matings result in
131 the transfer of sperm and/or seminal secretions, whether mating duration affects this
132 transfer, and whether gynes can adjust mating duration according to male quality or mating
133 status.

134

135 **ACKNOWLEDGEMENTS**

136 We thank C. Haussy for helping with colony collection, R. Péronnet for helping in rearing the
137 ants, and A. L. Cronin for linguistic help.

138

139 **REFERENCES**

140 Baer B (2011) The copulation biology of ants (Hymenoptera: Formicidae). *Myrmecol News*
141 14:55-68

142 Boomsma JJ (2007) kin selection versus sexual selection: why the ends do not meet. *Curr*
143 *Biol* 17:673-683

144 Boomsma JJ (2013) Beyond promiscuity: mate-choice commitments in social breeding. *Philos*
145 *Trans R Soc B* 368:2-21

146 Boomsma JJ, Baer B, Heinze J (2005) The evolution of male traits in social insects. *Annu Rev*
147 *Entomol* 50:395-420

148 Chéron B, Cronin AL, Doums C, Fédérici P, Haussy C, Tirard C, Monnin T (2011) Unequal
149 resource allocation among colonies produced by fission in the ant *Cataglyphis cursor*.
150 *Ecology*. 92: 1448-1458

151 Cronin A, Monnin T, Haussy C, Doums C (2011) Opportunities for mate choice in the fission-
152 performing ant *Cataglyphis cursor*. *Ecol Entomol* 36:522-525

153 Davidson DW (1982) Sexual selection in harvester ants (Hymenoptera: Formicidae:
154 *Pogonomyrmex*). *Behav Ecol Sociobiol* 10:245-250

155 den Boer SPA, Baer B, Boomsma JJ (2010) Seminal fluid mediates ejaculate competition in
156 social insects. *Science* 327:1506-1509

157 Doums C, Cronin AL, Ruel C, Fédérici P, Haussy C, Tirard C, Monnin T (2013) Facultative use
158 of thelytokous parthenogenesis for queen production in the polyandrous ant
159 *Cataglyphis cursor*. *J Evol Biol* 26:1431-1444

160 Franks NR, Hölldobler B (1987) Sexual competition during colony reproduction in army ants.
161 *Bio J Linn Soc* 30:229–243

162 Helft F, Monnin T, Doums C (2015) First evidence of inclusive sexual selection in the ant
163 *Cataglyphis cursor*: worker aggressions differentially affect male access to virgin
164 queens. *Ethology* 121:1-10

165 Lenoir A, Querard L, Pondicq N, Berton F (1988) Reproduction and dispersal in the ant
166 *Cataglyphis cursor* (Hymenoptera, Formicidae). *Psyche* 95:21–44

167 Oppelt A, Heinze J (2007) Dynamics of sperm transfer in the ant *Leptothorax gredleri*.
168 Naturwissenschaften 94:781-786


169 Pearcy M, Aron S, Doums C, Keller L (2004) Conditional use of sex and parthenogenesis for
170 workers and queen production in ants. Science 306:25–28

171 Sunamura E, Hoshizaki S, Sakamoto H, Fujii T, Nishisue K, Suzuki S, Terayama Y, Tatsuki S
172 (2011) Workers select mates for queens: a possible mechanism of gene flow
173 restriction between supercolonies of the invasive Argentine ant.
174 Naturwissenschaften 98:361-368

175

176 Figure 1: Timing of mating attempts and duration of matings. The x-axis shows the duration
 177 of the mating tests in min. On the y-axis, labels indicate colony identity, as well as gyne
 178 identity for colonies which provided multiple gynes. The shorter bars indicate mating
 179 attempts. A few that were largely overlapping have been slightly moved sideways for clarity.
 180 The taller bars indicate actual mating (copulation), with their width indicating the duration
 181 (given in s). W- males are in white and W+ males are in black.

182
 183



184