



HAL
open science

Genesis of the Pieniny Klippen Belt in the Carpathians: Possible effects of a major paleotransform fault in the Neo-Tethyan domain

Jean Chorowicz

► **To cite this version:**

Jean Chorowicz. Genesis of the Pieniny Klippen Belt in the Carpathians: Possible effects of a major paleotransform fault in the Neo-Tethyan domain. *Comptes Rendus Géoscience*, 2015, 10.1016/j.crte.2015.10.003. hal-01239459v1

HAL Id: hal-01239459

<https://hal.sorbonne-universite.fr/hal-01239459v1>

Submitted on 7 Dec 2015 (v1), last revised 3 May 2016 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Genesis of the Pieniny Klippen Belt in the Carpathians: possible effects of a major paleotransform fault in the Neo-Tethyan domain

Jean Chorowicz

1- Sorbonne Universités, UPMC Univ Paris 06, CNRS Institut des Sciences de la Terre Paris (iSTeP), 4 place Jussieu 75005 Paris, France; jean.chorowicz@upmc.fr

Keywords: Alps, Carpathians, Dinarides, paleotransform, Pieniny Klippen Belt, paleogeography

ABSTRACT

The Pieniny Klippen Belt is a narrow, discontinuous zone rich in olistostromes and olistoliths (klippen) in the Western Carpathians. This paper based on prior works including tectonic and stratigraphic evidences suggests that the PKB rocks were deposited from the Triassic to the Early Paleogene along the eastern footwall side of a 'SKICZ' paleotransform fault zone. This transform was then separating the continental crust of the Austro-Alpine zone in the west and a Carpathian Embayment Ocean in the east. It was only during the Late Paleogene-Early Miocene that the PKB rocks were integrated into the accretionary prism that formed at the front of the eastward extruded ALCAPA units. This interpretation also supports the existence of a major paleotransform fault zone in the Neo-Tethys during the Triassic-Early Paleogene. This paleotransform had been previously suggested to explain the observed reversal in obduction and subduction at the junction between the Eastern-Southern Alps and the Carpathians-Dinarides.

1. Introduction

The Pieniny Klippen Belt (PKB) in the Western Carpathians (Fig. 1) is a particular, narrow, segmented isopic zone, rich in Mesozoic to Cenozoic olistostromes and olistoliths (klippen). It has been the focus of recent studies but its paleogeographic origin is still unclear (e.g., Golonka et al., 2015; Jankowski et al., 2012; Plašienka, 2012; Ślaczka et al., 2012). This contributes to make difficult the paleogeographic reconstruction of the whole Alpine Neo-Tethyan domain (e.g., Aubouin et al., 1970; Săndulescu 1988; Dercourt et al., 2000; Golonka 2004; Schmid et al., 2008; Stampfli and Hochard, 2009).

On the one hand, Chorowicz (1977) has suggested that a Split-Karlovac-Wien paleotransform fault zone might have existed in the Neo-Tethys between Split and Vienna (Wien) and was forming the western end of the Carpathians (Fig. 1). On the other hand, Ratschbacher et al. (1991a, b) and Wölfler et al. (2011) have studied the Oligocene-Miocene lateral extrusion of the Eastern Alps inner zones (Austro-Alpine) towards the Western Carpathians. They suggested that the Austro-Alpine units prior to their extrusion were forming a compressed wedge which eastern boundary was unconstrained.

The present paper explores the relations between the paleotransform fault zone and the genesis and paleogeographic location of the Pieniny Klippen Belt. The first section briefly reports the particularities of the Pieniny Klippen Belt. The second section discusses the process of lateral extrusion of the inner Eastern Alps (Austro-Alpine) and discriminates the SK and CZ segments of the paleotransform fault zone (Fig. 1). A discussion section then examines the links between the PKB facies and the paleogeographic setting of the paleotransform fault. It is proposed that a segment of the paleotransform is the initial location of the PKB.

2. Pieniny Klippen Belt

The peri-Eurasia Alpine branch in Central Europe comprises the arched Western and the Eastern Alps that are in line with the Carpathians and the Balkans (Fig. 1). Further south, the peri-Adria branch comprises the Apennines, the Southern Alps and the Dinarides-Hellenides. The Carpathians form around the Pannonian basin an arch that is interpreted to result from post-orogenic extension in the backarc regions of southward subduction zones (Jolivet et al., 2008). The subduction would have been that of a 'Carpathian Embayment Ocean'.

The Pieniny Klippen Belt (PKB, Fig. 1) in the Northwestern Carpathians presently is a subvertical, narrow - up to 20km wide -, 600km long, strike-slip fault zone which separates the inner and the outer nappes (e.g., Golonka et al., 2015). The PKB emerges in the west from beneath the Neogene sediments of the Vienna basin and ends to the east against the eastern alongstrike continuation of the Balaton line. The 'black flysch', which crops out along the eastern continuation of the Balaton line, may belong to the PKB (e.g., Schmid et al, 2008). The Inner Northwest Carpathians, in between the Balaton line and the PKB, were compressed during the Late Paleogene-Neogene together with the Inner Southeast Carpathians composed of the Tisza and Dacia units (Fig.1). This compression pushed the Outer Carpathian flysch nappes onto the Miocene deposits of the Carpathian foredeep that covers the Eurasia platform (e.g., Csontos and Vörös, 2004; Săndulescu, 1975).



Figure 1. Geostructural frame of the Alpine belts in the Central Europe-Mediterranean region, showing present day location of the SKICZ (Split-Karlovac-Initial PKB-Crustal-Zone) segments of the paleotransform, drawn in blue colour: SK, Split-Karlovac; PKB: Pieniny Klippen Belt; CZ: Crustal Zone. Bn: Bosnian zone; Dlm: Dalmatian zone; Hk: High Karst zone; Sr: Serbian zone. 1: front of the Alpine belts; 2: Pieniny Klippen Belt (PKB); 3: SK and CZ end segments of the SKICZ paleotransform zone (dashed line in depth, full line Krakow-Sandomierz fault at surface); 4: vergence of obducted ophiolites; 5: the Split-Karlovac-Wien (SKW) line suggested in 1977 by Chorowicz.

The lithostratigraphic units of the PKB have been divided into several successions (e.g., Golonka and Krobicki, 2004). The matrix is shale, marls and flysch, spanning from the Triassic to the Neogene. The most typical feature of the PKB units is that they include olistostromes and olistolites (klippen), some of kilometre size. Deformation began during the Late Cretaceous (Nemčok and Nemčok, 1994) but mainly occurred from Eocene to Neogene times (Ratschbacher et al., 1993).

3. Lateral extrusion of Austro-Alpine units

The nappes of the Inner Northwest Carpathians are similar to those of the Eastern Alps inner (Austro-Alpine) zones (Săndulescu, 1988) and they together presently form the **Alps-Carpathians-Pannonia** - or 'ALCAPA' - system (Csontos and Vörös, 2004; Fig.1). The ALCAPA system is interpreted to result from the Oligocene to Middle Miocene lateral extrusion of fragments of Austro-Alpine units away from the zone of convergence between Adria and Eurasia continents. This event is generally considered to be a compressional eastward tectonic escape (Wölfler et al., 2011) but it might have combined extensional collapse driven by gravity (Ratschbacher et al., 1991b). The latter authors have illustrated in map view the pre-extrusion situation during the Early Paleogene time and suggested that the Austro-Alpine continental block was separated from both the Eurasia and the Adria continents by EW-striking major contacts whereas their eastern boundary was an unconstrained margin. The 'pre-Gosau' compression phase in the Albian to Late Cretaceous (Schuller, 2004) thickened the Austro-Alpine crust that became rheologically weak 30-50 Ma later (Oligocene). The weakened unconstrained margin collapsed toward the low domain (Ratschbacher et al., 1991b).

The present Carpathian foreland was originally a passive margin bordering an oceanic lithosphere that formed the 'Carpathian Embayment - or Magura - Ocean', belonging to the Eurasian plate (e.g., Royden et al, 1982; Csontos and Vörös 2004; Wölfler et al., 2011). SW-ward slab subduction roll-back of the Carpathian Embayment oceanic lithosphere would be responsible for the current arched shape of the Carpathians (e.g., Nemčok, 1993). Subduction was active at least during the Middle Miocene, as evidenced by back arc volcanism (Szabó et al., 1992). Schmid et al. (2008) compiled evidences for the Carpathian Embayment Ocean to be in place since at least the Jurassic.

Cretaceous collisions occurred on either side of the Carpathian Ocean Embayment. In the east, the Dacia and Tisza blocks were afterward both pushed into the Carpathian Embayment during the Cenozoic (Fig. 1; Csontos and Vörös 2004). ALCAPA is presently separated from Tisza-Dacia by the Balaton line and its mid-Hungarian eastern continuation that are reported to have accommodated differential horizontal movements (e.g. Márton et al., 2007). During the Middle Oligocene-Neogene time, the ALCAPA and Dacia-Tisza units were docked against the Eurasia continent foreland and this induced progressive folding and thrusting of the Outer Carpathians flysch units.

4. Split-Karlovac and Vienna segments

The Split-Karlovac zone in the Dinarides (SK, Fig. 1) is a narrow NNW-trending faulted area across which the otherwise regular arrangement of the NW-trending folds and thrusts is modified (Chorowicz, 1975). The SK faulted region comprises isopic zones that are shown in Fig. 2 and characterized in Table 1. Some units are offset and/or differ in their lithostratigraphic successions and styles across the SK fault zone. While the outermost Dalmatian, Inner Dalmatian and Outer Karst zones are unmodified across the SK fault zone, except for a sigmoid change in their strike, the High Karst carbonate platform is intersected by faults (Figs. 1 and 2) since the Permian (Chorowicz, 1977). The High Karst zone forms a nappe sheet that rests on the Outer Karst of the Gračac window. The Inner Karst zone extends across the High Karst and includes Late Jurassic pelagic deposits (*Lemeš* levels). The Prekarst zone is transitional toward the NE with the pelagic inner Bosnian zone (Blanchet et al., 1969), which is dextrally offset by about 100km across the SK fault zone. The Outer and Bosnian units of the Dinarides extend up NW-ward into similar zones in the Southern Alps (Cousin, 1981). Overall, the lateral offsets across the SK fault zone increases toward the inner zones whereas they die out southward in the Adria micro-continent.




Figure 2. Structural map of the Dinarides between Split and Karlovac according to Chorowicz (1977), completed from Schmid et al. (2008) and Lužar-Oberiter et al. (2012). The isopic zones are shortly characterized in Table 1.

	Zones	Facies	Initiation of Flysch	Paleogeography
Inner zones	<i>Sava</i>	continental crust + Late Cretaceous ophiolites and back-arc ocean	Late Cretaceous-Paleogene	Suture fragments of the Adria distal margin
	<i>Serbian (Western Vardar)</i>	ophiolites, mélanges	Senonian	obducted Jurassic back-arc marginal basin
	<i>Bosnian</i>	dolomitic Mesozoic	1- Late Jurassic; 2- Late Cretaceous	basin of the distal margin of Adria
	<i>Prekarst</i>	slope and deep-marine Mesozoic carbonates, stratigraphic gaps	Late Cretaceous-Paleogene	Mesozoic transitional platform-slope
Outer zones	<i>Inner Karst</i>	similar to the High Karst but pelagic Late Jurassic <i>Lemeš</i> levels		High Karst areas affected by SK paleoactivity
	<i>High Karst</i>	marine Late Palaeozoic, marine Mesozoic platform carbonates	Early Eocene	carbonate platform
	<i>Outer Karst</i>	volcano-sedimentary Ladinian; marine Mesozoic platform and slope carbonates.	Late Lutetian-Early Priabonian	transitional platform slope
	<i>Budva</i>	hidden by High Karst nappes but revealed by tectonic 'slice' of pelagic Paleocene limestone		deep-water basin
	<i>Inner Dalmatian</i>	marine carbonate rich in pelagic facies and turbidites	Early Priabonian	transitional platform slope
	<i>Dalmatien</i>	marine Mesozoic platform carbonate, bauxite, Lutetian transgression	Late Priabonian	carbonate platform

Table 1. Main characteristics of the isopic zones in the Split-Karlovac (SK) fault zone in the Croatia-Bosna border region of Fig. 2.

The NW-striking ophiolitic nappes that extend along the Dinarides and the Hellenides form two parallel groups related to the Vardar and the Pindos domains (Smith and Spray, 1984). They both verge to the SW (Fig. 1). The vergence is conversely to the north for the ophiolitic Penninic nappes that crop out in major windows in the Eastern Alps of Austria, thus opposite to that of the Dinarides-Hellenides. This reversal in thrust vergence (Laubscher, 1971) is related to opposing subduction polarities as shown in tomographic images (Lippitsch et al., 2003). These observations suggest that a transform fault zone might have linked the Eastern Alps and the Dinarides. Location of the transform was about that of the northward continuation of the Split-Karlovac line. The SK line was likely a segment of the inferred paleo-transform for it was accomodating vertical motions since at least the Permian, and since the Late Eocene it was compressed during the Alpine shortening phases (Chorowicz, 1977). It can be interpreted that the SK fault zone has formed the southern tip segment of the paleo-transform, as such accomodating vertical motions before collision phases.

Further north, the virgation of the Alpine-Carpathian front near Vienna seems to be guided by a NE-striking fault zone (Beranek and Dudek, 1972) that forms the hidden eastern boundary of the Bohemian massif. This fault zone seems to have initiated in the Permian (Suk, 1984). Zatopek (1979) and Malkovsky (1987) have evidenced a vertical step in the Moho discontinuity across the NE fault zone, which they call 'crustal zone' (CZ, Fig. 1). Their map of the Moho and tomographic images in horizontal section at 50 km in depth (Ustaszewski et al., 2008) obtained by the method of Bijwaard and Spakman (2000) suggest that the CZ fault zone extends at depth under the Carpathian nappes, then crops out further northeast in the Carpathian foreland where it forms the so-called Krakow-Sandomierz fault (Fig. 1; Púchy and Varga, 1973; Chorowicz et al., 1999). The overall CZ fault zone dies out NE-ward and may be the former prolongation of the paleotransform fault zone. It might therefore represent the continuation of that transform fault within the Eurasia continent. Similar to the SK segment further south in the Adria continent, the CZ fault zone would have been the northern tip of the paleo-transform and as such would have originally accomodated mainly vertical motions.

5. Discussion

Time and space distribution of PKB facies

The PKB olistostrome and olistolith deposits were formed either because an orogenic front was advancing towards the foreland (e.g., Golonka et al., 2015) or because an accretionary prism was forming above a subduction (e.g., Birkenmajer, 1986). They might also have formed during rifting and be subsequently integrated into a subduction-related accretionary prism (e.g., Cieszkowski et al., 2009).

Analysis of the distribution of the PKB facies through time and space allows distinguishing: a) clast samples, representative of the history of the uplifted areas; b) matrix and flysch deposits, characterizing lower sedimentary basins. To examine whether the PKB facies are related to the time evolution of the transform fault zone, they must be analyzed over different time periods, mainly the Triassic-Jurassic-Early Cretaceous, then the Late Cretaceous-Early Paleogene, and finally the Late Paleogene-Early Neogene. In doing so, the PKB deposits should be moved back to their original inferred position along a ~NS-trending transform fault.

1- From the Triassic to the Early Cretaceous times, the diversified olistostromes and olistoliths are consistent with the PKB being at eastern foot of transform fault scarps. Clasts are found for instance in the 'black flysch' of Early Jurassic age (Birkenmajer et al., 2008). Conglomerate yields small clasts of metamorphic, magmatic and sedimentary rocks of the Eurasian continent, which basement was affected by the Variscan orogeny and was partly covered by a Mesozoic carbonate platform. By contrast, west of the transform fault, stratigraphy indicates that a shallow marine pelagic environment first developed during the Middle Triassic to Early Dogger (Mišík, 1993); then a deeper basin developed during the Middle to Late Jurassic-Early Cretaceous. During the Late Jurassic-Early Cretaceous, Penninic obductions and the 'pre-Gosau' compression in the Austro-Alpine domain altogether created new reliefs west of the transform fault. In contrast, east of the transform fault the Carpathian Embayment Ocean remained a deep basin.

2- During the Late Cretaceous-Early Paleogene, the eastern boundary of the newly formed Austro-Alpine and West-Carpathian belt remained an unconstrained margin (Ratschbacher et al., 1991b). Clasts in the PKB reveal that Austro-Alpine facies were uplifted west of the transform fault and finally a carbonate platform was created during the Maastrichtian-Paleocene. In Poland and Slovakia, facies of the Late Cretaceous-Early Paleogene flysch of the PKB and the Magura nappe (e.g., Cieszkowski et al., 2009) show that the PKB was lying along the deep Carpathian Embayment (Golonka and Krobicki, 2004). In Western Ukraine (Ślączka et al., 2006), the Late Cretaceous-Early Paleogene matrix shows facies attesting for a deep marine origin facies east of the transform, whereas clasts reveal: - lower Jurassic (Hettangian?) clastics; - Middle Jurassic crinoidal limestone; - Bajocian to Tithonian red nodular limestones of Ammonitico Rosso-type with thin radiolarite intercalations; - Berriasian Calpionella-bearing limestones; - Berriasian basaltic pillow lavas. The latter magmatism is of mid-ocean-ridge basalt (MORB) type which suggests the existence of an oceanic crust (Golonka and Bocharova, 2000). There are also evidences of subduction-related calc-alkaline volcanics in the area during the Berriasian (Krobicki et al., 2003). Occurrence of Jurassic ophiolites west of the transform were also suggested from heavy mineral analyses (e.g., Aubrecht et al., 2009).

3- During the Late Paleogene-Early Neogene time, olistostromes are more likely due to brittle faulting on faults that destructed the pre-existing fold-and-thrust structures (e.g., Ratschbacher et al., 1993). The PKB was integrated into an accretionary prism at the front of the eastward-extruded Austro-Alpine and West-Carpathian units which now form the ALCAPA.

Paleogeographic location of the paleotransform zone segments

The paleogeography of the transform zone can be best understood at the Oxfordian (155 Ma) because that period predated the first Late Jurassic-Early Cretaceous obductions in the concerned area (Fig. 3). A paleogeographic reconstruction must however be regarded as purely speculative (e.g., Schmid et al., 2008). The proposal is that the SK segment in Adria and the CZ segment in Eurasia were both trending ~NNE-SSW relative to Eurasia during the Oxfordian (Fig. 3). The two segments were almost aligned. Moving the PKB back to its original position 'I' ('I' for Initial position) between the SK and the CZ segments makes 'I' becoming almost N-S. The SK, the original 'I' and the CZ segments were thus altogether forming a single N-S fault zone, dubbed here the SKICZ transform fault zone. At the Oxfordian time, the original PKB ('I') transform segment was possibly linking several oceans (to the west, the Penninic Ocean and the inferred Insubric Ocean; to the east, the Carpathian Embayment and the Vardar oceans) and was likely

accommodating a transcurrent motion between accretion or/and subduction zones. This ‘I’ transcurrent fault zone segment possibly also included during the Oxfordian elongate fault compartments along the boundary of the Austro-Alpine/Western Carpathians continental fragments and the Carpathian Embayment Ocean (Fig. 3), for example the Czorsztyn Submarine Ridge (Mišík, 1993). The two SK and CZ segments were forming the tips of the transform fault zone and as such they were sustaining vertical movements. The SKICZ paleotransform fault zone was active from the Permian to at least the Late Jurassic–Early Cretaceous. During that entire period, it kept a ~N-S overall trend. It afterward was partly the eastern unconstrained boundary of the Austro-Alpine belt during the Late Cretaceous–Early Paleogene. It is only later that it was divided and rotated up to its present attitude.




Figure 3. Location of the SKICZ paleotransform during the Oxfordian (155 Ma). The purpose of this map is purely paleogeographic; it does not explicitly concern the plate geodynamics, and the map does not show the zones of accretion, subduction, obduction, back-arc basins and other transforms but the SKICZ. Drawings are superimposed over extracts of Figures from Stampfli et al. (2001), Stampfli and Hochard (2009) and Bonev and Stampfli (2011) that provide the main geometry of the European, African and Adria continents and the paleolocation of mega-units. Dark green: oceanic lithosphere; white surfaces: continental lithosphere; blue lines, the SKICZ paleotransform comprising - dashed line labelled as ‘I’ in the text, transcurrent movements, - full lines, SK and CZ, with vertical movements. The Insubric line is sometimes considered to represent a former major suture between the Eurasia and Adria continents (e.g., Dewey and

Bird, 1970). AA: Austro-Alpine; Ana: Anatolia; Bri: Briançonnais; CzR: Czorsztyn ridge; Dac: Dacia; Dal: Dalmatia; Hel: helvetic; Kar: High Karst; Li: Liguria; Lyc: Lycia; Men: Menderes; Moe: Moesia; Pel: Pelagonian; Pen: Pennic; Pie: Piemont; Tau: Taurus; Tis: Tisza; WCa: West-Carpathians.

Presently, the SK and CZ segments almost meet northwest of Zagreb and strike at right angle to each other (Fig. 1). Paleomagnetic data (e.g., Muttoni et al., 2001) show that Adria and Africa have rotated from the Permian to the Late Cretaceous counterclockwise almost 90° relative to Eurasia. Since then, the Africa/Eurasia convergence has been ~N-S (Platt et al., 1989). These rotations and northward convergence have modified the trends of the two SK and CZ segments, up to their present attitude.

Some authors have already suggested the existence of paleotransform fault zones in the Neo-Tethys (e.g., Aubouin et Dercourt, 1975; Chorowicz et Geysant, 1976; Chorowicz et al., 1980; Debelmas and Săndulescu, 1987; Lyberis et al., 1982; Royden et al., 1982; Smith and Spray, 1984; Trümpy, 1988; Robertson and Shallo, 2000). In particular, Stampfli et al. (2001) have drawn a transform zone quite similar to the SKICZ between the Vardar Ocean opening in the north and the Paleo-Tethys subduction in the south, active during the Late Permian-Anisian. The transform would have then extend at the western border of the Vardar Ocean during the Carnian, Sinemurian and Oxfordian times. Csontos and Vörös (2004) have suggested a transform line between the Pindos and the Vardar oceans in the Late Permian, Carnian, Sinemurian and Oxfordian times. Debelmas and Săndulescu (1987) have associated the PKB with a Penninic transform fault zone that would have been EW-trending with respect to Eurasia and active in the Early Cretaceous. Ustaszewski et al. (2008) proposed that the transform needed to explain the observed reversal in obduction and subduction polarities between the Eastern Alps and the Dinarides, is the Balaton line and its eastern continuation. This inferred transform is thus fairly close to the SKICZ line suggested in the present study. Although both interpretations are reasonable, the SKICZ better reconstructs the original position of the PKB.

6. Conclusions

The Pieniny Klippen Belt (PKB) in the Carpathians was originally a segment of a large transform fault in the Neo-Tethys, the SKICZ (Split-Karlovac-Initial PKB-Crustal-Zone) paleotransform. The transform mechanism was active from the Permian to the Early Paleogene. During this time interval, clasts depositing in the PKB were originated from the west, from zones of uplifts west of the SKICZ fault 'I' segment, essentially from the Austro-Alpine continental lithosphere. It is only afterward, during the Late Paleogene-Early Neogene that these clasts were included into the accretionary prism that formed along the free border of the Austro-Alpine units as the latter were laterally extruding eastward.

The paleotransform acted as a transcurrent transform fault zone linking opposite subduction and obduction polarities (Alpine polarity vs. Dinaridic polarity). In the Late Jurassic, the paleotransform formed the western boundary of the Carpathian Embayment and of the Vardar oceans, against the free border of the future ALCAPA units. The SK and CZ fault zones in Adria and Eurasia continents respectively were the fault sections developed at the tips of the paleotransform. As such, they mainly accommodated vertical motions. The SKICZ paleotransform was a major tectonic structure in the Western Neo-Tethys whose existence and functioning are consistent with several of the models and paleogeographic maps that have been proposed to explain the formation of the Alpine belts.

Acknowledgements

Jean-François Stephan, 40 years friend of mine, knew about such project of paper. Together with Marc Tardy, he encouraged me to achieve this work and to submit it for publication. I am indebted to both of them for helpful discussions. I am also thankful to the Pierre et Marie Curie Paris 6 University for providing a position of associate (émérite) professor and for giving me access to all facilities necessary for research activity. This paper has been reviewed by Bernard Mercier de Lépinay, invited Associate Editor, and by Isabelle Manighetti, Associate Editor, who largely helped to improve the wording and clarity of the paper.

Bibliographical reference list

Aubouin, J., Dercourt, J., 1975. Les transversales dinariques dérivent-elles de paléofailles transformantes? C.R. Acad. Sci. (D) 218, 347-350.

- Aubouin, J., Blanchet, R., Cadet, J.-P., Celet, P., Charvet, J., Chorowicz, J., Cousin, M., Rampnoux, J.-P., 1970. Essai sur la géologie des Dinarides. *Bull. Soc. Géol. Fr.* 12, 1060–1095.
- Aubrecht, R., Méres, Š., Sýkora, M., Mikuš, T., 2009. Provenance of the detrital garnets and spinels from the Albian sediments of the Czorsztyn Unit (Pieniny Klippen Belt, Western Carpathians, Slovakia). *Geologica Carpathica* 60(6), 463–483.
- Beranek, B., Dudek, A., 1972. The contribution of deep seismic sounding to the studies of deep fault tectonics. *Int. Geol. Congr.* 24th 3, Tectonics, 16-24.
- Bijwaard, H., Spakman, W., 2000. Non-linear global P-wave tomography by iterated linearized inversion. *Geophysical Journal International* 141, 71-82.
- Birkenmajer, K., 1986. Stages of structural evolution of the Pieniny Klippen Belt, Carpathians. *Stud. geol. pol.* 88, 7-32.
- Birkenmajer, K., Gedl, P., Myczyński, R., Tiscza, J., 2008. “Cretaceous black flysch” in the Pieniny Klippen Belt, West Carpathians: a case of geological misinterpretation. *Cretaceous Research* 29(3), 535-549.
- Blanchet, R., Cadet, J.-P., Charvet, J., Rampnoux, J.-P., 1969. Sur l’existence d’un important domaine de flysch Tithonique–Cretacé inférieur en Yougoslavie. L’unité de flysch Bosniaque. *Bull. Soc. Géol. Fr.*, ser. 7, 11, 871–880.
- Bonev, N., Stampfli, G., 2011. Alpine tectonic evolution of a Jurassic subduction-accretionary complex: Deformation, kinematics and ⁴⁰Ar/³⁹Ar age constraints on the Mesozoic low-grade schists of the Circum-Rhodope Belt in the eastern Rhodope-Thrace region, Bulgaria-Greece. *Journal of Geodynamics*, 52, pp. 143-167.
- Chorowicz J., 1975. Le mécanisme de la structure transversale Split-Karlovac, dans les Dinarides yougoslaves. *C.R. Acad. Sc. (D)* 280, 2313-2316.
- Chorowicz, J., 1977. Etude géologique des Dinarides le long de la structure transversale Split-Karlovac (Yougoslavie). *Société Géologique du Nord, Publication* 1, 331 p.
- Chorowicz, J., Geysant, J., 1976. La paléofaille transformante Split-Karlovac-Vienne. Introduction à un modèle d’évolution des chaînes alpines de l’Europe moyenne. *Rev. de Géog. Phys. et de Géol. Dyn.* XVI(1), 127-141.
- Chorowicz, J., Cadet, J.P., Stephan, J.F., 1980. Le secteur transversal de Scutari-Peć: apport de l’étude de la fracturation à partir des données Landsat. *Bull. Soc. géol. France*, ser. 7 21(3), 217-228.
- Chorowicz, J., Mering, C., Raymond, D., Szczuski, R., François, R. 1999. Lithofacies change revealed by SAR ERS imagery in the Przemysl sigmoid, Carpathians: implication of a NE-striking paleofault on the North Tethyan passive margin. *Tectonophysics*, 304, 187-200.
- Cieszkowski, M., Golonka, J., Krobicki, M., Ślaczka, A., Oszczytko, N., Waśkowska, A., Wendorff, M., 2009. The Northern Carpathians plate tectonic evolutionary stages and origin of olistoliths and olistostromes. *Geodinamica Acta* 22(1-3), 101-126.
- Cousin, M., 1981. Les rapports Alpes-Dinarides. Les confins de l’Italie et de la Yougoslavie. *Société Géologique du Nord, Publication* 5(1), 521 p.
- Csontos L., Vörös, A., 2004. Mesozoic plate tectonic reconstruction of the Carpathian region. *Paleogeography Paleoclimatology Paleoecology* 210, 1–56.
- Debelmas, J., Săndulescu, M., 1987. Transformante nord-pennique et problèmes de corrélation palinspastique entre les Alpes et les Carpathes. *Bull. Soc. géol. France* 8, 3(2), 403-405.
- Dercourt, J., Gaetani, M., Vrielynck, B., Barrier, E., Biju-Duval, B., Brunet, M.F., Cadet, J.P., Crasquin, S., Săndulescu, M. (Eds.), 2000. Atlas Peri-Tethys, Palaeogeographical maps. CCGM/CGMW, 24 maps and explanatory notes I-XX, pp. 1–269.
- Dewey, J.F., Bird, J.M., 1970. Mountain belts and the new global tectonics. *Journ. Geoph. Res.*, 75, 2625-2647.
- Golonka, J., 2004. Plate tectonic evolution of the southern margin of Eurasia in the Mesozoic and Cenozoic. *Tectonophysics* 381, 235–273.
- Golonka, J., Bocharova, N.Y., 2000. Hot spots activity and the break-up of Pangea. *Paleogeography, Paleoclimatology, Paleoecology* 161, 49–69.
- Golonka, J., Krobicki, M., 2004. Jurassic paleogeography of the Pieniny and Outer Carpathian basins. *Rivista Italiana di Paleontologia e Stratigrafia* 110(1), 5–14.

- Golonka, J., Krobicki, M., Waskowska, A., Cieszkowski, M., Ślącza, A., 2015. Olistostromes of the Pieniny Klippen Belt, Northern Carpathians. *Geol. Mag.* 152(2), 269–286.
- Jankowski, L., Kopctowski, R., Rylko, W., Danysh, V., Tsarnenko, P.T., Hnylko, O., 2012. Lithostratigraphic correlation of the outer Carpathian borderlands of Poland, Ukraine, Slovakia and Romania. *Biuletyn Państwowego Instytutu Geologicznego* 449, 87-98.
- Jolivet, L., Augier, R., Faccenna, C., Negro, F., Rimmeles, G., Agard, P., Robin, C., Rossetti, F., Crespo-Blanc, A., 2008. Subduction, convergence and the mode of backarc extension in the Mediterranean region. *Bull. Soc. géol. Fr.*, 179(6), 525-550.
- Krobicki, M., Golonka, J., Aubrecht, R., 2003. Pieniny Klippen Belt: general geology and geodynamic evolution. In: Golonka, J., Lewandowski, M. (Eds.), *Geology, geophysics, geothermics and deep structure of the West Carpathians and their basement*. Publications of the Institute of Geophysics, Polish Academy of Sciences (Warszawa), Monographie M-28, 363, pp. 25-33.
- Laubscher, H., 1971. Das Alpen-Dinariden-Problem und die Palinspastik der südlichen Tethys. *Geologische Rundschau* 60, 813–833.
- Lippitsch, R., Kissling, E., Ansorge, J., 2003. Upper mantle structure beneath the Alpine orogen from high resolution teleseismic tomography. *J. Geophys. Res.*, 108(B8), 2376, doi:10.1029/2002JB002016.
- Lužar-Oberiter, B., Mikes, T., Dunkl, I., Babić, L., von Eynatten, H., 2012. Provenance of Cretaceous synorogenic sediments from the NW Dinarides (Croatia). *Swiss J. Geosci.*, 105, pp. 377-399.
- Lyberis, N., Chorowicz, J., Papamarinopoulos, S., 1982. La transversale du Kastanotikos (Grèce): télédétection, données de terrain et géophysiques. *Bull. Soc. géol. France* (7), 24(1), 73-85.
- Malkovsky, M., 1987. The Mesozoic and Tertiary basins of the Bohemian Massif and their evolution. *Tectonophysics* 137, 31–42.
- Márton, E., Tischler, M., Csontos, L., Fügenschuh, B., Schmid, S., 2007. The contact zone between the ALCAPA and Tisza-Dacia mega-tectonic units of Northern Romania in the light of new paleomagnetic data. *Swiss Journal of Geosciences* 100, 109–124.
- Mišík, M., 1993. The Czorsztyn submarine ridge (Jurassic-Lower Cretaceous, Pieniny Klippen Belt): an exemple of a pelagic swell. *Mitt. Österr. Geol. Ges.* 86, 133-140.
- Muttoni, G., Garzanti, E., Alfonsi, L., Cirilli, S., Germani, D., Lowrie, W., 2001. Motion of Africa and Adria since the Permian: paleomagnetic and paleoclimatic constraints from northern Libya. *Earth and Planetary Science Letters* 192, 159-174.
- Nemčok, M., 1993. Transition from convergence to escape: field evidence from the West Carpathians. *Tectonophysics* 217, 117-142.
- Nemčok, M., Nemčok, J., 1994. Late Cretaceous deformation of the Pieniny Klippen Belt, West Carpathians. *Tectonophysics* 239(1–4), 81–109.
- Plašienka, D., 2012. Jurassic syn-rift and Cretaceous synorogenic, coarse-grained deposits related to opening and closure of the Vahic (South Penninic) Ocean in the Western Carpathians, an overview. *Geological Quarterly* 56(4), 601–628.
- Platt, J.P., Behrmann, J.H., Cunningham, P.C., Dewey, J.F., 1989. Kinematics of the Alpine arc and the motion history of Adria, *Nature* 337, 158-161.
- Púchy, R., Varga, J., 1973. Tectonic map of the Carpathian–Balkan mountain system and adjacent areas (1:1,000,000). Geological Institute in Bratislava, and UNESCO.
- Ratschbacher, L., Froisch, W., Linzer, H.G., Merle, O., 1991a. Lateral extrusion in the Eastern Alps, part 2: Structural analysis. *Tectonics* 10(2) 257-271.
- Ratschbacher, L., Merle, O., Davy, P., Cobbold, P., 1991b. Lateral extrusion in the Eastern Alps, Part 1: boundary conditions and experiments scaled for gravity. *Tectonics* 10(2), 245-256.
- Ratschbacher, L., Frisch, W., Linzer, H.G., Sperner, B., Meschede, M., Decker, K., Nemčok, M., Nemčok, J., Grygar, R., 1993. The Pieniny Klippen Belt in the Western Carpathians of northeastern Slovakia: structural evidence for transpression. *Tectonophysics* 226, 471-483.
- Robertson, A.H.F., Shallo, M., 2000. Mesozoic–Tertiary tectonic evolution of Albania in its regional Eastern Mediterranean context. *Tectonophysics* 316, 197–254.
- Royden, L.H., Horváth, F., Burchfiel, B. C., 1982. Transform faulting, extension, and subduction in the Carpathian Pannonian region. *Geol. Soc. Am. Bull.*, 93, 717-725.
- Săndulescu, M., 1975. Essai de synthèse structurale des Carpathes. *Bulletin de la Société Géologique de France* 17/3, 299–358.

- Săndulescu, M., 1988. Cenozoic tectonic history of the Carpathians. In: Royden, L.H., Horváth, F. (Eds.), *The Pannonian Basin, a study in basin evolution*. American Association of Petroleum Geologists Memoir 35, pp. 17–25.
- Schmid, S.M., Bernoulli, D., Fügenschuh, B., Matenco, L., Schefer, S., Schuster, R., Tischler, M., Ustaszewski, K., 2008. The Alpine–Carpathian–Dinaridic orogenic system: compilation and evolution of tectonic units. *Swiss Journal of Geosciences* 101, 139–183.
- Schuller, V., 2004. Evolution and geodynamic significance of the Upper Cretaceous Gosau basin in the Apuseni Mountains (Romania). *Tübinger Geowissenschaftliche Arbeiten Reihe A* 70, 112 p.
- Ślączka, A., Kruglov, S., Golonka, J., Oszczypko, N., Popadyuk, I., 2006. The General Geology of the Outer Carpathians, Poland, Slovakia, and Ukraine. In: Golonka, J., Picha, F. (Eds.), *The Carpathians and their Foreland: Geology and Hydrocarbon Resources*, American Association of Petroleum Geologists Memoir 84, pp. 221–258.
- Ślączka, A., Renda, P., Cieszkowski, M., Golonka, J., Nigro, F., 2012. Sedimentary basins evolution and olistoliths formation: The case of Carpathian and Sicilian regions. *Tectonophysics* 568, 306–19.
- Smith, A.G., Spray, J.G., 1984. A half-ridge transform model for the Hellenic-Dinaric ophiolites. In: Dixon, J.E., Robertson, A.H.F. (Eds.), *The Geological Evolution of the Eastern Mediterranean*. Geological Society London Special Publication, 17, pp. 589–603.
- Stampfli, G.M., Hochard, C., 2009. Plate tectonics of the Alpine realm. In: Murphy, J.B., Keppie, J.D., Hynes, A.J. (Eds.), *Ancient orogens and modern analogues*. Special Publications 327, Geological Society, London, pp. 89–111.
- Stampfli, G.M., Mosar, J., Favre, P., Pillevuit, A., Vannay, J.-C., 2001. Permo-Mesozoic evolution of the western Tethyan realm: The NeoTethys/east Mediterranean basin connection. In: Cavazza, W., Robertson, A.H.F., Ziegler, P. (Eds.), *PeriTethyan Rift/Wrench Basins and Passive Margins*. *Memoires du Muséum National d'Histoire Naturelle, Paris, PeriTethys Memoir* 6, pp. 51–108.
- Suk, M., 1984. Geological history of the Czech Socialist Republic. Geological Survey, the Czechoslovak academy of sciences, Prague.
- Szabó, C., Harangi, S., Csontos, L., 1992. Review of Neogene and Quaternary volcanism of the Carpathian–Pannonian region. *Tectonophysics* 208, 243–256.
- Trümpy, R., 1988. A possible Jurassic–Cretaceous transform system in the Alps and the Carpathians. Geological Society of America, Special paper 218, 93–109.
- Ustaszewski, K., Schmid, S.M., Fügenschuh, B., Tischler, M., Kissling, E., Spakman, W., 2008. A map-view restoration of the Alpine–Carpathian–Dinaridic system for the Early Miocene. *Swiss J. Geosci.* 101, Supplement 1, 273–294.
- Wölfler, A., Kurz, W., Fritz, H., Stüwe, K., 2011. Lateral extrusion in the Eastern Alps revisited: refining the model by thermochronological, sedimentary and seismic data. *Tectonics* 30(4), TC4006, doi:10.1029/2010TC002782.
- Zatopek, A., 1979. On geodynamical aspects of geophysical synthesis in Central Europe. Geodynamic investigations in Czechoslovakia. Final Rep., Veda, Bratislava, 91–104.