



HAL
open science

The coupling of non-linear supersymmetry to supergravity

Ignatios Antoniadis, Chrysoula Markou

► **To cite this version:**

Ignatios Antoniadis, Chrysoula Markou. The coupling of non-linear supersymmetry to supergravity. European Physical Journal C: Particles and Fields, 2015, 75 (12), pp.582. 10.1140/epjc/s10052-015-3783-0 . hal-01242943

HAL Id: hal-01242943

<https://hal.sorbonne-universite.fr/hal-01242943>

Submitted on 14 Dec 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

The coupling of non-linear supersymmetry to supergravity

Ignatios Antoniadis^{1,2,a}, Chrysoula Markou^{1,b} 

¹ LPTHE, UMR CNRS 7589, Sorbonne Universités, UPMC Paris 6, 75005 Paris, France

² Albert Einstein Center for Fundamental Physics, Institute for Theoretical Physics, University of Bern, Sidlestrasse 5, 3012 Bern, Switzerland

Received: 4 October 2015 / Accepted: 5 November 2015

© The Author(s) 2015. This article is published with open access at Springerlink.com

Abstract We study the coupling of non-linear supersymmetry to supergravity. The goldstino nilpotent superfield of global supersymmetry coupled to supergravity is described by a geometric action of the chiral curvature superfield \mathcal{R} subject to the constraint $(\mathcal{R} - \lambda)^2 = 0$ with an appropriate constant λ . This constraint can be found as the decoupling limit of the scalar partner of the goldstino in a class of $f(\mathcal{R})$ supergravity theories.

1 Introduction

Studies of non-linear supersymmetric actions have been revived recently due to potential interesting applications in particle physics [1] and cosmology [2–4] and their realization in particular string compactifications [5,6]. Indeed, effective actions with non-linear supersymmetry parametrize in a model independent way the effects of supersymmetry breaking at low energies compared to the mass of the sgoldstino (supersymmetric partner of the goldstino) which is in general of the order of the supersymmetry breaking scale at the ‘hidden’ sector. The goldstino on the other hand, although part of the massive gravitino providing its longitudinal degrees of freedom, is always in the low-energy spectrum since it becomes massless in the absence of gravity and interacts with a strength fixed by the supersymmetry breaking scale, in contrast to the transverse gravitino components, which become free and decouple. In the string theory context, non-linear supersymmetry appears naturally on the world-volume of D-branes realizing the broken supersymmetries of the bulk. It can then even remain an exact symmetry of certain vacua, if for instance the orientifold projection respects it [5,7].

At the global level, a convenient way to write off-shell non-linear supersymmetric actions is by utilizing a nilpotent chiral superfield [8–11]. In analogy to ordinary non-linear sigma

models, the constraint eliminates the sgoldstino component, playing the role of the radial Higgs mode, and replaces it by a goldstino bilinear. In the absence of matter fields, the low-energy action (i.e. without higher order super-derivatives) is completely determined in terms of the goldstino decay constant (or equivalently the supersymmetry breaking scale), and reproduces [11, 12] the Volkov–Akulov action [13] on-shell. Indeed, the most general Kähler potential is canonical and the superpotential is linear in the nilpotent goldstino superfield. In the presence of matter, the simple nilpotent constraint may change if some matter fields have superheavy superpartners (of the order of the sgoldstino mass) [14], but it remains valid if all other extra fields belong to ordinary linear supermultiplets [1].

The coupling to supergravity is straightforward since the constraint does not involve any derivatives [2]. The superpotential now admits also a constant piece, allowing for an arbitrary cosmological constant of any sign and space-time to be anti-de Sitter, de Sitter or flat. In flat space, the gravitino mass is given by the usual relation in terms of the supersymmetry breaking scale and the Planck mass. In the unitary gauge, the action is reduced to the ordinary $N = 1$ supergravity with a mass term for the gravitino that has absorbed the goldstino. The theory has an alternative geometric formulation in terms of the chiral curvature superfield \mathcal{R} , which obeys an appropriate quadratic constraint $(\mathcal{R} - \lambda)^2 = 0$ with λ a constant [2, 15, 16].

In this work, we first show the equivalence of the two formulations of non-linear supersymmetry coupled to supergravity by computing explicitly the two actions in components. An alternative way to obtain the constraint is to add it in the action with an independent coupling-coefficient ρ and take the limit $\rho \rightarrow \infty$. The resulting \mathcal{R}^2 supergravity contains besides the graviton and the gravitino the degrees of freedom of a chiral multiplet that should play the role of the goldstino multiplet. It turns out, however, that this theory does not have a minimum in flat space for finite ρ , while, starting from a de Sitter minimum, the decoupling limit of

^a e-mail: antoniad@lpthe.jussieu.fr

^b e-mail: chrysoula@lpthe.jussieu.fr

the sgoldstino, and thus of non-linear supersymmetry, does not exist. We then study a general class of $f(\mathcal{R})$ $N = 1$ supergravity theories [17, 18] that satisfy the required limit in flat space; the mass of the extra complex scalar goes to infinity in a Minkowski minimum of the scalar potential and the geometric constraint for the chiral curvature is recovered.

The structure of the paper is as follows. In Sect. 2, we show the equivalence between the two formulations of non-linear supersymmetry coupled to $N = 1$ supergravity using first a formal argument with superfields and then by comparing the two effective actions in components. We find two possible values for the constant λ entering the geometric constraint: $\lambda = 0$ and $\lambda = 6W_0$ with W_0 a constant superpotential. In Sect. 3, we recover this constraint in the sgoldstino decoupling limit of a particular class of $f(\mathcal{R})$ supergravity theory that we construct as a Taylor series expansion around a flat space minimum of the corresponding scalar potential. Finally, in ‘‘Appendix A’’ we give some details for the derivation of the solution of the geometric constraint, while in ‘‘Appendix B’’ we show why an \mathcal{R}^2 supergravity does not have a stable supersymmetry breaking vacuum that reproduces the constraint in a suitable limit.

2 Two equivalent Lagrangians

In the constrained superfield formalism of non-linear supersymmetry, the goldstino is described by the fermionic component of a chiral superfield, X , that satisfies the nilpotent constraint $X^2 = 0$ [8–11]. The scalar component (sgoldstino) is then eliminated by the constraint and is replaced by a goldstino bilinear. The most general low-energy (without superderivatives) Lagrangian, invariant (upon space-time integration) under global supersymmetry, is then given by

$$\mathcal{L}_{\text{VA}} = [X\bar{X}]_D + ([fX]_F + \text{h.c.}), \quad (2.1)$$

where $f \neq 0$ is a complex parameter. The subscripts D and F denote D- and F-term densities, integrated over the full or the chiral superspace, respectively, and correspond to the Kähler potential and superpotential of $N = 1$ supersymmetry. It can be shown [11, 12] that \mathcal{L}_{VA} is equivalent to the Volkov–Akulov Lagrangian [13] on-shell.

The coupling to supergravity in the superconformal context [19, 20], (2.1), takes the form

$$\mathcal{L} = -[(1 - X\bar{X})S_0\bar{S}_0]_D + \left(\left[\left(fX + W_0 + \frac{1}{2}TX^2 \right) S_0^3 \right]_F + \text{h.c.} \right), \quad (2.2)$$

where we have used the superconformal tensor calculus [21], [22] with S_0 being the superconformal compensator superfield. We have also used a Lagrange multiplier T in order to impose the constraint $X^2 = 0$ explicitly in \mathcal{L} , while the

factor $\frac{1}{2}$ is put merely for convenience. W_0 is a complex constant parameter whose importance will appear shortly. The Kähler potential corresponding to (2.2) is given by

$$K(X, \bar{X}) = -3\ln(1 - X\bar{X}) = -3 \left[-X\bar{X} - \frac{(-X\bar{X})^2}{2} + \dots \right] = 3X\bar{X}. \quad (2.3)$$

We would now like to find a geometrical formulation of (2.2), that is, to eliminate X and write an equivalent Lagrangian that contains only superfields describing the geometry of spacetime, such as the superspace chiral curvature \mathcal{R} [2, 15, 16]. To this aim, we observe that the following Kähler potential K' :

$$K' = -3\ln(1 + X + \bar{X}) = -3 \left(X + \bar{X} - \frac{(X + \bar{X})^2}{2} + \dots \right) = 3X\bar{X} - 3(X + \bar{X}), \quad (2.4)$$

is related to the Kähler potential K via a Kähler transformation of the type

$$\begin{aligned} K &\rightarrow K' = K - 3(X + \bar{X}) \\ W &\rightarrow W' = e^{3X}W. \end{aligned} \quad (2.5)$$

This tells us that \mathcal{L} is equivalent to \mathcal{L}' , where

$$\mathcal{L}' = -[(1 + X + \bar{X})S_0\bar{S}_0]_D + \left(\left[\left(fX + W_0 + \frac{1}{2}TX^2 \right) e^{3X}S_0^3 \right]_F + \text{h.c.} \right). \quad (2.6)$$

Using the constraint $X^2 = 0$, we have

$$\begin{aligned} \mathcal{L}' &= -[(1 + X + \bar{X})S_0\bar{S}_0]_D \\ &+ \left(\left[\left(fX + W_0(1 + 3X) + \frac{1}{2}TX^2 \right) S_0^3 \right]_F + \text{h.c.} \right) \\ &= -[S_0\bar{S}_0]_D + \left(\left[\left(\lambda X + W_0 - X\frac{\mathcal{R}}{S_0} + \frac{1}{2}TX^2 \right) S_0^3 \right]_F + \text{h.c.} \right), \end{aligned} \quad (2.7)$$

where we have set $\lambda = f + 3W_0$ and we have used the identity [22]

$$[X \cdot \mathcal{R} \cdot S_0^2]_F = [S_0\bar{S}_0(X + \bar{X})]_D + \text{total derivatives}. \quad (2.8)$$

In (2.7), X enters only in F-terms without derivatives and can be thus integrated out. Solving the equation of motion for X , we have

$$\lambda - \frac{\mathcal{R}}{S_0} + TX = 0 \quad \Rightarrow \quad X = \frac{\frac{\mathcal{R}}{S_0} - \lambda}{T}, \quad (2.9)$$

and substituting it back into (2.7), we get

$$\begin{aligned} \mathcal{L}' &= -[S_0 \bar{S}_0]_D + \left[\left(-\frac{1}{2T} \left(\frac{\mathcal{R}}{S_0} - \lambda \right)^2 + W_0 \right) S_0^3 \right]_F + \text{h.c.} \\ &= \left[\left(-\frac{1}{2} \frac{\mathcal{R}}{S_0} + W_0 - \frac{1}{2T} \left(\frac{\mathcal{R}}{S_0} - \lambda \right)^2 \right) S_0^3 \right]_F + \text{h.c.}, \end{aligned} \tag{2.10}$$

where we have used again the identity (2.8). We can now view $\frac{1}{T}$ as a Lagrange multiplier that imposes the constraint

$$\left(\frac{\mathcal{R}}{S_0} - \lambda \right)^2 = 0. \tag{2.11}$$

Consequently, we have established an equivalence between the constrained Lagrangians (2.2) and (2.10); they both describe the coupling of non-linear supersymmetry to supergravity, with \mathcal{L}' providing its geometric formulation with the use of a constraint imposed on \mathcal{R} instead of X . This constraint was proposed in [2] for $\lambda = 0$. In what follows we will confirm the equivalence by writing these Lagrangians in terms of component fields.

2.1 Constraining a chiral superfield X

In the following we use the method and conventions of [23] except from a factor of 1/6 which we omit in the expression of \mathcal{R} but introduce at the Lagrangian level. We also set the gravitational coupling $\kappa^2 = 8\pi G_N$ (given here in natural units) to be equal to one, in accordance with the usual convention. After gauge-fixing the superconformal symmetry by using the convenient gauge $S_0 = 1$, the Lagrangian (2.2) can be written as follows:

$$\begin{aligned} \mathcal{L} &= \int d^2\Theta 2\mathcal{E} \left\{ \frac{3}{8} \left(\bar{\mathcal{D}}\bar{\mathcal{D}} - \frac{8}{6}\mathcal{R} \right) e^{-K/3} + W \right\} + \text{h.c.} \\ &\text{with } W(X) = fX + W_0 \text{ and } X^2 = 0, \end{aligned} \tag{2.12}$$

where \mathcal{D} is the super-covariant derivative and \mathcal{E} the chiral superfield density that is constructed from the vielbein e_a^m :

$$\mathcal{E} = \frac{1}{2} e \{ 1 + i\Theta\sigma^a\bar{\psi}_a - \Theta\Theta[\bar{M} + \bar{\psi}_a\bar{\sigma}^{ab}\bar{\psi}_b] \}. \tag{2.13}$$

Here ψ_a is the gravitino, Θ the fermionic coordinates of the curved superspace and $\sigma^a = (-1, \sigma)$, $\sigma_\alpha^{ab} = \frac{1}{4}(\sigma_{\alpha\dot{\alpha}}^a \bar{\sigma}^{b\dot{\alpha}\beta} - \sigma_{\alpha\dot{\alpha}}^b \bar{\sigma}^{a\dot{\alpha}\beta})$ with σ the Pauli matrices. Note that the Lagrange multiplier T in (2.2) has been used to impose the constraint $X^2 = 0$, which can be solved, fixing the scalar component (sgoldstino) in terms of the goldstino G and the auxiliary field F of X [11].

We now substitute X , \mathcal{E} and \mathcal{R} with their respective expressions in component fields:

$$\begin{aligned} X &= \frac{G^2}{2F} + \sqrt{2}\Theta G + (\Theta\Theta)F \equiv A + \sqrt{2}\Theta G + (\Theta\Theta)F \\ \mathcal{R} &\equiv -M - \Theta B - (\Theta\Theta)C \\ \Xi &\equiv \left(\bar{\mathcal{D}}\bar{\mathcal{D}} - \frac{8}{6}\mathcal{R} \right) \bar{X} \equiv -4\bar{F} + \frac{4}{3}M\bar{A} + \Theta D + (\Theta\Theta)E. \end{aligned} \tag{2.14}$$

The exact components of \mathcal{R} and Ξ are computed, for example, in [23] (our convention for \mathcal{R} differs by 1/6 with respect to [23]). M and b_a are the auxiliary fields of the $N = 1$ supergravity multiplet in the old-minimal formulation. Then

$$\begin{aligned} &-\frac{3}{4} \left[\mathcal{E} \left(\bar{\mathcal{D}}\bar{\mathcal{D}} - \frac{8}{6}\mathcal{R} \right) \bar{X} X \right]_F \\ &= -\frac{3}{8} [2\mathcal{E}\Xi X]_F = -\frac{3}{8} e \left(EA - 4F\bar{F} + \frac{4}{3}MF\bar{A} - \frac{\sqrt{2}}{2}(DG) \right) \\ &\quad + \frac{3}{16} ie(y\sigma^a\bar{\psi}_a) + \frac{3}{8} e [\bar{M} + \bar{\psi}_a\bar{\sigma}^{ab}\bar{\psi}_b] \left[-4A\bar{F} + \frac{4}{3}MA\bar{A} \right], \end{aligned} \tag{2.15}$$

where

$$y = \sqrt{2}G \left(-4\bar{F} + \frac{4}{3}M\bar{A} \right) + DA. \tag{2.16}$$

This expression is simplified significantly if we choose to use the unitary gauge, setting $G = 0$ and thus $A = y = 0$:

$$-\frac{3}{4} \left[\mathcal{E} \left(\bar{\mathcal{D}}\bar{\mathcal{D}} - \frac{8}{6}\mathcal{R} \right) \bar{X} X \right]_F = \frac{3}{2} e F \bar{F}. \tag{2.17}$$

Moreover, also in the unitary gauge, one can compute

$$[2\mathcal{E}(fX + W_0)]_F = efF - e[\bar{M} + \bar{\psi}_a\bar{\sigma}^{ab}\bar{\psi}_b]W_0. \tag{2.18}$$

Now, using the property

$$\begin{aligned} (\bar{\psi}_a\bar{\sigma}^{ab}\bar{\psi}_b)^\dagger &= \frac{1}{4} [\bar{\psi}_a(\bar{\sigma}^a\sigma^b - \bar{\sigma}^b\sigma^a)\bar{\psi}_b]^\dagger \\ &= \frac{1}{4} [\psi_b(\sigma^b\bar{\sigma}^a - \sigma^a\bar{\sigma}^b)\psi_a] = \psi_a\sigma^{ab}\psi_b, \end{aligned} \tag{2.19}$$

the Lagrangian (2.12) in terms of the component fields becomes

$$\begin{aligned} \mathcal{L} &= -\frac{1}{2}eR - \frac{1}{3}eM\bar{M} + \frac{1}{3}eb^ab_a \\ &\quad + \frac{1}{2}e\epsilon^{abcd}(\bar{\psi}_a\bar{\sigma}_b\bar{\mathcal{D}}_c\psi_d - \psi_a\sigma_b\bar{\mathcal{D}}_c\bar{\psi}_d) \\ &\quad + efF - eW_0[\bar{M} + \bar{\psi}_a\bar{\sigma}^{ab}\bar{\psi}_b] \\ &\quad + e\bar{f}\bar{F} - e\bar{W}_0[M + \psi_a\sigma^{ab}\psi_b] + 3eF\bar{F}, \end{aligned} \tag{2.20}$$

where R is the Ricci scalar. The equations of motion for the auxiliary fields b^a, M, F are then

$$\begin{aligned} b^a &= 0, \\ M &= -3W_0, \bar{M} = -3\bar{W}_0, \\ F &= -\frac{\bar{f}}{3}, \bar{F} = -\frac{f}{3}. \end{aligned} \tag{2.21}$$

Substituting it back into (2.20) we get

$$\begin{aligned} \mathcal{L} &= -\frac{1}{2}eR + \frac{1}{2}e\epsilon^{abcd}(\bar{\psi}_a\bar{\sigma}_b\tilde{\mathcal{D}}_c\psi_d - \psi_a\sigma_b\tilde{\mathcal{D}}_c\bar{\psi}_d) \\ &\quad - eW_0\bar{\psi}_a\bar{\sigma}^{ab}\bar{\psi}_b - e\bar{W}_0\psi_a\sigma^{ab}\psi_b + 3e|W_0|^2 - \frac{1}{3}e|f|^2. \end{aligned} \tag{2.22}$$

In this form, it is obvious that the Lagrangian reduces to the usual $N = 1$ supergravity, together with a gravitino mass term:

$$m_{3/2} = |W_0|. \tag{2.23}$$

Imposing that the cosmological constant (i.e. the vacuum expectation value of the scalar potential) vanishes, one finds

$$3|W_0|^2 - \frac{1}{3}|f|^2 = 0 \Rightarrow |f|^2 = 9|W_0|^2. \tag{2.24}$$

This means that $W_0 \neq 0$, which justifies the use of the constant piece W_0 in the superpotential in \mathcal{L} . Then the final form of \mathcal{L} is

$$\begin{aligned} \mathcal{L} &= -\frac{1}{2}eR + \frac{1}{2}e\epsilon^{abcd}(\bar{\psi}_a\bar{\sigma}_b\tilde{\mathcal{D}}_c\psi_d - \psi_a\sigma_b\tilde{\mathcal{D}}_c\bar{\psi}_d) \\ &\quad - eW_0\bar{\psi}_a\bar{\sigma}^{ab}\bar{\psi}_b - e\bar{W}_0\psi_a\sigma^{ab}\psi_b. \end{aligned} \tag{2.25}$$

It is important to notice that the use of the constrained superfield X is what has generated the gravitino mass term: the final form of the Lagrangian in flat space is just the pure $N = 1$ supergravity, but with a massive gravitino. The use of the unitary gauge $G = 0$ results in the gravitino absorbing the goldstino and becoming massive, in analogy with the well-known Brout–Englert–Higgs mechanism.

2.2 Constraining the superspace curvature superfield \mathcal{R}

After gauge-fixing the superconformal symmetry by imposing $S_0 = 1$, the Lagrangian (2.10) can be written as follows:

$$\begin{aligned} \mathcal{L}' &= -\int d^2\Theta \mathcal{E}(\mathcal{R} - 2W_0) + \text{h.c.}, \\ (\mathcal{R} - \lambda)^2 &= 0. \end{aligned} \tag{2.26}$$

\mathcal{L}' then yields

$$\begin{aligned} \mathcal{L}' &= -\frac{1}{2}eR - \frac{1}{3}eM\bar{M} + \frac{1}{3}eb^a b_a \\ &\quad + \frac{1}{2}e\epsilon^{abcd}(\bar{\psi}_a\bar{\sigma}_b\tilde{\mathcal{D}}_c\psi_d - \psi_a\sigma_b\tilde{\mathcal{D}}_c\bar{\psi}_d) \\ &\quad - eW_0[\bar{M} + \bar{\psi}_a\bar{\sigma}^{ab}\bar{\psi}_b] - e\bar{W}_0[M + \psi_a\sigma^{ab}\psi_b]. \end{aligned} \tag{2.27}$$

Now let us solve the constraint which is the second of Eq. (2.26). To this aim, we substitute the second of Eq. (2.14) into the constraint and find the set of the following equations:

$$\begin{aligned} (M + \lambda)^2 &= 0, \\ (M + \lambda)B_\alpha &= 0, \\ 4(M + \lambda)C &= (BB), \end{aligned} \tag{2.28}$$

where

$$\begin{aligned} B_\alpha &= \sigma_{\alpha\dot{\alpha}}^a \bar{\sigma}^{b\dot{\beta}\beta} \psi_{ab\beta} - i\sigma_{\alpha\dot{\alpha}}^a \bar{\psi}_a^{\dot{\alpha}} M + i\psi_{a\alpha} b^a \\ &\quad \text{with } \psi_{ab} \equiv \tilde{\mathcal{D}}_a\psi_b - \tilde{\mathcal{D}}_b\psi_a \\ C &= -\frac{1}{2}R + \mathcal{O}\{M, b_a, \psi_a\} \neq 0. \end{aligned} \tag{2.29}$$

Equation (2.28) yield

$$M = -\lambda \quad \text{and} \quad b^a = 0. \tag{2.30}$$

Indeed, B in this case depends only on the gamma-trace or the divergence of the gravitino, $\bar{\sigma}^a\psi_a$ and $\tilde{\mathcal{D}}^a\psi_a$ (using the Clifford algebra property of sigma-matrices $(\sigma^a\bar{\sigma}^b + \sigma^b\bar{\sigma}^a)_\alpha^\beta = -2\eta^{ab}\delta_\alpha^\beta$), which can be put to zero by an appropriate gauge choice. Alternatively, one can show that B vanishes on-shell (see ‘‘Appendix A’’).

Using (2.30), Eq. (2.27) becomes

$$\begin{aligned} \mathcal{L}' &= -\frac{1}{2}eR - \frac{1}{3}e|\lambda|^2 + \frac{1}{2}e\epsilon^{abcd}(\bar{\psi}_a\bar{\sigma}_b\tilde{\mathcal{D}}_c\psi_d - \psi_a\sigma_b\tilde{\mathcal{D}}_c\bar{\psi}_d) \\ &\quad + eW_0\bar{\lambda} + e\bar{W}_0\lambda - eW_0\bar{\psi}_a\bar{\sigma}^{ab}\bar{\psi}_b - e\bar{W}_0\psi_a\sigma^{ab}\psi_b. \end{aligned} \tag{2.31}$$

Substituting now $\lambda = f + 3W_0$, one finds that the cosmological constant is given by $3e|W_0|^2 - \frac{1}{3}e|f|^2$ and the Lagrangian (2.31) is identical to (2.22). Note that the vanishing of the cosmological constant

$$-\frac{1}{3}e|\lambda|^2 + eW_0\bar{\lambda} + e\bar{W}_0\lambda = 0 \tag{2.32}$$

gives two possible solutions for λ :

$$\lambda = 6W_0 \quad \text{and} \quad \lambda = 0, \tag{2.33}$$

corresponding to $f = \pm 3W_0$, which solves the condition (2.24).

3 Without imposing direct constraints

In this section, we would like to start with a regular \mathcal{R}^2 supergravity and recover the constraint in an appropriate limit where the additional (complex) scalar arising from \mathcal{R}^2 becomes superheavy and decouples from the low-energy spectrum. Indeed, by analogy with ordinary General Relativity in the presence of an R^2 -term (with R the scalar curvature), an \mathcal{R}^2 supergravity can be re-written as an ordinary Einstein $N = 1$ supergravity coupled to an extra chiral multiplet.¹ Let us then consider the Lagrangian

$$\bar{\mathcal{L}} = \left[\left(-\frac{1}{2} \frac{\mathcal{R}}{S_0} + W_0 + \frac{1}{2} \rho \left(\frac{\mathcal{R}}{S_0} - \lambda \right)^2 \right) S_0^3 \right]_F + \text{h.c.}, \tag{3.1}$$

where ρ is a real parameter. In the limit $|\rho| \rightarrow \infty$, one would naively expect to recover the constraint $(\mathcal{R} - \lambda)^2 \rightarrow 0$, and thus (3.1) should be reduced to (2.10). In principle, one could linearize (3.1) with the use of a chiral superfield S and then demonstrate that in the limit $|\rho| \rightarrow \infty$, \mathcal{L} , \mathcal{L}' and $\bar{\mathcal{L}}$ are all equivalent. If this were true, one would expect that S corresponds to the goldstino superfield and that supersymmetry is non-linearly realized (in the limit $|\rho| \rightarrow \infty$), as is the case for the chiral nilpotent superfield X . In other words, the mass of the scalar component of S would approach infinity as $|\rho| \rightarrow \infty$ and would, therefore, decouple from the spectrum. However, upon computing the scalar potential and the scalar mass matrix corresponding to (3.1), we found that this is not the case. This means that the parameter space (λ, W_0, ρ) does not allow for a supersymmetry breaking minimum that realizes the goldstino decoupling and the equivalence between $\bar{\mathcal{L}}$ with \mathcal{L} and \mathcal{L}' . The detailed analysis can be found in ‘‘Appendix B’’.

To solve this problem, we start with a more general class of $f(\mathcal{R})$ supergravity actions. More precisely, we modify $\bar{\mathcal{L}}$ with the addition of a suitable term that is suppressed by ρ in the limit $|\rho| \rightarrow \infty$ ²:

$$\mathcal{L}'' = \left[\left(-\frac{1}{2} \frac{\mathcal{R}}{S_0} + W_0 + \frac{1}{2} \rho \left(\frac{\mathcal{R}}{S_0} - \lambda \right)^2 + \frac{1}{\rho} \left(S \frac{\mathcal{R}}{S_0} - F(S) \right) \right) S_0^3 \right]_F + \text{h.c.}, \tag{3.2}$$

where S is a chiral superfield coupled to gravity and $F(S)$ is a holomorphic function of the superfield S . This extra term has already been studied in the literature and is known as

¹ Note that \mathcal{R}^2 supergravity is not the supersymmetrization of R^2 gravity which is described by a D-term $\mathcal{R}\mathcal{R}$, bringing about two chiral multiplets to be linearized [21,22].

² In principle, we may replace $1/\rho$ by $1/\hat{\rho}(\rho)$ with $|\hat{\rho}(\rho)| \rightarrow \infty$ when $\rho \rightarrow \infty$. One can show, however, that our results do not change and thus we make the simple choice $\hat{\rho} = \rho$.

$f(\mathcal{R})$ supergravity [17,18]. Indeed, S can be integrated out by its equation of motion at finite ρ :

$$\mathcal{R} = F' S_0, \tag{3.3}$$

where $F' = \frac{\partial F}{\partial S}$. This equation can be in principle solved to give S as a function of \mathcal{R} and replacing it back in (3.2) one finds an $f(\mathcal{R})$ theory.

We will now study the physical implications of \mathcal{L}'' in the limit $\rho \rightarrow \infty$ so as to confirm the equivalence between \mathcal{L} , \mathcal{L}' and \mathcal{L}'' (without loss of generality, we take ρ positive). We first use Eq. (3.3) to replace \mathcal{R} in terms of S in the third term of (3.2), instead of doing the reverse as described above. Using then the identity (2.8), we get

$$\mathcal{L}'' = - \left[\left(1 - \frac{1}{\rho} (S + \bar{S}) \right) S_0 \bar{S}_0 \right]_D + \left\{ \left[\left(W_0 + \frac{1}{2} \rho (F' - \lambda)^2 - \frac{1}{\rho} F \right) S_0^3 \right]_F + \text{h.c.} \right\}. \tag{3.4}$$

We now fix the gauge according to $S_0 = 1$ and set ϕ to be the lowest component of S . Then the Kähler potential and the superpotential corresponding to \mathcal{L}'' are given by

$$K = -3 \ln \left(1 - \frac{1}{\rho} (\phi + \bar{\phi}) \right) \\ W = W_0 + \frac{1}{2} \rho (F' - \lambda)^2 - \frac{1}{\rho} F, \tag{3.5}$$

where now $F' = \frac{\partial F}{\partial \phi}$. (We use the same symbols K and W as in Sect. 2 as there is no confusion.)

It follows that

$$\exp(K) = \frac{\rho^3}{(\rho - \phi - \bar{\phi})^3} \tag{3.6}$$

and

$$g_{\phi\bar{\phi}} = \frac{\partial}{\partial \phi} \frac{\partial}{\partial \bar{\phi}} K = \frac{3}{(\rho - \phi - \bar{\phi})^2}, \quad g^{\phi\bar{\phi}} = \frac{(\rho - \phi - \bar{\phi})^2}{3}. \tag{3.7}$$

Also

$$D_\phi W = \partial_\phi W + K_\phi W = \rho F'' (F' - \lambda) - \frac{1}{\rho} F' \\ + \frac{3}{\rho - \phi - \bar{\phi}} \left(W_0 + \frac{1}{2} \rho (F' - \lambda)^2 - \frac{1}{\rho} F \right). \tag{3.8}$$

Putting everything together, we find that the scalar potential V is given by

$$V = \exp(K) \left[g^{\phi\bar{\phi}} (D_\phi W) (\bar{D}_{\bar{\phi}} \bar{W}) - 3 \bar{W} W \right] = \frac{\rho^2}{3(\rho - \phi - \bar{\phi})^2} \tilde{V}, \tag{3.9}$$

where

$$\begin{aligned} \tilde{V} = & \rho^4 |F''(F' - \lambda)|^2 + \rho^3 \\ & \times \left[-(\phi + \bar{\phi}) |F''(F' - \lambda)|^2 \right. \\ & \left. + \frac{3}{2} |F' - \lambda|^2 (F''(\bar{F}' - \bar{\lambda}) + \text{h.c.}) \right] \\ & + \rho^2 [-\bar{F}' F''(F' - \lambda) + 3\bar{W}_0 F''(F' - \lambda) + \text{h.c.}] \\ & + \rho [(\phi + \bar{\phi}) \bar{F}' F''(F' - \lambda) - 3\bar{F}' F''(F' - \lambda) \\ & - \frac{3}{2} F'(\bar{F}' - \bar{\lambda})^2 + \text{h.c.}] + \rho^0 [|F'|^2 - 3F' \bar{W}_0 - 3\bar{F}' W_0] \\ & + \rho^{-1} [-(\phi + \bar{\phi}) |F'|^2 + 3F' \bar{F}' + 3\bar{F}' F']. \end{aligned} \tag{3.10}$$

For $\rho \rightarrow \infty$, the leading behavior of V is given by

$$V = \frac{\rho^4}{3} |F''(F' - \lambda)|^2. \tag{3.11}$$

It is positive definite with a minimum at zero when $F' = \lambda$ or $F'' = 0$. In the following, we will analyze the case $F' = \lambda$; its curvature defines the (canonically normalized) scalar mass given by

$$m_\phi = \frac{\rho^3}{3} (F'')^2 \tag{3.12}$$

which goes to infinity at large ρ and ϕ decouples. At the minimum $F' = \lambda$, the potential at large ρ becomes constant, proportional to $|\lambda|^2 - 3\lambda \bar{W}_0 - 3\bar{\lambda} W_0$. This term vanishes precisely if Eq. (2.32), or equivalently (2.24), holds. We conclude that in the model (3.2) the cosmological constant can be tuned to zero (in the limit $\rho \rightarrow \infty$) by using the same condition as for the model (2.10). As shown in Sect. 2.2, this is the case for two possible values of λ :

$$\boxed{\lambda = 6W_0} \quad \text{or} \quad \boxed{\lambda = 0}. \tag{3.13}$$

Now let us investigate the minimum of the potential at finite but large ρ . We shall construct the solution as a power series in $1/\rho$ around the asymptotic field value of the minimum $\phi = \phi_0$ that solves $F' = \lambda$. Simple inspection of the potential (3.10) shows that it is sufficient to consider only even powers in $1/\rho$:

$$\begin{aligned} \phi = & \phi_0 + \frac{\phi_1}{\rho^2} \\ F'(\phi) = & F'(\phi_0) + (\phi - \phi_0) F''(\phi_0) \\ & + \frac{1}{2} (\phi - \phi_0)^2 F'''(\phi_0) + \dots \end{aligned} \tag{3.14}$$

or equivalently,

$$F'(\phi) = \lambda + \frac{c}{\rho^2} + \frac{d}{\rho^4} + \dots \tag{3.15}$$

where

$$c = \phi_1 F_0'', \quad d = \frac{1}{2} \phi_1^2 F_0'''. \tag{3.16}$$

We then compute the derivative of \tilde{V} with respect to ϕ and keep only the terms that do not vanish in the limit $\rho \rightarrow \infty$:

$$\begin{aligned} \tilde{V}_\phi = \frac{\partial \tilde{V}}{\partial \phi} = & \rho^4 (\bar{F}'' F''' |F' - \lambda|^2 + |F''|^2 F''(\bar{F}' - \bar{\lambda})) \\ & - \rho^3 (\phi + \bar{\phi}) |F''|^2 F''(\bar{F}' - \bar{\lambda}) + \rho^2 [F''^2 (3\bar{W}_0 - \bar{F}')] \\ & - |F''|^2 (\bar{F}' - \bar{\lambda}) + F'''(F' - \lambda) (3\bar{W}_0 - \bar{F}')] \\ & + \rho F''^2 [\bar{F}'(\phi + \bar{\phi}) - 3\bar{F}] + \rho^0 F'' [\bar{F}' - 3\bar{W}_0]. \end{aligned} \tag{3.17}$$

This expression vanishes if every coefficient at each order vanishes.

We now substitute the expansion (3.14), (3.15) into \tilde{V}_ϕ (ignoring orders that vanish as ρ^{-2} and higher) and impose each coefficient to be set to zero so as to have an extremum. Assuming for simplicity that $W_0, \lambda, \phi_0, c, d$ are real, we find the following constraints on the function F :

$$\begin{aligned} cF_0'' = & \lambda - 3W_0, \\ F_0 = & 2W_0\phi_0, \\ c^2 F_0''' = & \frac{2}{3} (\lambda - 3W_0), \end{aligned} \tag{3.18}$$

which yield

$$\begin{aligned} F(\phi) = & 2\phi_0 W_0 + \lambda(\phi - \phi_0) + \frac{\lambda - 3W_0}{2c} (\phi - \phi_0)^2 \\ & + \frac{1}{3!} \frac{2(\lambda - 3W_0)}{3c^2} (\phi - \phi_0)^3 + \dots \\ = & 2\phi_0 W_0 + \lambda(\phi - \phi_0) \pm \frac{3W_0}{2c} (\phi - \phi_0)^2 \\ & \pm \frac{1}{3!} \frac{2W_0}{c^2} (\phi - \phi_0)^3 + \dots, \end{aligned} \tag{3.19}$$

where in the second line above, we used the two possible values of λ (3.13), $\lambda = 6W_0$ for the + sign and $\lambda = 0$ for the - sign, for which the potential vanishes at the minimum.

At the minimum, the F-auxiliary term of S, \mathcal{F}_ϕ , is given by

$$\begin{aligned} \langle |\mathcal{F}^\phi| \rangle = & \left\langle \left| e^{K/2} \sqrt{g^{\phi\bar{\phi}}} \bar{D}_{\bar{\phi}} \bar{W} \right| \right\rangle \\ \xrightarrow{\rho \rightarrow \infty} & \frac{\rho^2}{\sqrt{3}} \langle |F''(F' - \lambda)| \rangle + (\text{subleading terms}) \\ = & \frac{1}{\sqrt{3}} \langle |F_0'' c| \rangle + \mathcal{O}(1/\rho^2) = \frac{1}{\sqrt{3}} |\lambda - 3W_0| \\ = & \sqrt{3} |W_0| \neq 0, \end{aligned} \tag{3.20}$$

where in the third line we used $\lambda = 0$ or $\lambda = 6W_0$. We conclude that supersymmetry is spontaneously broken in this limit along the direction of ϕ , which can be identified with the scalar superpartner of the goldstino that becomes superheavy and decouples. The supersymmetry breaking scale remains finite and is given by $f = 3|W_0|$. Therefore, we identify the fermionic component of S with the goldstino and ϕ with its superpartner, the sgoldstino. According to (3.12), the latter decouples from the spectrum in the limit $\rho \rightarrow \infty$, which is equivalent to imposing the nilpotent constraint for the goldstino superfield $X^2 = 0$ on \mathcal{L} . Finally, the gravitino mass is given by

$$m_{3/2} = \langle |W|e^{K/2} \rangle \rightarrow |W_0| \quad \text{as } \rho \rightarrow \infty, \quad (3.21)$$

which completes the proof of equivalence between \mathcal{L} , \mathcal{L}' and \mathcal{L}'' .

Acknowledgments I.A. would like to thank E. Dudas for enlightening discussions. C.M. would like to thank the laboratory LPTHE of the UPMC for partial financial support as well as for the warm hospitality in regard to the Master’s Thesis. C.M. is also grateful to the Greek State Scholarships Foundation (IKY) for partially funding her Master’s Studies through the programme “IKY Scholarships”.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. Funded by SCOAP³.

Appendix A

Here, we derive the equation of motion for the gravitino from (2.31):

$$\frac{1}{2}\epsilon^{abcd}\sigma_b\tilde{\mathcal{D}}_c\bar{\psi}_d = -\bar{W}_0\sigma^{ab}\psi_b. \quad (A.1)$$

Contracting (A.1) with $\tilde{\mathcal{D}}_a$, we obtain the following equation:

$$\sigma^{ab}\tilde{\mathcal{D}}_a\psi_b = 0. \quad (A.2)$$

Moreover, contracting the hermitian conjugate

$$\frac{1}{2}\epsilon^{abcd}\bar{\sigma}_b\tilde{\mathcal{D}}_c\psi_d = W_0\bar{\sigma}^{ab}\bar{\psi}_b \quad (A.3)$$

of (A.1) with σ_a , we have

$$\epsilon^{abcd}\sigma_a\bar{\sigma}_b\tilde{\mathcal{D}}_c\psi_d \sim \epsilon^{abcd}\sigma_{ab}\tilde{\mathcal{D}}_c\psi_d \sim \sigma^{cd}\tilde{\mathcal{D}}_c\psi_d = 0, \quad (A.4)$$

where we have used (A.2) and

$$\epsilon^{abcd}\sigma_{ab} = -2i\sigma^{cd}. \quad (A.5)$$

Consequently,

$$\sigma_a\bar{\sigma}^{ab}\bar{\psi}_b = 0 \Rightarrow \sigma^a\bar{\psi}_a = 0, \quad (A.6)$$

where we have used the identity

$$\sigma^a\bar{\sigma}^b\sigma^c - \sigma^c\bar{\sigma}^b\sigma^a = 2i\epsilon^{abcd}\sigma_d. \quad (A.7)$$

Now let us consider B_α of Eq. (2.29). Its last term $i\psi_a b^a$ vanishes due to the equation of motion for b^a , while its second term vanishes due to Eq. (A.6). B_α ’s first term is

$$\begin{aligned} \sigma^a\bar{\sigma}^b\psi_{ab} &= \sigma^a\bar{\sigma}^b(\tilde{\mathcal{D}}_a\psi_b - \tilde{\mathcal{D}}_b\psi_a) = (\sigma^a\bar{\sigma}^b - \sigma^b\bar{\sigma}^a)\tilde{\mathcal{D}}_a\psi_b \\ &= 4\sigma^{ab}\tilde{\mathcal{D}}_a\psi_b = 0, \end{aligned} \quad (A.8)$$

where we have used the definition

$$\sigma^{ab} \equiv \frac{1}{4}(\sigma^a\bar{\sigma}^b - \sigma^b\bar{\sigma}^a) \quad (A.9)$$

and the relation (A.2). Consequently $B_\alpha = 0$ on-shell, which justifies the solution $M = -\lambda$ and $b^a = 0$ we chose in Sect. 2.2.

Appendix B

We will now demonstrate why the Lagrangian

$$\tilde{\mathcal{L}} = \left[\left(-\frac{1}{2}\frac{\mathcal{R}}{S_0} + W_0 + \frac{1}{2}\rho \left(\frac{\mathcal{R}}{S_0} - \lambda \right)^2 \right) S_0^3 \right]_F + \text{h.c.} \quad (B.1)$$

does not reproduce (2.10) with the constraint (2.11) in the limit $\rho \rightarrow \infty$. We first set

$$\begin{aligned} a &= W_0 + \frac{1}{2}\rho\lambda^2, \\ b &= 1 + 2\rho\lambda, \end{aligned} \quad (B.2)$$

assuming again reality of all parameters for simplicity. We then introduce a chiral superfield

$$S = A + \sqrt{2}\Theta\chi + (\Theta\Theta)F$$

(A and F are not the same as in the previous sections), such that

$$\begin{aligned} \bar{\mathcal{L}} &= \left[\left(a - \frac{1}{2} b \frac{\mathcal{R}}{S_0} + \frac{1}{2} \rho \frac{\mathcal{R}^2}{S_0^2} \right) S_0^3 \right]_F + \text{h.c.} \\ &= \left[\left(a - \frac{1}{2} b \frac{\mathcal{R}}{S_0} + \frac{S}{S_0} \frac{\mathcal{R}}{S_0} - \frac{1}{2\rho} \frac{S^2}{S_0^2} \right) S_0^3 \right]_F + \text{h.c.} \end{aligned} \tag{B.3}$$

It follows that $b > 0$ in order to have canonical gravity for a metric tensor with signature $(-+++)$. It is obvious from (B.3) that we have linearized our initial theory (B.1), which now describes the coupling of supergravity to a chiral superfield S that satisfies the equation of motion

$$S = \rho \mathcal{R}. \tag{B.4}$$

Next, using the identity (2.8) and fixing the gauge at $S_0 = 1$, we have

$$\bar{\mathcal{L}} = -[b - S - \bar{S}]_D + \left(\left[a - \frac{1}{2\rho} S^2 \right]_F + \text{h.c.} \right), \tag{B.5}$$

and the corresponding Kähler potential and superpotential are

$$K = -3 \ln(b - A - \bar{A}), \quad W = a - \frac{1}{2\rho} A^2. \tag{B.6}$$

The scalar potential V is given by

$$V = e^K \left[g^{A\bar{A}} (D_A W) (\bar{D}_{\bar{A}} \bar{W}) - 3 \bar{W} W \right]. \tag{B.7}$$

Note that positivity of the kinetic terms implies that $b - 2A_R > 0$, where we have set $A = A_R + iA_I$. We now compute

$$\begin{aligned} e^K &= \frac{1}{(b - A - \bar{A})^3}, \quad g_{A\bar{A}} = \frac{\partial}{\partial A} \frac{\partial}{\partial \bar{A}} K \\ &= \frac{3}{(b - A - \bar{A})^2}, \quad g^{A\bar{A}} = \frac{(b - A - \bar{A})^2}{3}, \end{aligned} \tag{B.8}$$

and

$$D_A W = \partial_A W + K_A W = -\frac{A}{\rho} + \frac{3}{b - A - \bar{A}} \left(a - \frac{1}{2\rho} A^2 \right). \tag{B.9}$$

Putting everything together, we get

$$\begin{aligned} V &= \frac{A\bar{A}}{3\rho^2(b - A - \bar{A})} - \frac{A}{\rho(b - A - \bar{A})^2} \left(a - \frac{1}{2\rho} \bar{A}^2 \right) \\ &\quad - \frac{\bar{A}}{\rho(b - A - \bar{A})^2} \left(a - \frac{1}{2\rho} A^2 \right) \\ &= \frac{1}{\rho^2(b - 2A_R)^2} \left\{ \frac{1}{3} (A_R^2 + A_I^2) (b + A_R) - 2a\rho A_R \right\}. \end{aligned} \tag{B.10}$$

The range of A_R is given by

$$-b \leq A_R < \frac{b}{2}, \quad b > 0, \tag{B.11}$$

so that the scalar potential is bounded from below.

To find the minimum of the potential, we demand that

$$\left\langle \frac{\partial V}{\partial A_R} \right\rangle = \left\langle \frac{\partial V}{\partial A_I} \right\rangle = 0. \tag{B.12}$$

The second of the equations above gives

$$\langle A_I (b + A_R) \rangle = 0. \tag{B.13}$$

If $\langle A_R \rangle = -b$, then

$$\left\langle \frac{\partial V}{\partial A_R} \right\rangle = 0 \Rightarrow \langle A_I^2 \rangle = -2a\rho - b^2 \xrightarrow{\rho \rightarrow \infty} -\infty, \tag{B.14}$$

so this case is rejected. Consequently $\langle A_I \rangle = 0$. Then

$$\left\langle \frac{\partial V}{\partial A_R} \right\rangle = 0 \Rightarrow \langle (b + 2A_R)(A_R^2 - 2bA_R + 6a\rho) \rangle = 0, \tag{B.15}$$

which yields three solutions whose compatibility with the condition (B.11) is given in Table 1. Only the solutions $\langle A_R \rangle = -\frac{b}{2}$ and $\langle A_R \rangle = b - \sqrt{b^2 - 6a\rho}$ are compatible with the range of A_R . Now we would like to check whether one of them is compatible with the condition

$$\langle V \rangle = 0. \tag{B.16}$$

Equation (B.16) has two solutions whose compatibility with the condition (B.11) is given in Table 2.

Table 1 Possible values of $\langle A_R \rangle$ for $\langle A_I \rangle = 0$

$\langle A_R \rangle$	$-b \leq A_R < \frac{b}{2}, b > 0$
$-\frac{b}{2}$	True always
$b + \sqrt{b^2 - 6a\rho}, b^2 - 6a\rho \geq 0$	Never true
$b - \sqrt{b^2 - 6a\rho}, b^2 - 6a\rho \geq 0$	True if $b^2 > 8a\rho$ and $b^2 \geq -2a\rho$

Table 2 Possible values of $\langle A_R \rangle$ for $\langle V \rangle = 0$ and $\langle A_I \rangle = 0$

$\langle A_R \rangle$	$-b \leq A_R < \frac{b}{2}, b > 0$
0	True always
$-\frac{b}{2} + \frac{\sqrt{b^2 + 24a\rho}}{2}, b^2 + 24a\rho \geq 0$	True if $b^2 > 8a\rho$
$-\frac{b}{2} - \frac{\sqrt{b^2 + 24a\rho}}{2}, b^2 + 24a\rho \geq 0$	True if $a\rho \leq 0$

Table 3 The (canonically normalized) scalar squared-mass matrix for $\langle A_R \rangle = -\frac{b}{2}$, $\langle A_I \rangle = 0$ and $\langle V \rangle = 0$

$g^{A\bar{A}} \frac{\partial^2 V}{\partial \phi_i \partial \phi_j}$	A_R	A_I
A_R	$-\frac{1}{\rho^2} \frac{b}{9}$	0
A_I	0	$\frac{1}{\rho^2} \frac{b}{9}$

It is straightforward to see that the solution $\langle A_R \rangle = b - \sqrt{b^2 - 6a\rho}$ is compatible with (B.16) only if $b^2 = 8a\rho$ (for $\langle A_R \rangle \neq 0$) or if $a = 0$ (for $\langle A_R \rangle = 0$). The first case is rejected, since then $\langle A_R \rangle = b/2$ and the metric $g_{A\bar{A}}$ diverges. The second case is rejected, because then $\langle D_A W \rangle = 0$ and there is thus no spontaneous supersymmetry breaking. On the other hand, the solution $\langle A_R \rangle = -\frac{b}{2}$ is compatible with (B.16) for $b^2 + 24a\rho = 0$. It can also lead to spontaneous supersymmetry breaking, as

$$\left\langle e^{K/2} \sqrt{g^{A\bar{A}}} \bar{D}_{\bar{A}} \bar{W} \right\rangle \sim ab^{-3/2} \neq 0 \quad \text{for finite } \rho. \quad (\text{B.17})$$

However, it is easy to see that the scalar squared-masses corresponding to A_R and A_I have opposite signs and thus the point ($\langle A_R \rangle = -\frac{b}{2}$, $\langle A_I \rangle = 0$) is a saddle point of the potential and not a minimum; see Table 3. Moreover, all the eigenvalues of the scalar mass matrix approach 0 as $|\rho| \rightarrow \infty$ and thus the extra scalar (sgoldstino) does not decouple. We conclude that neither of the two solutions for $\langle A_R \rangle$ can be used to tune the cosmological constant to zero for every value of ρ , consistently with the decoupling of the extra scalar.

Instead, we can investigate what happens if the condition (B.16) holds for the potential only in the limit $\rho \rightarrow \infty$. For both possible solutions

$$\begin{aligned} \langle A_I \rangle = 0, \quad \langle A_R \rangle = b - \sqrt{b^2 - 6a\rho} \approx \rho\lambda - 1 + 3\frac{W_0}{\lambda}, \quad \lambda \neq 0, \\ \langle A_I \rangle = 0, \quad \langle A_R \rangle = -\frac{b}{2} = -\frac{1}{2} - \rho\lambda, \end{aligned} \quad (\text{B.18})$$

we find that $V \rightarrow 0$ for $\rho \rightarrow \infty$; however, none of the eigenvalues of the scalar mass matrix approaches ∞ at $\rho \rightarrow \infty$ (they approach 0 instead), which is again incompatible with the sgoldstino decoupling. We conclude that the parameter space of the model (B.1) does not allow for the realization of the non-linear supersymmetry coupled to gravity. Thus, (B.1) has to be modified suitably, which is what we proposed in Sect. 3.

References

1. I. Antoniadis, E. Dudas, D.M. Ghilencea, P. Tziveloglou, Non-linear MSSM. Nucl. Phys. B **841**, 157 (2010). [arXiv:1006.1662](#) [hep-ph]
2. I. Antoniadis, E. Dudas, S. Ferrara, A. Sagnotti, The Volkov–Akulov–Starobinsky supergravity. Phys. Lett. B **733**, 32 (2014). [arXiv:1403.3269](#) [hep-th]
3. S. Ferrara, R. Kallosh, A. Linde, Cosmology with nilpotent superfields. JHEP **1410**, 143 (2014). [arXiv:1408.4096](#) [hep-th]
4. E.A. Bergshoeff, D.Z. Freedman, R. Kallosh, A. Van Proeyen, *Pure de Sitter Supergravity*. [arXiv:1507.08264](#) [hep-th]
5. I. Antoniadis, E. Dudas, A. Sagnotti, Brane supersymmetry breaking. Phys. Lett. B **464**, 38 (1999). [arXiv:hep-th/990802](#)
6. I. Antoniadis, M. Tuckmantel, Nonlinear supersymmetry and intersecting D-branes. Nucl. Phys. B **697**, 3 (2004). [arXiv:hep-th/0406010](#)
7. R. Kallosh, F. Quevedo, A.M. Uranga, *String Theory Realizations of the Nilpotent Goldstino*. [arXiv:1507.07556](#) [hep-th]
8. M. Rocek, Linearizing the Volkov–Akulov model. Phys. Rev. Lett. **41**, 451 (1978)
9. U. Lindstrom, M. Rocek, Constrained local superfields. Phys. Rev. D **19**, 2300 (1979)
10. R. Casalbuoni, S. De Curtis, D. Dominici, F. Feruglio, R. Gatto, Nonlinear realization of supersymmetry algebra from supersymmetric constraint. Phys. Lett. B **220**, 569 (1989)
11. Z. Komargodski, N. Seiberg, From linear SUSY to constrained superfields. JHEP **0909**, 066 (2009). [arXiv:0907.2441](#) [hep-th]
12. S.M. Kuzenko, S.J. Tyler, On the Goldstino actions and their symmetries. JHEP **1105**, 055 (2011). [arXiv:1102.3043](#) [hep-th]
13. D.V. Volkov, V.P. Akulov, Is the neutrino a Goldstone particle? Phys. Lett. B **46**, 109 (1973)
14. I. Antoniadis, E. Dudas, D.M. Ghilencea, Goldstino and sgoldstino in microscopic models and the constrained superfields formalism. Nucl. Phys. B **857**, 65 (2012). [arXiv:1110.5939](#) [hep-th]
15. C. Markou, *The coupling of Nonlinear Supersymmetry to Supergravity*. Master’s Thesis, Ecole Polytechnique Paris and ETH Zurich (2015)
16. E. Dudas, S. Ferrara, A. Kehagias, A. Sagnotti, *Properties of Nilpotent Supergravity*. [arXiv:1507.07842](#) [hep-th]
17. S.J. Gates Jr, S.V. Ketov, Superstring-inspired supergravity as the universal source of inflation and quintessence. Phys. Lett. B **674**, 59 (2009). [arXiv:0901.2467](#) [hep-th]
18. S.V. Ketov, Scalar potential in $F(R)$ supergravity. Class. Quantum Gravity **26**, 135006 (2009). [arXiv:0903.0251](#) [hep-th]
19. R. Kallosh, L. Kofman, A.D. Linde, A. Van Proeyen, Superconformal symmetry, supergravity and cosmology. Class. Quantum Gravity **17**, 4269 (2000) (Class. Quantum Gravity **21**, 5017, 2004). [arXiv:hep-th/0006179](#)
20. D.Z. Freedman, A. Van Proeyen, *Supergravity* (Cambridge University Press, Cambridge, 2012)
21. S. Cecotti, Higher derivative supergravity is equivalent to standard supergravity coupled to matter. I. Phys. Lett. B **190**, 86 (1987)
22. S. Ferrara, R. Kallosh, A. Van Proeyen, On the supersymmetric completion of $R + R^2$ gravity and cosmology. JHEP **1311**, 134 (2013). [arXiv:1309.4052](#) [hep-th]
23. J. Wess, J. Bagger, *Supersymmetry and Supergravity* (Princeton University Press, Princeton, 1992)