



HAL
open science

The internal consistency of the North Sea carbonate system

Lesley A. Salt, Helmuth Thomas, Yann Bozec, Alberto V. Borges, Hein J.W. de Baar

► **To cite this version:**

Lesley A. Salt, Helmuth Thomas, Yann Bozec, Alberto V. Borges, Hein J.W. de Baar. The internal consistency of the North Sea carbonate system. *Journal of Marine Systems*, 2016, 157, pp.52-64. 10.1016/j.jmarsys.2015.11.008 . hal-01246273

HAL Id: hal-01246273

<https://hal.sorbonne-universite.fr/hal-01246273v1>

Submitted on 18 Dec 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The internal consistency of the North Sea carbonate system.

Lesley A. Salt^{a,1*}, Helmuth Thomas^b, Yann Bozec^c, Alberto V. Borges^d, and Hein J. W. de Baar^a

^aRoyal Netherlands Institute for Sea Research, Landsdiep 4, 1797 SZ, Texel, Netherlands. Hein.de.Baar@nioz.nl

^bDepartment of Oceanography, Dalhousie University, Halifax, B3H 4J1, Nova Scotia, Canada. Email: helmuth.thomas@dal.ca

^cSorbonne Universités, UPMC Univ Paris 06, CNRS, Adaptation et Diversité en Milieu Marin (AD2M UMR7144), Station Biologique de Roscoff, 29680 Roscoff, France. Email: bozec@sb-roscoff.fr

^dUnité d'Océanographie Chimique, Institut et Physique B5, Université de Liège, 4000 Liège, Belgium. Email: alberto.borges@ulg.ac.be

¹Sorbonne Universités, UPMC Univ Paris 06, CNRS, Adaptation et Diversité en Milieu Marin (AD2M UMR7144), Station Biologique de Roscoff, 29680 Roscoff, France

*Corresponding author: Email: lesley.salt@sb-roscoff.fr. Tel: +33 (0)2 98 29 28 17.

Abstract

In 2002 (February) and 2005 (August) the full suite of carbonate system parameters (Total Alkalinity (A_T), Dissolved Inorganic Carbon (DIC), pH, and partial pressure of CO_2 (pCO_2)) were measured on two re-occupations of the entire North Sea basin, with three parameters (A_T , DIC, pCO_2) measured on four additional re-occupations, covering all four seasons, allowing an assessment of the internal consistency of the carbonate system. For most of the year there is a similar level of internal consistency, with A_T being calculated to within $\pm 6 \mu mol kg^{-1}$ using DIC and pH, DIC to $\pm 6 \mu mol kg^{-1}$ using A_T and pH, pH to ± 0.008 using A_T and pCO_2 , and pCO_2 to $\pm 8 \mu atm$ using

DIC and pH, with the dissociation constants of Millero et al. (2006). In spring, however, we observe a significant decline in the ability to accurately calculate the carbonate system. Lower consistency is observed with an increasing fraction of Baltic Sea water, caused by the high contribution of organic alkalinity in this water mass, not accounted for in the carbonate system calculations. Attempts to improve the internal consistency by accounting for the unconventional salinity-borate relationships in freshwater and the Baltic Sea, and through application of the new North Atlantic salinity-boron relationship (Lee et al., 2010), resulted in no significant difference in the internal consistency.

Keywords: *carbon dioxide, North Sea, internal consistency, total alkalinity*

1 Introduction

Increasing atmospheric CO₂ has been partly mitigated by oceanic uptake, which currently accounts for 25-30% of the total CO₂ emissions (Sabine et al., 2004; Canadell et al., 2007; Khatiwala et al., 2009; Le Quéré et al., 2015). The uptake of CO₂ in surface waters causes a shift in the chemistry of seawater, notably increases in [H⁺] and a decrease in [CO₃²⁻] leading to the phenomenon known as ocean acidification (OA) (Caldeira and Wickett, 2003, 2005; Raven, 2005). Recent findings of Wootton et al. (2008), Provost et al. (2010) and Borges and Gypens (2010) indicate that ocean acidification is proceeding in coastal areas more rapidly than has been predicted. The associated changes related to OA have been shown to have a variety of effects on marine phytoplankton (Fabry et al., 2008). The coastal zone is responsible for a disproportionately large amount of primary production per surface area as compared to the open ocean (Wollast, 1998) and is responsible for a similarly large proportion of carbon export (Dunne et al., 2007). As such, understanding how these waters respond to acidification is vital for accurately predicting the future consequences of continued CO₂ increase.

The North Sea has been identified as an effective continental shelf-pump of CO₂ (Thomas et al., 2004) and has thus undergone intensive study with respect to the CO₂

system. High temporally resolved datasets are necessary to understand the large, observed seasonal variability of the carbonate system in the North Sea (Frankignoulle and Borges, 2001; Bozec et al., 2006; Prowe et al., 2009). The use of Voluntary Observing Ships (VOS) to collect underway pCO₂ data have been used extensively in the North Sea (Omar et al., 2010), complemented by measurements of one other of the measurable carbonate variables (total alkalinity (A_T), dissolved inorganic carbon (DIC), and pH). From at least two of the measured variables (a combination pair of A_T, DIC, pH or pCO₂), the carbonate system can then be fully solved and quantified using thermodynamic relationships (Park, 1969), which can be integrated in computer programs, such as CO2SYS (Lewis and Wallace, 1998). These calculated parameters, however, are prone to errors, which have been well documented (Millero et al., 1993; Lee et al., 2000; Koeve et al., 2012; Hoppe et al., 2012). These errors are due to the accuracy of the two measured variables, and to the reliability of the first and second dissociation constants of carbonic acid (K₁, K₂). The assessment of the reliability of dissociations constants can be achieved with internal consistency studies.

The first internal consistency studies took place in the open ocean and focused on the reliability of the various sets of carbonic acid dissociation constants, which are used to calculate the distribution of inorganic carbon species. Acid dissociation constants describe the interaction between acids (HA), their dissociated proton (H⁺) and the conjugate base (A⁻), shown in equation 1.

$$K = \{H^+\} \{A^-\} / \{HA\}, \quad (\text{equation 1})$$


where {} denote activities. The aforementioned determinations of K₁ and K₂ of carbonic acid were performed analytically on artificial seawater (ASW) (Roy et al., 1993; Goyet and Poisson, 1989; Hansson, 1973) and natural seawater (NSW) (Mehrbach et al., 1973) and the data were later refitted using improved computer-generated models (Dickson and Millero, 1987). These constants are technically *apparent* dissociation constants (denoted by K'), as they are based on concentrations (indicated by []), not activities, so that

$$K' = \{H^+\} [A^-] / [HA], \quad (\text{equation 2})$$

This approach also means that the K' values must be determined across the range of

salinities, temperatures and pressures at which they are to be applied and that the measurements take place in the same medium as the solution to which the K' value is to be applied. Assessments and comparisons of each set of constants have taken place in several locations, including the northwest European shelf (Ribas-Ribas et al., 2014), equatorial Pacific (Millero et al., 1993) and the North Atlantic (Lee et al., 1997), and spanning across several ocean basins (Wanninkhof et al., 1999; Millero et al., 2002). At sea studies have been complemented by further work using laboratory set-ups (Lee et al., 1996; Lueker et al., 2000) and more recently investigations into inconsistencies in culture experiments (Hoppe et al., 2012; Koeve et al., 2012).

The conclusions from these studies is that the first and second dissociation constants of carbonic acid (K'_1 and K'_2) calculated by Mehrbach et al. (1973), (refit by Dickson and Millero, 1987), are the most reliable when $p\text{CO}_2$ is being used in the calculations or is being calculated (Wanninkhof et al., 1999; Lee et al., 1996). These constants were the only ones determined in NSW and the difference in accuracy has been attributed to the lack of boric acid in the ASW (Mojica Prieto and Millero, 2002). In solution the majority of dissolved boron (BT = total boron) is present in two forms; boric acid and borate, following the equilibrium relationship below:



where $K_B = [\text{B(OH)}_4^-][\text{H}^+]/[\text{B(OH)}_3]$ (equation 4)

In NSW boric acid interacts with HCO_3^- and CO_3^{2-} to lower the activities, and concentrations of HCO_3^- , and increase the activities and concentrations of CO_3^{2-} , thus lowering pK_1 and increasing pK_2 . Calculations of $p\text{CO}_2$ depend directly on the ratio of K_2/K_1 , thus errors in this ratio will be evident in a lack of internal consistency of $p\text{CO}_2$. In contrast, the lack of use of this ratio in calculations with pH, A_T or DIC results in other dissociation constants being more appropriate, notably those of Roy et al., (1993) or Goyet and Poisson (1989) (Lee et al., 1997; Clayton et al. 1995). Despite this, in an attempt to homogenize the way carbon chemists carry out these calculations, it is now the recommended standard procedure to use the constants of Mehrbach et al. (1973), refit by Dickson and Millero (1987), on the Total pH scale (Dickson et al., 2007).

Beyond the scope of dissociation constants, additional errors have been associated with systematical differences in equipment (Körtzinger et al., 2000; Ribas-Ribas et al., 2014), high DIC/A_T ratios (>1.10)(Lee et al., 1996), the interference of fatty acids (Gripenberg, 1960), and contribution from organic acids and bases to A_T (Hernandez-Ayon et al., 2007; Koeve et al., 2012; Ulfsbo, et al., 2015). The latter errors are only thought to be a problem where dissolved organic carbon (DOC) reaches particularly high concentrations (Koeve et al., 2012), thus this is not an issue for open ocean measurements, although presents a major problem for calculations in freshwaters (Abril et al., 2015). Similarly, the contributions of fatty acids and high DIC/A_T ratios in the open ocean are sufficiently low that internal consistency remains within the same order of magnitude as the measurement precision. In culture experiments, the differences between measured and calculated carbonate parameters have been much larger, and thus coastal areas where high primary production takes place, could also be susceptible to large discrepancies in calculations of the carbonate system. Furthermore, in the original work of the recommended constants of Mehrbach et al., (1973), the constants were only determined to salinities above 26. The salinity range was extended in further work by Millero et al. (2006), however, the results have not yet been applied to further internal consistency studies. Coastal regions can also exhibit non-oceanic-like, chemical relationships with salinity, for example, the atypical borate-salinity relationship in the Baltic Sea (Kremling, 1972), which is not taken into account when performing carbonate system calculations and can lead to further error. Here we present the first internal consistency study performed in coastal waters with seasonal resolution over a range of salinities and biogeochemical regimes.

2 Methods

2.1 Hydrography

The hydrographic properties in the surface waters of the North Sea demonstrate the presence of 3 different water masses, acknowledged in the literature as North Atlantic water, Baltic water, and German Bight water, with the resulting composite water mass known as Central North Sea Water (Kempe and Pegler, 1991; Bozec et al., 2006; Salt

et al., 2013). In the northern North Sea (nNS), North Atlantic water flows over the Shetland Shelf bringing characteristic warm and high saline waters into the North Sea. The inflowing water follows an anti-clockwise trajectory around the North Sea, with additional North Atlantic inflow occurring through the English Channel (Otto et al., 1990). In the southeast the Skagerrak water mass, a mixture of Central North Sea water and Baltic water, is easily identifiable through much lower salinities and temperatures than the central North Sea. The German Bight water comes from the coastal region bordering northwest continental Europe in the southeastern North Sea, where the riverine influence is greatest in the North Sea.

The North Sea can be divided into two biogeochemical regimes, within which behavior is determined by depth (Thomas et al., 2005; Prowe et al., 2009; Omar et al., 2010). The nNS (north of 56°N) is deeper than the south (50-400 m) and experiences more ocean-like behavior, e.g. higher salinity and seasonal stratification. The southern North Sea (sNS) is much shallower (<50 m) and remains well-mixed throughout the year. Both regions receive nutrient and organic carbon inputs from terrestrial sources, facilitating high primary production that leads to overall annual autotrophy in the nNS and near balanced trophic status in the sNS (Bozec et al., 2005; Schiettecatte et al. 2007).

2.2 Measurements and comparisons

The data used in this study comes from 6 separate re-occupations of the North Sea, which took place in August 2001, November 2001, February 2002, May 2002, August 2005 and August 2008. All four carbonate parameters were measured on two of these cruises; February 2002, and August 2005, with a precision of $\pm 2-3 \mu\text{mol A}_T \text{ kg}^{-1}$ for A_T , $\pm 1-2 \mu\text{mol DIC kg}^{-1}$ for DIC, ± 0.001 for pH, and $\pm 1 \mu\text{atm}$ for pCO_2 , respectively. The remaining cruises measured A_T , DIC, and pCO_2 with the same level of precision.

2.2.1 Carbonate system measurements

For all of the aforementioned cruises, A_T was measured by potentiometric titration to the second end-point (pK_2) of carbonic acid, using 0.1M hydrochloric acid as the

titrant. For the cruises taking place in August 2001, November 2001, February 2002, May 2002 and August 2005 the samples for A_T were filtered on GF/F shortly after sampling, prior to analysis. The resultant acid volumes and corresponding e.m.f values are fitted to a Gran titration plot using least squares (Gran, 1952). An uncertainty of $\pm 2-3 \mu\text{mol } A_T \text{ kg}^{-1}$ was found as the combined precision for all cruises. Samples for DIC were not filtered in order to prevent gas exchange with the atmosphere. Any sediments or particles collected in the samples would likely settle to the bottom of the sample bottle before analysis and thus interference in the measurements should be minimal. All of the DIC data collected in the North Sea was analyzed by the coulometric method of Johnson et al. (1993), with an overall uncertainty of $\pm 1-2 \mu\text{mol DIC kg}^{-1}$ for all cruises. For quality control and calibration of both DIC and A_T , Certified Reference Material (CRM), of known DIC and A_T , acquired from Prof. Andrew Dickson (Scripps Institute of Oceanography, USA) were used.

In 2002 and 2005, discrete samples were taken from the CTD and analyzed for pH potentiometrically on the total hydrogen ion scale, calibrated using TRIS (2-amino-2-hydroxymethyl-1,3-propanediol) and AMP (2-aminopyridine) buffers prepared at a salinity of 35 according to Dickson (1993). The accuracy of this method is strongly dependent on the sensitivity of the pH meter and how accurately the two buffer solutions are made up. For the measurements used in this work a precision of ± 0.001 pH units was attained. For all North Sea cruises $p\text{CO}_2$ was measured continuously, underway, from surface waters, providing the greatest coverage of data. The instrument used continuous equilibration with a headspace and infrared detection with an accuracy of $\pm 1 \mu\text{atm}$ (Körtzinger et al., 1996). The calibration of $p\text{CO}_2$ measurements was carried out regularly (every 2-3 hours) using National Oceanic and Atmospheric Administration (NOAA, Global Monitoring Division, Carbon Cycle Greenhouse Gases Group) calibrated gas standards. For all cruises a minimum of 3 calibration gases were used, spanning a concentration range from 0 ppm to 750 ppm.

The concentrations of silicate and phosphate are also required to account for their

contribution to A_T , which can be of the order of up to $1 \mu\text{mol } A_T \text{ kg}^{-1}$ in the North Sea. The inorganic nutrients nitrate (NO_3), ammonium (NH_4), dissolved silicate (H_4SiO_4), and phosphate (PO_4), were analyzed following the methods of Grasshof et al. (1983). Reference nutrient samples (RMNS Kanso, Japan) were analyzed in every run as a control, giving a precision of ± 0.2 , 0.1 , 0.2 and 0.01 for the four nutrients, respectively, for each cruise. The salinity and temperature values were recorded by CTD (Conductivity-Temperature-Depth) and salinity values were calibrated post-cruise to salinity samples taken onboard.

2.3 Calculations

The internal consistency was assessed using the measured in-situ salinity, temperature, silicate, phosphate and pressure values. The KSO_4 of Dickson (1990) is used and pH values are given throughout on the total hydrogen ion scale. Not all station positions were occupied and measured in all four seasonal cruises that are compared here. In order to overcome this, the percentage of data points considered internally consistent is often used, and given in tables 3 and 4, to allow direct comparison between the seasons without the loss of data. The calculations were performed using the CO2SYS program (Lewis and Wallace, 1998), adapted for MATLAB (van Heuven, 2011; http://cdiac.ornl.gov/ftp/co2sys/CO2SYS_calc_MATLAB_v1.1/). All calculated parameters as a function of two input parameters were explored, thus consisting of: $A_T(\text{DIC}, \text{pH})$, $A_T(\text{DIC}, \text{pCO}_2)$, $A_T(\text{pH}, \text{pCO}_2)$, $\text{DIC}(A_T, \text{pH})$, $\text{DIC}(A_T, \text{pCO}_2)$, $\text{DIC}(\text{pH}, \text{pCO}_2)$, $\text{pH}(\text{DIC}, A_T)$, $\text{pH}(\text{DIC}, \text{pCO}_2)$, $\text{pH}(A_T, \text{pCO}_2)$, $\text{pCO}_2(\text{DIC}, A_T)$, $\text{pCO}_2(\text{DIC}, \text{pH})$, and $\text{pCO}_2(A_T, \text{pH})$, for the following seven different sets of carbonic acid dissociation constants: Roy et al. (1993)(R), Goyet & Poisson (1989)(GP), Hansson (1973), refit by Dickson and Millero (1987)(H), Mehrbach et al. (1973), refit by Dickson and Millero (1987)(M), Hansson & Mehrbach (1973), refit by Dickson and Millero (1987)(HM), Mojico-Prieto & Millero (2002)(MM) and Millero et al. (2006)(ML). In the remaining text, for simplicity, the constants will be referred to by the abbreviations (R, GP, H, M, HM, MM and ML). The parameterizations of these constants, along with the medium and salinity ranges for which they were determined, are shown in Table 1.

2.4 Assessment

The internal consistency of the carbonate system can be considered in a number of different ways. Here, we assess how accurate and precise it is by comparing measured parameters to calculated parameters. Comparing the measured and calculated values, an accurate but imprecise system would be indicated by an average difference (Δ) close to 0, but with a high level of variance, denoted by the standard deviation (σ). A precise but inaccurate system, however, is indicated by a small σ , but with a bias, indicated by an average Δ varying from 0. Thus, an inaccurate and imprecise system would have a high standard deviation and a large average Δ , whilst an accurate and precise system would show the inverse. The measured and calculated values of each parameter, for each set of apparent dissociation constants, was fitted to the form $Y = a.X + b$, by means of *total least squares* using Matlab 2012®, where X is the calculated value and Y is the observed value. By fitting the data using total least squares, we take into account the errors in both the x and y variables.

2.5 Error Propagation

When calculating the carbonate parameters in CO2SYS, the uncertainties associated with the observations and calculations are carried through (propagated). Errors are introduced via the measurement techniques of the input parameters, the measurement techniques of the output parameters against which we compare our calculations, and the determinations of the K_1 and K_2 values of carbonic acid. The measurement errors carried through in the calculations depend on the input parameter combinations. These margins vary insignificantly over the range of conditions (e.g. temperature, salinity) observed in the North Sea. The error propagation for each calculated parameter, with all variations of input parameter, was calculated following equation 5.

$$(\epsilon z)^2 = \{\delta z / \delta x\}^2 \cdot (\epsilon x)^2 + \{\delta z / \delta y\}^2 \cdot (\epsilon y)^2 \quad (\text{equation 5})$$

The partial derivatives were determined seasonally, using average temperature and salinity values, and determined over the range of values observed. The determined errors for each set of input parameters for each season are given in brackets in the

standard deviation section of table 2a and 2b. The error calculation performed here does not take into account errors in the acid dissociation constants, as we later compare the performance of these constants. When the term ‘internally consistent’ is used, it refers to whether the observed values and associated measurement errors fit within the limits of the calculated propagated error.

2.6 Total Borate Calculations

To independently calculate the total borate contribution from each water mass, the individual water mass fractions were first calculated, following Friis et al. (2003). To calculate the Elbe contribution we used the salinity and A_T data from the stations closest to the German Bight (at latitudes $\geq 54.4^\circ\text{N}$ and longitudes $\geq 8.0^\circ\text{E}$) to form a regression line with the Central North Sea end member. We then extrapolated this to a salinity of 0 to get a representative freshwater end member. This assumes that all the fresh water present comes from the same source, however, due to other rivers in the region having a similar catchment lithology we assume that the chemical signal of the water will be similar. This resulted in a 0-salinity end member of $2616 \mu\text{mol } A_T \text{ kg}^{-1}$ in February 2002, and $1678 \mu\text{mol } A_T \text{ kg}^{-1}$ in August 2005. These values comparing well with recent seasonal A_T data from the Elbe (Amann et al., 2012), which reports A_T values of $\sim 2500 \mu\text{mol } A_T \text{ kg}^{-1}$ in winter and $\sim 1750 \mu\text{mol } A_T \text{ kg}^{-1}$ in summer at salinities < 1 , thus we assume our estimation is sufficient for the purpose of the calculation.

The borate alkalinity (BAlk) was then calculated for the water mass end members from total boron concentrations. The total boron concentrations (BT) were calculated from (a) the equations of Barth (1998) (equations 6 and 7) for the Elbe (German Bight) salinity-boron relationship, (b) Gripenberg (1960) (equation 8) for the Baltic salinity-boron ratio, (c) Uppström, (1974) for the North Atlantic ratio and also the revised North Atlantic ratio of Lee et al. (2010). At a salinity of 35 the two seasonal linear relationships of Barth (1998), for summer (September, 1994) and winter (February, 1995), yield an approximate difference of $6 \mu\text{mol kg}^{-1}$ in the calculated $[\text{B}(\text{OH})_4^-]$. As such, for our calculations, equation 6 was used for the calculations of

August 2005 (summer), and equation 7 for the calculation of February 2002 (winter). The water mass fractions were then multiplied by the theoretical BAlk contribution and summed to obtain the actual BAlk of the sample. All of the following equations (6 – 9) are given for total borate concentration (BT) in mol kg⁻¹ seawater.

$$BT = (0.000127/10.811) \cdot S + 1.66e^{-5} \text{ (for September, 1994) (equation 6)}$$

$$BT = (0.000136/10.811) \cdot S + 1.57e^{-5} \quad \text{(for February, 1995) (equation 7)}$$

The Baltic Sea has demonstrated a number of chemical relationships varying from standard ocean behaviour. The chlorinity-boron relationship has been documented by Gripenberg, (1960), Dyrssen and Uppström (1974), and Kremling, (1970), who all noted a positive anomaly in the boron/chlorinity ratio. There is little variation in their findings, with the ratio varying from 0.22 (Kremling, 1970) to 0.244 (Gripenberg, 1960). Here, we use the following equation from the data presented by Gripenberg (1960):

$$BT = (0.000244/10.811) \cdot (S/1.80655) \quad \text{(equation 8)}$$

For the North Atlantic fraction the traditional seawater relationship of Uppström (1974) was used:

$$BT = (0.000232/10.811) \cdot (S/1.80655) \quad \text{(equation 9)}$$

Furthermore, Lee et al. (2010) recently found boron-salinity relationships significantly different to those reported by Uppström (1974), which were based on measurements solely from the Pacific Ocean. Here, we also apply this new boron-salinity equation (equation 10), to test whether this improves our internal consistency in the North Sea.

$$BT = (0.0001336/10.811) \cdot S \quad \text{(equation 10)}$$

3 Results and Discussion

3.1 A seasonal comparison

The sampling program of the North Sea is shown in Figure 1, with the different distinct, biogeochemical regions labelled. The two seasons in which data was collected of all four measurable carbonate parameters were winter (February) and summer (August). These two data sets allow us to compare the internal consistency in two very different conditions, in terms of values of the carbonate variables, and water temperature. As the North Sea is a temperate coastal sea, it is characterized by high primary production during summer and very low, or no, production in winter. The seasonal cycle is fully assessed in the works of Bozec et al. (2006) and Thomas et al. (2009), thus here the discussion is kept brief. The average DIC in February 2002 was $64 \mu\text{mol DIC kg}^{-1}$ higher than that in August. The average A_T changed by just $2 \mu\text{mol } A_T \text{ kg}^{-1}$, however, the various contributions of chemical species to A_T showed larger variability. The changes in A_T contributions in the North Sea from February to August are shown in Figures 2a and 2b, respectively. For average conditions in the North Sea, the HCO_3^- and CO_3^{2-} species were by far the greatest contributors to A_T supplying $\sim 2212 \mu\text{mol } A_T \text{ kg}^{-1}$, followed by borate ($\sim 68 \mu\text{mol } A_T \text{ kg}^{-1}$), hydroxide ($\sim 3 \mu\text{mol } A_T \text{ kg}^{-1}$), phosphate ($\sim 0.07 \mu\text{mol } A_T \text{ kg}^{-1}$) and silicate ($\sim 0.04 \mu\text{mol } A_T \text{ kg}^{-1}$). The largest changes between the summer and winter situation were in carbonate alkalinity ($\text{HCO}_3^- + 2 \cdot \text{CO}_3^{2-}$), which was significantly lower in August, when photosynthesis was ongoing. The contribution of silicate and phosphate were also much smaller then, due to lower concentrations resulting from uptake by phytoplankton. The borate contribution remains the second largest contributor to A_T for both seasons, but was slightly higher in August. Isolating the various contributors to this change, we find that the increases in temperature, salinity and pH from February to August all affect an individual increase of $\sim 3 \mu\text{mol kg}^{-1}$ in B(OH)_4^- (total increase $\sim 9 \mu\text{mol kg}^{-1}$).

The change in average pCO_2 , from February 2002 to August 2005, was just $8 \mu\text{atm}$, however, this is not representative of the entire North Sea, as the pCO_2 range was two times larger in August ($287 - 455 \mu\text{atm}$) compared to February ($315 - 396 \mu\text{atm}$). The range reflected the two biogeochemical regimes within the North Sea: the stratified

nNS, where stratification enables prolonged phytoplankton growth causing a large drawdown of CO₂, shows very low pCO₂ values in August, whereas in the sNS, very low pCO₂ values are observed only in spring, and a combination of remineralization and higher temperatures lead to much higher pCO₂ in August (Bozec et al., 2005; Thomas et al., 2004, 2005; Schiettecatte et al. 2007). In February, the water column is well-mixed throughout, and biological activity is lower thus leading to a more homogenous surface distribution of pCO₂.

The results from all internal consistency calculations using the 4 carbonate parameters, with various dissociation constants, in the February 2002 and August 2005 data sets are shown in Tables 2a and 2b, respectively. A regression slope of >1 indicates that the measured value is larger than the calculated values. A regression slope of 1.000 is achieved with several combinations in both seasons. Overall, in February, when the three parameters involved are DIC, A_T and pCO₂, (e.g. DIC(A_T,pCO₂), A_T(DIC,pCO₂) or pCO₂(DIC,A_T)) we find that the constants of M are the most suitable, whereas when DIC, A_T and pH are involved the constants of GP outperform the others. The largest deviations are consistently observed when the input parameters are pH and pCO₂, with MM producing the most accurate and precise reproduction of measurements. We also observe that the observed standard deviations of differences are largely comparable with the propagation error, with the exception of pH, in which we observe notably larger discrepancies than can be accounted for from the propagation error. In August the constants of ML are the best choice when the parameter combination is A_T, DIC and pCO₂ and, additionally, when A_T, DIC and pH are all involved. However, for calculations using both pH and pCO₂ the constants of GP and H are the most accurate.

In February and August, the variability in the standard deviations between the dissociation constants (e.g. comparing $\sigma \Delta A_{T,R}$ with $\sigma \Delta A_{T,ML}$) were relatively minor compared to the differences obtained with various input parameter combinations (e.g. comparing $\sigma \Delta A_{T,(pH,pCO_2)}$ with $\sigma \Delta A_{T,(DIC,pH)}$ etc.). As such, beyond assessing the parameter combinations with the largest degree of noise, the standard deviations are

not further discussed in this section. The results for both February and August are consistent, showing that the best regression coefficients for A_T were attained with the inputs (DIC, pCO_2), for DIC it was (A_T , pCO_2) and for pCO_2 it was (DIC, A_T). For A_T and DIC, this is in line with smaller error propagations, as similarly found by Dickson and Riley (1974), however, the much improved result for calculating pCO_2 using (DIC, A_T) is indicative of the improvements in measurement accuracy obtained with these parameters with the use of CRMs. When calculating pH, a regression coefficient of 1.000 was attained with all combination parameters. Similarly, consistency was found in the parameter input combinations resulting in the greatest precision, with A_T being best represented by (DIC, pH), DIC by (A_T , pH), pH by (A_T , pCO_2) and pCO_2 by (DIC, pH). The combination pair resulting in the lowest residuals varied between the two seasons, possibly indicating a bias in some of the measurements. The prevalence of better results being obtained when using pH as an input parameter, indicate that these measurements are reliable, however, the difference in the resulting residuals from e.g. DIC(A_T , pH) and DIC(A_T , pCO_2), are still directly comparable.

The results from February show a slight overall decrease in precision, represented by a higher standard deviation, compared to that obtained in August, however, the average differences show a similar level of accuracy for DIC and pCO_2 and a mildly lower bias in February for A_T and pH. The standard deviations of the differences between the calculated and measured parameters (Δ) for February 2002 and August 2005 (shown in brackets, respectively) are approximately double to those found in the open ocean by Lee et al. (1997), of $\pm 3 \mu\text{mol } A_T \text{ kg}^{-1}$ in A_T ($\pm 7, 6 \mu\text{mol } \text{kg}^{-1}$), $\pm 3 \mu\text{mol } \text{DIC } \text{kg}^{-1}$ in DIC ($\pm 7, 6 \mu\text{mol } \text{kg}^{-1}$), and $\pm 1.3\%$ in pCO_2 ($\pm 10, 8 \mu\text{atm}$; $\pm 3\%, 2\%$).

3.1.1 The effects of salinity on dissociation constants

Different salinity and temperature parameterizations used in the determination of pK_1 and pK_2 (Table 1) are largely responsible for the deviations in the calculations between dissociation constants. Plotting the difference in ΔA_T between results from different dissociation constants (e.g. the differences between the residuals using the constants of R and the residuals using the constants of M: $\Delta A_T^R - \Delta A_T^M$; or ΔDIC , ΔpH

and $\Delta p\text{CO}_2$) shows an exponential-like increase, or decrease, around a salinity of ~ 34.5 (Fig. 3), indicating that at higher salinities the choice of dissociation constants becomes more significant. We compared the differences in residuals of all dissociation constants to those of M, and found the same relationship to that shown in Fig. 3 for the constants of R, GP, H and HM. The differences in ΔA_T between M and MM showed a gradual linear increase with salinity, with a slope of 0.3 ($R^2=0.9$, $p\text{-value}<0.01$), indicating that across the range of 5 salinity units, there is an increase of $1.5 \mu\text{mol } A_T \text{ kg}^{-1}$ difference in ΔA_T . The residuals resulting from the use of constants ML showed no discernible difference to those of M over the salinity range investigated. At lower salinities the ratio of K_1/K_2 is larger (Mosley et al. 2010; Zeebe and Wolf-Gladrow, 2001) leading to relatively greater HCO_3^- concentrations compared to CO_2 and CO_3^{2-} . This ratio is used directly in the calculation of pH and A_T from DIC and $p\text{CO}_2$. Due to the different parameterizations of K_1 and K_2 for the different constants, the difference in this ratio between constants may become larger or smaller at lower salinities. This will lead to larger or smaller differences between the measured and calculated parameters. As such, at lower salinities, one would expect, overall, lower variation between the calculations from the different dissociation constants, as is observed here.

To investigate the performance of the six dissociation constants across a range of different salinities, the North Sea data from August ('05) and February ('02) were split into two groups; low salinity waters, with salinities less than 34.5, and high salinity waters, with salinities greater than or equal to 34.5. The 1 standard deviation (σ) of ΔA_T , ΔDIC , ΔpH and $\Delta p\text{CO}_2$ is larger, for all parameters and in both seasons, for low salinity waters compared to high salinity waters. In February, in high salinity waters, the standard deviation of ΔA_T is $2 \mu\text{mol } A_T \text{ kg}^{-1}$ lower than that of the total dataset; ΔDIC is $1 \mu\text{mol DIC kg}^{-1}$ lower, ΔpH is 0.002 lower and $\Delta p\text{CO}_2$ is $2 \mu\text{atm}$ lower. In the less saline waters there is an increase in the standard deviation of ΔA_T , ΔDIC , ΔpH and $\Delta p\text{CO}_2$ by $2 \mu\text{mol } A_T \text{ kg}^{-1}$, $1 \mu\text{mol DIC kg}^{-1}$, 0.003, and $2 \mu\text{atm}$, respectively. The same pattern is observed in August, however the differences are marginally smaller.

Furthermore, in February, there is a clear dichotomy between the different dissociation constants in low and high salinity waters (Fig. 4). As previously mentioned, in winter, when DIC, A_T , and pH are involved in calculations (that is to say, the calculation of $\text{pH}(\text{DIC}, A_T)$ or $\text{DIC}(A_T, \text{pH})$ or $A_T(\text{DIC}, \text{pH})$), the constants of GP produce the least deviation between measured and calculated parameters. However, we find that in high salinity ($S \geq 34.5$) waters the constants of M become more appropriate and in the lower salinity waters ($S < 34.5$) the constants of R outperform the others. Despite these changes, the constants of MM and ML show little variation in performance between the two salinity environments. For August, there was no difference between the most suitable constants for the entire dataset compared to the different salinity regimes regarding measurements with DIC, A_T and pH, with the constants of GP still most closely representing the measurements. With regard to measurements of pH, pCO_2 and DIC, however, in the lower salinity waters the constants of H outperformed those of GP. Similarly to February, the constants of MM and ML also perform well and show the best consistency across the entire salinity range, with the constants of ML generally displaying a greater level of accuracy. The salinity range does not change between the two seasons, however, in February, 2002 a 3.5 fold higher Elbe river discharge was recorded compared to August, 2005 (GKSS, 2012, <http://coast.gkss.de/staff/kappenberg/elbe/abfluss/elbe.abfluss>). A larger proportion of freshwater in the North Sea could explain why a greater difference is observed between the two salinity environments during this time.

From this application of the various dissociation constants of carbonic acid, we show that the range of salinity directly affects the robustness of carbonate system calculations with different constants. Specifically, the constants R, H, GP and HM show much larger errors in mediums with a larger freshwater constituent compared to those of MM and ML. The larger range of salinities for which these latter constants were determined is likely to contribute to more accurate determinations of K_1 and K_2 , thus also leading to more accurate K_1/K_2 ratios. Based on these results, for datasets that cover a large range of salinities, for consistency, one can achieve the best overall results using the constants of ML in carbonate system calculations. Despite this, and bearing in mind that the current recommended dissociation constants for carbonate

system measurements are those of M, all further calculations and comparisons are performed using this set of constants.

3.1.2 The Role of Borate Alkalinity

The ratio of K_1/K_2 varies between different parameterizations of carbonic acid dissociation constants, partly depending on whether the constants were determined in artificial seawater (ASW) or natural seawater (NSW). In NSW interactions between borate and carbonate cause an increase in K_1 and a decrease in K_2 (Mojico-Prieto and Millero, 2002), which does not occur in ASW due to the lack of boric acid. As such, we suggest that finding the constants of R, GP and H to perform comparably with those of M in less saline waters may be indicative of the lower borate concentrations in freshwater and thus the errors in pK_2 , arising from the lack of boric acid in their ASW medium, are minimized. Figure 2a and 2b clearly show that borate is the second largest non-carbon contributor to A_T , constituting on average ~4% (Zeebe and Wolf-Gladrow, 2001), thus deviations from the true borate concentration can potentially cause substantial errors in calculations involving A_T . Additionally, the K_B of the boric acid system is very close to ocean pH, thus any miscalculation of $[H^+]$ can also lead to significant errors in the calculation of $[B(OH)_3]$ and $[B(OH)_4^-]$.

Borate has a long residence time in the oceans leading to a conservative relationship with salinity, thus in carbonate system calculations the total boron contribution is calculated using the linear relationship to chlorinity as found by Uppström (1974), and the borate then calculated using a thermodynamically derived K_B . The data used by Uppström (1974) is based on data obtained from the Pacific Ocean. However, recent findings suggest that this relationship is not consistent in all the worlds oceans (Lee et al., 2010). Lee et al. (2010) recently found that this relationship is not as ubiquitous in the oceans as originally thought and has made adjustments which, on provisional tests, led to an increase in the internal consistency of carbonate system calculations in the North Atlantic. The Uppström (1974) borate-salinity relationship was substituted in CO2SYS for the new Lee et al. (2010) relationship, to see if this improved the internal consistency. The new calculations led to a maximum difference

in borate alkalinity of $4 \mu\text{mol A}_T \text{ kg}^{-1}$, during both seasons. Despite this change, the influence on internal consistency was almost imperceptible, and resulted in no significant improvement.

In the North Sea, the Baltic input through the Skagerrak, and the Elbe input in the German Bight, have also revealed non-conventional salinity-borate relationships (Kremling, 1972; Barth, 1998). To evaluate the potential error introduced into calculations of the carbonate system in the North Sea by unaccounted-for borate, the borate concentrations in the Elbe- and Baltic-influenced areas of the North Sea were calculated using the water mass fractions and borate-salinity equations from the literature (see section 2.5). This 'pre-calculated' borate contribution to alkalinity (BAlk_{pre}) was then implemented into CO2SYS, replacing the normal salinity-borate calculation (Uppström et al., 1974). For comparative purposes BAlk was also calculated using the traditional Uppström (1974) equation but using various $[\text{H}^+]$ determinations, including that from direct pH measurements, using DIC and A_T as input parameters and using A_T and pCO_2 as input parameters (DIC/ pCO_2 produce the same results as A_T/pCO_2 , hence have not been included). The BAlk_{pre} calculations showed little change from the BAlk concentrations, the difference showing a normal distribution with 0 as the mean of the absolute values and σ of $3 \mu\text{mol kg}^{-1}$. The equations of Barth (1998) are very similar to the oceanic relationships, thus no significant difference was observed in the German Bight between the two calculations. The greatest difference occurs in the Skagerrak where the Uppström et al. (1974) ratio underestimates the BAlk compared to the equation derived from Gripenberg's data. The differences in calculation resulted in a maximum difference in BAlk of $8 \mu\text{mol kg}^{-1}$ in August 2005. Despite these differences, using the new calculated BAlk_{pre} values in CO2SYS did not improve the internal consistency; on the contrary the changes caused greater deviations between the measured and calculated parameters, reducing the number of internally consistent stations for all input combinations. Thus we conclude that the non-conservative salinity-borate relationships in the North Sea are not limiting factors for the internal consistency.

When the $[\text{H}^+]$ from pH measurements is used to calculate BAlk directly, and this

value is implemented into the thermodynamic calculations, the overall internal consistency is again unimproved. On the other hand, when the BAlk calculated using $[H^+]$ from (DIC, A_T) is implemented, the internal consistency improves notably if DIC or A_T are being calculated, for both seasons. The number of internally consistent (IC) stations in February and August, for A_T (DIC,pH) and DIC(A_T ,pH), increased by 11% (7 stations) and 36% (29 stations). The internal consistency of A_T (DIC,pCO₂) increased by 8% and 19% for February and August, respectively, and DIC(A_T ,pCO₂) increased by 6% and 16% stations, respectively. It should be noted that the $[H^+]$ calculated from (DIC, A_T) is only used to calculate the BAlk, whereas the $[H^+]$ used in the other CO2SYS calculations is calculated from the stated input parameters. As such, we can conclude that it is the effect of miscalculation of $[H^+]$ on the BAlk concentration that contributes significantly to a lack of internal consistency for (A_T , pCO₂) and (DIC, pCO₂). This can be illustrated by Figure 5(a), where the change in ΔA_T between the measured and calculated values are plotted against the change in $[H^+]$ between that calculated from (DIC, A_T) and (DIC, pCO₂). When the $[H^+]$ from (DIC, A_T) is used to calculate BAlk the ΔA_T values are shifted towards the x-axis (0), indicative of the improvement in internal consistency. The reason for such a bias is unknown, and may be caused by an undetected bias in one of the measured parameters.

3.2 Seasonal comparison (A_T , DIC, and pCO₂)

3.2.1 Seasonal Internal Consistency

The results from the internal consistency study from all 6 data sets, using just three carbonate parameters (A_T , DIC and pCO₂) show a clear dominance of the constants of M and ML outperforming the others. Both of these sets of constants show comparable results. The results of the seasonal calculations with the use of M constants are given in Table 3. The statistics indicate that the internal consistency of the North Sea follows a seasonal pattern, with May – the most biologically productive time period – producing less consistent results and the rest of the year displaying a good level of consistency. These findings are in agreement with those of Koeve et al. (2012), who similarly noted significant inconsistencies in the carbonate system in waters of high

biological activity.

Additionally, the largest fraction of Baltic water found in the Skagerrak is present in May (17%). Further analysis shows that during this season there is a significant ($p=0.008$, $R^2=0.55$), positive correlation between the Baltic fraction and the deviation between calculated and measured parameters (Figure 5(b)). Despite this, in May, the internal consistency could also not be improved by changing the BAlk calculation as done above (see section 3.1.2). However, Gripenberg (1960) concluded, from back titrations of alkalinity, that whilst borate did not directly affect the accuracy of the A_T titrations, the interactions between boric acid and organic material could lead to underestimated alkalinity values. Recent work by Ulfsbo et al. (2015) similarly find that in the Baltic, dissolved organic matter can contribute up to $30 \mu\text{mol } A_T \text{ kg}^{-1}$ to alkalinity, thus rendering the A_T equation in CO2SYS invalid. It thus seems likely that the lack of internal consistency observed in Baltic-influenced waters in May stems from additional organic material in the water at this time of year, with possibly further errors introduced from the interaction of this organic material with boric acid.

The North Sea receives a high input of allochthonous dissolved organic matter, for which analysis of colored dissolved organic matter (CDOM) indicates that it contains a high proportion of terrigenous sources (Stedmon et al., 2009). Thus may contain terrestrial “titratable” compounds, which, are also not accounted for in our calculations. It has been suggested that fatty acids and humic substances may contribute to measured A_T and in the Baltic Sea there is a higher than average concentration of both. The contribution of fatty acids to A_T has been fairly extensively examined by Osterroht (1993) and maximum contributions of $<1 \mu\text{mol } A_T \text{ kg}^{-1}$ to alkalinity were found, which is insufficient to cause the inconsistencies observed here. In contrast, more recently, the presence of humic substances has been shown to contribute to large excess A_T in the Baltic Sea (Ulfsbo et al., 2015). The contribution of organic matter to A_T has been observed elsewhere, as well. Muller et al. (2008), using the technique of Hernandez-Ayon et al. (2005) on fjord surface waters, compared the A_T estimation of DIC to the measured value and found that organic

alkalinity can contribute from 2-22 $\mu\text{mol A}_T \text{ kg}^{-1}$ in regions of high organic matter. These errors are sufficiently large to account for the noted decrease in internal consistency observed in the North Sea in May, compared to other seasons.

As the North Sea consists of two different biogeochemical regimes, the nNS and the sNS, we investigated whether these two environments contributed towards a difference in the internal consistency of the carbonate system. The North Sea was subsequently divided along the 56°N latitude and the internal consistency re-examined in both areas, applying just the constants of M (Table 4). In May there is a significant increase in the standard deviation of ΔDIC and ΔpCO_2 in the nNS compared to the sNS, however, ΔA_T shows no difference. As previously mentioned there is greater freshwater influence in the sNS. However, the difference observed between the nNS and sNS is larger than the difference previously mentioned between oceanic and estuarine waters. In the nNS larger blooms of phytoplankton are maintained for long periods due to seasonal, thermal stratification, which is in contrast to the sNS where rapid remineralization takes place in the permanently well-mixed water column. Errors have previously been associated with intense biological activity (Kim and Lee, 2009; Hoppe et al., 2012). Thus we are inclined to suggest that this may have a similar effect here. However, it is interesting that the errors in ΔA_T are not affected on a north-south latitudinal gradient, like ΔpCO_2 and ΔDIC .

The different regimes in the nNS and sNS also influence the dominant phytoplankton species present in each location. Calcifying organisms and the presence of coccolithophores are well documented in the nNS, where the development of coccolithophore blooms is controlled by the North Atlantic influence and the stratification of the water column (Houghton et al., 1991; Burkill et al., 2002). The presence of coccolithophores in waters sampled for DIC could conceivably lead to a degree of CaCO_3 dissolution during the DIC titration. This would introduce a degree of noise to the DIC values in the nNS, and possibly account for some of the larger errors observed in this region. Furthermore, Fraga & Alvarez-Salgado (2005) quantified the change in A_T during photosynthesis and found that the differences are

caused predominantly by CaCO_3 formation. Calcification influences the A_T/DIC (X) ratio of the waters as it removes one mole of DIC per 2 moles of A_T (Wolf-Gladrow et al., 2007). Lee et al. (1996, 1997) found that high X ratios (>1.10) can impact the sensitivity of pK_2 , thus affecting the consistency between measured and calculated carbonate parameters. The range of X (A_T/DIC) found in the North Sea varies from 1.06 - 1.16 throughout the year, with the smallest range (1.06 – 1.09) occurring in February and the largest range (1.07 - 1.16) found in May. During May, X also shows a positive linear correlation with ΔDIC , ΔA_T and ΔpCO_2 with R^2 values of 0.6, 0.6 and 0.7, respectively (Figure 6). The size of the error introduced by varying X can be up to $5 \mu\text{mol } A_T \text{ kg}^{-1}$, as found by Lee et al. (1996), which would contribute a large proportion of the errors we observe in the North Sea.

4 Conclusions

On a basin-wide scale in the North Sea we find that the carbonic acid dissociation constants of Mehrbach et al., (1973, refit by Dickson and Millero, 1987), and the constants of Millero et al., (2006) provide the best internal consistency. In the case of the former constants, this is in agreement with previous findings (Lee et al., 1996; Wanninkhof et al., 1999). In regions of lower than average ocean salinity, there is markedly less deviation between the internal consistency of the different constants and the constants of ML outperform those of M. At high salinities these two sets of constants are comparable in performance, thus over datasets with a wide variety of salinities, we recommend the use of ML constants for more consistent results.

In the North Sea in general, we find that the internal consistency of the carbonate system is much less than that found in the open ocean, by approximately 2-fold, which is unsurprising considering the additional input of terrestrial compounds in coastal areas, and generally higher productivity. Application of the new borate-salinity relationship noted by Lee et al. (2010), and further accounting for the non-conventional salinity-borate relationships in the water masses contributing to the North Sea, did not result in any improvement in the internal consistency. The internal consistency is relatively constant throughout the year, with the exception of May,

where there is a marked decrease in our ability to accurately calculate the carbonate system. Waters in which the Baltic Sea constituent is larger than 2% show lower internal consistency in May driven by the presence of excess organic alkalinity unaccounted for in carbonate system calculations. In the entire North Sea there is a significant correlation between the internal consistency and X (A_T/DIC), similar to that found in previous studies. Higher X values in the nNS compared to the sNS are likely to contribute to the larger errors observed in this area through the increased sensitivity on pK_2 .

Acknowledgements

We thank the captains and crews of the Research Vessel *Pelagia* and an anonymous reviewer for constructive comments on a previous version of the manuscript. This program was supported by the Research Council for Earth and Life Sciences (ALW) of the Netherlands Organization for Scientific Research (NWO), project number 817.01.004 and contributes to CARBO-OCEAN, an integrated project of the European Union (contract no. 511176-2). This work contributes to IGBP/IHDP LOICZ. AVB is a senior research associate at the FRS-FNRS.

References

Abril G., Bouillon, S., Darchambeau, F., Teodoru, C. R., Marwick, T. R., Tamooch, F., Omengo, F. O., Geeraert, N., Deirmendjian, L., Polsenaere, P., and Borges, A. V., 2015. Technical Note: Large overestimation of pCO_2 calculated from pH and alkalinity in acidic, organic-rich freshwaters. *Biogeosciences*, 12(1), 67-78.

Amann, T., Weiss, A., and Hartmann, J., 2012. Carbon dynamics in the freshwater part of the Elbe estuary, Germany: Implications of improving water quality. *Estuar. Coast. Shelf Sci.* 107, 112-121.

Barth, S., 1998. $^{11}B/^{10}B$ variations of dissolved boron in a freshwater-seawater mixing plume (Elbe Estuary, North Sea). *Mar. Chem.* 62, 1-14.

Borges, A. V. and Gypens, N., 2010. Carbonate chemistry in the coastal zone responds

more strongly to eutrophication than to ocean acidification. *Limnol. Oceanogr.* 55, 346-353.

Bozec, Y., Thomas, H., Elkalay, K., and de Baar, H. J. W., 2005. The continental shelf pump for CO₂ in the North Sea - evidence from summer observation. *Mar. Chem.* 93, 131-147.

Bozec, Y., Thomas, H., Schiettecatte, L. -S., Borges, A. V., Elkalay, K., and de Baar, H. J. W., 2006. Assessment of the processes controlling seasonal variations of dissolved inorganic carbon in the North Sea. *Limnol. Oceanogr.* 51, 2,746-2,762.

Brewer, P. G., and Goldman, J. C., 1976. Alkalinity changes generated by phytoplankton growth. *Limnol. Oceanogr.* 21,108-117.

Burkill, P. H., Archer, S. D., Robinson, C., Nightingale, P. D., Groom, S. B., Tarran, G. A. and Zubkov, M. V., 2002. Dimethyl sulphide biogeochemistry within a coccolithophore bloom (DISCO): an overview. *Deep-Sea Res. Pt. II* 49, 2,863-2,885.

Caldeira, K. and Wickett, M. E., 2003. Anthropogenic carbon and ocean pH. *Nature* 425, 365.

Caldeira, K. and Wickett, M. E., 2005. Ocean model predictions of chemistry changes from carbon dioxide emissions to the atmosphere and ocean. *J. Geophys. Res.* 110, C09S04, doi:10.1029/2004JC002671.

Canadell, J. G., Le Quere, C., Raupach, M. R., Field, C. B., Buitenhuis, E. T., Ciais, P., Conway, T. J., Gillett, N. P., Houghton R. A. and Marland G., 2007. Contributions to accelerating atmospheric CO₂ growth from economic activity, carbon intensity, and efficiency of natural sinks. *P. Natl. Acad. Sci.* 104, 18,866-18,870.

Clayton, T. D., Byrne, R. H., Breland, J. A., Feely, R. A., Millero, F. J., Campbell, D. M., Murphy, P. P., Lamb, M. F., 1995. The role of pH measurements in modern oceanic CO₂-system characterizations: precision and thermodynamic consistency. *Deep-Sea Res. Pt. I* 42, 411-431.

Dickson, A. G., and Millero, F. J., 1987. A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep-Sea Res. Pt. I* 34, 1,733-1,743.

Dickson, A. G., 1990. Standard potential of the reaction: $\text{AgCl(s)} + 1/2\text{H}_2(\text{g}) = \text{Ag(s)} + \text{HCl(aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic seawater from 273.15 to 318.15K. *J. Chem. Thermodyn.* 22,113-127.

Dickson, A. G., 1993. pH buffers for sea water media based on the total hydrogen ion concentration scale. *Deep-Sea Res. A.* 40, 107-118.

Dickson, A. G., Sabine, C.L. and Christian, J. R., 2007. Guide to best practices for ocean CO_2 measurements. PICES Special Publication 3.

Dunne, J. P., Sarmiento J. L. and Gnanadesikan, A., 2007. A synthesis of global particle export from the surface ocean and cycling through the ocean interior and on the seafloor. *Global Biogeochem. Cy.* 21, GB4006, doi:10.1029/2006GB002907.

Fabry, V. J., Seibel, B. A., Feely, R. A. and Orr, J. C., 2008. Impacts of ocean acidification on marine fauna and ecosystem processes, ICES. *J. Mar. Sci.* 65, 414-432.

Fraga, F. and Alvarez Salgado, X. A., 2005. On the variation of alkalinity during phytoplankton photosynthesis. *Cienc. Mar.* 31(4), 627-639.

Frankignoulle, M. and Borges, A.V., 2001. European continental shelf as a significant sink for atmospheric carbon dioxide. *Global Biogeochem. Cy.* 15, 569-576.

Friis, K., Körtzinger, A. and Wallace, D. W. R., 2003. The salinity normalization of marine inorganic carbon chemistry data. *Geophys. Res. Lett.* 30, 1085, doi:10.1029/2002GL015898.

Goyet, C. and Poisson, A., 1989. New determination of carbonic acid dissociation

constants in seawater as a function of temperature and salinity. *Deep-Sea Res. I* 36, 1,635-1,654.

Gran, G., 1952. Determination of the equivalence point in potentiometric titrations of seawater with hydrochloric acid. *Oceanol. Acta*, 5, 209-218.

Gripenberg, S., 1960. On the Alkalinity of Baltic Waters. *ICES J. Mar. Sci.* 26, 5-20.

Hansson, I., 1973. A new set of acidity constants for carbonic acid and boric acid in sea water. *Deep-Sea Res. I* 20, 461-478.

Hernandez-Ayon, J. M., Zirino, A., Dickson, A. G., Camiro-Vargas, T., and Valenzuela-Espinoza, E., 2007. Estimating the contribution of organic bases from microalgae to the titration alkalinity in coastal seawaters. *Limnol. Oceanogr-meth.* 5, 225-232.

van Heuven, S., Pierrot, D., Lewis, E. and Wallace, D.W.R., 2011. MATLAB Program developed for CO₂ system calculations. ORNL/CDIAC-105b. Carbon dioxide information analysis center. Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge.

Hoppe, C. J. M., Langer, G., Rokitta, S. D., Wolf-Gladrow, D. A., and Rost, B., 2012. Implications of observed inconsistencies in carbonate chemistry measurements for ocean acidification studies. *Biogeosciences* 9, 2401-2405.

Houghton, S. D., 1991. Coccolith sedimentation and transport in the North Sea. *Mar. Geol.* 99, 267-274.

Hydes, D. and Hartman, S., 2012. Seasonal and inter-annual variability in alkalinity in Liverpool Bay (53.5°N, 3.5°W) and in major river inputs to the North Sea. *Ocean Dynam.* 62, 321-333.

Johnson, K. M., Wills, K. D., Butler, D. B., Johnson, W. K., and Wong, C. S., 1993. Coulometric total carbon dioxide analysis for marine studies: maximizing the

performance of an automated gas extraction system and coulometric detector. *Mar. Chem.* 44, 167-187.

Kempe, S. and Pegler, K., 1991. Sinks and sources of CO₂ in coastal seas: the North Sea. *Tellus B* 43, 224-235.

Khatiwala, S., Primeau, F., and Hall, T., 2009. Reconstruction of the history of anthropogenic CO₂ concentrations in the ocean. *Nature* 462, 346 – 349.

Kim, H. -C. and Lee, K., 2009. Significant contribution of dissolved organic matter to seawater alkalinity. *Geophys. Res. Lett.* 36, L20603.

Koeve, W. and Oschlies, A., 2012. Potential impact of DOC accumulation on fCO₂ and carbonate ion computations in ocean acidification experiments. *Biogeosciences* 9, 3,787-3,798.

Körtzinger, A., Thomas, H., Schneider, B., Gronau, N., Mintrop, L. and Duinker, J. C., 1996. At-sea intercomparison of two newly designed underway pCO₂ systems – encouraging results. *Mar. Chem.* 52,133-145.

Körtzinger, A., Mintrop, L., Wallace, D. W. R., Johnson, K. M., Neill, C., Tilbrook, B., Towler, P., Inoue, H. Y., Ishii, M., Shaffer, G., Torres Saavedra, R. F., Ohtaki, E., Yamashita, E., Poisson, A., Brunet, C., Schauer, B., Goyet, C., and Eiseid, G., 2000. The international at-sea intercomparison of fCO₂ systems during the R/V Meteor Cruise 36/1 in the North Atlantic Ocean. *Mar. Chem.* 72, 171-192.

Kremling, K., 1972. Untersuchungen ueber die chemische Zusammensetzung des Meerwassers aus der Ostsee. *Kieler Meeresforschungen*, XXVIII, H.2, 99-118.

Le Quéré, C., Moriarty, R., Andrew, R. M., Peters, G. P., Ciais, P., Friedlingstein, P., Jones, S. D., Sitch, S., Tans, P., Arneeth, A., Boden, T. A., Bopp, L., Bozec, Y., Canadell, J. G., Chini, L. P., Chevallier, F., Cosca, C. E., Harris, I., Hoppema, M., Houghton, R. A., House, J. I., Jain, A. K., Johannessen, T., Kato, E., Keeling, R. F., Kitidis, V., Klein Goldewijk, K., Koven, C., Landa, C. S., Landschützer, P., Lenton,

A., Lima, I. D., Marland, G., Mathis, J. T., Metzl, N., Nojiri, Y., Olsen, A., Ono, T., Peng, S., Peters, W., Pfeil, B., Poulter, B., Raupach, M. R., Regnier, P., Rodenbeck, C., Saito, S., Salisbury, J. E., Schuster, U., Schwinger, J., Seferian, R., Segschneider, J., Steinhoff, T., Stocker, B. D., Sutton, A. J., Takahashi, T., Tilbrook, B., van der Werf, G. R., Viovy, N., Want, Y. -P., Wanninkhof, R., Wiltshire, A., and Zeng, N., 2015. Global carbon budget 2014. *Earth Syst. Sci. Data* 7, 47-85.

Lee, K. and Millero, F., 1995. Thermodynamic studies of the carbonate system in seawater. *Deep-Sea Res. Pt. I* 42, 2035-2061.

Lee, K., Millero, F. J. and Campbell, D. M., 1996. The reliability of the thermodynamic constants for the dissociation of carbonic acid in seawater. *Mar. Chem.* 55, 233-245.

Lee, K., Millero, F. and Wanninkhof, R., 1997. The carbon dioxide system in the Atlantic Ocean. *J. Geophys. Res.* 102, 15,693–15,707.

Lee, K., Millero, F. J., Byrne, R. H., Feely, R. A. and Wanninkhof, R., 2000. The recommended dissociation constants for carbonic acid in seawater. *Geophys. Res. Lett.* 27, 229-232.

Lewis, E. L. and Wallace, D. W. R., 1998. Program developed for CO₂ system calculations, ORNL/CDIAC-105. Carbon dioxide information analysis center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge.

Lueker, T. J., Dickson, A. G. and Keeling, C. D., 2000. Ocean pCO₂ calculated from dissolved inorganic carbon, alkalinity, and equations for K₁ and K₂: validation based on laboratory measurements of CO₂ in gas and seawater at equilibrium. *Mar. Chem.* 70, 105-119.

Mehrbach, C., Culberson, C. H., Hawley, J. E., and Pytkowicz, R. M., 1973. Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure. *Limnol. Oceanogr.* 18, 897-907.

Millero, F. J., Byrne, R. H., Wanninkhof, R., Feely, R., Clayton, T., Murphy, P. and Lamb, M. F., 2002. The internal consistency of CO₂ measurements in the equatorial Pacific. *Mar. Chem.* 44, 269-280.

Millero, F. J., Pierrot, D., Lee, K., Wanninkhof, R., Feely, R., Sabine, C. L., Key, R. M. and Takahashi, T., 2002. Dissociation constants for carbonic acid determined from field measurements. *Deep-Sea Res. Pt. I* 49, 1705-1723.

Millero, F. J., Graham, T. B., Huang, F., Bustos-Serrano, H. and Pierrot, D., 2006. Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Mar. Chem.* 100, 80-94.

Mojica-Preito F. J. and Millero, F. J., 2002. The values of pK₁ and pK₂ for the dissociation of carbonic acid in seawater. *Geochim. Cosmochim. Acta* 66, 2,529-2,540.

Mosley, L. M., Peake, B. M. and Hunter, K. A., 2010. Modelling of pH and inorganic carbon speciation in estuaries using the composition of the river and seawater end members. *Environ. Model. Software* 25, 1,658-1,663.

Muller F. L. L. and Bleie, B., 2008. Estimating the organic acid contribution to coastal seawater alkalinity by potentiometric titrations in a closed cell. *Anal. Chim. Acta* 619, 183-191.

Omar, A. M., Olsen, A., Johannessen, T., Hoppema, M., Thomas, H. and Borges A. V., 2010. Spatiotemporal variations of fCO₂ in the North Sea. *Ocean Sci.* 6, 77-89, doi:10.5194/os-6-77-2010.

Osterroht, C., 1993. Extraction of dissolved fatty acids from sea water. *Fresenius J. Anal. Chem.* 345, 773-779.

Otto, L., Zimmerman, J. T. F., Furnes, G. K., Mork, M., Saetre, R., and Becker, G., 1990. Review of the physical oceanography of the North Sea. *Neth. J. Sea Res.* 26, 161-238.

- Park, K. P., 1969. Oceanic CO₂ System: An evaluation of ten methods of investigation. *Limnol. Oceanogr.* 14, 179-186.
- Provost, P., van Heuven, S., Soetaert, K., Laane, R. W. P. M. and Middelburg, J. J., 2010. Seasonal and long-term changes in pH in the Dutch coastal zone. *Biogeosciences* 7, 3,869-3,878.
- Prowe, A. E. F., Thomas, H., Pätsch, J., Kühn, W., Bozec, Y., Schiettecatte, L. -S., Borges, A. V., and de Baar, H. J. W., 2009. Mechanisms controlling the air-sea CO₂ flux in the North Sea. *Cont. Shelf Res.* 29, 1801-1808.
- Raven, J., 2005. Ocean acidification due to increasing atmospheric carbon dioxide. Document No. 12/05, The Royal Society, London.
- Ribas-Ribas, M., Rérolle, V. M. C., Bakker, D. C. E., Kitidis, V., Lee, G. A., Brown, I., Achterberg, E. P. and Tyrrell, T., 2014. Intercomparison of carbonate chemistry measurements on a cruise in northwestern European shelf seas. *Biogeosciences* 11, 4,339-4,355.
- Roy, R. N., Roy, L. N., Vogel, K. M., Porter-Moore, C., Pearson, T., Good, C. E., Millero, F. J. and Campbell, D. M., 1993. Determination of the ionization constants of carbonic acid in seawater. *Mar. Chem.* 44, 249-259.
- Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C. S., Wallace, D. W. R., Tilbrook, B., Millero, F. J., Peng, T. -H., Kozyr, A., Ono, T. and Rios, A. F., 2004. The Oceanic Sink for Anthropogenic CO₂. *Science* 305, 367-371.
- Salt, L. A, Thomas, H., Prowe, F. A. E, Borges, A. V., Bozec, Y., and de Baar, H. J. W., 2013. Variability of North Sea pH and CO₂ in response to North Atlantic Oscillation forcing. *J. Geophys. Res. Biogeosci.* 118, 1-9, doi:10.1002/2013JG002306.

Schiettecatte, L. -S., Thomas, H., Bozec, Y. and Borges, A.V., 2007. High temporal coverage of carbon dioxide measurements in the Southern Bight of the North Sea. *Mar. Chem.* 106, 161-173.

Stedmon, C. A., Osburn, C. L., and Kragh, T., 2010. Tracing water mass mixing in the Baltic-North Sea transition zone using the optical properties of coloured dissolved organic matter. *Estuar. Coast. Shelf Sci.* 87, 156-162.

Thomas, H., Bozec, Y., Elkalay, K., and de Baar, H. J. W., 2004. Enhanced Open Ocean Storage of CO₂ from Shelf Sea Pumping. *Science* 304, 1005 - 1008.

Thomas, H., Bozec, Y., Elkalay, K., de Baar, H. J. W., Borges, A. V. and Schiettecatte, L. -S., 2005. Controls of the surface water partial pressure of CO₂ in the North Sea. *Biogeosciences* 2, 323-334.

Thomas, H., Schiettecatte, L. -S., Suykens, K., Koné, Y. J. M., Shadwick, E. H., Prowe, A. E. F., Bozec, Y., de Baar, H. J. W. and Borges, A. V., 2009. Enhanced ocean carbon storage from anaerobic alkalinity generation in coastal sediments. *Biogeosciences* 6, 267-274.

Ulfso, A., Kulinski, K., Anderson, L.G., and Turner, D.R., 2015. Modelling organic alkalinity in the Baltic Sea using a Humic-Pitzer approach. *Mar. Chem.* 168, 18-26.

Uppström, L. R., 1974. The boron/chlorinity ratio of deep-sea water from the Pacific Ocean. *Deep-Sea Res.* 21, 161-162.

Wanninkhof, R., Lewis, E., Feely, R. A. and Millero, F. J., 1999. The optimal carbonate dissociation constants for determining surface water pCO₂ from alkalinity and total inorganic carbon. *Mar. Chem.* 65, 291 - 301.

Wolf-Gladrow, D. A., Zeebe, R. E., Klaas, C., Körtzinger, A., and Dickson, A. G., 2007. Total alkalinity: The explicit conservative expression and its application to biogeochemical processes. *Mar. Chem.* 106, 287-300.

Wollast, R., 1998. Evaluation and comparison of the global carbon cycle in the coastal zone and in the open ocean. In: Brink, K. H., Robinson, A. R. (Eds.), *The Sea*. John Wiley & Sons, New York, pp. 213-252.

Wootton, J. T., Pfister, C. A. and Forester, J. D., 2008. Dynamic patterns and ecological impacts of declining ocean pH in a high-resolution multi-year dataset. *P. Natl Acad. Sci.* 105, 18,848-18,853.

Zeebe, R. E. and Wolf-Gladrow, D., 2001. *CO₂ in Seawater: Equilibrium, Kinetics, Isotopes*. Elsevier, Amsterdam.

Table Captions

Table 1. Table of parameterizations of the carbonic acid dissociation constants, with temperature and salinity range for each set, and the medium in which measurements were performed.

Table 2. Calculated statistics for all calculated values of parameters compared to the measured values during February 2002 (a), and for August 2005 (b). Both datasets were tested using the dissociation constants of Roy et al. (1993)(R), Goyet & Poisson (1989)(GP), Hansson (1973), refit by Dickson and Millero (1987)(H), Mehrbach et al. (1973), refit by Dickson and Millero (1987)(M), Hansson & Mehrbach (1973), refit by Dickson and Millero (1987)(HM), Mojico-Prieto & Millero (2002)(MM) and Millero et al. (2006)(ML).

Table 3. Derived linear regressions, using total least squares, and standard deviations of differences between measured and calculated parameters (A_T , DIC, and pCO_2) with the percentage of internally consistent stations for all six cruises. The calculations were performed using the acid dissociation constants of Mehrbach et al., (1973), refit by Dickson and Millero, (1987).

Table 4. Calculated internal consistency statistics for all data sets divided into northern North Sea stations (nNS)(latitude $>56^{\circ}$ N) and southern North Sea stations (sNS)(latitude $\leq 56^{\circ}$ N).

Figure Captions

Figure 1. Map of the North Sea region, with the stations sampled marked in black. The 50m depth contour is marked, which separates the southern North Sea from the northern North Sea.

Figure 2. The distribution of total alkalinity in the North Sea and respective contributions to total alkalinity by (i) carbonate species, (ii) borate, (iii) silicate, (iv) phosphate and (v) ammonia in February 2002 (panel a) and in August 2005 (panel b).

Figure 3. The difference in ΔA_T ($\mu\text{mol } A_T \text{ kg}^{-1}$) across the measured salinity range. The ΔA_T^R and ΔA_T^M are calculated using (DIC, $p\text{CO}_2$) as input parameters.

Figure 4. Radar plot showing (a) the regression coefficients for $A_T(\text{DIC}, p\text{CO}_2)$ and (b) $p\text{CO}_2(\text{DIC}, A_T)$ in different salinity environments, and their respective standard deviation (c), (d) of the fit.

Figure 5. (a) The change in ΔA_T caused by the difference in BAlk calculated using the $[\text{H}^+]$ from (DIC, A_T)(red) and (DIC, $p\text{CO}_2$)(black). (b) Regression of ΔDIC with the calculated water mass fraction of Baltic water, selectively displaying those with a fraction of 2% or higher.

Figure 6. The change in ΔDIC and ΔA_T (both in $\mu\text{mol kg}^{-1}$) and $\Delta p\text{CO}_2$ (in μatm) observed with a varying ratio of A_T/DIC (X) during May 2002.

Constants	pK_1 and pK_2 parameterization:	Temperature range	Salinity range	Media
R	$\ln K_1 = 2.83655 - 2307.1266/T - 1.5529413 \cdot \ln(T) + (-0.20760841 - 4.0484/T) \cdot S^{0.5} + 0.08468345 \cdot S - 0.00654208 \cdot S^{1.5}$	0 - 45	5 - 45	Artificial SW
	$\ln K_2 = -9.226508 - 3351.6106/T - 0.2005743 \cdot \ln(T) + (-0.106901773 - 23.9722/T) \cdot S^{0.5} + 0.1130822 \cdot S - 0.00846934 \cdot S^{1.5}$			
GP	$pK_1 = 812.27/T + 3.356 - 0.00171 \cdot S \cdot \ln(T) + 0.000091 \cdot S^2$	-1 - 40	10 - 50	Artificial SW
	$pK_2 = 1450.87/T + 4.604 - 0.00385 \cdot S \cdot \ln(T) + 0.000182 \cdot S^2$			
H	$pK_1 = 851.4/T + 3.237 - 0.0106 \cdot S + 0.000105 \cdot S^2$	2 - 35	20 - 40	Artificial SW
	$pK_2 = -3885.4/T + 125.844 - 18.141 \cdot \ln(T) - 0.0192 \cdot S + 0.000132 \cdot S^2$			
M	$pK_1 = 3670.7/T - 62.008 + 9.7944 \cdot \ln(T) - 0.0118 \cdot S + 0.000116 \cdot S^2$	2 - 35	26 - 43	Natural SW
	$pK_2 = 1394.7/T + 4.777 - 0.0184 \cdot S + 0.000118 \cdot S^2$			
HM	$pK_1 = 845/T + 3.248 - 0.0098 \cdot S + 0.000087 \cdot S^2$	2 - 35	20 - 40	Artificial SW
	$pK_2 = 1377.3/T + 4.824 - 0.0185 \cdot S + 0.000122 \cdot S^2$			
MM	$pK_1 = -43.6977 - 0.0129037 \cdot S + 1.364e^{-4} \cdot S^2 + 2885.378/T + 7.045159 \cdot \ln(T)$	0 - 45	5 - 42	Natural SW
	$pK_2 = -452.0940 + 13.142162 \cdot S - 8.101e^{-4} \cdot S^2 + 21263.61/T + 68.483143 \cdot \ln(T) + (-581.4428 \cdot S + 0.259601 \cdot S^2/T - 1.967035 \cdot S \cdot \ln(T))$			

ML	$\begin{aligned} \text{pK}_1 = & -126.34048 + \\ & 6320.813/\text{T} + 19.568224.\ln(\text{T}) \\ & + 13.4038.\text{S}^{0.5} + 0.03206.\text{S} - \\ & 5.242\text{e}^{-5}.\text{S}^2 + (-530.659.\text{S}^{0.5} - \\ & 5.8210.\text{S})/\text{T} + (- \\ & 2.0664.\text{S}^{0.5}).\ln(\text{T}) \end{aligned}$	0 - 50	1 - 50	Natural SW
	$\begin{aligned} \text{pK}_2 = & -90.18333 + 5143.692/\text{T} \\ & + 14.613358.\ln(\text{T}) + \\ & 21.3728.\text{S}^{0.5} + 0.1218.\text{S} - \\ & 3.688\text{e}^{-4}.\text{S}^2 + (-788.289.\text{S}^{0.5} - \\ & 19.189.\text{S})/\text{T} + (- \\ & 3.374.\text{S}^{0.5}).\ln(\text{T}) \end{aligned}$			

Table 1. Table of parameterizations of the carbonic acid dissociation constants, with temperature and salinity range for each set, and the medium in which measurements were performed.

Table 2a

			R	GP	H	M	HM	MM	ML	
A_T	Regress (slope)	DIC,pH	0.999	1.000	0.998	1.001	0.999	0.999	1.001	
		DIC,pCO₂	0.997	0.997	0.995	1.000	0.997	0.998	0.999	
		pH,pCO₂	1.022	1.035	1.042	1.013	1.031	1.009	1.027	
	R² (p-value)	DIC,pH	0.93 (<0.01)	0.93 (<0.01)	0.93 (<0.01)	0.93 (<0.01)	0.93 (<0.01)	0.93 (<0.01)	0.93 (<0.01)	0.93 (<0.01)
		DIC,pCO₂	0.92 (<0.01)	0.92 (<0.01)	0.93 (<0.01)	0.93 (<0.01)	0.92 (<0.01)	0.92 (<0.01)	0.93 (<0.01)	
		pH,pCO₂	0.03 (0.19)	0.03 (0.20)	0.06 (0.18)	0.06 (0.05)	0.04 (0.13)	0.05 (0.07)	0.06 (0.05)	
	Average ΔAT (σ) (μmol kg⁻¹)	DIC,pH (±2)	3 (8)	1(8)	4 (7)	-2 (7)	2 (8)	3 (8)	2 (7)	
		DIC,pCO₂ (±2)	8 (9)	8 (9)	12 (9)	1 (8)	8 (9)	6 (8)	6 (8)	
		pH,pCO₂ (±43)	-50 (63)	-79 (62)	-93 (62)	-32 (60)	-70 (61)	-23 (61)	48 (60)	
DIC	Regress (slope)	TA,pH	1.001	1.000	1.002	0.999	1.001	1.002	0.999	
		TA,pCO₂	1.003	1.003	1.005	1.000	1.003	1.002	1.001	

		pH,pCO₂	1.024	1.036	1.045	1.013	1.033	1.011	1.027
	R²	TA,pH	0.89 (<0.01)	0.90 (<0.01)	0.90 (<0.01)	0.90 (<0.01)	0.90 (<0.01)	0.90 (<0.01)	0.90 (<0.01)
	(p-value)	TA,pCO₂	0.88 (<0.01)	0.88 (<0.01)	0.90 (<0.01)	0.90 (<0.01)	0.89 (<0.01)	0.89 (<0.01)	0.90 (<0.01)
		pH,pCO₂	0.06 (0.07)	0.06 (0.07)	0.06 (0.05)	0.07 (0.03)	0.06 (0.06)	0.07 (0.04)	0.07 (0.04)
	Average ΔDIC (σ) (μmol kg⁻¹)	TA,pH (±8)	-3 (7)	-1 (7)	-4 (7)	2 (7)	-2 (7)	-3 (7)	-2(7)
		TA,pCO₂ (±7)	-7 (8)	-7 (8)	-11 (7)	-1 (7)	-7 (8)	-5 (7)	6 (7)
		pH,pCO₂ (±42)	-50 (59)	-76 (58)	-92 (57)	-29 (55)	-68 (57)	-25 (56)	44 (55)
pH	Regress (slope)	DIC,A_T	1.001	1.000	1.001	0.999	1.000	1.001	1.000
		A_T,pCO₂	0.999	0.998	0.998	0.999	0.998	1.000	0.999
		DIC,pCO₂	0.999	0.998	0.998	0.999	0.998	0.999	0.999
	R² (p-value)	DIC,A_T	0.23 (<0.01)	0.23 (<0.01)	0.25 (<0.01)	0.24 (<0.01)	0.24 (<0.01)	0.25 (<0.01)	0.25 (<0.01)
		A_T,pCO₂	0.53 (<0.01)	0.53 (<0.01)	0.53 (<0.01)	0.57 (<0.01)	0.55 (<0.01)	0.56 (<0.01)	0.56 (<0.01)
		DIC,pCO₂	0.49 (<0.01)	0.49 (<0.01)	0.49 (<0.01)	0.53 (<0.01)	0.51 (<0.01)	0.52 (<0.01)	0.53 (<0.01)

pCO₂	Average ΔpH (σ)	DIC,A_T(±0.008)	-0.006 (0.017)	0.001 (0.017)	-0.008 (0.017)	0.004 (0.017)	-0.004 (0.017)	-0.007 (0.017)	0.004 (0.017)
		A_T,pCO₂(±0.001)	0.009 (0.011)	0.014 (0.011)	0.016 (0.011)	0.006 (0.010)	0.012 (0.011)	-0.004 (0.011)	0.011 (0.011)
		DIC,pCO₂(±0.001)	0.010 (0.012)	0.015 (0.012)	0.018 (0.012)	0.006 (0.011)	0.014 (0.011)	0.005 (0.011)	0.011 (0.011)
	Regress (slope)	DIC,A_T	0.959	0.959	0.936	0.994	0.957	0.969	0.980
		DIC,pH	0.975	0.963	0.956	0.985	0.967	0.987	0.973
		A_T,pH	0.976	0.963	0.957	0.985	0.967	0.989	0.972
	R² (p-value)	DIC,A_T	0.14 (<0.01)	0.14 (<0.01)	0.14 (<0.01)	0.17 (<0.01)	0.16 (<0.01)	0.17 (<0.01)	0.18 (<0.01)
		DIC,pH	0.52 (<0.01)	0.52 (<0.01)	0.53 (<0.01)	0.57 (<0.01)	0.54 (<0.01)	0.56 (<0.01)	0.56 (<0.01)
		A_T,pH	0.52 (<0.01)	0.52 (<0.01)	0.53 (<0.01)	0.56 (<0.01)	0.54 (<0.01)	0.55 (<0.01)	0.56 (<0.01)
Average ΔpCO₂ (σ)	DIC,A_T(±8)	15 (18)	15 (18)	24 (17)	2 (17)	16 (17)	11 (17)	7 (17)	
	DIC,pH(±4)	9 (10)	14 (11)	17 (11)	5 (10)	12 (10)	5 (10)	10 (10)	

	(μatm)	A_T,pH (± 3)	9 (11)	14 (11)	16 (11)	6 (10)	12 (11)	4 (10)	10 (10)
--	---------------------	--	--------	---------	---------	--------	---------	--------	---------

Table 2b

			R	GP	H	M	HM	MM	ML	
A_T	Regress	DIC,pH	0.992	0.994	0.996	0.997	0.995	0.995	0.998	
		DIC,pCO₂	0.994	0.994	0.995	1.001	0.996	0.999	1.000	
		pH,pCO₂	0.977	0.991	0.999	0.962	0.988	0.960	0.976	
	R² (p-value)	DIC,pH	0.96 (<0.01)	0.96 (<0.01)	0.97 (<0.01)	0.97 (<0.01)	0.97 (<0.01)	0.97 (<0.01)	0.97 (<0.01)	0.97 (<0.01)
		DIC,pCO₂	0.94 (<0.01)	0.94 (<0.01)	0.94 (<0.01)	0.94 (<0.01)	0.94 (<0.01)	0.94 (<0.01)	0.94 (<0.01)	0.94 (<0.01)
		pH,pCO₂	0.18 (<0.01)	0.19 (<0.01)	0.19 (<0.01)	0.22 (<0.01)	0.22 (<0.01)	0.20 (<0.01)	0.21 (<0.01)	
	Average ΔAT (σ) (μmol kg⁻¹)	DIC,pH (±2)	18 (7)	14 (7)	10 (6)	7 (6)	11 (6)	12 (6)	5 (6)	
		DIC,pCO₂ (±3)	14 (9)	13 (9)	11 (9)	1 (9)	9 (9)	3 (9)	0 (9)	
		pH,pCO₂ (±55)	53 (52)	19 (51)	1 (51)	86 (52)	28 (51)	96 (52)	55 (51)	
DIC	Regress	A_T,pH	1.008	1.006	1.004	1.003	1.005	1.005	1.002	
		A_T,pCO₂	1.006	1.005	1.004	0.999	1.004	1.001	1.000	

pCO₂	Average ΔpH (σ)	DIC,A_T (±0.007)	-0.031 (0.012)	-0.024 (0.012)	-0.018 (0.012)	-0.012 (0.012)	-0.020 (0.012)	-0.021 (0.012)	-0.009 (0.012)
		A_T,pCO₂ (±0.001)	0.009 (0.009)	-0.003 (0.008)	0.000 (0.008)	-0.014 (0.008)	-0.005 (0.008)	-0.015 (0.008)	-0.009 (0.008)
		DIC,pCO₂ (±0.001)	-0.006 (0.009)	-0.001 (0.009)	0.002 (0.009)	-0.014 (0.009)	-0.003 (0.009)	-0.015 (0.009)	-0.009 (0.009)
	Regress	DIC,A_T	0.939	0.942	0.952	1.004	0.959	0.984	1.000
		DIC,pH	1.014	1.000	0.994	1.035	1.006	1.036	1.021
		A_T,pH	1.022	1.006	0.998	1.038	1.011	1.041	1.023
	R² (p-value)	DIC,A_T	0.89 (<0.01)	0.89 (<0.01)	0.89 (<0.01)	0.89 (<0.01)	0.89 (<0.01)	0.89 (<0.01)	0.89 (<0.01)
		DIC,pH	0.96 (<0.01)	0.96 (<0.01)	0.96 (<0.01)	0.96 (<0.01)	0.96 (<0.01)	0.96 (<0.01)	0.96 (<0.01)
		A_T,pH	0.97 (<0.01)	0.97 (<0.01)	0.97 (<0.01)	0.97 (<0.01)	0.97 (<0.01)	0.97 (<0.01)	0.97 (<0.01)
Average ΔpCO₂ (σ)	DIC,A_T (±6)	22 (15)	22 (14)	17 (14)	-2 (13)	15 (14)	5 (13)	0 (13)	
	DIC,pH (±4)	-5 (8)	-1 (8)	2 (8)	-12 (8)	-2 (8)	-13 (8)	-8 (8)	

	(μatm)	A_T,pH (± 3)	-8 (8)	-3 (8)	0 (8)	-13 (8)	-4 (8)	-14 (8)	-8 (8)
--	-------------------------------------	--	--------	--------	-------	---------	--------	---------	--------

Table 3

Input combination		Aug '01	Nov '01	Feb '02	May '02	Aug '05	Aug '08
A_T(DIC,pCO₂)	Regress	0.999	1.001	1.000	0.993	1.001	1.000
	Std	29	8	8	27	9	9
	IC (%)	15.5	47.7	43.5	9.4	34.9	50.2
DIC(A_T,pCO₂)	Regress	1.000	0.999	1.000	1.007	1.000	1.000
	Std	24	7	7	23	7	8
	IC (%)	17	48	44	9	39	52
pCO₂(DIC,A_T)	Regress	0.976	1.004	0.994	0.906	1.004	0.998
	Std	41	15	17	41	13	15
	IC (%)	26	50	40	20	49	57

	Std	24	8	8	12	7	7
	IC (%)	17.6	41.7	38.5	10.3	44.1	56.8
<hr/>							
	Regress	0.96	1.00	1.00	0.97	1.00	1.00
pCO₂(DIC,A_T)	Std	46	17	19	19	14	15
	IC (%)	23.5	47.2	38.5	25.6	52.9	59.5
<hr/>							
	# stations	34	36	26	39	34	37




Figure 1





Figure 2a

**Figure 2b**




Figure 3




Figure 4




Figure 5




Figure 6

Highlights

- The internal consistency of the carbonate system in the North Sea is notably less than the open ocean.
- The consistency of the carbonate system follows a seasonal pattern, with the largest errors in May.
- The carbonic acid dissociation constants of Millero et al. (2006) are most consistent across a large salinity range (0-36).