



**HAL**  
open science

## A note on the fate of the Landau–Yang theorem in non-Abelian gauge theories

Matteo Cacciari, Luigi del Debbio, José R. Espinosa, Antonio D. Polosa,  
Massimo Testa

► **To cite this version:**

Matteo Cacciari, Luigi del Debbio, José R. Espinosa, Antonio D. Polosa, Massimo Testa. A note on the fate of the Landau–Yang theorem in non-Abelian gauge theories. *Physics Letters B*, 2016, 753 (10), pp. 476-481. 10.1016/j.physletb.2015.12.053 . hal-01250179

**HAL Id: hal-01250179**

**<https://hal.sorbonne-universite.fr/hal-01250179>**

Submitted on 4 Jan 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# A note on the fate of the Landau–Yang theorem in non-Abelian gauge theories



Matteo Cacciari<sup>a,b,c,d</sup>, Luigi Del Debbio<sup>e,d</sup>, José R. Espinosa<sup>f,g</sup>, Antonio D. Polosa<sup>h,\*</sup>, Massimo Testa<sup>h</sup>

<sup>a</sup> Université Paris Diderot, F-75013 Paris, France

<sup>b</sup> Sorbonne Universités, UPMC Univ Paris 06, UMR 7589, LPHE, F-75005 Paris, France

<sup>c</sup> CNRS, UMR 7589, LPHE, F-75005 Paris, France

<sup>d</sup> CERN, PH-TH, CH-1211 Geneva 23, Switzerland

<sup>e</sup> Higgs Centre for Theoretical Physics, School of Physics & Astronomy, University of Edinburgh, Edinburgh EH9 3FD, United Kingdom

<sup>f</sup> ICREA, Institució Catalana de Recerca i Estudis Avançats, Barcelona, Spain

<sup>g</sup> IFAE, Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Spain

<sup>h</sup> Dipartimento di Fisica and INFN, 'Sapienza' Università di Roma, Piazzale Aldo Moro 2, I-00185 Roma, Italy

## ARTICLE INFO

### Article history:

Received 15 October 2015

Received in revised form 17 December 2015

Accepted 17 December 2015

Available online 21 December 2015

Editor: A. Ringwald

## ABSTRACT

Using elementary considerations of Lorentz invariance, Bose symmetry and BRST invariance, we argue why the decay of a massive color-octet vector state into a pair of on-shell massless gluons is possible in a non-Abelian  $SU(N)$  Yang–Mills theory, we constrain the form of the amplitude of the process and offer a simple understanding of these results in terms of effective-action operators.

© 2015 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP<sup>3</sup>.

## 1. Introduction

The Landau–Yang theorem states that a massive vector (i.e. spin 1) particle cannot decay into two on-shell massless photons. The proofs of Landau [1] and Yang [2] show that one can reach this conclusion under very general conditions, using only Lorentz invariance, gauge invariance (in the form of transversality of photon polarization vectors) and the Bose symmetry of the photons.

One can also consider the case of a massive color-octet vector state decaying into two on-shell massless gluons. This process is of phenomenological interest not only for models predicting the existence of colored massive vector particles, but also for heavy quarkonium physics, e.g. for the hadroproduction of a  $J/\psi$  particle or its decay into hadrons.<sup>1</sup> Evidence that this amplitude vanishes at tree-level has been given many times in Quantum Chromodynamics (QCD), usually by explicitly calculating the two-gluon decay of a quark–antiquark pair projected onto a massive color-octet spin-1,  $S$ -wave state that we denote as  $(Q\bar{Q})_{J=1}^{(8)}$ , see e.g.

Refs. [3–9]. In these papers it was generally understood, albeit with some exceptions, that the proofs of Landau and Yang could not be immediately extended to the color-octet case, because of additional terms in the amplitude induced by the antisymmetric character of the color quantum numbers. However, for a number of years no attempt was apparently made to study the color-octet case in more detail or to explicitly check if the vanishing of the amplitude at tree level still held at higher orders.

This situation began to change only recently, when two calculations found non-zero results for the one-loop amplitude of the transition between a massive color-octet vector state and two massless gluons: Ref. [10] calculated the next-to-leading order corrections to gluon fusion production of massive color-octet vector bosons (colorons), while Ref. [11] reported that the  $(Q\bar{Q})_{J=1}^{(8)} \rightarrow gg$  amplitude is also different from zero at one-loop level.

These results indicate that a colored version of the Landau–Yang theorem breaks down once quantum corrections are taken into account. In this note we analyze in depth the origin of the cancellation at tree level and the structure of the amplitude in full generality, we explore where the proof of a would-be Landau–Yang theorem for colored states fails, and derive constraints on the form of the amplitude for the decay of a massive vector color-octet in two massless gluons. We also identify higher-dimension effective operators whose presence can explain the non-vanishing results at

\* Corresponding author.

E-mail address: [antonio.polosa@roma1.infn.it](mailto:antonio.polosa@roma1.infn.it) (A.D. Polosa).

<sup>1</sup> In both cases the two gluons will not be exactly on-shell, but their off-shellness will be at most of the order of a few hundred MeV, and its effects therefore suppressed by the ratio with the much larger scale set by the  $J/\psi$  mass or transverse mass.

one-loop and beyond, and show how the LY theorem (or its failure) can be understood in a very direct and simple way at the operator level.

## 2. The Abelian case

It is instructive to first reconsider the original Landau–Yang theorem, highlighting the difficulties in extending it to the case of a color-octet state. We denote by

$$M(1, 2) \equiv \langle \gamma(k_1, \epsilon_1) \gamma(k_2, \epsilon_2) | V(P, \epsilon) \rangle, \quad (1)$$

the amplitude for the decay  $V \rightarrow \gamma\gamma$ , where  $V$  is a colorless massive spin-1 state and the  $\gamma$ 's are photons.  $k_1$  and  $k_2$  are the photons 4-momenta,  $P$  is the 4-momentum of  $V$ , and momentum conservation dictates  $P = k_1 + k_2$ . Relying only on Lorentz invariance and Bose symmetry ( $M(1, 2) = M(2, 1)$ ), one can write

$$\begin{aligned} M \sim & A(\epsilon_1 \cdot \epsilon_2)[\epsilon \cdot (k_1 + k_2)] \\ & + B[(\epsilon_1 \cdot \epsilon)(\epsilon_2 \cdot k_1) + (\epsilon_2 \cdot \epsilon)(\epsilon_1 \cdot k_2)] \\ & + C[\epsilon \cdot (k_1 + k_2)](\epsilon_1 \cdot k_2)(\epsilon_2 \cdot k_1), \end{aligned} \quad (2)$$

where  $\epsilon_1 \equiv \epsilon_1^*(k_1)$ ,  $\epsilon_2 \equiv \epsilon_2^*(k_2)$  and  $\epsilon \equiv \epsilon(P)$  are the polarization 4-vectors of the two photons and of the massive vector state respectively.  $A, B, C$  are coefficients that can be determined in perturbation theory: they only depend on scalar products of momenta, and therefore are constants in the decay of a massive vector particle. Terms that vanish because of transversality,  $\epsilon_1 \cdot k_1 = \epsilon_2 \cdot k_2 = 0$ , have not been included. Instead, for the sake of introducing the discussion of the non-Abelian case that will follow, we have kept in Eq. (2) also the  $A$  and  $C$  terms, even if in the Abelian case these terms are trivially equal to zero as a consequence of the transversality of the massive vector polarization,  $\epsilon(P) \cdot P = 0$ .

It is convenient to consider the  $B$  term in Eq. (2) after a Lorentz transformation to the rest frame of the massive state  $V$ . The photon polarizations transform under the Lorentz transformation, yielding in general non vanishing time components for  $\epsilon_{1,2}^\mu$ . However, the gauge invariance of the amplitude  $M$  in Eq. (2) is equivalent to the invariance of  $M$  under the transformations

$$\epsilon_j^\mu(k_j) \rightarrow \tilde{\epsilon}_j^\mu(k_j) = \epsilon_j^\mu(k_j) + \beta_j k_j^\mu, \quad (j = 1, 2), \quad (3)$$

where  $\beta_j$  are arbitrary constants. This means that we are allowed to use  $\tilde{\epsilon}_{1,2}$  in place of  $\epsilon_{1,2}$  in the expression of  $M$ . In particular, we choose  $\beta_j$  in such a way that

$$\tilde{\epsilon}_j^\mu(k_j, \pm) = \frac{1}{\sqrt{2}}(0, \mp 1, -i, 0). \quad (4)$$

Therefore, only polarizations transverse to the  $\hat{z}$  direction will eventually appear in  $M$ . Taking the 3-momenta  $\mathbf{k}_1$  and  $\mathbf{k}_2$  along  $\hat{z}$ , in the rest frame of  $V$ , we simply have  $\mathbf{k}_1 = -\mathbf{k}_2$  so that

$$\tilde{\epsilon}_{1(2)} \cdot k_{2(1)} = 0. \quad (5)$$

$M$  in Eq. (2) is therefore zero and  $V \rightarrow \gamma\gamma$  is forbidden, leading to a proof of the Landau–Yang theorem in the Abelian case.<sup>2</sup> The treatment of gauge invariance in the non-Abelian case, discussed in the next section, will require instead more care.

<sup>2</sup> If the initial particle were a  $1^+$  axial-vector, the only non-trivial parity conserving term would be proportional to  $\epsilon_{\mu\nu\rho\sigma} \epsilon^\mu \epsilon_1^\nu \epsilon_2^\rho (k_1 - k_2)^\sigma$ . In the rest frame of the massive vector particle  $k_1 - k_2 = (0, 0, 0, 2E)$ ,  $E$  being the photon energy. Thus  $\epsilon, \epsilon_1, \epsilon_2$  can only have 0, 1, 2 indices and permutations. Since  $\epsilon_{1,2}$  are transverse polarizations, and  $\epsilon^0 \sim |\mathbf{P}|/M$  (where  $\mathbf{P}$  and  $M$  are the massive vector momentum and mass), we get  $\epsilon_{\mu\nu\rho\sigma} \epsilon^\mu \epsilon_1^\nu \epsilon_2^\rho (k_1 - k_2)^\sigma = 0$ . This illustrates that the Landau–Yang theorem applies as well to positive parity vectors.

Before considering the non-Abelian case, it is instructive to understand the Landau–Yang result at the level of operators in the Lagrangian (or effective action) that describes the interaction between a color-singlet massive vector  $V^\mu$  and the electromagnetic tensor  $F^{\mu\nu}$ . This approach takes care of gauge invariance automatically and further cancellations can be shown by using the field equations of motion, as illustrated below. The only non-trivial operator<sup>3</sup> leading to the relevant Lorentz structures for the  $V \rightarrow \gamma\gamma$  decay written in Eq. (2) is:

$$\Delta\mathcal{L} = a(\partial^\mu V_\nu)F_{\mu\rho}F^{\rho\nu}, \quad (6)$$

where  $a$  is some coefficient with dimension  $\text{mass}^{-2}$ . Other operators with more derivatives just modify the momentum-dependent form factors of the vertices without changing its Lorentz structure.

Integrating by parts judiciously, the operator in Eq. (6) can be rewritten as

$$\Delta\mathcal{L} = -aV_\nu(\partial^\mu F_{\mu\rho})F^{\rho\nu} - \frac{a}{4}(\partial^\nu V_\nu)F_{\mu\rho}F^{\mu\rho}. \quad (7)$$

The second term, which can be written in this form thanks to the symmetry of  $F_{\mu\rho}F^{\mu\rho}$  under photon exchange, can be dropped due to  $\partial^\nu V_\nu = 0$ . In the first, we can use the equation of motion of the electromagnetic field to replace  $\partial^\mu F_{\mu\rho}$  by a sum over all electromagnetic currents to which the photon couples in the theory. This replacement, equivalent to a field redefinition, does not change the physics [12], but makes clear the fact that this Lagrangian does not contribute to the  $V \rightarrow \gamma\gamma$  decay with on-shell photons. (The modified Lagrangian in terms of electromagnetic currents will give the right description for processes with virtual photons coupled to such currents in the final state.)

## 3. Non-Abelian case: tree-level cancellation

As mentioned in the Introduction the  $V^a \rightarrow g^b g^c$  amplitude is known to vanish at tree level. The purpose of this section is to trace the origin of the tree-level cancellation, showing it explicitly.

We choose to work with a colored vector field  $V^a$  rather than in full QCD and projecting a quark–antiquark pair onto a spin-one color-octet state. We therefore introduce a pure-gluon  $SU(N)$  Lagrangian, to which we add a massive colored vector field  $V_\mu^a$ , in the adjoint representation of  $SU(N)$ , interacting with the gluons in a gauge invariant way:

$$\mathcal{L}_{\text{VF}} = -\frac{1}{4}V_{\mu\nu}^a V^{a\mu\nu} - \frac{1}{2}M^2 V_\mu^a V^{a\mu} - \frac{1}{4}F_{\mu\nu}^a F^{a\mu\nu} + \mathcal{L}_1. \quad (8)$$

Considering operators of dimension  $d = 4$ , there is only one operator that can contribute to the interactions:

$$\begin{aligned} \mathcal{L}_1 &= \frac{g'}{2}V_{\mu\nu}^a F^{a\mu\nu} \\ &= g' \partial_\mu V_\nu^a F^{a\mu\nu} - g g' f^{abc} A_\mu^c V_\nu^b (\partial^\mu A^{a\nu} - \partial^\nu A^{a\mu}). \end{aligned} \quad (9)$$

In this expression

$$V_{\mu\nu}^a \equiv D_\mu^{ab} V_\nu^b - D_\nu^{ab} V_\mu^b, \quad (10)$$

$D_\mu^{ab}$  denotes the covariant derivative in the adjoint representation

$$D_\mu^{ab} = \partial_\mu \delta^{ab} - g f^{abc} A_\mu^c, \quad (11)$$

<sup>3</sup> Operators like  $(\partial^\mu V_\mu)F_{\nu\rho}F^{\rho\nu}$  are trivial in the sense that they can be removed by using  $\partial^\mu V_\mu = 0$ .




Fig. 1. Polar ( $A_1$ ) and direct ( $A_2$ ) contributions to the  $V^a \rightarrow g^b g^c$  amplitude.

and

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f^{abc} A_\mu^b A_\nu^c. \quad (12)$$

Using the interaction term between  $V$  and the gluons introduced in Eq. (9) we can extract the relevant contributions from the two vertices directly relevant to our calculation,  $V \rightarrow g$  and  $V \rightarrow gg$ , and use them to evaluate the two amplitudes in Fig. 1.

From the first term in the r.h.s. of Eq. (9) we get, after integration by parts<sup>4</sup>

$$V^a \rightarrow g^a: \quad -g' M^2 \epsilon_\nu, \quad (13)$$

where  $M$  is the mass of  $V$ , whereas from the second term we get

$$V^a \rightarrow g^b g^c: \quad igg' f^{abc} (\epsilon_2 \cdot \epsilon_1) [\epsilon \cdot (k_1 - k_2)]. \quad (14)$$

In both cases, polarization vectors for legs that will eventually be external ones have already been included.

In order to complete the amplitude  $A_1$  in Fig. 1 we need to propagate the gluon with  $1/(k_1 + k_2)^2 = 1/M^2$  from the  $V$  state to the triple-gluon vertex. This leads to

$$A_1 = -igg' M^2 \epsilon_\nu \frac{1}{M^2} f^{abc} (k_1 - k_2)^\nu (\epsilon_1 \cdot \epsilon_2). \quad (15)$$

The amplitude  $A_2$  is instead given simply by the expression for the  $V \rightarrow gg$  vertex in Eq. (14). The sum  $A_1 + A_2$  shows a full cancellation.

In Ref. [3] the same conclusion was reached by computing explicitly the  $Q\bar{Q} \rightarrow gg$  amplitude with off-shell gluons after projecting the  $Q\bar{Q}$  pair onto an  $\ell = 0$  (i.e.  $S$ -wave, obtained by setting the relative momentum between  $Q$  and  $\bar{Q}$  equal to zero), spin 1, color-octet state, and later setting  $k_1^2 = k_2^2 = 0$ . The result found in Ref. [3] with off-shell gluons has the form

$$M^{bca} \sim f^{bca} \{ D (\epsilon_1 \cdot \epsilon_2) [\epsilon \cdot (k_1 - k_2)] + E [(\epsilon_1 \cdot \epsilon) (\epsilon_2 \cdot k_1) - (\epsilon_2 \cdot \epsilon) (\epsilon_1 \cdot k_2)] \} \quad (16)$$

with

$$D = E/2 = 1 + \frac{M^2}{k_1^2 + k_2^2 - M^2}. \quad (17)$$

In this expression the second term comes from the sum of the two amplitudes with a propagating quark, whereas the first one originates from the amplitude with the tri-linear gluon vertex. In the on-shell limit  $k_1^2 = k_2^2 = 0$  and one finds  $D = E = 0$ .

We have therefore shown that the vanishing of the amplitude for the decay of a massive color-octet vector bosons into two massless gluon is due to a cancellation between the diagram with the triple-gluon vertex and the rest of the amplitude, whose forms are themselves dictated by gauge invariance. When working in full QCD and with a projected  $Q\bar{Q}$  pair instead of the vector boson  $V$ ,

the cancellation takes place between the antisymmetric combination of the Abelian diagrams and the triple-gluon vertex diagram.

As we will show below, this delicate cancellation does not survive quantum corrections.

#### 4. Non-Abelian case: full analysis

We consider now the process where a massive color-octet vector state decays into two massless colored gluons,  $V^a \rightarrow g^b g^c$ , in full generality. Lorentz invariance and Bose symmetry lead to the matrix element

$$M^{bca} \sim \epsilon^\rho \epsilon_1^\mu \epsilon_2^\nu A_{\mu\nu\rho}^{bca} = f^{bca} \left\{ D (\epsilon_1 \cdot \epsilon_2) [\epsilon \cdot (k_1 - k_2)] + E [(\epsilon_1 \cdot \epsilon) (\epsilon_2 \cdot k_1) - (\epsilon_2 \cdot \epsilon) (\epsilon_1 \cdot k_2)] + \frac{F}{k_1 \cdot k_2} [\epsilon \cdot (k_1 - k_2)] (\epsilon_1 \cdot k_2) (\epsilon_2 \cdot k_1) \right\}, \quad (18)$$

where  $f^{abc}$  are the color-SU( $N$ ) structure constants. We have now denoted by  $\epsilon_1 \equiv \epsilon_1^{b*}(k_1)$ ,  $\epsilon_2 \equiv \epsilon_2^{c*}(k_2)$  and  $\epsilon \equiv \epsilon^{a*}(P)$  the polarizations of the massless colored gluons. Note the minus sign between  $k_1$  and  $k_2$  momenta in the  $D$  and  $F$  terms, as well as the minus sign in the  $E$  term, compared to Eq. (2). Since  $\epsilon \cdot (k_1 - k_2) \neq 0$ , the  $D$  and  $F$  terms cannot be dropped as was done in the Abelian case. An additional term proportional to the symmetric tensor  $d^{bca}$ , rather than to the antisymmetric one  $f^{bca}$ , is identical to the two-photon case considered previously, and will therefore not be discussed further.

In the following we exploit the BRST symmetry [13] of the gauge-fixed action to constrain the form of the amplitude in Eq. (18). We will show that, differently from the Abelian case, we cannot conclude that all its terms are simultaneously zero.

The tensor  $A_{\mu\nu\rho}^{bca}(k_1, k_2)$  can be readily extracted from Eq. (18):

$$A_{\mu\nu\rho}^{bca}(k_1, k_2) = f^{bca} \left[ D g_{\mu\nu} (k_1 - k_2)_\rho + E (g_{\mu\rho} k_{1\nu} - g_{\nu\rho} k_{2\mu}) + \frac{F}{k_1 \cdot k_2} (k_1 - k_2)_\rho k_{2\mu} k_{1\nu} \right]. \quad (19)$$

As shown in the Appendix, BRST invariance implies that this tensor satisfies

$$k_1^\mu \epsilon_2^\nu A_{\mu\nu\rho}^{bca}(k_1, k_2) = \epsilon_1^\mu k_2^\nu A_{\mu\nu\rho}^{bca}(k_1, k_2) = 0, \quad (20)$$

which, together with Eq. (19), gives

$$0 = k_1^\mu \epsilon_2^\nu A_{\mu\nu\rho}^{bca} = f^{bca} \left[ (D + E + F) (k_1 \cdot \epsilon_2) k_{1\rho} - (D + F) (k_1 \cdot \epsilon_2) k_{2\rho} - E (k_1 \cdot k_2) \epsilon_{2\rho} \right], \quad (21)$$

where we set  $k_1^2 = 0$ , i.e. on-shell gluon, and used the transversality condition  $k_2 \cdot \epsilon_2 = 0$ . From Eq. (21) we conclude that BRST invariance requires  $E = 0$  and  $D + F = 0$ . However the latter relation can be satisfied even if  $D \neq 0$  and  $F \neq 0$ .

As discussed in the Abelian case, once BRST invariance is enforced, we can work with transverse polarizations and hence we know that the  $E$  and  $F$  terms will not contribute to the amplitude because terms of the form  $\epsilon_1 \cdot k_2$  and  $\epsilon_2 \cdot k_1$  vanish in an appropriate frame. However, we also see that the  $D$  term can survive in the non-Abelian case, consistently with the non-vanishing results at one loop obtained in Refs. [10,11].

<sup>4</sup> We omitted null terms like  $\partial^\nu V_\nu^a = 0$  and terms proportional to  $k_1 \cdot \epsilon_2$  and  $k_2 \cdot \epsilon_1$  that do not contribute to the final amplitude because of the general transversality considerations made in Section 2.

The explicit tree-level calculation performed in Section 3 shows the cancellation of the  $V^a$  gluonic decay amplitude expected from (9). As done above for the Abelian case, it is interesting to understand this result directly in terms of the operators that can appear in the Lagrangian. The  $d = 4$  operator in (9) generates the Lorentz structures of the terms  $E$  and  $D$  in Eq. (18). By themselves, these terms do not contribute to the  $V^a \rightarrow g^b g^c$  decay, as we have seen. At the operator level, that result is immediate to see in analogy to what we did for the Abelian case. Simple integration by parts yields

$$\frac{g'}{2} V_{\mu\nu}^a F^{a\mu\nu} = -g' V_\nu^b (D_\mu^{ba} F^{a\mu\nu}), \quad (22)$$

and then, use of the equation of motion for the gluons to replace  $D_\mu^{ba} F^{a\mu\nu}$  by colored quark currents shows that the decay amplitude into on-shell gluons is zero.

The operator that can generate both  $D$  and  $F$  terms (in the combination  $D + F$ ), and thus contribute to the gluonic decay of  $V^a$ , is

$$\Delta\mathcal{L}_6 = f^{bca} V_{\mu\nu}^a F^{b\nu\rho} F_\rho^{c\mu}, \quad (23)$$

a  $d = 6$  operator that can be radiatively generated as a finite correction to the effective action already at 1-loop order. Consistently with the results of our previous discussion, attempts to reduce this operator to equations of motion or trivially vanishing terms, as we did for the Abelian operator (6), fail in this case due to the non-Abelian nature of the gauge symmetry: no expression like (7) exists in this case. Note that the additional  $d = 6$  operator,  $D^{ab\mu} V_{\mu\nu}^b D_\rho^{ac} F^{c\rho\nu}$  will not contribute to the two-gluon decay of  $V^a$ , as can be shown by direct use of the gluonic equation of motion. The relevance of these  $d = 6$  operators for technicolor phenomenology and this particular gluonic decay amplitude has been studied in [14,15].

## 5. Landau–Yang in $\ell = 1$ scattering states

Up to this point we have considered the decay of a colored massive vector particle. Now we wish to analyze the Landau–Yang selection rule for the  $Q(p_1)\bar{Q}(p_2) \rightarrow g(k_1)g(k_2)$  annihilation in  $P$ -wave, with quarks treated as spinless for simplicity. This configuration constitutes a stand-in for that of spin-1/2 quarks whose projection onto a given spin and angular momentum state leads to a  $J = 1$  vector state for the  $Q\bar{Q}$  pair.

We start by considering the amplitude

$$M = \epsilon_{1\mu}\epsilon_{2\nu} A^{\mu\nu}(k_1, k_2, p_1) = \langle k_1, \epsilon_1; k_2, \epsilon_2 | p_1, p_2 \rangle, \quad (24)$$

with on shell particles. The  $A^{\mu\nu}$  tensor can be decomposed in terms of form factors as

$$\begin{aligned} A^{\mu\nu}(k_1, k_2, p_1) = & A_1^{(\mp)} (p_1^\mu k_2^\nu \pm p_1^\nu k_1^\mu) + \\ & + A_2^{(\mp)} (p_1^\mu k_1^\nu \pm p_1^\nu k_2^\mu) + \\ & + A_3^{(-)} (k_1 \cdot k_2) g^{\mu\nu} + A_4^{(-)} k_1^\nu k_2^\mu + \\ & + A_5^{(-)} k_1^\mu k_2^\nu, \end{aligned} \quad (25)$$

where  $A_i$  are functions of the Mandelstam variables  $s = (p_1 + p_2)^2$  and  $t = (p_1 - k_1)^2$ . The form factors  $A_i^{(\pm)}$  are symmetric/antisymmetric with respect to  $k_1 \leftrightarrow k_2$  exchange. In Eq. (25) we assume that the gluon configuration is color-odd so that the final state is Bose-symmetric.

BRST identities imply that if we saturate Eq. (25) with  $k_{1\mu}\epsilon_{2\nu}$  we get

$$\begin{aligned} 0 = & k_{1\mu}\epsilon_{2\nu} A^{\mu\nu} = \\ = & [A_2^{(-)} + A_2^{(+)}] (k_1 \cdot p_1) (k_1 \cdot \epsilon_2) + \\ & + [A_2^{(-)} - A_2^{(+)}] (k_1 \cdot k_2) (p_1 \cdot \epsilon_2) + \\ & + [A_3^{(-)} + A_4^{(-)}] (k_1 \cdot k_2) (k_1 \cdot \epsilon_2). \end{aligned} \quad (26)$$

From this equation we deduce that  $A_2^{(-)} = A_2^{(+)} = 0$ , while due to the transversality of the gluon polarizations, the  $A_1^{(\mp)}$  and the  $A_5^{(-)}$  contributions to the physical amplitude vanish.

One can further see that, in the threshold limit,  $\mathbf{p}_1 = \mathbf{p}_2 = 0$ , we have  $A_i^{(-)} = 0$  because of antisymmetry, and therefore  $A_{3,4}^{(-)} = 0$  at threshold. However this is not sufficient to imply the Landau–Yang selection rule. In fact the initial state  $|p_1, p_2\rangle$  is to be projected onto the  $P$ -wave in the center of mass according to

$$|p_1, p_2; \ell = 1, m\rangle = \int d\Omega_{\hat{\mathbf{p}}_1} Y_m^{\ell=1}(\hat{\mathbf{p}}_1) |p_1, p_2\rangle, \quad (27)$$

where  $\mathbf{p}_1 + \mathbf{p}_2 = 0$  and  $\hat{\mathbf{p}}_1 = \mathbf{p}_1 / \|\mathbf{p}_1\|$  is the unit vector along  $\mathbf{p}_1$ , and the transition amplitude is given by

$$M \propto \epsilon_{1\mu}\epsilon_{2\nu} \int d\Omega_{\hat{\mathbf{p}}_1} Y_m^{\ell=1}(\hat{\mathbf{p}}_1) A^{\mu\nu}. \quad (28)$$

The  $P$ -wave condition selects contributions to  $A^{\mu\nu}$  that overall contain one power of  $p_1$  which originates from the  $t$  dependence of  $A_i$ 's which, close to threshold, we parameterize as

$$A_i^{(-)} \simeq (\mathbf{p}_1 \cdot \mathbf{k}_1) B_i, \quad (29)$$

with  $B_i$  constrained by (26). Replacing in Eq. (28) we find

$$\begin{aligned} M \propto & B_3 \epsilon_{1\mu}\epsilon_{2\nu} g^{\mu\nu} \int d\Omega_{\hat{\mathbf{p}}_1} Y_m^{\ell=1}(\hat{\mathbf{p}}_1) (\mathbf{p}_1 \cdot \mathbf{k}_1) \propto \\ \propto & B_3 (\epsilon_1 \cdot \epsilon_2) k_1^m, \end{aligned} \quad (30)$$

where  $\sqrt{2}k_1^{\pm 1} = k_{1x} \mp ik_{1y}$ ,  $k_1^0 = k_{1z}$  and we used Eq. (5).

The result in Eq. (30) shows that the form of the amplitude for the process is very constrained. However, one cannot say anything about the value of  $B_3$ , because the BRST identity in Eq. (26) only allows one to conclude that the combination  $B_3 + B_4$  must vanish. This is the same kind of roadblock that was met in Section 4, and it shows that the Landau–Yang selection rule cannot be extended to non-Abelian gluons also in the case in which the initial state is not a single particle but a scattering continuum.

## 6. Conclusions

In this note we have considered the process where a massive, color-octet vector state decays into two on-shell massless gluons. The well-known result that this amplitude vanishes at tree level in QCD has been shown in general terms to be due to a cancellation between the color-antisymmetric, Abelian part of the amplitude and its non-Abelian part.

Using considerations of Lorentz invariance and Bose symmetry we have also written down the most general expression for this amplitude, and employed BRST invariance to constrain its form. We have shown that, at variance with the Abelian case, one cannot conclude that the amplitude vanishes to all orders. This is consistent with recent evidence that the amplitude does indeed not vanish at one-loop. We have explained this result in terms of the emergence of higher-dimension operators, radiatively generated in the effective action already at one-loop order. In addition, we offer a novel way of understanding the Landau–Yang theorem (or its failure) directly in terms of the relevant operators, manipulated by integration by parts or using field equations of motion. We

have also considered in detail a situation where the massive vector state is given by the projection of a spinless quark–antiquark pair onto an  $\ell = 1$  angular momentum state, reaching the same conclusions.

## 7. Note added

While our note was being finalized Ref. [16] appeared, also discussing the failure of the Landau–Yang theorem in a non-Abelian gauge theory and reporting that an explicit calculation shows that the  $(Q\bar{Q})_{J=1}^{(8)} \rightarrow gg$  amplitude at one loop is non-vanishing. This paper also prompted a revised version of Ref. [17] that now features a discussion of the Landau–Yang theorem in QCD.

## Acknowledgements

A.D.P. thanks the CERN Theory group for hospitality while this work was performed. M.C. is supported in part by the ILP LABEX (ANR-10-LABX-63) financed by French state funds managed by the ANR within the Investissements d’Avenir programme under reference ANR-11-IDEX-0004-02. L.D.D. is supported by STFC, grant ST/L000458/1, and the Royal Society, Wolfson Research Merit Award, grant WM140078. The work of J.R.E. is supported by the grants FPA2013-44773-P and FPA2014-55613-P from the Spanish Ministry for Economy and Competitiveness (MINECO), by the grant 2014SGR1450 from the Generalitat de Catalunya and by the Severo Ochoa excellence program of MINECO under the grant SO-2012-0234.

## Appendix A

We now detail the derivation of the identities in Eq. (20). The BRST transformations are defined by

$$\delta_{\text{BRST}} A_\mu^a \equiv \theta \delta A_\mu^a = \theta (D_\mu)^{ab} c^b, \quad (31)$$

$$\delta_{\text{BRST}} B^a \equiv \theta \delta B^a = 0, \quad (32)$$

$$\delta_{\text{BRST}} \bar{c}^a \equiv \theta \delta \bar{c}^a = \theta B^a, \quad (33)$$

$$\delta_{\text{BRST}} c^a \equiv \theta \delta c^a = -\frac{1}{2} g \theta f^{abc} c^b c^c, \quad (34)$$

where  $\theta$  is an anticommuting parameter and  $B^a$  are Lagrange multipliers enforcing the gauge fixing conditions. The theory that we are considering also contains a colored massive vector field  $V_\mu^a(x)$ , whose BRST variation is given by

$$\delta V_\mu^a(x) = -\frac{g}{2} f^{abc} c^b V_\mu^c. \quad (35)$$

The Lagrangian density for the gauge-fixed theory,

$$\mathcal{L} = \mathcal{L}_{\text{VF}} - \delta (\bar{c}^a \partial_\mu A^{a\mu}) - \frac{1}{2\alpha} B^a B^a, \quad (36)$$

satisfies  $\delta \mathcal{L} = 0$  and therefore the expectation value of any BRST variation vanishes,

$$\langle \delta O \rangle = 0, \quad (37)$$

where  $\langle \dots \rangle$  denotes the vacuum  $T$ -product.

In particular, from

$$\begin{aligned} \delta (\bar{c}^a(x) A_\nu^b(y) V_\rho^c(z)) &= \\ &= B^a(x) A_\nu^b(y) V_\rho^c(z) + \bar{c}^a(x) (D_\nu)^{bd} c^d(y) V_\rho^c(z) + \\ &\quad - \frac{g}{2} f^{cde} \bar{c}^a(x) A_\nu^b(y) c^d(z) V_\rho^e(z), \end{aligned} \quad (38)$$

we have

$$\begin{aligned} 0 &= \langle T(B^a(x) A_\nu^b(y) V_\rho^c(z)) \rangle + \\ &\quad + \langle T(\bar{c}^a(x) (D_\nu)^{bd} c^d(y) V_\rho^c(z)) \rangle + \\ &\quad - \frac{g}{2} f^{cde} \langle T(\bar{c}^a(x) A_\nu^b(y) c^d(z) V_\rho^e(z)) \rangle. \end{aligned} \quad (39)$$

Integrating over the Nakanishi–Lautrup field  $B^a$  yields:

$$\begin{aligned} \langle T(B^a(x) A_\nu^b(y) V_\rho^c(z)) \rangle &= \\ &= \alpha \partial_x^\mu \langle T(A_\mu^a(x) A_\nu^b(y) V_\rho^c(z)) \rangle, \end{aligned} \quad (40)$$

hence

$$\begin{aligned} \alpha \partial_x^\mu \langle T(A_\mu^a(x) A_\nu^b(y) V_\rho^c(z)) \rangle &= \\ &= -\langle T(\bar{c}^a(x) (D_\nu)^{bd} c^d(y) V_\rho^c(z)) \rangle + \\ &\quad + \frac{g}{2} f^{cde} \langle T(\bar{c}^a(x) A_\nu^b(y) c^d(z) V_\rho^e(z)) \rangle. \end{aligned} \quad (41)$$

The tensor  $A_{\mu\nu\rho}^{bca}$  in Eq. (18) is given by the LSZ reduction formalism in the form

$$\begin{aligned} A_{\mu\nu\rho}^{bca}(k_1, k_2) &\propto \\ &\propto \lim_{(k_1+k_2)^2 \rightarrow M^2} \lim_{k_1^2 \rightarrow 0} \lim_{k_2^2 \rightarrow 0} k_1^2 k_2^2 [(k_1+k_2)^2 - M^2] \times \\ &\quad \int dx dy \langle T(A_\mu^a(x) A_\nu^b(y) V_\rho^c(0)) \rangle e^{ik_1 x} e^{ik_2 y}. \end{aligned} \quad (42)$$

Eqs. (41) and (42) imply

$$\begin{aligned} k_1^\mu A_{\mu\nu\rho}^{abc}(k_1, k_2) &\propto \\ &\propto \lim_{(k_1+k_2)^2 \rightarrow M^2} \lim_{k_1^2 \rightarrow 0} \lim_{k_2^2 \rightarrow 0} k_1^2 k_2^2 [(k_1+k_2)^2 - M^2] \times \\ &\quad \int dx dy \langle T(\bar{c}^a(x) (D_\nu)^{bd} c^d(y) V_\rho^c(0)) \rangle e^{ik_1 x} e^{ik_2 y}, \end{aligned} \quad (43)$$

where the second term in the r.h.s. of Eq. (41) does not contribute because there are no single particle poles in the channel of the composite operator  $f^{cde} c^d(z) V_\rho^c(z)$ . As for the first term, the massless ghost can contribute a term proportional to the momentum  $k_{2\nu}$  in the channel of the operator  $(D_\nu)^{bd} c^d(y)$ . Eq. (43) can therefore be written as

$$k_1^\mu A_{\mu\nu\rho}^{abc}(k_1, k_2) \propto k_{2\nu} A_\rho^{abc}, \quad (44)$$

so that

$$k_1^\mu \epsilon_2^\nu \epsilon^\rho A_{\mu\nu\rho}^{abc}(k_1, k_2) = (k_2 \cdot \epsilon_2) \epsilon^\rho A_\rho^{abc} = 0, \quad (45)$$

because  $k_2 \cdot \epsilon_2 = 0$ . This proves Eq. (20).

## References

- [1] L.D. Landau, Dokl. Akad. Nauk Ser. Fiz. 60 (1948) 207; L.D. Landau, Phys. Abstr. A 52 (1949) 125.
- [2] C.N. Yang, Phys. Rev. 77 (1950) 242.
- [3] P.L. Cho, A.K. Leibovich, Phys. Rev. D 53 (1996) 150, arXiv:hep-ph/9505329.
- [4] W.K. Tang, M. Vanttinen, Phys. Rev. D 53 (1996) 4851, arXiv:hep-ph/9506378.
- [5] J.P. Ma, Nucl. Phys. B 460 (1996) 109, arXiv:hep-ph/9510333.
- [6] W.K. Tang, M. Vanttinen, Phys. Rev. D 54 (1996) 4349, arXiv:hep-ph/9603266.
- [7] A. Petrelli, M. Cacciari, M. Greco, F. Maltoni, M.L. Mangano, Nucl. Phys. B 514 (1998) 245, arXiv:hep-ph/9707223.
- [8] W. Beenakker, et al., J. High Energy Phys. 1310 (2013) 120, arXiv:1304.6354 [hep-ph].
- [9] P. Bärnreuther, M. Czakon, P. Fiedler, J. High Energy Phys. 1402 (2014) 078, arXiv:1312.6279 [hep-ph].
- [10] R.S. Chivukula, A. Farzinia, J. Ren, E.H. Simmons, Phys. Rev. D 87 (9) (2013) 094011, arXiv:1303.1120 [hep-ph].

- [11] J.P. Ma, J.X. Wang, S. Zhao, *Phys. Lett. B* 737 (2014) 103, arXiv:1405.3373 [hep-ph].
- [12] S.R. Coleman, J. Wess, B. Zumino, *Phys. Rev.* 177 (1969) 2239; C. Arzt, *Phys. Lett. B* 342 (1995) 189, arXiv:hep-ph/9304230.
- [13] C. Becchi, A. Rouet, R. Stora, *Ann. Phys.* 98 (2) (1976) 287–321; I.V. Tyutin, Lebedev Institute preprint (1975, unpublished).
- [14] R.S. Chivukula, A. Grant, E.H. Simmons, *Phys. Lett. B* 521 (2001) 239, arXiv:hep-ph/0109029.
- [15] Y. Bai, W.Y. Keung, *Int. J. Mod. Phys. A* 30 (2015) 1550120, arXiv:1407.6355 [hep-ph].
- [16] W. Beenakker, R. Kleiss, G. Lustermsans, arXiv:1508.07115 [hep-ph].
- [17] V. Pleitez, arXiv:1508.01394 [hep-ph].