



Emergence of ITD tuning in the MSO with a realistic periphery model

Pierre Yger, Victor Benichoux, M Stimberg, R Brette

► To cite this version:

Pierre Yger, Victor Benichoux, M Stimberg, R Brette. Emergence of ITD tuning in the MSO with a realistic periphery model. BMC Neuroscience, 2015, 16 (Suppl 1), pp.P20 10.1186/1471-2202-16-S1-P20 . hal-01250244

HAL Id: hal-01250244

<https://hal.sorbonne-universite.fr/hal-01250244>

Submitted on 4 Jan 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

POSTER PRESENTATION

Open Access

Emergence of ITD tuning in the MSO with a realistic periphery model

P Yger^{1,2,3,4*}, V Benichoux^{1,2,3,4}, M Stimberg^{1,2,3,4}, R Brette^{1,2,3,4}

From 24th Annual Computational Neuroscience Meeting: CNS*2015
Prague, Czech Republic. 18-23 July 2015

To localize sounds in the environment, animals mostly rely on spectro-temporal cues originating from the physical disparities of the sound waveforms impacting the two ears. Among those, the Interaural Time Difference (ITD) has been shown to be crucial in mammals for locating low-frequency sounds, and is known to be processed by neurons in a particular structure, the Medial Superior Olive (MSO). While it is classically considered that the emergence of ITD selectivity in a neuron of the MSO is simply due to differences in the axonal delays originating from the two ears and impinging those neurons (the so called “delay-line” model [1]), experimental evidence shows that the best delay (the ITD at which the neuron’s firing rate is maximum) is also dependent on the frequency of the sound [2].

To investigate more on this challenging experimental observation, we developed a realistic periphery model to mimic cochlear inputs from auditory nerves fibers onto the MSO. Using known plasticity rules such as Spike Timing Dependent Plasticity to structure the wiring from those connections onto the MSO, we extend the work that has already been performed [3], and study the emergence of binaural tuning in the MSO in a realistic scenario. The system is trained with binaural sounds such as white noise, and then, as in most experimental papers, we tested the ITD selectivity of cells in the MSO by presenting pure tones at various frequencies.

Finally, we discuss, from a coding point of view the potential implications raised by the frequency dependence of the best delay. As pointed out by recent work [4], with such a frequency-dependent best delay, neurons in the MSO should be seen as coding for a

particular position in space, rather than for just a fixed delay difference.

Acknowledgment

This work is funded with a Fellowship Grant from the FRM (Fondation pour la Recherche Médicale), and the European Research Council (StG 240132)

Authors' details

¹Institut d'Etudes de la Cognition, Ecole Normale Supérieure, Paris, France.
²Sorbonne Universités, UPMC Univ. Paris 06, UMR S 968, Institut de la Vision, Paris, F-75012, France. ³INSERM, U968, Paris, F-75012, France. ⁴CNRS, UMR 7210, Paris, F-75012, France.

Published: 18 December 2015

References

- Jeffress LA: A place theory of sound localization. *J Comp Physiol Psychol* 1948, **41**(1):35-39.
- Joris P, Yin TC: A matter of time: internal delays in binaural processing. *Trends Neurosci* 2007, **30**(2):70-78.
- Fontaine B, Brette R: Neural Development of Binaural Tuning through Hebbian Learning Predicts Frequency-Dependent Best Delays. *J Neurosci* 2011, **31**(32):11692-11696.
- Benichoux V, Fontaine B, Karino S, Joris P, Brette R: Frequency-dependent time differences between the ears are matched in neural tuning (in revision).

doi:10.1186/1471-2202-16-S1-P20

Cite this article as: Yger et al.: Emergence of ITD tuning in the MSO with a realistic periphery model. *BMC Neuroscience* 2015 **16**(Suppl 1):P20.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: pierre.yger@inserm.fr

¹Institut d'Etudes de la Cognition, Ecole Normale Supérieure, Paris, France
Full list of author information is available at the end of the article