



HAL
open science

Photosynthetic pigments of oceanic Chlorophyta belonging to prasinophytes clade VII

Adriana Lopes dos Santos, Priscillia Gourvil, Francisco Rodríguez, José Luis Garrido, Daniel Vaultot

► **To cite this version:**

Adriana Lopes dos Santos, Priscillia Gourvil, Francisco Rodríguez, José Luis Garrido, Daniel Vaultot. Photosynthetic pigments of oceanic Chlorophyta belonging to prasinophytes clade VII. *Journal of Phycology*, 2015, 52 (1), pp.148-155. 10.1111/jpy.12376 . hal-01258694

HAL Id: hal-01258694

<https://hal.sorbonne-universite.fr/hal-01258694>

Submitted on 19 Jan 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Received Date : 25-Apr-2015
Revised Date : 17-Sep-2015
Accepted Date : 13-Nov-2015
Article type : Note

Article Type: Research Note

Photosynthetic pigments of oceanic Chlorophyta belonging to prasinophytes clade VII.

Adriana Lopes dos Santos¹, Priscillia Gourvil¹, Francisco Rodríguez², José Luis Garrido³, and Daniel Vaultot¹¹.

¹Sorbonne Universités, UPMC Univ. Paris 06, CNRS, UMR 7144, Station Biologique, Place Georges Teissier, 29680 Roscoff, France

²Instituto Español de Oceanografía (IEO), Centro Oceanográfico de Vigo, Subida a Radio Faro 36390 Vigo, España.

³Instituto de Investigaciones Marinas(CSIC). Av. Eduardo Cabello, 6. 36208 Vigo, España.

Editorial Responsibility: K. Valentin (Associate Editor)

Revised version

Submitted to Journal of Phycology, September 18, 2015

Abstract

The ecological importance and diversity of pico/nano-planktonic algae remains poorly studied in marine waters, in part because many are tiny and without distinctive morphological features. Amongst green algae, Mamiellophyceae such as *Micromonas* or *Bathycoccus* are dominant in coastal waters while prasinophytes clade VII, yet not formerly described, appear

¹ Author for correspondence : vaultot@gmail.com

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/jpy.12376-15-065

to be major players in open oceanic waters. The pigment composition of 14 strains representative of different sub-clades of clade VII was analyzed using a method that improves the separation of loroxanthin and neoxanthin. All the prasinophytes clade VII analyzed here showed a pigment composition similar to that previously reported for RCC287 corresponding to pigment group prasino-2A. However we detected in addition astaxanthin for which it is the first report in prasinophytes. Among the strains analyzed the pigment signature are qualitatively similar within sub-clades A and B. In contrast, RCC3402 from sub-clade C (*Picocystis*) lacks loroxanthin, astaxanthin and antheraxanthin, but contains alloxanthin, diatoxanthin and monadoxanthin, that are usually found in diatoms or cryptophytes. For sub-clades A and B, loroxanthin was lowest at highest light irradiance suggesting a light-harvesting role of this pigment in clade VII as in *Tetraselmis*.

Keywords : phytoplankton, picoplankton, prasinophytes, pigments, HPLC

The paraphyletic group of prasinophytes is an assemblage of free-living unicellular microalgae present in both marine and freshwater habitats (Leliaert et al. 2012). Molecular phylogenetic, ultra-structural, and biochemical approaches have helped taxonomists to re-organize gradually the group into new classes and clades (Guillou et al. 2004, Marin and Melkonian 2010, Subirana et al. 2013, Lemieux et al. 2014a). Currently the prasinophytes are divided into nine groups known as clades I to IX, based on phylogenetic analyses of the nuclear 18S (nuclear-encoded small subunit rRNA) gene (Fawley et al. 2000, Guillou et al. 2004, Viprey et al. 2008). These clades may correspond to true classes, or be composed of a small number of species or of environmental sequences only. For example, Chlorodendrophyceae (Massjuk 2006) known previously as prasinophytes clade IV was

recently raised to the class level and added to the “core of chlorophytes” (Fucikova et al. 2014). Clade V corresponds to the order Pycnococaceae with two major species, *Pseudoscourfieldia marina* and *Pycnococcus provasolii* which are probably two forms of a single life cycle (Fawley et al. 1999, Guillou et al. 2004). Clades VIII and IX are composed entirely by environmental sequences without representatives in culture (Viprey et al. 2008). Clade II, previously corresponding to the order Mamiellales, was raised recently to the class level as Mamiellophyceae (Marin and Melkonian 2010) and contains three important genera of marine pico-phytoplankton: *Micromonas* (Butcher 1952), *Bathycoccus* (Eikrem and Throndsen 1990) and *Ostreococcus* (Chrétiennot-Dinet et al. 1995).

In coastal waters, Mamiellophyceae appear largely dominant, especially within the pico-plankton, with the genus *Micromonas* making the highest contribution and followed to a lesser extent by *Bathycoccus* (Throndsen, J. and Kristiansen 1991, Not et al. 2004, Collado-Fabri et al. 2011, Balzano et al. 2012). In contrast in the open ocean, another group of prasinophytes, clade VII, with cell size in the 3 to 5 μm range, has been found to make an important contribution to the pico-plankton community in regions such as the Equatorial Pacific and Mediterranean Sea (Moon-van der Staay et al. 2000, Viprey et al. 2008, Shi et al. 2009). The distribution of clade VII in typically oceanic mesotrophic waters makes this an interesting group. Prasinophyte clade VII contains several cultured strains mostly from tropical and sub-tropical waters but also from temperate regions. Although it has not been described formerly yet. Guillou et al. (2004) divided this group into three well-supported sub-clades, A, B and C, the latter being formed by *Picocystis salinarum*, a small species found in saline lakes (Lewin et al. 2000, Roesler et al. 2002, Krienitz et al. 2012).

Traditionally, pigment signature has been used to determine the taxonomy of algae groups present in the water column (Jeffrey et al. 2011). This approach has been largely superseded by molecular approaches (Liu et al. 2009) but pigments remain an important

phenotypic characteristic that allowed to point out the importance of green algae in specific regions of Pacific Ocean, Mediterranean Sea or Arctic Ocean (Obayashi and Tanoue 2002, Miki et al. 2008, Gutiérrez-Rodríguez et al. 2010, Coupel et al. 2014). The study of pigments in different types of prasinophytes has revealed a diversity of photosynthetic signatures in this group. Prasinophytes can be divided into three major groups based on their carotenoid composition (Egeland et al. 1997, Garrido et al. 2009). Group 1 contains the basic set of carotenoids present in Chlorophyceae: neoxanthin, violaxanthin, lutein, zeaxanthin, antheraxanthin and β - β -carotene. Group 2 consists of the basic set of carotenoids plus loroxanthin (2A) and siphonaxanthin (2B). Group 3 contains prasinoxanthin (3A) and uriolide, micromonal, micromonol and dihydrolutein (3B) in addition to the main pigments found in group 1 (Jeffrey et al. 2011).

Within clade VII, only three strains have been analyzed until now: two isolates of *Picocystis salinarum* (subclade C) from saline lakes (Lewin et al. 2000, Roesler et al. 2002) and the marine strain RCC287 (subclade A; Latasa et al. 2004). A large number of clade VII strains are available from the Roscoff Culture Collection (<http://roscoff-culture-collection.org/>) originating from a range of environment. The aim of this study was to determine the phenotypic characteristics of this important group of marine green algae by analyzing the pigment composition of fourteen strains belonging to the three sub-clades (A, B, C) of prasinophytes clade VII isolated from a range of oceanic location and depths (Table 1). We also assessed the effect of three light irradiances on pigment composition for a subset of these strains.

Twelve strains belonging to clade VII (Table 1) were grown at 22°C in 25 cm² culture flasks with 50 ml of K seawater medium (Keller et al. 1987) under 140 $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ in continuous light. Two other strains, added later, were grown under the same conditions except for light (100 $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ in 12:12 Light:Dark cycle). A subset

of nine strains was also grown at two other light levels (14 and 65 $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$). All strains were acclimated to the light conditions during at least five generations. Prior to sample collection, cell concentration was determined by flow cytometry using a Becton Dickinson Accuri C6. Approximately 50 ml of cultures were collected in late exponential or early stationary phase by filtration onto glass fiber GF/F filters (Whatman, Maidstone, UK) without vacuum. Total time for filtration did not exceed 10 min and filters were removed as soon as the passage of liquid through it was undetectable. Total volume filtered was recorded. Filters were protected from light at all processing stages, immediately frozen in liquid nitrogen and stored at -80°C . Pigments were analyzed within one month. Frozen filters were extracted with 3 mL of 90% acetone in screw cap glass tubes with polytetrafluoroethylene (PTFE) lined caps, placed in an ice-water bath. After 15 min, filters were homogenized using a stainless steel spatula for filter grinding. Tubes were placed in an ultrasonic bath with water and ice for 5 min. The slurries were then centrifuged 5 minutes at 4,500 r.p.m. and supernatants filtered through 13 mm diameter polypropylene syringe filters (MS PTFE, 0.22 μm pore size) to remove cell and filter debris. Before injection 1 mL of each sample extract was added with 0.4 mL of Milli-Q water to avoid peak distortion. Pigments extracted from clade VII strains were analyzed using a modification of Zapata et al. (2000) method, described by Garrido et al. (2009) to improve the separation of loroxanthin and neoxanthin (Table S1 in the Supporting Information). Pigment extracts of RCC3402 (*Picocystis*) were also analyzed employing a polymeric octadecyl silica column as described by Garrido and Zapata (1997). All graphs and analyses were performed with the R software using the ggplot2 and FactoMineR libraries (R Development Core Team 2013).

Intracellular chlorophyll (Chl) *a* content ranged from 4 to 26 fg per cell in most strains except for RCC996 (VIIA) and RCC3402 (*Picocystis* - clade VIIC) for which it was much higher (Table 2). This range agreed with values previously determined for marine

microalgae in the same size range (Simon et al. 1994). More recently, in a field survey, Giovagnetti et al (2013) found 20-60 fg per cell in nanophytoplankton (>3 μm). Brunet et al (2006) estimated a range of 17-168 fg per cell in picoeukaryotes from the DCM and finally, DuRand et al. (2002) as well as Not et al. (2004) reported 25 fg per cell in the picoplanktonic species *Micromonas pusilla*.

All the prasinophytes clade VIIA and B analyzed here showed a very similar pigment composition (Table 2). It did not seem to change drastically between sub-clades A and B, nor with the depth of isolation (Fig. 1). This composition is similar to that reported for RCC287 by Latasa et al. (2004) corresponding to pigment group prasino-2A. We did not observe strong differences (Fig. 1; Table 2) with the data of Latasa et al. (2004): in particular the ratios obtained for zeaxanthin and lutein were very similar in both studies despite the slight difference in light levels (100 vs. 140 $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$ in our study): zeaxanthin, 0.042 (w/w) versus 0.043 (w/w) and lutein, 0.382 (w/w) versus 0.363 (w/w). However their study used a less resolutive method and did not report the presence of loroxanthin and astaxanthin in RCC287. For loroxanthin this is probably due to the co-elution of this pigment with neoxanthin in the analytic method employed by these authors.

In our study, only RCC1124 and RCC1871 (both from sub-clade A) did not contain loroxanthin within strains belonging sub-clades A and B (Table 2). Violaxanthin and lutein were the most abundant carotenoids for sub-clades A and B. Astaxanthin came as third for most other A and B strains except for RCC1871 (sub-clade A) and RCC2339 (sub-clade B) for which it was neoxanthin and β - β -carotene, respectively. *Picocystis* (RCC3402, clade VIIC) had a clearly distinct carotenoid profile compared to sub-clades A and B. It did not contain loroxanthin, astaxanthin and antheraxanthin but instead diatoxanthin, alloxanthin and monadoxanthin (Fig. 1; Table 2). For this strain, β - β -carotene, monadoxanthin and diatoxanthin were the most abundant carotenoids, respectively (Fig. 1; Table 2) and the ratio

of accessory pigments to Chl *a* was much lower than in clades VIIA and B (Fig. 1). The presence in *Picocystis* of these pigments usually found in cryptophytes or diatoms (Takaichi 2011), that are parts of the so-called red lineage (by opposition to the green lineage to which clade VII belongs, Falkowski et al. 2004) was also reported by Lewin et al. (2000) and Roesler et al. (2002) as well as found in *Coccomyxa*, a green alga belonging to the Chlorophyceae (Crespo et al. 2009).

We analyzed the influence of irradiance (14, 65 and 140 $\mu\text{mol photons} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$) on pigment composition of nine strains of prasinophytes VIIA and B (Fig. 2; Table S2 in the Supporting Information). Accessory chlorophylls and carotenoids involved in light harvesting tend to increase relative to Chl *a* at low light, while photoprotective carotenoids increase at high light (Schlüter et al. 2000, Henriksen et al. 2002, Brunet et al. 2011a). In our study, Chl *b* ratios increased slightly at low light, as expected, except for RCC3376 that showed a very slightly lower ratio at low light than at high light (0.78 vs. 0.81; Fig. 2; Table S2). A similar slight decrease was also observed by Garrido et al. (2009) for the green alga *Tetraselmis suecica*.

The increase at low light of neoxanthin, β - ϵ carotene and loroxanthin points to a light harvesting role for these pigments in most of the strains (Fig. 2; Table S2). The changes can be subtle, as in the case of neoxanthin or drastic, as observed for loroxanthin (Fig. 2). Neoxanthin has been found to be associated with light harvesting complexes in the Mamiellophyceae *Mantoniella squamata* (Wilhelm and Lenarz-Weiler 1987). A major light harvesting role could be suggested for loroxanthin in clades VII A and B in agreement with what observed Garrido et al. (2009) in another Chlorophyta *Tetraselmis*. Interestingly, two strains lacking loroxanthin (RCC1124 and RCC1871) have been isolated from temperate North Atlantic Ocean waters, in contrast to the other strains from subclade A which originate from tropical waters (Table 1).

The increase of astaxanthin (from 2 to 4-fold depending on the strains) with light intensity suggests that this carotenoid has a photoprotective role (Fig. 2), as previously demonstrated in the Chlorophyceae *Haematococcus pluvialis* (Wang et al. 2003, Gao et al. 2012). Among all strains, RCC3374 showed the most impressive accumulation of astaxanthin which contributed up to 42% of the total carotenoid pool under high light conditions (Fig. 2). In comparison, *H. pluvialis* can accumulate 86 - 90% of astaxanthin in the total carotenoid pool after sixteen days cultures of under stress conditions (Sarada et al. 2002).

The photoprotective role attributed to lutein (Jahns and Holzwarth 2012) seems to happen also in these species. Its contribution to total carotenoids increased sharply from low to medium light and stabilized at the highest irradiance (Fig. 2; Table S2). Such increase under high light conditions has been previously reported by Böhme et al. (2002) in the Mamiellophyceae *M. squamata*. These authors suggested that lutein played an important role as intermediate of biosynthesis for light harvesting pigments after light shifts from HL to LL. This role was coherent with its loose binding to the LHC apoprotein, also observed for the violaxanthin cycle (VAZ) carotenoids. However, lutein and loroxanthin are xanthophylls derived from β - ϵ carotene, and both have also been suggested also to take part in photoprotective mechanisms (non-photochemical quenching, NPQ) to prevent photo-oxidative damage in high light conditions in the green alga *Chlamydomonas reinhardtii* (Niyogi et al. 1997).

As for lutein, the content of the photoprotective xanthophyll cycle involving violaxanthin, antheraxanthin and zeaxanthin (VAZ cycle) relative to Chl *a* increased from low to medium light and then stabilized (Fig. 2; Table S2). However the evolution of individual pigments differed among strains. For example, zeaxanthin did not change much in RCC287 and RCC857 while it increased several-fold in other strains (e.g., RCC719, Fig. 2).

A relationship between strain origin and pigment composition is unlikely according to a Principal Component Analysis (PCA) based on pigments to Chl *a* ratios (Fig. 3). The first two components explained more than 50% (dimension 1 and 2, 33.1 % and 20.4 %, respectively) of the variance. Pigments contributing positively to dimension 1 included some which may have a photoprotective role (lutein, zeaxanthin, antheraxanthin and astaxanthin) while pigments suggested to be involved in light harvesting, such as loroxanthin contributed negatively to this axis. Pigments with moderate response to light, such as Chl *b* and neoxanthin, contributed to dimension 2. Strains distributed along dimension 1 according to the light treatment, irrespectively of their sub-clade, latitude or depth of isolation (surface vs. DCM). The use of HPLC data to assess the role of individual pigments as light-harvesting or photoprotective must be considered with caution. Photoacclimation processes operate at different scales (from seconds to several days) and pigment changes are influenced by multiple factors (genetics, ecology, physiology). Despite all this, some common patterns can be found when pigment data are given in terms of their ratios to Chl *a*. Light-harvesting pigments and Chl *a* content increase under low irradiance, and tend to co-vary under variable light conditions. In turn, photoprotective pigments are synthesized under light stress and increase their ratios to Chl *a* in higher light irradiance (Brunet et al. 2011b). The behavior of pigments analyzed in clade VII resembled that expected for light-harvesting or photoprotective ones, but without a more complete dataset (biochemistry, photosynthetic dynamics, etc.) this cannot be stated unambiguously. The discovery of loroxanthin (a putative light harvesting pigment) and astaxanthin (with a suggested photoprotective role) in prasinophytes clades VIIA and B prompts the need to reexamine the pigment composition of other members of this diverse and ancient group using improved analytical protocols.

Recent phylogenetic results pointed clade VII A/B as a sister group of the core of Chlorophyta (Guillou et al. 2004, Leliaert et al. 2012, Lemieux et al. 2014a, 2014b). The presence of astaxanthin as in core Chlorophyta while it is absent in other prasinophytes may reflect another common feature between clade VII and core Chlorophyta. Moreover while Guillou et al. (2004) included *Picocystis* into clade VII based on the phylogenetic analysis of 18S rRNA gene, the recent analysis of chloroplast genomes (Lemieux et al. 2014a) has shown widely divergent traits between *Picocystis* and sub-clade VIIA. The divergent carotenoid composition of *Picocystis* (absence of loroxanthin, astaxanthin, and antheraxanthin, and confirmation of the presence of red lineage pigments such as diatoxanthin and monadoxanthin) reinforce these phylogenetic analyses and point out the interest of pigments as phenotypic markers.

Acknowledgments

Financial support for this work was provided by the European Union program MaCuMBA (FP7-KBBE-2012-6-311975). This work is a contribution of Microalgas Nocivas (IEO), Unidad Asociada al IIM (CSIC). We acknowledge the constructive remarks of two referees that helped improving our manuscript.

References

- Balzano, S., Marie, D., Gourvil, P. & Vaulot, D. 2012. Composition of the summer photosynthetic pico and nanoplankton communities in the Beaufort Sea assessed by T-RFLP and sequences of the 18S rRNA gene from flow cytometry sorted samples. *ISME J.* 6:1480–98.
- Böhme, K., Wilhelm, C. & Goss, R. 2002. Light regulation of carotenoid biosynthesis in the prasinophycean alga *Mantoniella squamata*. *Photochem. Photobiol. Sci.* 1:619–28.
- Brunet, C., Casotti, R., Vantrepotte, V., Corato, F. & Conversano, F. 2006. Picophytoplankton diversity and photoacclimation in the Strait of Sicily (Mediterranean Sea) in summer. I. Mesoscale variations. *Aquat. Microb. Ecol.* 44:127–41.

- Brunet, C., Johnsen, G., Lavaud, J. & Roy, S. 2011a. Pigments and photoacclimation processes. In Roy, S., Llewellyn, C. A., Egeland, E. S. & Johnsen, G. [Eds.] *Phytoplankton Pigments*. Cambridge University Press, pp. 445–71.
- Brunet, C., Johnsen, G., Lavaud, J. & Roy, S. 2011b. Pigments and photoacclimation processes. In Roy, S., Llewellyn, C. A., Egeland, E. S. & Johnsen, G. [Eds.] *Phytoplankton Pigments*. Cambridge University Press, pp. 445–71.
- Butcher, R. 1952. Contributions to our knowledge of the smaller marine algae. *J. Mar. Biol. Assoc. U.K.* 19:175–91.
- Chrétiennot-Dinet, M.-J., Courties, C., Vaquer, A., Neveux, J., Claustre, H., Lautier, J. & Machado, M.C. 1995. A new marine picoeucaryote: *Ostreococcus tauri* gen. et sp. nov. (Chlorophyta, Prasinophyceae). *Phycologia*. 34:285–92.
- Collado-Fabri, S., Vaultot, D. & Ulloa, O. 2011. Structure and seasonal dynamics of the eukaryotic picophytoplankton community in a wind-driven coastal upwelling ecosystem. *Limnol. Oceanogr.* 56:2334–46.
- Coupe, P., Matsuoka, A., Ruiz-Pino, D., Gosselin, M., Claustre, H., Marie, D., Tremblay, J.E. et al. 2014. Pigment signatures of phytoplankton communities in the Beaufort Sea. *Biogeosciences Discuss.* 11:14489–530.
- Crespo, C., Rodríguez, H., Segade, P., Iglesias, R. & García-Estévez, J.M. 2009. *Coccomyxa* sp. (Chlorophyta: Chlorococcales), a new pathogen in mussels (*Mytilus galloprovincialis*) of Vigo estuary (Galicia, NW Spain). *J. Invertebr. Pathol.* 102:214–9.
- Durand, M.D., Green, R.E., Sosik, H.M. & Olson, R.J. 2002. Diel Variations in Optical Properties of *Micromonas pusilla* (Prasinophyceae). *J. Phycol.* 38:1132–42.
- Egeland, E.S., Guillard, R.R.L. & Liaaen-Jensen, S. 1997. Additional carotenoid prototype representatives and a general chemosystematic evaluation of carotenoids in Prasinophyceae (Chlorophyta). *Phytochemistry*. 44:1087–97.
- Eikrem, W. & Throndsen, J. 1990. The ultrastructure of *Bathycoccus* gen. nov. and *B. prasinos* sp. nov., a non-motile picoplanktonic alga (Chlorophyta, Prasinophyceae) from the Mediterranean and Atlantic. *Phycologia*. 29:344–50.
- Falkowski, P.G., Schofield, O., Katz, M.E., van de Schootbrugge, B. & Knoll, A. 2004. Why is the land green and the ocean red? In Thierstein, H. & Young, J. [Eds.] *Coccolithophorids*. Springer-Verlag, Berlin, pp. 429–53.
- Fawley, M.W., Qin, M. & Yun, Y. 1999. The relationship between *Pseudoscurfieldia marina* and *Pycnococcus provasolii* (Prasinophyceae, Chlorophyta): Evidence from 18S rDNA sequence data. *J. Phycol.* 35:838–43.
- Fawley, M.W., Yun, Y. & Qin, M. 2000. Phylogenetic analyses of 18S rDNA sequences reveal a new coccoid lineage of the Prasinophyceae (Chlorophyta). *J. Phycol.* 36:387–93.
- Fucikova, K., Leliaert, F., Cooper, E.D., Åkaloud, P., D'Hondt, S., De Clerck, O., Gurgel, C.F.D. et al. 2014. New phylogenetic hypotheses for the core Chlorophyta based on chloroplast sequence data. *Front. Ecol. Evol.* 2.
- Gao, Z., Meng, C., Zhang, X., Xu, D., Zhao, Y., Wang, Y., Lv, H. et al. 2012. Differential expression of carotenogenic genes, associated changes on astaxanthin production and photosynthesis features induced by JA in *H. pluvialis*. *PLoS One*. 7:e42243.
- Garrido, J.L., Rodríguez, F. & Zapata, M. 2009. Occurrence of loroxanthin, loroxanthin decenoate, and

- loroxanthin dodecenoate in *Tetraselmis* species (Prasinophyceae, Chlorophyta). *J. Phycol.* 45:366–74.
- Garrido, J.L. & Zapata, M. 1997. Reversed-phase high-performance liquid chromatographic separation of mono- and divinyl chlorophyll forms using pyridine-containing mobile phases and a polymeric octadecylsilica column. *Chromatographia.* 44:43–9.
- Giovagnetti, V., Brunet, C., Conversano, F., Tramontano, F., Obernosterer, I., Ridame, C. & Guieu, C. 2013. Assessing the role of dust deposition on phytoplankton ecophysiology and succession in a low-nutrient low-chlorophyll ecosystem: A mesocosm experiment in the mediterranean sea. *Biogeosciences.* 10:2973–91.
- Guillou, L., Eikrem, W., Chrétiennot-Dinet, M.-J., Le Gall, F., Massana, R., Romari, K., Pedrós-Alió, C. et al. 2004. Diversity of picoplanktonic prasinophytes assessed by direct nuclear SSU rDNA sequencing of environmental samples and novel isolates retrieved from oceanic and coastal marine ecosystems. *Protist.* 155:193–214.
- Gutiérrez-Rodríguez, A., Latasa, M., Estrada, M., Vidal, M. & Marrasé, C. 2010. Carbon fluxes through major phytoplankton groups during the spring bloom and post-bloom in the Northwestern Mediterranean Sea. *Deep. Res. Part I Oceanogr. Res. Pap.* 57:486–500.
- Henriksen, P., Riemann, B., Kaas, H., Sorensen, H.L.H.M. & Sørensen, H.M.H.L. 2002. Effects of nutrient-limitation and irradiance on marine phytoplankton pigments. *J. Plankton Res.* 24:835–58.
- Jahns, P. & Holzwarth, A.R. 2012. The role of the xanthophyll cycle and of lutein in photoprotection of photosystem II. *Biochim. Biophys. Acta - Bioenerg.* 1817:182–93.
- Jeffrey, S., Wright, S. & Zapata, M. 2011. Phytoplankton Pigments: Characterization, Chemotaxonomy and Applications in Oceanography. In Roy S, Egeland ES, Johnsen G, L. C. [Ed.]. Cambridge University Press, Cambridge., pp. 3–77.
- Keller, M.D., Selvin, R.C., Claus, W. & Guillard, R.R.L. 1987. Media for the culture of oceanic ultraphytoplankton. *J. Phycol.* 23:633–8.
- Krienitz, L., Bock, C., Kotut, K. & Luo, W. 2012. *Picocystis salinarum* (Chlorophyta) in saline lakes and hot springs of East Africa. *Phycologia.* 51:22–32.
- Latasa, M., Scharek, R., Le Gall, F. & Guillou, L. 2004. Pigment suites and taxonomic groups in Prasinophyceae. *J. Phycol.* 40:1149–55.
- Leliaert, F., Smith, D.R., Moreau, H., Herron, M.D., Verbruggen, H., Delwiche, C.F. & De Clerck, O. 2012. Phylogeny and molecular evolution of the green algae. *CRC. Crit. Rev. Plant Sci.* 31:1–46.
- Lemieux, C., Otis, C. & Turmel, M. 2014a. Six newly sequenced chloroplast genomes from prasinophyte green algae provide insights into the relationships among prasinophyte lineages and the diversity of streamlined genome architecture in picoplanktonic species. *BMC Genomics.* 15:857.
- Lemieux, C., Otis, C. & Turmel, M. 2014b. Chloroplast phylogenomic analysis resolves deep-level relationships within the green algal class Trebouxiophyceae. *BMC Evol. Biol.* 14:211.
- Lewin, R.A., Krienitz, L., Oerlickei, R.G., Takeda, H. & Hepperle, D. 2000. *Picocystis salinarum* gen. et sp. nov. (Chlorophyta) - a new picoplanktonic green alga. *Phycologia.* 39:560–5.
- Liu, H., Probert, I., Uitz, J., Claustre, H., Aris-Brosou, S., Frada, M., Not, F. et al. 2009. Haptophyta rule the waves: extreme oceanic biodiversity in non-calcifying haptophytes explains the 19-Hex paradox. *Proc. Natl. Acad. Sci.* 106:12803–8.
- Marin, B. & Melkonian, M. 2010. Molecular phylogeny and classification of the Mamiellophyceae class. nov. (Chlorophyta) based on sequence comparisons of the nuclear- and plastid-encoded rRNA Operons.

Protist. 161:304–36.

- Massjuk, N.P. 2006. Chlorodendrophyceae class. nov. (Chlorophyta, Viridiplantae) in the Ukrainian flora: I. The volume, phylogenetic relations and taxonomical status. *Ukr. Bot. J.* 63:601–6014.
- Miki, M., Ramaiah, N., Takeda, S. & Furuya, K. 2008. Phytoplankton dynamics associated with the monsoon in the Sulu Sea as revealed by pigment signature. *J. Oceanogr.* 64:663–73.
- Moon-van der Staay, S., Van der Staay, G.W.M., Guillou, L., Vaulot, D., Claustre, H. & Medlin, L. 2000. Abundance and diversity of prymnesiophytes in the picoplankton community from the equatorial Pacific Ocean inferred from 18S rDNA sequences. *Limnol. Oceanogr.* 45:98–109.
- Niyogi, K.K., Björkman, O. & Grossman, A.R. 1997. The roles of specific xanthophylls in photoprotection. *Proc. Natl. Acad. Sci. U. S. A.* 94:14162–7.
- Not, F., Latasa, M., Marie, D., Cariou, T., Vaulot, D. & Simon, N. 2004. A single species *Micromonas pusilla* (Prasinophyceae) dominates the eukaryotic picoplankton in the western English Channel. *Appl. Environ. Microbiol.* 70:4064–72.
- Obayashi, Y. & Tanoue, E. 2002. Growth and mortality rates of phytoplankton in the northwestern North Pacific estimated by the dilution method and HPLC pigment analysis. *J. Exp. Mar. Bio. Ecol.* 280:33–52.
- R Development Core Team 2013. R: A Language and Environment for Statistical Computing.
- Roesler, C.S., Culbertson, C.W., Etheridge, S.M., Goericke, R., Kiene, R.P., Miller, L.G. & Oremland, R.S. 2002. Distribution, production, and ecophysiology of *Picocystis* strain ML in Mono Lake, California. *Limnol. Oceanogr.* 47:440–52.
- Sarada, R., Tripathi, U. & Ravishankar, G. 2002. Influence of stress on astaxanthin production in *Haematococcus pluvialis* grown under different culture conditions. *Process Biochem.* 37:623–7.
- Schlüter, L., Møhlenberg, F., Havskum, H. & Larsen, S. 2000. The use of phytoplankton pigments for identifying and quantifying phytoplankton groups in coastal areas: Testing the influence of light and nutrients on pigment/chlorophyll a ratios. *Mar. Ecol. Prog. Ser.* 192:49–63.
- Shi, X.L., Marie, D., Jardillier, L., Scanlan, D.J. & Vaulot, D. 2009. Groups without cultured representatives dominate eukaryotic picophytoplankton in the oligotrophic South East Pacific Ocean. *PLoS One.* 4.
- Simon, N., Barlow, R.G., Marie, D., Partensky, F. & Vaulot, D. 1994. Characterization of oceanic photosynthetic picoeukaryotes by flow cytometry analysis. *J. Phycol.* 30:922–35.
- Subirana, L., Péquin, B., Michely, S., Escande, M.L., Meilland, J., Derelle, E., Marin, B. et al. 2013. Morphology, genome plasticity, and phylogeny in the genus *Ostreococcus* reveal a cryptic species, *O. mediterraneus* sp. nov. (Mamiellales, Mamiellophyceae). *Protist.* 164:643–59.
- Takaichi, S. 2011. Carotenoids in Algae: Distributions, biosyntheses and functions of carotenoids in algae. *Mar. Drugs.* 9:1101–18.
- Thronsen, J. and Kristiansen, S. 1991. *Micromonas pusilla* (Prasinophyceae) as part of pico- and nanoplankton communities of the Barents Sea. *Polar Res.* 10:201–7.
- Viprey, M., Guillou, L., Ferréol, M. & Vaulot, D. 2008. Wide genetic diversity of picoplanktonic green algae (Chloroplastida) in the Mediterranean Sea uncovered by a phylum-biased PCR approach. *Environ. Microbiol.* 10:1804–22.
- Wang, B., Zarka, A., Trebst, A. & Boussiba, S. 2003. Astaxanthin accumulation in *Haematococcus pluvialis* (Chlorophyceae) as an active photoprotective process under high irradiance. *J. Phycol.* 39:1116–24.
- Wilhelm, C. & Lenarz-Weiler, I. 1987. Energy transfer and pigment composition in three chlorophyll b-

containing light-harvesting complexes isolated from *Mantoniella squamata* (Prasinophyceae), *Chlorella fusca* (Chlorophyceae) and *Sinapis alba*. *Photosynth. Res.* 13:101–11.

Zapata, M., Rodriguez, F. & Garrido, J.L. 2000. Separation of chlorophylls and carotenoids from marine phytoplankton: a new HPLC method using a reversed phase C-8 column and pyridine-containing mobile phases. *Mar. Ecol. - Prog. Ser.* 195:29–45.

List of tables

Table 1. Characteristics of the strains used in this study. RCC refers to the Roscoff Culture Collection (<http://www.roscoff-culture-collection.org/>).

RCC	Sub-clade	Strain name	Other names	Ocean origin	Region origin	Latitude	Depth Isolation (m)
15	A	CCMP 1205		NA Pacific	NA	NA	NA
287	A	NOUM15	NOUM97015	Ocean	West Equatorial Pacific	0°	120
719	A	IndianOcean_45-8		Indian Ocean	East Equatorial Indian	12°S	76
856	A	Biosope_42 A2	CCMP3325	Ocean Pacific	Marquesas islands	8°S	10
857	A	Biosope_40 A2		Ocean Pacific	Marquesas islands	8°S	10
996	A	Biosope_46 B4S		Ocean Pacific	South East Pacific	9°S	100
998	A	Biosope_46 C3S	NIES2676, CCMP3334	Ocean	South East Pacific	9°S	100
1124	A	PAP_AD	PAP_Ludwig_AI	Atlantic Ocean	North Atlantic, PAP site	49°N	10
1871	A	RA090205-09		Atlantic Ocean	North Atlantic, English Channel	49°N	0
3374	A	CCMP 2152	A7831	Pacific Ocean	Hawaii	23°N	NA
3376	A	CCMP 2113	A9533	Ocean Pacific	Central Equatorial Pacific	9°N	85
2337	B	JST MH335	MH335, NIES2756	Ocean Pacific	Iki Island	34°N	0
2339	B	JST MH340	MH340, NIES2758, CCMP3360	Ocean Pacific	Iki Island	34°N	0
3402	C	CCMP 1897	SFBB	Ocean	San Francisco Bay	38°N	0

Remark : Ordered by clade and then RCC number

Table 2. Concentration of Chl *a* per cell, ratios of pigment to Chl *a* concentration and contribution to total carotenoids (in italics) for fourteen strains of prasinophytes clade VII (Chlorophyta) grown at 140 $\mu\text{E}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$.

Strain	sub-clade	Light $\mu\text{E}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$	fg Chl <i>a</i> /cell	Chl <i>b</i> /Chl <i>a</i>	Chlide <i>a</i> /Chl <i>a</i>	Chlide <i>b</i> /Chl <i>a</i>	Sum of carotenoids /Chl <i>a</i>	Carotenoids																			
								Loroxanthin		Neoxanthin		Violaxanthin		Astaxanthin		Antheraxanthin		Zeaxanthin		Lutein <i>a</i>		$\beta\beta$ -carotene		$\beta\epsilon$ -carotene		Alloxanthin	
								/Chl <i>a</i>	%	/Chl <i>a</i>	%	/Chl <i>a</i>	%	/Chl <i>a</i>	%	/Chl <i>a</i>	%	/Chl <i>a</i>	%	/Chl <i>a</i>	%	/Chl <i>a</i>	%	/Chl <i>a</i>	%	/Chl <i>a</i>	%
RCC15	A	140	20.34	0.901	0.066	0.018	1.321	0.058	4.37	0.003	0.20	0.327	24.77	0.165	12.47	0.043	3.25	0.103	7.79	0.376	28.45	0.093	7.03	0.154	11.66	0	0.00
RCC287	A	140	4.99	0.986	0.106	0	1.533	0.024	1.59	0.167	10.91	0.572	37.29	0.210	13.68	0.024	1.56	0.043	2.79	0.363	23.68	0.079	5.17	0.051	3.32	0	0.00
RCC287*	A	100	Nd	1.313	Nd	Nd	0.759	Nd	Nd	0.074	9.75	0.131	17.26	Nd	Nd	0.051	6.72	0.042	5.53	0.382	50.33	0.053	6.98	0.026	3.43	0	0.00
RCC719	A	140	14.89	0.683	0.000	0	1.928	0.035	1.80	0.086	4.44	0.617	31.97	0.264	13.71	0.000	0.00	0.478	24.79	0.193	10.02	0.149	7.73	0.107	5.53	0	0.00
RCC856	A	140	23.37	0.860	0.000	0	2.077	0.113	5.42	0.079	3.79	0.291	14.02	0.570	27.44	0.042	2.02	0.310	14.94	0.461	22.20	0.136	6.53	0.075	3.63	0	0.00
RCC857	A	140	4.10	1.000	0.000	0	1.399	0.026	1.89	0.122	8.74	0.534	38.15	0.187	13.38	0.024	1.73	0.094	6.69	0.291	20.83	0.064	4.55	0.056	4.03	0	0.00
RCC996	A	140	51.57	0.931	0.074	0	1.259	0.043	3.39	0.095	7.52	0.119	9.46	0.191	15.17	0.039	3.06	0.245	19.45	0.364	28.88	0.074	5.85	0.091	7.21	0	0.00
RCC998	A	140	26.07	0.783	0.000	0	1.902	0.014	0.75	0.117	6.15	0.877	46.10	0.157	8.24	0.058	3.04	0.168	8.84	0.285	14.98	0.131	6.90	0.095	4.99	0	0.00
RCC1124	A	140	8.61	0.960	0.099	0	1.471	0	0.00	0.116	7.87	0.525	35.66	0.148	10.07	0.020	1.39	0.086	5.83	0.399	27.14	0.082	5.59	0.095	6.45	0	0.00
RCC1871	A	100	3.30	1.196	0.000	0	1.296	0	0.00	0.119	9.20	0.460	35.45	0.064	4.92	0.015	1.15	0.091	7.03	0.444	34.28	0.025	1.93	0.078	6.03	0	0.00
RCC3374	A	140	4.14	0.726	0.229	0	1.841	0.012	0.67	0.082	4.47	0.272	14.78	0.778	42.25	0.041	2.25	0.118	6.39	0.332	18.06	0.146	7.92	0.059	3.22	0	0.00
RCC3376	A	140	4.08	0.813	0.105	0	1.526	0.008	0.53	0.112	7.33	0.572	37.47	0.324	21.26	0.054	3.56	0.129	8.47	0.048	3.14	0.194	12.68	0.085	5.57	0	0.00
RCC2337	B	140	4.37	0.882	0.394	0.236	2.186	0.031	1.40	0.137	6.27	0.504	23.07	0.302	13.82	0.051	2.34	0.154	7.03	0.706	32.30	0.240	10.97	0.061	2.80	0	0.00
RCC2339	B	140	14.58	0.624	0.000	0	1.520	0.026	1.74	0.078	5.13	0.613	40.35	0.045	2.98	0.024	1.56	0.074	4.84	0.302	19.86	0.330	21.72	0.028	1.83	0	0.00
RCC3402	C	100	60.40	0.283	0.000	0	0.583	0	0.00	0.039	6.69	0.035	6.03	0	0.00	0.003	0.53	0.018	3.17	0.071	12.12	0.129	22.06	0.015	2.59	0.047	8.13

Nd: Not determined

*Values reported by Latasa et al. 2004

List of supplementary tables

Table S1. Chromatographic retention times and spectral characteristics of the major pigments for strains of prasinophytes clade VII.

Table S2. Ratios of pigment to Chl *a* concentration and contribution to total carotenoids (in italics) for nine strains of prasinophytes clade VII under three light intensities.

List of Figures

Figure 1. A. Cumulative ratios of Chl *b* and five major carotenoids (lutein *a*, violaxanthin, zeaxanthin, astaxanthin, alloxanthin + monadoxanthin + diatoxanthin) to Chl *a* for 14 strains of prasinophytes clades VII at 140 or 100 $\mu\text{E}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ (see Table 2). Strains are ordered by sub-clades (A, B, C) and depth of isolation (surface, deep chlorophyll maximum-DCM). RCC287a correspond to the composition reported by Latasa et al. (2004) for this strain. B. Same as A but with relative abundance of Chl *b* and five major carotenoids.

Figure 2. Change in pigment to Chl *a* ratios for Chl *b* and nine major carotenoids in nine strains of prasinophytes clade VII under three light intensities. Solid lines correspond to sub-clade VIIA and dashed lines to VIIB. Open symbols correspond to surface strains, closed ones to DCM strains and grey to unknown depth of isolation.

Figure 3. Principal component analysis using the pigment to Chl *a* ratios as variables for the strains grown at 3 light levels (Table S2). Top. Variables. Bottom. Samples. Circles correspond to clade VIIA and triangles to clade VIIB. Closed symbols correspond to low light, grey symbols to medium light and open symbols to high light.





