



HAL
open science

Genetic profiles of cervical tumors by high-throughput sequencing for personalized medical care

Etienne Muller, Baptiste B. Brault, Allyson M. Holmes, Angelina Legros, Emmanuelle Jeannot, Maura Campitelli, Antoine Rousselin, Nicolas Goardon, Thierry Frebourg, Sophie Krieger, et al.

► **To cite this version:**

Etienne Muller, Baptiste B. Brault, Allyson M. Holmes, Angelina Legros, Emmanuelle Jeannot, et al.. Genetic profiles of cervical tumors by high-throughput sequencing for personalized medical care. *Cancer Medicine*, 2015, 4 (10), pp.1484-1493. <10.1002/cam4.492>. <hal-01259401>

HAL Id: hal-01259401

<https://hal.sorbonne-universite.fr/hal-01259401v1>

Submitted on 20 Jan 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

ORIGINAL RESEARCH

Genetic profiles of cervical tumors by high-throughput sequencing for personalized medical care

Etienne Muller^{1,2}, Baptiste Brault¹, Allyson Holmes³, Angelina Legros¹, Emmanuelle Jeannot⁴, Maura Campitelli⁵, Antoine Rousselin¹, Nicolas Goardon¹, Thierry Frébourg^{2,6}, Sophie Krieger^{1,2,7}, Hubert Crouet⁸, Alain Nicolas³, Xavier Sastre⁴, Dominique Vaur^{1,2} & Laurent Castéra^{1,2}

¹Department of Cancer Biology and Genetics, CCC François Baclesse, Caen, France

²Inserm U1079, Rouen, France

³Recombination and Genetic Instability, UMR 3244, Institut Curie, Paris, France

⁴Biopathology Department, Institut Curie, Paris, France

⁵Department of Radiotherapy, Institut Curie, Paris, France

⁶Department of Genetics, University Hospital, Rouen, France

⁷Caen University, Caen, France

⁸Gynecology Oncology Department, CCC François Baclesse, Caen, France

Keywords

Cervix, diagnosis, NGS, panel, targeted therapie

Correspondence

Laurent Castéra, Department of Cancer Biology and Genetics, Centre François Baclesse, 3 Avenue du Général Harris, 14076 Caen, France.

Tel: +33 231 455 154; Fax: +33 231 455 053;

E-mail: l.castera@baclesse.unicancer.fr

Funding Information

This work was conducted as a part of the Integrated Actions for Research Program (PAIR-HPV) and was supported by grants from INCa (French Cancer National Institute), Fondation ARC (Association pour la recherche sur le cancer) and the Ligue Contre Le Cancer and the French National League Against Cancer (LNCC).

Received: 15 April 2015; Revised: 2 June 2015; Accepted: 4 June 2015

Cancer Medicine 2015; 4(10):1484–1493

doi: 10.1002/cam4.492

Abstract

Cancer treatment is facing major evolution since the advent of targeted therapies. Building genetic profiles could predict sensitivity or resistance to these therapies and highlight disease-specific abnormalities, supporting personalized patient care. In the context of biomedical research and clinical diagnosis, our laboratory has developed an oncogenic panel comprised of 226 genes and a dedicated bioinformatic pipeline to explore somatic mutations in cervical carcinomas, using high-throughput sequencing. Twenty-nine tumors were sequenced for exons within 226 genes. The automated pipeline used includes a database and a filtration system dedicated to identifying mutations of interest and excluding false positive and germline mutations. One-hundred and seventy-six total mutational events were found among the 29 tumors. Our cervical tumor mutational landscape shows that most mutations are found in *PIK3CA* (E545K, E542K) and *KRAS* (G12D, G13D) and others in *FBXW7* (R465C, R505G, R479Q). Mutations have also been found in *ALK* (V1149L, A1266T) and *EGFR* (T259M). These results showed that 48% of patients display at least one deleterious mutation in genes that have been already targeted by the Food and Drug Administration approved therapies. Considering deleterious mutations, 59% of patients could be eligible for clinical trials. Sequencing hundreds of genes in a clinical context has become feasible, in terms of time and cost. In the near future, such an analysis could be a part of a battery of examinations along the diagnosis and treatment of cancer, helping to detect sensitivity or resistance to targeted therapies and allow advancements towards personalized oncology.

Introduction

Cancer treatment faces a major evolution since the advent of targeted therapies. It is now possible to specifically treat tumors presenting with a clearly identified genetic alteration(s). It is the case of nonsmall-cell lung cancers associated with *EGFR* mutations that responds to specific tyrosine kinase inhibitors [1]. This type of therapy can

only be effective in the case of a known molecular target. It requires the characterization of various alterations (commonly called actionable mutations) that a tumor may accumulate. Establishment of such genetic profiles would allow sensitivity, resistance, and toxicity predictions for such therapies [2]. Sequences from many human genomes have shown a wide interindividual genetic heterogeneity [3]. Variations among individuals are attributed to

“germline” mutations and constitute the individual’s inherited genetic characteristics. These mutations, however, should be distinguished from other mutations that may occur during the individual’s life, and also, might only be observed in one organ or some tissue. These events are called somatic mutations and have been shown to provoke various oncogenic processes [2] (they are called “driver” mutations).

Many research teams, brought together in international consortiums, are engaged in the characterization of mutations causing tumorigenesis. The Catalog Of Somatic Mutations In Cancer (COSMIC) Project [4], for example, aims at creating a database gathering all somatic mutations already described. The International Cancer Genome Consortium plans to build a comprehensive catalog of somatic mutations in tumors of 50 different types/subtypes of cancer, with additional epigenomic and transcriptomic information, with the aim of highlighting differences and common abnormalities across tumor types [5]. These works on tumor genetic profiling have demonstrated that tumors of different types can share protumorigenic signaling pathways, defined by common driver mutations which could also be actionable [6]. Consequently, it makes sense to explore the presence or absence of mutations within these driver genes, to identify biomarkers for sensibility or resistance to treatments, and to indicate those patients most likely to benefit from targeted therapies.

The number of genes to test, in order to establish a cancer gene map, is huge [7]. This approach is made possible by a new generation of sequencing devices (Next Generation Sequencing, NGS), able to analyze the equivalent of several entire genomes [8]. However, sequencing a whole genome can be quite long, regardless of analysis time for such an extensive quantity of data. Another approach consists of limiting the analysis to exomes or several hundreds of genes, using targeted methods. Using a rational approach, which targets a panel consisting of a few hundred genes, considered to be “actionable” and/or “driver”, it is possible to sequence and characterize a tumor, depending on its major molecular characteristics. The main advantages of this approach lie in the reduced delay for reporting the results and limited costs, compatible with a diagnostic use, while maintaining sufficient sequencing quality to detect somatic variations [9].

The sensitivity of these methods depends on bioinformatic tools specifically developed to highlight somatic mutations in cancer, by comparing healthy tissue and tumor DNA from the same individual (paired-mode). These tools verify that a somatic mutation found in tumor does not actually correspond to germline mutation or systematic false-positive mutation, found in healthy tissues [10]. In some cases, it is difficult to obtain the matched

healthy tissue in a clinical context, depending on legislative (local or national), ethical or logistical considerations; a constraint that must be overcome. On the other hand, the detection of mutations strongly depends on algorithms and their adjusted parameters used. In addition, their sensitivity depends on the type and rate of mutation present in the tumor. To date, there is no global consensus for the use of a particular algorithm or mode of analysis [10].

We report on the genetic profiles of cervix uteri tumors which represents the fourth leading cause of death by cancer in females worldwide [11]. It is well-known that some human papillomavirus (HPV) types are the cause of this cancer and its development is linked to the stable insertion of the HPV genome into the tumor’s DNA [12]. Cancer of cervix uteri can go undetectable for years and years, and is associated with a 5-year survival rate of 68% [13], depending on clinical stage at presentation: the 5-year relative survival rate is 91%, 58%, and 17% for patients with localized, regional, and metastatic disease, respectively (<http://seer.cancer.gov/statfacts/html/cervix.html>). Moreover, up to one-third of patients will develop recurrent tumors mostly within 2 years after initial treatments [14]. Most recurrences being detected by imaging or medical examination, it is essential that highly specific, sensitive and less invasive markers are identified to predict response to treatment, disease progression, and to develop personalized therapies.

In our laboratory, we have developed the sequencing of a panel of genes associated with tumors of cervix uteri, including exons within 226 genes chosen for their actionable nature and their potential implication in cervical tumorigenesis (Table 1). To overcome the lack of matched healthy tissue, an average healthy tissue has been generated *in silico* from nonmatched healthy tissues at our disposal. The aim of the study was to prove that a tumor can be managed in a clinical context using NGS technology and automated bioinformatic pipeline and to identify potential actionable mutations that could have a direct link with treatments.

Material and Methods

Tumor samples

Twenty-five squamous cell carcinoma and four adenocarcinoma tumor samples were obtained under approved protocols from the Curie Institute in Paris, France (Table 2). Seven healthy tissue samples were obtained from cervical surgical specimen provided by the Anatomic Pathology laboratory of cancer center François Baclesse in Caen, France. All patients gave written informed consent before entering the study.

Table 1. Genes selected for cervix uteri cancer panel.

<i>ABL1</i>	<i>CDK8</i>	<i>FGFR3</i>	<i>LTK</i>	<i>NSD1</i>	<i>SMAD2</i>
<i>ABL2</i>	<i>CDKN1A</i>	<i>FGFR4</i>	<i>MAP2K1</i>	<i>NTRK1</i>	<i>SMAD3</i>
<i>AFF3</i>	<i>CDKN2A</i>	<i>FHIT</i>	<i>MAP2K2</i>	<i>NTRK2</i>	<i>SMAD4</i>
<i>AKT1</i>	<i>CEACAM5</i>	<i>FKBP9</i>	<i>MAP2K4</i>	<i>NTRK3</i>	<i>SMARCA4</i>
<i>AKT2</i>	<i>CEBPA</i>	<i>FLT1</i>	<i>MCL1</i>	<i>PAK3</i>	<i>SMARCB1</i>
<i>AKT3</i>	<i>CHEK1</i>	<i>FLT3</i>	<i>MDM2</i>	<i>PARP1</i>	<i>SMO</i>
<i>ALK</i>	<i>CHEK2</i>	<i>FLT4</i>	<i>MDM4</i>	<i>PAX5</i>	<i>SOCS1</i>
<i>APC</i>	<i>CREBBP</i>	<i>FOXP4</i>	<i>MED1</i>	<i>PDGFRA</i>	<i>SOX10</i>
<i>AR</i>	<i>CRKL</i>	<i>GATA1</i>	<i>MEN1</i>	<i>PDGFRB</i>	<i>SOX2</i>
<i>ARFRP1</i>	<i>CROCC</i>	<i>GLIS2</i>	<i>MET</i>	<i>PDPK1</i>	<i>SRC</i>
<i>ARID4A</i>	<i>CSF1R</i>	<i>GNA11</i>	<i>MGMT</i>	<i>PIK3CA</i>	<i>STK11</i>
<i>ATM</i>	<i>CSMD1</i>	<i>GNAQ</i>	<i>MITF</i>	<i>PIK3R1</i>	<i>SUFU</i>
<i>AURKA</i>	<i>CTLA4</i>	<i>GNAS</i>	<i>MLH1</i>	<i>PKHD1</i>	<i>TBX22</i>
<i>AURKB</i>	<i>CTNNB1</i>	<i>GPR124</i>	<i>MLL</i>	<i>PLCG1</i>	<i>TCF4</i>
<i>BCL2</i>	<i>CTTN</i>	<i>GUCY1A2</i>	<i>MLL2</i>	<i>PLEKHO2</i>	<i>TENM1</i>
<i>BCL2A1</i>	<i>DAPK1</i>	<i>HIC1</i>	<i>MLL3</i>	<i>PRDM9</i>	<i>TERT</i>
<i>BCL2L1</i>	<i>DDX3X</i>	<i>HNF1A</i>	<i>MPL</i>	<i>PRKDC</i>	<i>TET2</i>
<i>BCL2L2</i>	<i>E2F3</i>	<i>HOXA3</i>	<i>MRE11A</i>	<i>PTCH1</i>	<i>TGFBR2</i>
<i>BCL6</i>	<i>EGFR</i>	<i>HRAS</i>	<i>MSH2</i>	<i>PTEN</i>	<i>TIMP2</i>
<i>BRAF</i>	<i>EMSY</i>	<i>HSP90AA1</i>	<i>MSH6</i>	<i>PTK2</i>	<i>TIMP3</i>
<i>BRCA1</i>	<i>EPHA3</i>	<i>IDH1</i>	<i>MTOR</i>	<i>PTK2B</i>	<i>TNFAIP3</i>
<i>BRCA2</i>	<i>EPHA5</i>	<i>IDH2</i>	<i>MUC1</i>	<i>PTPN11</i>	<i>TNFRSF10C</i>
<i>CADM1</i>	<i>EPHA6</i>	<i>IGF1R</i>	<i>MYB</i>	<i>PTPRD</i>	<i>TNFRSF10D</i>
<i>CASP8</i>	<i>EPHA7</i>	<i>IGF2R</i>	<i>MYC</i>	<i>RAD51</i>	<i>TOP1</i>
<i>CASZ1</i>	<i>EPHB1</i>	<i>IKBKE</i>	<i>MYCL1</i>	<i>RAD51B</i>	<i>TP53</i>
<i>CAV1</i>	<i>EPHB4</i>	<i>IKZF1</i>	<i>MYCN</i>	<i>RAF1</i>	<i>TP63</i>
<i>CBL</i>	<i>EPHB6</i>	<i>INHBA</i>	<i>MYH15</i>	<i>RARA</i>	<i>TP73</i>
<i>CCND1</i>	<i>ERBB2</i>	<i>IRF6</i>	<i>NF1</i>	<i>RARB</i>	<i>TSC1</i>
<i>CCND2</i>	<i>ERBB3</i>	<i>IRS2</i>	<i>NF2</i>	<i>RASSF1</i>	<i>TSC2</i>
<i>CCNE1</i>	<i>ERBB4</i>	<i>JAK2</i>	<i>NFE2L2</i>	<i>RB1</i>	<i>TSHR</i>
<i>CCNL1</i>	<i>ESR1</i>	<i>JAK3</i>	<i>NFIB</i>	<i>REL</i>	<i>USP9X</i>
<i>CDC73</i>	<i>EZH2</i>	<i>KDR</i>	<i>NKX2-1</i>	<i>RET</i>	<i>VHL</i>
<i>CDH1</i>	<i>FAM123B</i>	<i>KEAP1</i>	<i>NOTCH1</i>	<i>RICTOR</i>	<i>WT1</i>
<i>CDH2</i>	<i>FANCC</i>	<i>KIAA0774</i>	<i>NOTCH2</i>	<i>ROS1</i>	<i>ZBTB7C</i>
<i>CDH20</i>	<i>FANCF</i>	<i>KIT</i>	<i>NOTCH3</i>	<i>RPS6KB1</i>	<i>ZNF668</i>
<i>CDH5</i>	<i>FBXW7</i>	<i>KLF5</i>	<i>NOTCH4</i>	<i>RPTOR</i>	<i>ZNF91</i>
<i>CDK4</i>	<i>FGFR1</i>	<i>KRAS</i>	<i>NPM1</i>	<i>RUNX1</i>	
<i>CDK6</i>	<i>FGFR2</i>	<i>LRP1B</i>	<i>NRAS</i>	<i>RUNX1T1</i>	

Sample preparation and next-generation sequencing

Sequencing methods were described previously [15], enabling analysis of two tumors and one healthy tissue sample in the same sequencing run. The SureSelect-targeted enrichment process (Agilent, Santa Clara, CA) was performed after combining the indexed samples, equimolarly. Libraries were then sequenced on a MiSeq (Illumina, San Diego, CA), using the paired-end 2 × 150 bp method (full protocol available on request).

Bioinformatic analysis

The CASAVA suite v1.8 (Illumina) was used for demultiplexing and generating fastq files. These raw data were then analyzed by a home-made pipeline (BAPT, Bioinformatic

Analysis Pipeline Toolkit), to automate alignment, variant-calling, and annotation steps. Reads were first mapped to a reference genome (hg19) using BWA algorithm [16] and output files were reworked following GATK [17] (Genome Analysis ToolKit, Broad Institute) best practices, especially to calculate quality scores and undergo local realignments. Five variant-callers were used to call single-nucleotide substitutions and insertion–deletion (indels) events: HaplotypeCaller [18], UnifiedGenotyper [18], LofreqStar [19], VarScan2 [20], and MuTect [21]. The BAM files from seven healthy samples were used by randomly selecting reads from each file, to create an average healthy synthetic file (using SAMtools [22]). This “synthetic” BAM file was used in paired-mode during the variant-calling step with the programs LofreqStar, MuTect, and VarScan2. All variants found by the variant callers were annotated with Alamut-Batch (Interactive Biosoftware, Rouen, France) and integrated into a database called CanDiD

Table 2. Demographic, histological, and biological characteristics.

Patient no.	Histological type	Tumor stage (FIGO)	Age at diagnosis	HPV type
1	SCC	IIB	40	HPV 18
2	SCC	IB	33	HPV 18
3	Adenocarcinoma	II	45	HPV 18
4	SCC	IB	37	HPV 18
5	SCC	IB	34	HPV 16
6	SCC	IB2	47	HPV 16
7	Adenocarcinoma	IIB	49	HPV 16
8	SCC	IVB	57	HPV 16
9	SCC	IB1	33	HPV 16
10	SCC	NA	NA	HPV45
11	SCC	IIB	60	HPV 16
12	SCC	IIB	42	HPV 18
13	SCC	IIB	47	HPV 16, 18
14	SCC	IIB	34	HPV 16
15	SCC	IIB	68	HPV 16
16	SCC	IIB	43	HPV 18
17	SCC	IIB	44	HPV 51
18	SCC	IIB	65	HPV 33
19	SCC	IIB	45	HPV 73
20	SCC	IIIB	43	HPV 16
21	SCC	IIB	53	HPV 16
22	Adenocarcinoma	IB2	42	HPV 16
23	Adenocarcinoma	IIB	54	HPV-
24	SCC	IIB	54	HPV 18
25	SCC	III	33	HPV 16
26	SCC	IIB	55	HPV 16
27	SCC	IV + metastasis	44	HPV 73
28	SCC	IB1	25	HPV 18
29	SCC	IB2	31	HPV-

FIGO, International Federation of Gynecology; HPV, human papillomavirus; SCC, squamous cell carcinoma; NA, not available.

(Cancer Diagnostic Database, PostgreSQL) (Fig. S1). Variants were retrieved from CanDiD database according to five criteria: (1) the variant was found on a list of preferred transcripts; (2) the variant was in targeted zones; (3) the variant was within the coding sequence or ± 10 bp within the intronic sequences; (4) the variant was in a canonical splicing mutation sites or when outside they induced a 15% decrease of MaxEntScan score and a 5% decrease of the SpliceSiteFinder score [23] (hereafter called splicing mutation); and (5) the variant was observed 10 times at most in the Exome Sequencing Project Database (ESP; <http://evs.gs.washington.edu/EVS/>). Identified variants passed through a home-made filtration system (scripted on Python programming language, available on request) which compares sequencing data from the five variant-callers and tumor samples versus healthy samples, in order to rule out false-positive and germline variants. Data from each variant were first compared to both tumor and healthy tissue samples, to evaluate whether the variant have a greater presence in the tumor than in the healthy samples. Variants were then

filtered based on their quality score (the main criteria being the PHRED score and depth of coverage). The main objective was to obtain, for each tumor, a list of variants with a high probability of being somatic. Only variants seen by at least 2 variant callers were further considered.

Results were annotated and interpreted in sequence with: (1) two mutation impact prediction systems SIFT [24], Polyphen [25], (2) InterPro Database [26], (3) Clinical Trials.gov Database [27], (4) dbSNP [28], (5) COSMIC [4], (6) TARGET [29].

In the present study, all genes that constitute a target for the US Food and Drug Administration-approved targeted therapies were called actionable genes, regardless of the type of cancer. Similarly, a missense mutation was classified as deleterious if scored as “deleterious” by SIFT algorithm or “Probably Damaging” by Polyphen algorithm. Inactivating mutations (PTC, premature codon termination and splicing mutations) were directly considered as deleterious.

Results

Tumor samples from 29 patients with cervical cancer (stade I B to IV B), including 25 squamous cell carcinomas and 4 adenocarcinomas, have been sequenced (Table 2), as well as seven samples from healthy cervical biopsy (“control” samples). The time from DNA extraction to acquisition of most likely somatic mutations was about height working days (Fig. 1). The sequencing process produced an average of 8,078,023 reads per sample, with an average sequencing depth of 268 \times and >92% of nucleotides covered ≥ 50 -fold. Consequently, only variants with an allele ratio of more than 5% were called. After data processing in the BAPT pipeline, 11,267 variants were included in the CanDiD database. After extraction from the database, 2746 missense mutations were collected from the various variant callers (Fig. 2A). After passing through the filtration system, only 220 mutations were retained (Fig. 2B; Table S2). Most of the mutations deleted by the filtration system were germline mutations detected by HaplotypeCaller or UnifiedGenotyper (from the control samples) or false-positive selected by Varscan due to its lack of specificity. Among all mutations found in the 29 tumor samples, 41% (91/220) were identified by all variant callers, assuring accuracy of the mutations detected. Only variants detected by two or more variant-callers were selected leading to 156 missense mutations selected, among which 29 are already referenced in the COSMIC database. Seven nonsense mutations, nine small insertion/deletions (indels) inducing PTC and four splicing mutations were also identified. Altogether, they represent an average of 6.1 mutations per tumor sample. For each patient, 75 percent of mutations were considered




Figure 1. Sequencing workflow's major steps. After DNA extraction (Day 1), DNA molecules are sheared by sonification, and DNA fragments are ligated to adapters containing a patient-specific index (Day 2). DNA from 2 patients and 1 control are pooled equimolarly together during the multiplexing step (Day 3). Regions of interest (exons from 226 genes) are retrieved by a targeted enrichment system with biotinylated baits (Day 4–5). Then DNA is sequenced in an Illumina MiSeq (Day 6) and sequencing raw data are processed by the bioinformatic pipeline (Day 7–8), to extract the most likely somatic variations.




Figure 2. Representation of missense mutations found by each variant caller. (A) Mutations extracted directly from CANDID database (total: 2746). (B) mutations remaining after passing through filtration system (total: 221). (C) Proportion of mutations detected by at least 2 variant-callers classified as deleterious by SIFT or POLYPHEN.

deleterious (by SIFT or POLYPHEN algorithm) (Fig. 2C, Table S2).

No somatic missense mutation was found in the tumor suppressor gene *TP53*, which is consistent with molecular characteristics of cervix uteri tumors [30]. Similarly, missense mutations were found in the oncogene *PIK3CA* ($n = 8$), the *KRAS* gene implicated into anti-EGFR therapies resistance ($n = 4$) and in the *FBXW7* gene, recently described in the cervix uteri tumors mutational landscape ($n = 3$) (Table S2). Other missense mutations were found in tumor suppressor genes, including the susceptibility gene for breast cancer *BRCA2* ($n = 6$) or the RAS family negative regulator *NF1* ($n = 1$). Twenty inactivating

mutations (PTC and splicing mutations) were also identified in tumor suppressor genes including *RBI* ($n = 1$), *NF1* ($n = 3$) or *MLH1* ($n = 2$) (Table 3).

Among other actionable genes, deleterious missense mutations were observed in *ALK* ($n = 4$), *AR* ($n = 1$), *EGFR* ($n = 3$), *ERBB2* ($n = 1$), *MET* ($n = 1$), *mTOR* ($n = 2$), *PDGFRA* ($n = 1$) and *RAF1* ($n = 2$) (Table 4). Considering only deleterious mutations, 48% ($n = 14/29$) had at least one mutation in an actionable gene (Fig. 3A and B).

Several clinical trials (www.clinicaltrials.gov database) are looking for mutations in specific genes to evaluate sensitivity or resistance to approved, or in clinical trials, targeted therapies, among which *PIK3CA*, *ALK*, *EGFR*,

Table 3. Inactivating mutations.

Coding effect	Gene	Coding DNA sequence ¹	Protein sequence ¹	Transcript	Patients tumor sample no.
Splicing mutation	<i>CBL</i>	c.1096-1G>A	p. ?	NM_005188	5
Potential splicing mutations	<i>mTOR</i>	c.2514+3G>C	p. ?	NM_004958	15
	<i>MYH15</i>	c.1993-14G>A	p. ?	NM_014981	14
	<i>NF1</i>	c.1392+5G>T	p. ?	NM_001042492	21
Nonsense mutations	<i>C11orf30</i>	c.3004C>T	p.Gln1002*	NM_020193	26
	<i>CEACAM5</i>	c.646G>T	p.Glu216*	NM_004363	26
	<i>EPHA5</i>	c.2458G>T	p.Gly820*	NM_004439	6
	<i>FBXW7</i>	c.1053G>A	p.Trp351*	NM_033632	26
	<i>MLH1</i>	c.1630C>T	p.Gln544*	NM_000249	20
		c.755C>T	p.Ser252*		5
	<i>RB1</i>	c.1399C>T	p.Arg467*	NM_000321	5
Frameshift mutations	<i>CASP8</i>	c.790dup	p.Val264Glyfs*13	NM_001080125	5
	<i>CHEK2</i>	c.1229del	p.Thr410Metfs*15	NM_001005735	11
	<i>CREBBP</i>	c.4477dup	p.Ile1493Asnfs*26	NM_004380	5
	<i>FGFR2</i>	c.962dup	p.Asn321Lysfs*21	NM_022970	5
	<i>FGFR4</i>	c.2396_2403del	p.Gly799Aspfs*12	NM_213647	7
	<i>NF1</i>	c.5907_5908del	p.Arg1970Serfs*6	NM_001042492	5
		c.2033del	p.Pro678Argfs*10		
	<i>NF2</i>	c.301del	p.Tyr101Ilefs*22	NM_181832	16
	<i>NOTCH2</i>	c.6909del	p.Ile2304Leufs*2	NM_024408	5

Mutations are classified by mutation type.

¹Nomenclature according HGVS guidelines (Human Genome Variation Society).

Table 4. Deleterious mutations found in actionable genes.

Gene	Drugs in relation with gene of interest	Mutation	Number of tumors	Transcript	Associated clinical trial
<i>ALK</i>	Crizotinib, Ceritinib	A1266T A1234V V1149L R1120Q	4	NM_004304	NCT01548144, NCT01744652
<i>AR</i>	Entuzalamide, Abiraterone	K809N	1	NM_000044	–
<i>EGFR</i>	Cetuximab, Panitumumab, Erlotinib, Gefetinib, Afatinib, Vandetanib	S511Y T259M A611T	3	NM_005228	NCT00770263
<i>ERBB2</i>	Trastuzumab, Pertuzumab, Lapatinib	L696F	1	NM_001005862	NCT01953926
<i>KRAS</i>	Resistance to Cetuximab and others	G12D G13D	4	NM_033360	–
<i>MET</i>	Cabozantinib	L342F	1	NM_001127500	–
<i>mTOR</i>	Temsirolimus, Everolimus	M813I	1	NM_004958	–
<i>PDGFRA</i>	Regorafenib	P441L	1	NM_006206	NCT02029001
<i>RAF1</i>	Regorafenib	C96F Q255H	2	NM_002880	–

Genes are linked to targeted therapies already approved by Food and Drug Administration in at least one indication. All potential drug targets of each therapy are considered.

mTOR, *PTCH1*, *SMO*, and *PIK3R1* are included (Table S2). Overall, 59% of patients (17/29) would be eligible for potential inclusion in a clinical trial (Fig. 3A and B), based on carrier gene mutations and tumor origin.

Discussion

The aim of the study was to demonstrate in a clinical context the feasibility of detecting somatic variations, in

cervical cancers, using a targeted sequencing approach. The use of a “synthetic healthy tissue sample” (see above Material and Methods section) from the sequencing data of several unpaired healthy tissue samples allows the use of somatic variant-callers (which compares healthy tissues from tumors to extract somatic mutations) to detect poorly represented mutations. This approach, however, probably tolerates a few germline mutations which cannot be differentiated from somatic mutations. In our study, some




Figure 3. Distribution of deleterious mutations among the 29 tumors. A gene is considered actionable if linked to targeted therapies approved by Food and Drug Administration. A mutation is classified as deleterious if considered as such by SIFT algorithm or POLYPHEN algorithm. (A) Proportion of tumors with a deleterious mutation on gene considered actionable (blue) and proportion of patients for whom their deleterious mutations could allow inclusion in a clinical trial (purple). (B) Representation of most mutated genes across tumors.

mutations found in *BRCA2* (hereditary predisposition for breast and ovarian cancer) or in *MLH1* (hereditary predisposition to Lynch syndrome) genes through the genetic characterization of tumors could be germline (Table S2). The identification of germline mutations in genes in relation with mendelian diseases must be taken into account and the appropriate ethical issues should be considered, a subject still of debate. Tools and methods developed in our laboratory should not be considered as an automatic interpretation system; they should be considered as help to ease work discerning the most relevant mutations. For instance, with our data, the number of mutations to analyze drops from about 388 to 6 for one tumor, which represents a huge time-saving for biologists who interpret data in a clinical context. Moreover, analysis time (± 8 working days) is consistent with clinical practice, and should improve patient healthcare in a comprehensive way.

No somatic alterations were detected in the tumor suppressor gene *TP53* which seems consistent throughout our model. In cervical carcinomas *TP53* inactivation is linked to virus integration, which results in the expression of the E6 oncoprotein, able to induce the *TP53* degradation via the ubiquitin pathway [30]. Previous studies have already identified genes frequently mutated

in cervical carcinomas, such as *PIK3CA* [31], *KRAS* [32] (the most mutated genes in cervix uteri cancer according to COSMIC data) or *FBXW7* [33]. *PIK3CA* E545K ($n = 4$) and E542K ($n = 2$) have strong oncogenic properties due to increased kinase activity [34], consequences in growth factor-independent cell proliferation, resistance to apoptosis, and increased invasion and cell migration. This gene represents a prime target for drug development, and even if there is no available approved treatment, several clinical trials are currently testing *PIK3CA* selective inhibitors (ClinicalTrials ID: NCT01708161; NCT01928459). Two major mutations were also found in *KRAS*, G12D ($n = 1$) and G13D ($n = 3$), well known in colorectal cancer to predict a resistance to *EGFR* antibodies [35]. *FBXW7* is a gene recently described as mutated in cervical carcinoma [33] with two missense mutations already described, R505G and R456C. There was also a novel mutation at a position already found mutated, R479Q. All of these point mutations are located in the WD repeats domain. This domain is used to form a complex in charge of degrading several products of proto-oncogenes (including *MYC*, *CCNE1*, *NOTCH1*, and *JUN*) [36]. We found a novel truncating mutation W351*; it occurs just upstream of the WD domain (UniProt ID: Q969H0). Therefore, pipeline implementation

seems consistent, since key elements from cervical tumorigenesis were found.

Considering all of the actionable genes as defined before (Table 4), deleterious mutations could influence sensitivity or resistance to treatments. *ALK* mutations V1149L and A1266T are located next to previously described mutations L1152R [37] and G1269A [38], which have been identified as resistance mutations to Crizotinib and other tyrosine kinase inhibitors, and could then have the same impact on treatment response. Another mutation (T259M) has been observed in a specific subpart of extracellular domain of *EGFR*, called the dimerization loop [39]. This loop is essential for receptor dimerization and hence activation. A previous study showed that mutations in this subpart were able to increase receptor oncogenic activity [40]. *MET* had also a new mutation at residue L342, which correspond to a specific region of SEMA domain, considered as *MET* ligand-binding domain [41] and useful to receptor dimerization and activation [42]. Other mutations classified as deleterious were detected in *mTOR* (M813I, splice mutation c.2514+3G>C) and *PDGFRA* (P441L) but their real impact on protein functionality or the effectiveness of related targeted therapies is still unknown.

Eighteen deleterious mutations identified from our cases reside in actionable genes (Table 3), although their impact is still unclear for many. Therefore, functional studies should be performed to identify potential drug sensitivities or resistances to these targets. Deleterious classification (SIFT and POLYPHEN algorithm) should be used with caution since, for instance, *PIK3CA* mutations E545K and E542K are classified as tolerated by SIFT despite the fact that they are known driver mutations.

Among driver genes investigated in our study, interesting mutations were found on *BRCA2*, with 2 variants of unknown signification, classified here as deleterious missense mutations (L2721R and A2770D) in the OB1 (oligonucleotide/oligosaccharide-binding 1) domain, which serves to bind single-strand DNA. Functional assays showed that mutations in this domain can be disease-causing [43]. Another mutation (V2066I) was observed in BRC repeat domains, essential for binding to RAD51 protein [44]. The new PARP inhibitor Olaparib, now available, could potentially be of interest in these *BRCA* mutated tumors. Unfortunately, for the moment this drug is mainly tested in breast and ovarian cancers. Altogether, the real new challenge for a tumor's genetic profiling is the correct interpretation of mutation.

Sixty-nine percent of patients could have had their treatment decision guided by the targeted sequencing of 226 genes, either by revealing sensitivity or resistance to available drugs or by directing patients toward treatment protocols or diagnostic procedures being tested (Fig. 3A).

Even though no therapeutic answer could be provided by our “oncopanel” for certain patients, they were all analyzed for HPV insertion status. The HPV insertion site constitutes another biomarker in *cis*, which can provide information for targeted therapies. HPV has therefore been found inserted into *MYC*, *KLF5*, *KLF12*, as well as *RBI*, and many other important and potentially actionable genes (A. Holmes, unpubl. data). Such tests might participate in the overall study of cancer biomarkers in both prognostic (assessing tumor outcomes and deciding whether or not to undergo treatment) and predictive (evaluating sensitivity and resistance to treatments to choose the most effective drug) fields [45]. The outcome of such studies strives toward personalized oncology, which is to provide the most appropriate treatment to the patient with the correct dosage. The overall benefits first and foremost concern patients, who receive the most effective and least toxic treatments. Such an approach in patient care has economic advantages too. The correct administration of the adequate drug instead of using other ineffective and expensive therapies appears to be cost-effective. It has been calculated that US health-care system could save up to \$740 million dollars by only treating patients with Cetuximab with a wild-type *KRAS* status [46]. Other studies suggest that pharmacogenomics can be a good way to improve cost-effectiveness [47]. However, more studies are needed in this field to assess the real economic impact of these therapies [48].

Interpretation of somatic mutations remains challenging because, except for a few mutations already described, the impact of each mutation requires functional studies. In some countries, the inability to obtain matched healthy tissue in a daily clinical practice makes detection of somatic mutations more difficult. On the other hand, sequencing a limited number of genes selected for their implication in tumor development is now feasible in a clinical setting (in term of costs and time), and the “oncopanel” approach allows a larger screening power for diagnosis or treatment orientation by identification of sensitivity or resistance to drugs.

Acknowledgments

We thank the biological resources center of Curie Institute, the sequencing platform SésAME (Séquençage pour la Santé, l'Agonomie, la Mer et l'Environnement), CCC François Baclesse, Caen, the Tumorothèque Caen Basse Normandie (TCBN), and Michel Henry-Amar for critical review of the manuscript.

Conflict of Interest

None declared.

References

- Paez, J. G., P. A. Jänne, J. C. Lee, S. Tracy, H. Greulich, S. Gabriel, et al. 2004. EGFR mutations in lung cancer: correlation with clinical response to gefitinib therapy. *Science* 304:1497–1500.
- Vogelstein, B., N. Papadopoulos, V. E. Velculescu, S. Zhou, L. A. Diaz, and K. W. Kinzler. 2013. Cancer genome landscapes. *Science* 339:1546–1558.
- McClellan, J., and M.-C. King. 2010. Genetic heterogeneity in human disease. *Cell* 141:210–217.
- Forbes, S. A., G. Tang, N. Bindal, S. Bamford, E. Dawson, C. Cole, et al. 2009. COSMIC (the Catalogue of Somatic Mutations in Cancer): a resource to investigate acquired mutations in human cancer. *Nucleic Acids Res.* 38:652–657.
- Hudson, T. J., W. Anderson, A. Aretz, A. D. Barker, C. Bell, R. R. Bernabé, et al. 2010. International network of cancer genome projects. *Nature* 464:993–998.
- Zack, T. I., S. E. Schumacher, S. L. Carter, A. D. Cherniack, G. Saksena, B. Tabak, et al. 2013. Pan-cancer patterns of somatic copy number alteration. *Nat. Genet.* 45:1134–1140.
- Futreal, P. A., L. Coin, M. Marshall, T. Down, T. Hubbard, R. Wooster, et al. 2004. A census of human cancer genes. *Nat. Rev. Cancer* 4:177–183.
- Metzker, M. L. 2009. Sequencing technologies—the next generation. *Nat. Rev. Genet.* 11:31–46.
- Rehm, H. L. 2013. Disease-targeted sequencing: a cornerstone in the clinic. *Nat. Rev. Genet.* 14:295–300.
- Xu, H., J. DiCarlo, R. V. Satya, Q. Peng, and Y. Wang. 2014. Comparison of somatic mutation calling methods in amplicon and whole exome sequence data. *BMC Genom.* 15:244.
- Jemal, A., F. Bray, M. M. Center, J. Ferlay, E. Ward, and D. Forman. 2011. Global cancer statistics. *CA Cancer J. Clin.* 61:69–90.
- Zur Hausen, H. 2009. Papillomaviruses in the causation of human cancers—a brief historical account. *Virology* 384:260–265.
- Howlander, N., A. M. Noone, M. Krapcho, D. Miller, S. F. Altekruse, C. L. Kosary, et al. 2014. SEER cancer statistics review, 1975–2011. Bethesda MD Natl. Cancer Inst. p.1975–2011.
- Leitao, M. M., and D. S. Chi. 2002. Recurrent cervical cancer. *Curr. Treat. Options Oncol.* 3:105–111.
- Castéra, L., S. Krieger, A. Rousselin, A. Legros, J.-J. Baumann, O. Bruet, et al. 2014. Next-generation sequencing for the diagnosis of hereditary breast and ovarian cancer using genomic capture targeting multiple candidate genes. *Eur. J. Hum. Genet.* 22:1305–1313.
- Li, H., and R. Durbin. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760.
- Auwerda, G. A., M. O. Carneiro, C. Hartl, R. Poplin, G. del Angel, A. Levy-Moonshine, et al. 2013. From FastQ data to high-confidence variant calls: the Genome Analysis Toolkit best practices pipeline. *Curr. Protoc. Bioinformatics* 11:11.10.1–11.10.33.
- DePristo, M. A., E. Banks, R. Poplin, K. V. Garimella, J. R. Maguire, C. Hartl, et al. 2011. A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat. Genet.* 43:491–498.
- Wilm, A., P. P. K. Aw, D. Bertrand, G. H. T. Yeo, S. H. Ong, C. H. Wong, et al. 2012. LoFreq: a sequence-quality aware, ultra-sensitive variant caller for uncovering cell-population heterogeneity from high-throughput sequencing datasets. *Nucleic Acids Res.* 40:11189–11201.
- Koboldt, D. C., Q. Zhang, D. E. Larson, D. Shen, M. D. McLellan, L. Lin, et al. 2012. VarScan 2: somatic mutation and copy number alteration discovery in cancer by exome sequencing. *Genome Res.* 22:568–576.
- Cibulskis, K., M. S. Lawrence, S. L. Carter, A. Sivachenko, D. Jaffe, C. Sougnez, et al. 2013. Sensitive detection of somatic point mutations in impure and heterogeneous cancer samples. *Nat. Biotechnol.* 31:213–219.
- Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, et al. 2009. The sequence alignment/map format and SAMtools. *Bioinformatics* 25:2078–2079.
- Houdayer, C., V. Caux-Moncoutier, S. Krieger, M. Barrois, F. Bonnet, V. Bourdon, et al. 2012. Guidelines for splicing analysis in molecular diagnosis derived from a set of 327 combined in silico/in vitro studies on BRCA1 and BRCA2 variants. *Hum. Mutat.* 33:1228–1238.
- Ng, P. C., and S. Henikoff. 2003. SIFT: predicting amino acid changes that affect protein function. *Nucleic Acids Res.* 31:3812–3814.
- Adzhubei, I. A., S. Schmidt, L. Peshkin, V. E. Ramensky, A. Gerasimova, P. Bork, et al. 2010. A method and server for predicting damaging missense mutations. *Nat. Methods* 7:248–249.
- Hunter, S., P. Jones, A. Mitchell, R. Apweiler, T. K. Attwood, A. Bateman, et al. 2011. InterPro in 2011: new developments in the family and domain prediction database. *Nucleic Acids Res.* 40:306–312.
- US NATIONAL INSTITUTES OF HEALTH, et al. 2012. *ClinicalTrials.gov*.
- Sherry, S. T., M.-H. Ward, M. Kholodov, J. Baker, L. Phan, E. M. Smigielski, et al. 2001. dbSNP: the NCBI database of genetic variation. *Nucleic Acids Res.* 29:308–311.
- Van Allen, E. M., N. Wagle, P. Stojanov, D. L. Perrin, K. Cibulskis, S. Marlow, et al. 2014. Whole-exome sequencing and clinical interpretation of formalin-fixed,

- paraffin-embedded tumor samples to guide precision cancer medicine. *Nat. Med.* 20:682–688.
30. Tommasino, M., R. Accardi, S. Caldeira, W. Dong, I. Malanchi, A. Smet, et al. 2003. The role of TP53 in cervical carcinogenesis. *Hum. Mutat.* 21:307–312.
 31. McIntyre, J. B., J. S. Wu, P. S. Craighead, T. Phan, M. Köbel, S. P. Lees-Miller, et al. 2013. *PIK3CA* mutational status and overall survival in patients with cervical cancer treated with radical chemoradiotherapy. *Gynecol. Oncol.* 128:409–414.
 32. Kang, S., H.-S. Kim, S. S. Seo, S.-Y. Park, D. Sidransky, and S. M. Dong. 2007. Inverse correlation between *RASSF1A* hypermethylation, *KRAS* and *BRAF* mutations in cervical adenocarcinoma. *Gynecol. Oncol.* 105:662–666.
 33. Ojesina, A. I., L. Lichtenstein, S. S. Freeman, C. S. Peadarallu, I. Imaz-Rosshandler, T. J. Pugh, et al. 2014. Landscape of genomic alterations in cervical carcinomas. *Nature* 506:371–375.
 34. Samuels, Y., Z. Wang, A. Bardelli, N. Silliman, J. Ptak, S. Szabo, et al. 2004. High frequency of mutations of the *PIK3CA* gene in human cancers. *Science* 304:554.
 35. Amado, R. G., M. Wolf, M. Peeters, E. Van Cutsem, S. Siena, D. J. Freeman, et al. 2008. Wild-type *KRAS* is required for panitumumab efficacy in patients with metastatic colorectal cancer. *J. Clin. Oncol.* 26:1626–1634.
 36. Welcker, M., and B. E. Clurman. 2008. FBW7 ubiquitin ligase: a tumour suppressor at the crossroads of cell division, growth and differentiation. *Nat. Rev. Cancer* 8:83–93.
 37. Sasaki, T., J. Koivunen, A. Ogino, M. Yanagita, S. Nikiforow, W. Zheng, et al. 2011. A novel *ALK* secondary mutation and *EGFR* signaling cause resistance to *ALK* kinase inhibitors. *Cancer Res.* 71:6051–6060.
 38. Kim, S., T. M. Kim, D.-W. Kim, H. Go, B. Keam, S.-H. Lee, et al. 2013. Heterogeneity of genetic changes associated with acquired crizotinib resistance in *ALK*-rearranged lung cancer. *J. Thorac. Oncol.* 8:415–422.
 39. Ogiso, H., R. Ishitani, O. Nureki, S. Fukai, M. Yamanaka, J.-H. Kim, et al. 2002. Crystal structure of the complex of human epidermal growth factor and receptor extracellular domains. *Cell* 110:775–787.
 40. Lee, J. C., I. Vivanco, R. Beroukhim, J. H. Huang, W. L. Feng, R. M. DeBiasi, et al. 2006. Epidermal growth factor receptor activation in glioblastoma through novel missense mutations in the extracellular domain. *PLoS Med.* 3:e485.
 41. Gherardi, E., M. E. Youles, R. N. Miguel, T. L. Blundell, L. Iamele, J. Gough, et al. 2003. Functional map and domain structure of *MET*, the product of the *c-met* protooncogene and receptor for hepatocyte growth factor/scatter factor. *Proc. Natl. Acad. Sci. USA* 100:12039–12044.
 42. Wickramasinghe, D., and M. Kong-Beltran. 2005. *Met* activation and receptor dimerization in cancer: a role for the *Sema* domain. *Cell Cycle* 4:683–685.
 43. Wu, K., S. R. Hinson, A. Ohashi, D. Farrugia, P. Wendt, S. V. Tavtigian, et al. 2005. Functional evaluation and cancer risk assessment of *BRCA2* unclassified variants. *Cancer Res.* 65:417–426.
 44. Carreira, A., and S. C. Kowalczykowski. 2011. Two classes of *BRC* repeats in *BRCA2* promote *RAD51* nucleoprotein filament function by distinct mechanisms. *Proc. Natl. Acad. Sci. USA* 108:10448–10453.
 45. Sawyers, C. L. 2008. The cancer biomarker problem. *Nature* 452:548–552.
 46. Shankaran, V., D. J. Bentrem, M. F. Mulcahy, C. L. Bennett, and A. B. Benson III. 2009. Economic implications of *Kras* testing in metastatic colorectal cancer (mCRC). In: Abstract. 15–17.
 47. Carlson, J. J., L. P. Garrison, S. D. Ramsey, and D. L. Veenstra. 2009. The potential clinical and economic outcomes of pharmacogenomic approaches to *EGFR*-tyrosine kinase inhibitor therapy in non-small-cell lung cancer. *Value Health* 12:20–27.
 48. Lange, A., A. Prenzler, M. Frank, H. Golpon, T. Welte, and J.-M. von der Schulenburg. 2014. A systematic review of the cost-effectiveness of targeted therapies for metastatic non-small cell lung cancer (NSCLC). *BMC Pulm. Med.* 14:192.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Figure S1. Somatic mutations analysis pipeline: BAPT.

Table S2. Somatic mutations found across the 29 patients.