



HAL
open science

Double KK excited states in highly charged sulphur

P Amaro, J P Marques, P Indelicato, T K Mukherjee, J K Saha, L C Tribedi,
J P Santos

► **To cite this version:**

P Amaro, J P Marques, P Indelicato, T K Mukherjee, J K Saha, et al.. Double KK excited states in highly charged sulphur. *Journal of Physics: Conference Series*, 2015, 635 (2), pp.022071 10.1088/1742-6596/635/2/022071 . hal-01265591

HAL Id: hal-01265591

<https://hal.sorbonne-universite.fr/hal-01265591>

Submitted on 1 Feb 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Double *KK* excited states in highly charged sulphur

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 J. Phys.: Conf. Ser. 635 022071

(<http://iopscience.iop.org/1742-6596/635/2/022071>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 134.157.80.136

This content was downloaded on 01/02/2016 at 11:19

Please note that [terms and conditions apply](#).

Double KK excited states in highly charged sulphur

P. Amaro*¹, J. P. Marques[†], P. Indelicato[‡], T. K. Mukherjee[#],

J. K. Saha^{||}, L. C. Tribedi[&], J. P. Santos*²

* Laboratório de Instrumentação, Engenharia Biomédica e Física da Radiação (LIBPhys-UNL), Departamento de Física, Faculdade de Ciências e Tecnologia, FCT, Universidade Nova de Lisboa, 2829-516 Caparica, Portugal

[†] BioISI - Biosystems & Integrative Sciences Institute, Faculdade de Ciências da Universidade de Lisboa, Campo Grande, C8, 1749-016 Lisboa, Portugal

[‡] Laboratoire Kastler Brossel, École Normale Supérieure, CNRS, Université P. et M. Curie – Paris 6, Case 74; 4, place Jussieu, 75252 Paris CEDEX 05, France

[#] Narula Institute of Technology, Agarpara, Kolkata-700109, India

^{||} Indian Association for the Cultivation of Science, Jadavpur, Kolkata-700032, India

[&] Department of Nuclear and Atomic Physics, Tata Institute of Fundamental Research, Colaba, Mumbai 400005, India

Synopsis Calculation of energies and radiative decays was performed for several highly charge states of sulphur in order to identify experimental lines associated to double excited KK atomic states.

Double KK excited states have several important applications in astrophysics as well as in plasma diagnostics [1, 2]. Besides its applicability, the fundamental interest of dominance of fluorescence over autoionization decay for some KK excited states has attract much attention, both from theoreticians and experimentalists [3, 4]. Recently, it was observed the $2p3d(^1P^0) \rightarrow 1s3d(^1D^e)$ in helium with sufficient accuracy to distinguish from the H-like K_α line, showing an enhanced fluorescence rate compared to the one assuming standard selection rules [5]. In this work, we continue the investigation of KK excited states for sulphur by performing a systematic identification of experimental lines, which includes both single K and double KK excited states. The relativistic calculations were made within the MCDF approach using the general relativistic MCDF code (MDFGME) [6]. Experimental measurements were performed at the Pelletron accelerator facility in TIFR Mumbai.

In these preliminary calculations, all transition energies and probabilities were computed in a monoconfiguration approach. Almost all possible combinations of excitations for K_α , K_β and K_γ transitions of He through Be charge states were considered. The positions of the experimental lines are displayed in Fig. 1 along with the decay rates of the computed transitions. Transitions from KK excited states were also considered in both He- and Li-like states. It can observed in Fig. 1 that all experimental lines are unambiguously identified, with the exception of one line next to the He-like K_α line. Lines near

the H-like K_α line can only be associated to the KK excited states of He or Li charge states. A precise identification can be done by calculating the respective autoionization rate.




Figure 1. Comparison of obtained theoretical lines with experimental lines.

Referências

- [1] A. B. C. Walker and Jr., H. R. Ruge, 1971 *Astrophys. J.* **164** 181
- [2] T. Fujimoto and T. Kato, 1981 *Astrophys. J.* **264** 994
- [3] T.W. Gorczyca *at al* 2000 *Phys.Rev.Lett.* **85** 1202
- [4] J. G. Lambourne *at al* 2003 *Phys.Rev.Lett.* **90** 153004
- [5] S. Kasthurirangan *at al* 2013 *Phys.Rev.Lett.* **24** 243201
- [6] P. Indelicato and J. P. Desclaux 1990 *Phys.Rev.A* **42** 5139-5149

¹E-mail: pdamaro@fct.unl.pt

²E-mail: jps@fct.unl.pt

