



HAL
open science

Nondipole Effects in Chiral Molecules Measured with Linearly Polarized Light

K P Bowen, O Hemmers, R Guillemin, W C Stolte, M N Piancastelli, D W Lindle

► **To cite this version:**

K P Bowen, O Hemmers, R Guillemin, W C Stolte, M N Piancastelli, et al.. Nondipole Effects in Chiral Molecules Measured with Linearly Polarized Light. *Journal of Physics: Conference Series*, 2015, 635 (11), pp.112053 10.1088/1742-6596/635/11/112053 . hal-01265686

HAL Id: hal-01265686

<https://hal.sorbonne-universite.fr/hal-01265686>

Submitted on 1 Feb 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Nondipole Effects in Chiral Molecules Measured with Linearly Polarized Light

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 J. Phys.: Conf. Ser. 635 112053

(<http://iopscience.iop.org/1742-6596/635/11/112053>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 134.157.80.136

This content was downloaded on 01/02/2016 at 13:10

Please note that [terms and conditions apply](#).

Nondipole Effects in Chiral Molecules Measured with Linearly Polarized Light

K.P. Bowen*¹, O. Hemmers*, R. Guillemin†, W.C. Stolte‡, M.N. Piancastelli§, D.W. Lindle*

*Department of Chemistry & Biochemistry, University of Nevada Las Vegas, Las Vegas, NV, USA

† Laboratoire de Chimie Physique – Matière et Rayonnement, Université Pierre et Marie Curie, CNRS UMR 7614, Paris Cedex 05, France

‡National Security Technologies, LLC (NSTec), Livermore Operations, Livermore, CA, USA

§ Department of Physics and Astronomy, Molecular and Condensed Matter Physics, Uppsala Universitet, Uppsala, Sweden

Synopsis: We present the first-ever measurement of nondipole chiral angular distribution parameters for C 1s photoemission from each enantiomer of camphor in the photon energy range 296-343eV using linearly polarized light. The angular distribution parameters are determined to be enantiomer-specific, suggesting a new form of linear dichroism.

Over the past two decades, it has been demonstrated that higher-order corrections to the dipole approximation are necessary for the description of light-matter interactions in the soft x-ray range. These so-called ‘nondipole effects’ present themselves as asymmetries in the angular distributions of photoelectrons. Chiral molecules, known to have asymmetries in photoelectron angular distributions when exposed to circularly polarized light, have been proposed to demonstrate a chiral-specific nondipole effect when exposed to linearly polarized light[1]. This chiral-specific contribution to the angular distribution of photoelectrons is characterized by the parameter ϵ , which introduces a “twisting” of the angular distribution in the plane containing the polarization and perpendicular to the incoming photon propagation. This study presents $\epsilon(h\nu)$ values, as well as the conventional angular distribution asymmetry parameter $\beta(h\nu)$, for C 1s photoemission from both (R)-Camphor and (S)-Camphor, in the range of 5-50eV above the C 1s threshold.

The measurements were performed using electron time-of-flight (TOF) spectroscopy at the Advanced Light Source (ALS) at Lawrence Berkeley National Lab (LBNL) in Berkeley, CA.

The individual parameter values with respect to photon energy are presented in figures 1 and 2, respectively. The large difference in $\epsilon(h\nu)$ between enantiomers implies a new form of linear dichroism.




Figure 1. The anisotropy parameter $\beta(h\nu)$ for C 1s photoemission from (R)-Camphor and (S)-Camphor




Figure 2. The chiral nondipole parameter $\epsilon(h\nu)$ for C 1s photoemission from (R)-Camphor and (S)-Camphor

References

- [1] A.N. Grum-Grzhimailo 2003 *J. Phys. B* **36** 2385

¹ Email: bowenk4@unlv.nevada.edu



