



**HAL**  
open science

## **K-shell width, fluorescence yield, and $K \beta / K \alpha$ intensity ratio calculation for Fe in the Dirac-Fock approach**

J P Marques, Matteo Guerra, F Parente, J P Santos, P Indelicato, J M Sampaio

### ► To cite this version:

J P Marques, Matteo Guerra, F Parente, J P Santos, P Indelicato, et al.. K-shell width, fluorescence yield, and  $K \beta / K \alpha$  intensity ratio calculation for Fe in the Dirac-Fock approach. *Journal of Physics: Conference Series*, 2015, 635 (9), pp.092094 10.1088/1742-6596/635/9/092094 . hal-01266818

**HAL Id: hal-01266818**

**<https://hal.sorbonne-universite.fr/hal-01266818v1>**

Submitted on 3 Feb 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## K-shell width, fluorescence yield, and $K_{\beta}/K_{\alpha}$ intensity ratio calculation for Fe in the Dirac-Fock approach

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 J. Phys.: Conf. Ser. 635 092094

(<http://iopscience.iop.org/1742-6596/635/9/092094>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

### Download details:

This content was downloaded by: mauroguerra

IP Address: 85.242.187.37

This content was downloaded on 12/09/2015 at 12:55

Please note that [terms and conditions apply](#).

## *K*-shell width, fluorescence yield, and $K_\beta/K_\alpha$ intensity ratio calculation for Fe in the Dirac-Fock approach

J. P. Marques<sup>\*1</sup>, M. Guerra<sup>†</sup>, F. Parente<sup>†</sup>, J. P. Santos<sup>†</sup>, P. Indelicato<sup>‡</sup>, J. M. Sampaio<sup>\*</sup>

<sup>\*</sup> BioISI - Biosystems & Integrative Sciences Institute, Faculdade de Ciências da Universidade de Lisboa, Campo Grande, C8, 1749-016 Lisboa, Portugal

<sup>†</sup> Laboratório de Instrumentação, Engenharia Biomédica e Física da Radiação (LIBPhys-UNL), Departamento de Física, Faculdade de Ciências e Tecnologia da Universidade Nova de Lisboa, Monte da Caparica, 2892-516 Caparica, Portugal

<sup>‡</sup> Laboratoire Kastler Brossel, École Normale Supérieure, CNRS, Sorbonne Universités, UPMC Univ. Paris 06, Case 74; 4, place Jussieu, 75252 Paris CEDEX 05, France

**Synopsis** *K*-shell fluorescence yield, width, and  $K_\beta/K_\alpha$  intensity ratio for Fe are calculated using the Dirac-Fock approach. Results are compared with available data.

The knowledge of accurate values of atomic decay rates and fluorescence yields is fundamental in quantitative analysis in X-ray spectroscopy, plasma physics, nuclear physics, dosimetry, radiation protection, among many other areas [1]. Despite the increasing number of theoretical and experimental work concerning the determination of fluorescence yields for several elements, the available data are still scarce. Iron is one of the most abundant elements in the universe and is of extreme importance in the analysis of stellar objects like stars and galaxies. Iron spectral lines are used in plasma diagnostics and help us to know physical properties such as temperature and electron density as well as for abundance studies [2].

In this work, we present a relativistic calculation of the *K*-shell width, fluorescence yield and  $K_\beta/K_\alpha$  intensity ratio in Fe using the multi-configuration Dirac-Fock (MCDF) code of Desclaux and Indelicato [3, 4]. The code was used in a single-configuration approach, with Breit interaction and the vacuum polarization terms included in the self-consistent field calculations, and other QED effects included as perturbations.

**Table 1.** *K*-shell fluorescence yield ( $\omega_K$ ), width (eV), and  $K_\beta/K_\alpha$  intensity ratio for iron

	$\omega_K$	width	$K_\beta/K_\alpha$
This work	0.378	1.14	0.134
McGuire [5]	0.386		
Krause [6]	0.34		
Campbell [7]		1.19	
Carreras <i>et al</i> [8]			0.14
Scofield [9]			0.139
Jankowsky <i>et al</i> [10]			0.134
Bé <i>et al</i> [11]			0.142

Since iron has an open outer shell ( $3d^6$ ), electron coupling leads to 348105 possible radiative and radiationless transitions from the initial *K* hole configuration. Preliminary results for the *K*-shell fluorescence yield, width and  $K_\beta/K_\alpha$  intensity ratio are consistent with the available data (Table 1).

This work was supported by FCT Project PEstOE/FIS/UI0303/2011, Portugal. We also thank the Allianz Program of the Helmholtz Association, contract No. EMMI HA-216 Extremes of Density and Temperature: Cosmic Matter in the Laboratory. BioISI is supported by the centre grant UID/MULTI/04046/2013 from FCT/MCTES/PIDDAC, Portugal.

### References

- [1] J. M. Sampaio *et al* 2014 *Phys. Rev. A* **89** 012512
- [2] H. Hartman *et al* 2009 *J. Phys. Conf. Ser.* **130** 012010
- [3] J. Desclaux 1975 *Comput. Phys. Commun.* **9** 31
- [4] P. Indelicato, J. P. Desclaux 1990 *Phys. Rev. A* **42** 5139
- [5] E. J. McGuire 1970 *Phys. Rev. A* **2** 273
- [6] M. O. Krause 1979 *J. Phys. Chem Ref. Data* **8** 307
- [7] J. Campbell, T. Papp 2001 *At. Data Nucl. Data Tables* **77** 1
- [8] A. Carreras *et al* 2003 *Rad. Phys. Chem* **67** 703
- [9] J. Scofield 1974 *Phys. Rev. A* **9** 1041
- [10] K. Jankowski, M. Polasik 1989 *J. Phys. B* **22** 2369
- [11] M. -M Bé *et al* 1998 *Appl. Radiat. Isot.* **49** 1367

<sup>1</sup>E-mail: [jmmarques@fc.ul.pt](mailto:jmmarques@fc.ul.pt)

