



HAL
open science

Single photon core ionization with core excitation: a new spectroscopic tool

F Penent, S Carniato, P Lablanquie, P Selles, J Palaudoux, L Andric, M Žitnik, K Bučar, E Shigemasa, M Nakano, et al.

► To cite this version:

F Penent, S Carniato, P Lablanquie, P Selles, J Palaudoux, et al.. Single photon core ionization with core excitation: a new spectroscopic tool. *Journal of Physics: Conference Series*, 2015, 635 (11), pp.112093 10.1088/1742-6596/635/11/112093 . hal-01266835

HAL Id: hal-01266835

<https://hal.sorbonne-universite.fr/hal-01266835>

Submitted on 3 Feb 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Single photon core ionization with core excitation: a new spectroscopic tool

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 J. Phys.: Conf. Ser. 635 112093

(<http://iopscience.iop.org/1742-6596/635/11/112093>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 134.157.80.136

This content was downloaded on 03/02/2016 at 13:24

Please note that [terms and conditions apply](#).

Single photon core ionization with core excitation: a new spectroscopic tool.

Penent F.^{*1}, Carniato S.^{*2}, Lablanquie P.^{*}, Selles P.^{*}, Palaudoux J.^{*}, Andric L.^{*}, Žitnik M.[&],
 Bučar K.[&], Shigemasa E.[£], Nakano M.[†], Hikosaka Y.[#] and Ito K.[†]

^{*} Sorbonne Université, UPMC and CNRS, LCP-MR (UMR 7614), 11 rue P. & M. Curie, 75231 Paris Cedex 05, France

[&] Jozef Stefan Institute, P.O. Box 3000, SI-1001 Ljubljana, Slovenia

[£] UVSOR Facility, Institute for Molecular Science, Okazaki 444-8585, Japan


[†] Institute of Materials Structure Science, Photon Factory, Tsukuba, Ibaraki 305-0801, Japan

[#] Department of Environmental Science, Niigata University, Niigata 950-2181, Japan

Synopsis. The simultaneous core ionization and core excitation process (or $K^{-2}V$ process) induced by absorption of a single photon provides the basis of a new spectroscopy that offers both advantages of X-ray Photoelectron Spectroscopy (XPS) and near-edge x-ray absorption fine structures (NEXAFS) spectroscopy

In the last few years, double-core-hole states (DCHs) have been extensively studied by single photon double-core ionization at synchrotron centers [1] and, in relation to X-ray two photon double-core spectroscopy [2], at X-ray free electron laser facilities (XFELs). Closely related to double core ionization, the single photon process where a core electron is ionized while another core electron is excited simultaneously, reveals interesting properties on the formation mechanisms of $K^{-2}V$ states and provides a new spectroscopic insight that offers the advantages of both XPS and NEXAFS spectroscopies.

The creation of $K^{-2}V$ states and their subsequent decay are the following:



Since one photoelectron and two Auger electrons are emitted, by detecting in coincidence these three electrons, it is possible to filter out the specific photoelectron spectrum associated with $K^{-2}V$ states. This can be done very efficiently by using a magnetic bottle time-of-flight electron spectrometer under single bunch operation at SOLEIL (France) and Photon Factory (Japan).

The study of the above process for the C_2H_{2n} ($n=1-3$) series [3] has revealed a NEXAFS signature in the spectra (Fig.1). Two competing channels appear with comparable intensities:

- The first one corresponds to the dipolar ionization of a core electron accompanied by monopolar excitation (direct shake-up) of the remaining core electron and can be interpreted as a "Super" K^{-1} satellite state.

- The second one corresponds to the dipolar excitation of a core electron to a vacant orbital accompanied by monopolar shake-off of the

other core electron (conjugate shake-up process) which has a NEXAFS-like character.

Close to the K^{-2} threshold, these two processes appear with comparable intensities. This can be understood as resulting from aborted double-core ionization where one of the outgoing electrons is recaptured in a vacant orbital. At higher photon energy, the conjugate shake-up process is unfavorable since the sudden approximation leads only to the direct process.




Figure 1: Experimental and theoretical $K^{-2}V$ spectra in C_2H_{2n} series at a photon energy $h\nu = 770\text{eV}$.

Examples on small molecules: H_2O [4], N_2 [5] and many others, have been obtained, which show how powerful this new spectroscopy is and allows also a better interpretation of resonances appearing in NEXAFS spectrum.

References

- [1] Lablanquie *et al* 2011, *Phys. Rev. Lett.* **106**, 063003
- [2] Santra *et al* 2009, *Phys. Rev. Lett.* **103**, 013002
- [3] Nakano *et al* 2013, *Phys. Rev. Lett.* **111**, 123001
- [4] Carniato *et al* 2015, *J. Chem. Phys.* **142**, 014307
- [5] Carniato *et al* 2015, *J. Chem. Phys.* **142**, 014308

¹ E-mail: francis.penent@upmc.fr

^{*2} E-mail: stephane.carniato@upmc.fr

