



# Experimental exposure to trace metals affects plumage bacterial community in the feral pigeon

Marion Chatelain, Adrien Frantz, Julien Gasparini, Sarah Leclaire

## ► To cite this version:

Marion Chatelain, Adrien Frantz, Julien Gasparini, Sarah Leclaire. Experimental exposure to trace metals affects plumage bacterial community in the feral pigeon. 2016. hal-01274875

**HAL Id: hal-01274875**

**<https://hal.sorbonne-universite.fr/hal-01274875>**

Preprint submitted on 16 Feb 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# **Experimental exposure to trace metals affects plumage bacterial community in the feral pigeon**

Chatelain M<sup>1\*</sup>, Frantz A<sup>1</sup>, Gasparini J<sup>1</sup> and Leclaire S<sup>1,2</sup>

<sup>1</sup> Sorbonne Universités, UPMC Univ Paris 06, UPEC, Paris 7, CNRS, INRA, IRD, Institut d'Ecologie et des Sciences de l'Environnement de Paris, F-75005, Paris, France

<sup>2</sup>Centre d'Ecologie Fonctionnelle et Evolutive, UMR 5175, CNRS, 1919 route de Mende, 34293 Montpellier, France

\*Corresponding author: Marion Chatelain, Address : Université Pierre et Marie Curie 7 quai St Bernard 75005 Paris France, E-mail : marion.chatelain@upmc.fr, Tel: +331442752

## Abstract

Bacteria are fundamental associates of animals, and recent studies have highlighted their major role in host behaviour, immunity or reproductive investment. Thus, any environmental factor modifying bacterial community may affect host fitness. In birds, trace metals emitted by anthropogenic activities accumulate onto the plumage where they may alter bacterial community and ultimately affect bird fitness. Although trace metals are current major environmental issues in urban habitats, their effects on feather bacterial community have never been investigated. Here, we supplemented feral pigeons (*Columba livia*), an emblematic urban species, with zinc and/or lead in drinking and bath water. As expected, lead and zinc supplementations modified plumage bacterial community composition. Zinc decreased bacterial load, while lead decreased bacterial richness and the frequency of preening behaviour in birds, known to regulate feather bacteria. Our results demonstrate for the first time the effects of common urban trace metals on plumage bacterial community and shed light on one of the mechanisms by which trace metals can affect bird fitness. Further studies are now needed to investigate how this effect modulates avian life history traits known to depend on plumage bacterial community.

Key words: urban ecology, birds, bacteria

## INTRODUCTION

Bacteria successfully colonize numerous and various habitats, including animal body parts that are in direct contact with the surroundings such as skin, feathers or fur, and the digestive tract. The normal flora, also called the microbiota, lives in symbiosis with its host, fulfilling essential functions for host metabolism, such as cellulose degradation, vitamins synthesis (Hill 1997) and inhibition of pathogens proliferation (Olsson et al. 1992, Oh et al. 2006, Balcázar et al. 2007). Recent studies have also revealed that bacteria may have surprising effects such as shaping host behaviour and investment in reproduction (Ezenwa et al. 2012, Jacob et al. 2015). However, we have just made the first steps in discovering the diversity and functions of these bacterial ecosystems, and it is now necessary to evaluate the factors that regulate microbiota.

The microbiota is influenced by numerous interconnected factors, including host behaviour, genotype and physiology (Mueller et al. 2006, Frank et al. 2011, Rosenthal et al. 2011, Hildebrand et al. 2013, Leclaire et al. 2014a), as well as its environment (Dotterud et al. 2008, Burkholder et al. 2008, Ruiz-de-Castañeda et al. 2011). Trace metals emitted by anthropogenic activities have well-known detrimental effects on animals (Jarup 2003, Hsu et al. 2006, Berglund et al. 2007, Eeva et al. 2009) and are thus of major environmental concern in most urban environments (Azimi et al. 2005, Scheifler et al. 2006, Roux and Marra 2007, Kekkonen et al. 2012). Some trace metals, such as lead, cadmium, zinc, copper, chrome and nickel, are known to have toxic effects on environmental microbial communities (Babich and Stotzky 1978, Giller et al. 1998). In contrast, although trace metals can be naturally ingested or inhaled by animals, and deposited on integuments, their effects on animal microbiota have been poorly investigated (but see Hojberg et al. 2005, Vahjen et al. 2010, Breton et al. 2013, Liu et al. 2014).

Plumage is a key point of interaction between birds and the microbial world, and, like other integuments, is highly colonized by bacteria (Burt Jr and Ichida 1999, Muza et al. 2000, Shawkey and Hill 2004, Whitaker et al. 2005). Although several studies have shown that keratinolytic bacteria can alter feather structure *in vitro* (Burt Jr and Ichida 1999, Shawkey et al. 2007), the *in vivo* effects of feather bacteria on their host have been scarcely studied. A few correlative and experimental studies in captive and free-living birds have shown, however, that feather bacteria may influence sexual signalization such as feather colouration or plumage condition (Shawkey et al. 2007, Gunderson et al. 2009, Kilgas et al. 2012, Leclaire et al. 2014b; but see Jacob et al. 2014), and affect bird immune system (Leclaire et al. 2015) and investment in reproduction (Jacob et al. 2015). Any change in feather microbiota is thus likely to have a strong impact on bird fitness. The plumage and the uropygial gland secretions spread onto it can accumulate metals (Pilastro et al. 1993, Frantz et al. 2012), which can affect plumage bacterial community by direct contact. Ingested metals also circulate in the bloodstream and accumulate in organs and bones (Pattee 1984, Dauwe et al. 2002, Scheifler et al. 2006, Kekkonen et al. 2012, Reid et al. 2012), and may induce noxious (Redig et al. 1991, Snoeijs et al. 2004, Dauwe et al. 2005, Eeva et al. 2009) or beneficial (Mertz 1981, Prasad 1998) effects on bird physiology; therefore, metals can affect bird ability to regulate their microbiota (Piault et al. 2008, Moreno-Rueda 2010, Leclaire et al. 2014b). Whatever the exact underlying mechanisms, further studies are clearly needed to evaluate the effects of trace metals on feather bacteria.

The feral pigeon (*Columba livia*) is an emblematic urban bird living in high density populations, with potentially elevated plumage bacteria transmission between individuals (reviewed in Archie and Theis 2011). Mate choice is crucial for this species which mate for

life (Johnston and Janiga 1995). In this species, bacterial load affects immunity (Leclaire et al. 2015), as well as feather condition and colouration (Leclaire et al. 2014b) which are two of the main criteria used in mate choice (Johnston and Janiga 1995). Consequently, trace metals, besides their direct effects on bird physiology, may further impact pigeon reproduction and fitness through their effects on plumage bacterial load and composition. Here we investigated the effects of an experimental exposure to lead and/or zinc on two traits that can affect each other: plumage bacteria composition and frequency of preening in the feral pigeon. Preening is a costly antimicrobial behaviour that can be directly influenced by feather bacterial load (Leclaire et al. 2014b) and bird health status (Piault et al. 2008, Moreno-Rueda 2010), which are both potentially affected by trace metals.

## **METHODS**

### **Subjects and housing**

Ninety six (48 males and 48 females) free-living adult feral pigeons (*Columba livia*) were caught during winter 2013 (February/March) in several pigeons' flocks within the Parisian agglomeration. Birds were all considered as adults because of their well-formed caruncle, the absence of juvenile plumage and the presence of iridescent neck feathers (Johnston and Janiga 1995). Pigeons were kept in 8 outdoor aviaries (3.10 m x 2.00 m x 2.40 m) at the CEREEP field station (Centre d'Ecologie Expérimentale et Prédictive-Ecotron Ile-de-France, UMS 3194, Ecole Normale Supérieure, Saint-Pierre-lès-Nemours, France). They were evenly distributed among aviaries according to their flock, sex and plumage eumelanin level (see below) in such a way that there was no confounding effect between aviaries and these variables (i.e. no statistically significant link between aviary and flock:  $\chi^2=71.09$ ,  $df=70$ ,  $P=0.441$ ; sex: 6

males and 6 females per aviary; or plumage eumelanin level:  $F_{1,80}=0.38$ ,  $P=0.537$ ). Birds were fed *ad libitum* with a mix of maize, wheat and peas and water was provided in a trough. The aviaries were enriched with a bowl of water used for bathing and with branches as perches. Birds were individually identified with a numbered plastic ring. Birds were genetically sexed following the protocol described by Griffiths et al. (1998). Before onset of treatment, birds were kept 2 to 7 weeks for acclimation. Alongside this experiment, we measured the effects of treatments (see below) on bird body mass condition, reproduction and immunity. At the end of the breeding season, all birds were released back to the wild at their site of capture. All experiments were carried out in strict accordance with the recommendations of the “European Convention for the Protection of vertebrate Animals used for Experimental and Other Scientific Purposes” and were conducted under the authorizations of the “Ministère de l’éducation nationale, de l’enseignement supérieur et de la recherche” (authorization N\_00093.02) and the “Direction Départementale des Services Vétérinaires de Seine et-Marne” (authorization N\_77-05).

### **Measurement of plumage colouration**

At capture, birds were first categorised as eumelanic (grey to black pigmented) or pheomelanic (red pigmented) which define their melanin type. Then, eumelanic birds were individually photographed in order to measure their eumelanin level. Eumelanin level was calculated as the percentage of black on the wing surface (number of black pixels/number of white pixels x 100) using the Gimp image retouching and editing software. This measure has been shown to be a reliable and repeatable estimation of melanin concentration in pigeons (Jacquin et al. 2011). In eumelanic pigeons, plumage eumelanin level ranged from 4.2 to 95.9%. Because of the small amount of pheomelanic birds (14 over 96) the measure of a pheomelanin level was not relevant.

## Treatments

Two weeks before the onset of treatment, aviaries were divided into 4 metal-exposure treatments; this means that there were 2 aviaries per treatment with 12 pigeons each (24 pigeons in total per treatment). For each treatment, the two aviaries were purposely spatially separated from one another. Aviaries were in direct contact along a linear transect and numbered from 1 to 8 (*lead* (1), *zinc* (2), *control* (3), *lead+zinc* (4), *lead* (5), *control* (6), *zinc* (7), *lead+zinc* (8)). Side-by-side aviaries were separated by wire mesh. Treatments consisted of water supplemented with lead (*lead* group; 1ppm lead acetate; aviaries 1 and 5), zinc (*zinc* group; 10 ppm zinc sulphate; aviaries 2 and 7), lead and zinc (*lead+zinc* group; 1ppm lead acetate and 10ppm zinc sulphate; aviaries 4 and 8), or control (*control* group; tap water with no metal added; aviaries 3 and 6). We chose these concentrations based on both lead blood concentrations measured in urban birds (ranging from 0,053 to 0,264ppm; Roux and Marra 2007) and the gastrointestinal absorption rate of lead in zebra finches (<10%) calculated from (Dauwe et al. 2002). Zinc concentrations were approximated using the zinc/lead concentration ratio in the environment and in bird feathers (on average, zinc was 10 times more concentrated than lead; Azimi et al. 2005, Frantz et al. 2012, Chatelain et al. 2014). Drinking troughs and baths were filled with the corresponding treated water every other day. Our supplementation treatments were validated by measuring lead and zinc concentrations in blood and feathers of the birds. Blood was sampled 10 weeks after the start of the experiment. Moreover, the fifth secondary remige of each bird was removed a first time and the regrowth feather was used for metal measurements. Both blood and feathers were digested using a previously described protocol (Chatelain et al. 2014) and lead and zinc concentrations were measured by mass spectrometry (ICP-MS) and by optic emission spectrometry (ICP-OES) respectively. Validity of analytical methods was checked by means



of a standard biological reference material (TMDA-64.2. Environment Canada). Lead and zinc concentrations in blood were the highest among birds exposed to lead (*lead* and *lead+zinc* groups) and birds exposed to zinc (*zinc* and *lead+zinc* groups) respectively (Table 1). These results ensured that metals added to water were ingested by the birds. In feathers, while lead concentrations were significantly the highest among birds exposed to lead (*lead* and *lead+zinc* groups), the increase in zinc concentration among birds exposed to zinc (*zinc* and *lead+zinc* groups) was not significant (Table 1). Zinc and lead measured in the feathers were respectively 80 and 1.5 times less concentrated than the ones measured in feathers of urban pigeons (Nam et al. 2004, Adout et al. 2007, Hoff Brait and AntoniosiFilho 2011, Frantz et al. 2012, Chatelain et al. 2014), suggesting that our experimental exposure corresponded to the lower range of urban exposure.

#### **Measurement of plumage bacterial load**

Plumage bacterial load was measured 20 weeks after onset of treatment. 4 hours after renewing the water of the bowls used for bathing, 10 birds (5 males and 5 females) were randomly sampled in each treatment. Each sampled bird was caught with a net that had been previously sprayed with 70% ethanol. Then, a whole flora agar slide (Hygialim, 3026091, Plate Count Agar +triphenyltetrazoliumchloride+Neutralizing) was put flat against the back of the bird for 10 seconds. The slides were then incubated at 37°C for 24h. Feather bacterial load was expressed as the number of bacterial colonies per slide.

#### **Molecular analysis of plumage bacterial communities**

Fifteen weeks after the start of the experiment, 91 adults were caught with a net previously sprayed with 70% ethanol (n.b. 5 pigeons died for unknown reasons before this measure, 2 from the *zinc* group, 2 from the *lead+zinc* group and 1 from the *control* group). After washing

her hands with alcohol, the experimenter cut a clump (10 feathers on average) of back feathers with sterilized scissors and pliers, avoiding the outermost feathers. The feathers were immediately placed in sterile 2ml plastic tubes and stored at -20°C until analysis.

We extracted DNA using the QiagenDNeasy® Blood and Tissue Kit and the standard protocol designed for the purification of total DNA from Gram-positive bacteria (Qiagen, Venlo, Netherlands; July 2006).

To characterize the bacterial communities present in each sample, we performed automated ribosomal intergenic spacer analyses (ARISA; Ranjard et al. 2000). This DNA fingerprinting method is based on the amplification of the internal transcribed spacer (ITS) region lying between the 16S and 23S ribosomal RNA genes in the ribosomal operon. The ITS region is extremely variable, in both sequence and length, for different bacterial species. Therefore, the DNA amplification profile obtained with ARISA allows straightforward estimation of bacterial diversity, avoiding biases inherent in classical culture-based techniques (Ranjard et al. 2000). We amplified the ITS using the FAM (6-carboxyfluorescein)-labeled primer S-D-Bact-1522-b-S-20 (5'-[6FAM] TGCGGCTGGATCCCCTCCTT-3') and the unlabeled primer L-D-Bact-132-a-A-18 (5'-CCGGGTTTCCCCATTCGG-3') (Ranjard et al. 2000). We performed the PCR amplification in 10 µL mixtures containing 200 µM each deoxynucleotide triphosphate, 0.20 µM each primer, 1.25 units of PerfectTaq DNA polymerase, 1× PCR buffer (5 Prime, GmbH, Hamburg, Germany), and 1 µL DNA extract, using the following protocol: initial denaturation at 94 °C for 3 min, 40 cycles consisting of denaturation at 94 °C for 30 s, annealing at 55 °C for 45 s, elongation at 72 °C for 1 min, and a final elongation at 72 °C for 10 min. We then mixed 1 µL of the PCR products with 15 µL of highly deionized formamide and 0.2 µL of Genescan 1200 LIZ size standard (Applied Biosystems, Foster City, CA). The mixtures were separated with a 24-capillary 3500XL DNA Analyzer (Applied Biosystems) using POP-7 polymer and the manufacturer's default electrophoresis

run settings. Data analysis and genotyping were performed with GeneMapper software (Applied Biosystems). For each sample, the sequencer produced an ARISA profile in which each peak corresponds to 1 phylotype or operational taxonomic unit (OTU). In the various samples, the sequencer detected ITS fragments ranging in size from 300 to 950base pairs.

For each individual, we calculated bacterial richness as the number of different OTUs. Because the probability to detect an OTU is likely affected by the amount of feathers used for DNA extraction, feathers were dried overnight and weighted to the nearest mg after DNA extraction. We estimated bacterial community dissimilarities between individuals using Jaccard distance based on presence/absence of OTUs.

## **Observations of preening behaviour**

An observer recorded a total of 95 independent behavioural sessions of 5 minutes each. The observer remained outside the aviary and waited a few minutes before starting her observations not to influence bird behaviour. The observed birds were chosen randomly but the observer switched to a new treatment for each new session to have a similar number of observations for all treatments: 25 observations (corresponding to 21 different individuals) among the *control* group, 29 observations (24 individuals) among the *lead* group, 22 observations (20 individuals) among the *zinc* group and 19 observations (16 individuals) along the *lead+zinc* group. The behaviour recording was performed with the JWatcher software. We recorded the time birds spent preening; during this behaviour, the plumage may be spread with preen secretions (Mardon et al. 2011), which have antimicrobial and antiparasite properties (e.g. in feral pigeons, house sparrows, eastern bluebirds and hoopoes; Moyer et al. 2003, Shawkey et al. 2007, Ruiz-Rodriguez et al. 2009, Waite et al. 2012, Czirják et al. 2013). Because preening is costly (Piault et al. 2008, Moreno-Rueda 2010), the time allocated to preening has been shown to be adjusted to bacterial load (Leclaire et al.

2014b), and to reflect birds' health status. For instance, juvenile apapanes (*Himationesanguinea*) experimentally infected with *Plasmodium relictum* spent less time preening (Yorinks and Atkinson 2000).

## **Statistical analyses**

Statistical analyses were performed using R (R.3.0.2; R Development Core Team). Final models were retained based on their AIC.

To test the effects of metal exposure on the composition of bacterial communities, we performed a PERMANOVA with 5000 permutations (i.e. nonparametric multivariate analysis of variance, Adonis function, VEGAN package in R (Oksanen et al. 2007), based on Jaccard distance for OTU presence/absence data. Zinc exposure, lead exposure, sex and their interactions were included as explanatory variables. Because spatial proximity may influence bacterial community similarity between individuals, we added the aviary as a covariate. Then, we ran similar analyses between each pair of treatments. Finally, we tested the differences of bacterial communities between aviaries among each metal treatment.

To investigate more precisely the effect of spatial proximity on bacterial communities' similarities, we compared a matrix of bacterial Jaccard distances between individuals to a matrix of spatial distances (scored as 0 for individuals inhabiting the same aviary, to 7 for the most distant individuals), considering a matrix of treatment membership (scored as 0 for individuals submitted to the treatment and 1 for individuals of different treatments) using partial mantel test with 5000 permutations.

We graphically represented similarities between individuals using a constrained redundancy analysis (RDA function in R) based on the Jaccard distance matrix.

We also tested plumage bacterial richness using a generalized linear model for Poisson distribution with zinc exposure, lead exposure, sex and their interactions as explanatory variables, the weight of feathers used for the analysis as a covariate and aviary as random effect.

To test the effects of metal exposure on plumage bacterial load, we performed Wilcoxon tests because our sample size was low (n=10 per treatment). First, we tested the effects of zinc and the effects of lead in two different tests; then, we performed Wilcoxon tests between each pair of treatments to test the effects of the interaction between zinc and lead exposure.

Finally, we investigated the amount of time birds allocated to preening by performing a generalized linear mixed model for Poisson distribution with zinc exposure, lead exposure, sex and their interactions as explicative variables and bird identity and aviary as random effects.

All the previously described models were performed on the totality of the birds, whatever their plumage colouration (i.e. eumelanic and pheomelanic birds). The same models were performed on eumelanic birds only (i.e. excluding pheomelanic pigeons). In these models, plumage eumelanin level and its interaction with the other considered parameters were added as explanatory variables.

## **RESULTS**

The composition of bacterial communities depended on the interaction between lead and zinc exposure ( $F_{1,82}=3.47$ ,  $P<0.001$ ; Fig. 1) and on aviaries ( $F_{1,82}=3.91$ ,  $P<0.001$ ). Each pairwise test between metal exposures was significant (Table 2). The composition of bacterial communities differed significantly between aviaries among each metal treatment but was less dissimilar among the *lead* group (zinc-exposure:  $F_{1,19}=3.48$ ,  $P<0.001$ , lead-exposure:  $F_{1,21}=1.70$ ,  $P=0.018$ , zinc and lead-exposure:  $F_{1,19}=4.48$ ,  $P<0.001$ , control:  $F_{1,20}=7.59$ ,  $P<0.001$ ). Moreover, there was a highly significant positive correlation between bacterial distance and spatial distance ( $r=0.30$ ,  $P<0.001$ ; Fig. 2).

Plumage bacterial richness depended on the interaction between zinc and lead-exposure ( $\chi^2=9.09$ ,  $df=80$ ,  $P=0.003$ ; Fig. 3): the *lead* group had lower bacterial richness than the *control* group ( $\chi^2=5.58$ ,  $df=42$ ,  $P=0.018$ ) and the *zinc+lead* group ( $\chi^2=12.63$ ,  $df=41$ ,  $P<0.001$ ). Moreover, the *lead+zinc* group tended to have higher bacterial richness than the *zinc* group ( $\chi^2=3.63$ ,  $df=40$ ,  $P=0.057$ ).

Plumage bacterial load was significantly lower among birds exposed to zinc (*zinc* and *lead+zinc* groups) than among the others (*lead* and *control* groups;  $W=280.5$ ,  $P=0.029$ ; Fig. 4). Although there was no significant difference between each pair of treatments ( $P>0.067$ ), the *zinc* group tended to have lower plumage bacterial loads than the *control* group ( $W=29$ ,  $P=0.072$ ) and the *lead* group ( $W=25$ ,  $P=0.067$ ). Lead did not significantly affect plumage bacterial load ( $W=181$ ,  $P=0.623$ ).

Finally, the time birds spent preening depended on the interaction between zinc and lead exposure ( $\chi^2=3.97$ ,  $df=92$ ,  $P=0.04$ ). We performed partial models to compare each pair of treatments. Although there was no significant difference between each pair of treatments

( $P > 0.101$ ), our results suggest that time spent preening was shorter among the *lead* group than among the other groups (*control*:  $65.00 \pm 15.18$ s, *lead*:  $34.21 \pm 9.69$ s, *zinc*:  $53.86 \pm 13.79$ s, *lead+zinc*:  $60.84 \pm 15.30$ s).

Sex and plumage eumelanin level were retained in none of the tested models.

## DISCUSSION

As expected, the composition of plumage bacterial community varied with metal exposure. The exposure to lead alone appears to induce the strongest effect. Plumage bacterial composition was more similar amongst the two aviaries hosting birds exposed to lead only than expected if considering a spatial effect only. In addition, the plumage bacterial composition of birds exposed to lead only were the most distant from the communities of the other treatments. Moreover, birds of the *lead* group had reduced plumage bacterial richness compared to birds of the *control* group. These results suggest that lead may select for lead-tolerant plumage bacteria. To the best of our knowledge, our study is the first to show that lead exposure has effects on plumage bacterial community. It is consistent with a previous study showing that lead alters the intestinal microbiome of mice (Breton et al. 2013). Moreover, birds of the *lead* group tended to preen less frequently than birds of the *control* group. Lead exposure decreases bird immunity (unpublished results) and reproductive success (but there is no effect of metal exposure on bird breeding success; Chatelain et al. in press). Lead, by decreasing bird condition, may affect bird ability to preen, a costly behaviour that helps to regulate feather microbiota (Piault et al. 2008, Moreno-Rueda 2010, Leclaire et al. 2014b). Because preening is adjusted to feather bacterial load (Leclaire et al. 2014b), the change in

preening frequency observed in our study may also non-exclusively result from the change in feather bacterial community caused by trace metal exposure. Whatever the mechanism underlying the differences in preening frequency, the tendency of lead to reduce bird control on its plumage bacterial community may change the dominant status of bacteria species and therefore induce the proliferation of species that were previously sensitive to preen secretions. High-throughput DNA sequencing would help identifying lead-tolerant bacteria species and therefore inferring their potential pathogenicity and propensity to degrade feathers. More analyses should also be conducted to identify the proximal mechanisms involved in lead toxicity. For instance, *in vitro* exposure of feathers to these metals would allow us to disentangle the direct and indirect effects that may induce these metals.

Like lead exposure, zinc exposure had toxic effects on the plumage bacterial community with birds exposed to zinc exhibiting lower bacterial load than control birds. Similarly, high doses of zinc decrease bacterial load and change bacterial community in the gastrointestinal tract of piglets (Højberg et al. 2005, Vahjen et al. 2010), and inhibit bacterial growth in sludge and sediment (Cabrero et al. 1998, Vega-López et al. 2007). Zinc is known to be essential to several metabolic functions of bacteria (Sugarman 1983). At high concentrations, zinc can however reduce protein and ATP content, interact with nucleic acids and enzyme active sites, decrease membrane health and eventually lead to cell necrosis (Martinez-Tabche and Gutierrez 2000, Vega-López et al. 2007). Although the concentration of zinc we used is within the natural range found in cities, it may be high enough to negatively affect feather bacteria and to decrease bacterial load. Zinc may also affect feather bacterial load indirectly through its immunostimulating effect (Smith 2003). In feral pigeons, zinc has a positive effect on the production of specific antibodies (unpublished data), and might, therefore, increase the bactericidal capacity of uropygial secretions.



343

344 Bacterial community composition, bacterial richness and the time birds spent preening  
345 depended on the interaction between lead and zinc exposure. More precisely, the toxic effects  
346 of lead exposure was not detected in birds exposed to both lead and zinc, suggesting that zinc  
347 may compensate lead toxicity. Zinc is known to reduce the absorption and retention of  
348 ingested lead (Cerklewski and Forbes 1976, El-Gazzar et al. 1978, Prasanthi et al. 2010),  
349 which may therefore reduce the negative effects of lead on bird condition. In addition, the  
350 negative effect of zinc exposure on bacterial load would be higher in birds exposed to zinc  
351 only than in birds exposed to both lead and zinc, suggesting again an interaction between lead  
352 and zinc exposure.

353

354 Our results showed a strong effect of spatial proximity on bird plumage bacterial community,  
355 with birds in closer aviaries showing more similar bacterial communities. In accordance with  
356 other studies (Bisson et al. 2007, 2009, Saag et al. 2011), they point out the relatively small  
357 spatial scale transmission of plumage bacteria. While bacteria are likely transmitted through  
358 close contacts (Kulkarni and Heeb 2007) and reciprocal delousing, some bacteria may be able  
359 to survive on non-feather substrates and, therefore, be transmitted through bathwater, perches,  
360 soil and the grids separating the aviaries (Bisson et al. 2007). Because pigeons live in high  
361 density but have limited movements within their local environment (Frantz et al. 2012), the  
362 plumage bacterial community of wild pigeons may, therefore, greatly vary between  
363 populations, which may lead to local coevolution and co-adaptation between the host and its  
364 bacterial community. The strong effect of spatial proximity on plumage bacterial community  
365 detected in our study may have decreased and increased bacterial community similarities  
366 between and within aviaries respectively; ideally, future studies should increase the number of  
367 replicates (aviaries) or house pigeons in individual and spatially distant cages.

368

369 Our experimental exposure of feral pigeons to naturally occurring concentrations of lead  
370 and/or zinc highlights, for the first time, the effects of some trace metals commonly  
371 encountered in urban areas on plumage bacterial community. The birds used in our study were  
372 captured in Paris, and had therefore been previously exposed to trace metals in their natural  
373 urban habitat. Consequently, plumage bacterial communities at the start of the experiment  
374 might have already been shaped by past metal exposure. Because there was no significant  
375 correlation between bird capture site (i.e. pigeon flock) and aviary, the potential initial  
376 differences in bacterial community between the birds would, however, have reduced the  
377 power of our analysis, and the significant differences between treatments observed in our  
378 study are therefore conservative.

379 Although our knowledge on plumage bacterial community composition and function is  
380 scarce, feather bacteria seem to play a role in bird immunity, reproduction and feather  
381 colouration and condition (Clayton 1999, Shawkey et al. 2007, Gunderson et al. 2009,  
382 Leclaire et al. 2014, 2015, Jacob et al. 2015 but see Jacob et al. 2014). Through their effects  
383 on plumage bacteria, trace metals may, for instance, affect thermoregulation and visual  
384 signals potentially involved in dominant status assessment and mate choice (Wolf 2000, Hill  
385 and McGraw 2006). Future studies should now investigate if and how the changes in plumage  
386 bacterial communities induced by trace metals affect bird fitness.

387

388

## 389 **ACKNOWLEDGMENTS**

390

391 We thank the “Mairie de Paris” (Thomas Charachon) for allowing the capture of birds and the  
392 Centre de Recherche en Ecologie Expérimentale et Prédictive (CEREEP) which provided

393 logistic support for the field work of this study. We are very thankful to T. Gayet, S. Pollet, S.  
394 Hasnaoui, F. Lorente, S. Perret and B. Decenci re for their help all along the field work.  
395 Molecular analyses were partly performed at the technical facilities of the labexCeMEB and  
396 were funded by an “ANR PDOC” grant (to SL) (No. ANR-13-PDOC-0002).

## REFERENCES

- Adout, A., Hawlena, D., Maman, R., Paz-Tal, O. and Karpas, Z. 2007. Determination of trace elements in pigeon and raven feathers by ICPMS. - *Int. J. Mass Spectrom.* 267: 109–116.
- Agusa, T., Matsumoto, T., Ikemoto, T., Anan, Y., Kubota, R., Yasunaga, G., Kunito, T., Tanabe, S., Ogi, H. and Shibata, Y. 2005. Body distribution of trace elements in black-tailed gulls from Rishiri Island, Japan: Age-dependent accumulation and transfer to feathers and eggs. - *Environ. Toxicol. Chem.* 24: 2107–2120.
- Archie, E. A. and Theis, K. R. 2011. Animal behaviour meets microbial ecology. - *Anim. Behav.* 82: 425–436.
- Azimi, S., Rocher, V., Muller, M., Moilleron, R. and Thevenot, D. R. 2005. Sources, distribution and variability of hydrocarbons and metals in atmospheric deposition in an urban area (Paris, France). - *Sci. Total Environ.* 337: 223–239.
- Babich, H. and Stotzky, G. 1978. Toxicity of zinc to fungi, bacteria, and coliphages: influence of chloride ions. - *Appl. Environ. Microbiol.* 36: 906–914.
- Balcázar, J. L., Vendrell, D., de Blas, I., Ruiz-Zarzuela, I., Gironés, O. and Múzquiz, J. L. 2007. In vitro competitive adhesion and production of antagonistic compounds by lactic acid bacteria against fish pathogens. - *Vet. Microbiol.* 122: 373–380.
- Berglund, A., Sturve, J., Forlin, L. and Nyholm, N. 2007. Oxidative stress in pied flycatcher (*Ficedula hypoleuca*) nestlings from metal contaminated environments in northern Sweden. - *Environ. Res.* 105: 330–339.

419 Bisson, I.-A., Marra, P. P., Burt, E. H., Sikaroodi, M. and Gillevet, P. M. 2007. A molecular  
 420 comparison of plumage and soil bacteria across biogeographic, ecological, and  
 421 taxonomic scales. - *Microb. Ecol.* 54: 65–81.

422 Bisson, I.-A., Marra, P. P., Burt Jr, E. H., Sikaroodi, M. and Gillevet, P. M. 2009. Variation  
 423 in Plumage Microbiota Depends on Season and Migration. - *Microb. Ecol.* 58: 212–  
 424 220.

425 Breton, J., Massart, S., Vandamme, P., De Brandt, E., Pot, B. and Follign , B. 2013.  
 426 Ecotoxicology inside the gut: impact of heavy metals on the mouse microbiome. -  
 427 *BMC Pharmacol. Toxicol.* 14: 62.

428 Burkholder, K. M., Thompson, K. L., Einstein, M. E., Applegate, T. J. and Patterson, J. A.  
 429 2008. Influence of stressors on normal intestinal microbiota, intestinal morphology,  
 430 and susceptibility to *Salmonella enteritidis* colonization in broilers. - *Poult. Sci.* 87:  
 431 1734–1741.

432 Burt Jr, E. H. and Ichida, J. M. 1999. Occurrence of feather-degrading bacilli in the plumage  
 433 of birds. - *The Auk* 116: 364–372.

434 Cabrero, A., Fernandez, S., Mirada, F. and Garcia, J. 1998. Effects of copper and zinc on the  
 435 activated sludge bacteria growth kinetics. - *Water Res.* 32: 1355–1362.

436 Caro, S. P., Balthazart, J. and Bonadonna, F. 2015. The perfume of reproduction in birds:  
 437 Chemosignaling in avian social life. - *Horm. Behav.* 68: 25–42.

438 Cerklewski, F. L. and Forbes, R. M. 1976. Influence of dietary zinc on lead toxicity in the rat.  
 439 - *J. Nutr.* 106: 689–696.

- 440 Chatelain, M., Gasparini, J., Jacquin, L. and Frantz, A. 2014. The adaptive function of  
441 melanin-based plumage colouration to trace metals. - Biol. Lett. 10: 20140164–  
442 20140164.
- 443 Chatelain, M., Gasparini, J. and Frantz, A. In press. Do trace metals select for darker birds in  
444 urban areas? An experimental exposure to lead and zinc. – Glob. Change Biol.
- 445 Clayton, D. H. 1999. Feather-busting bacteria. - The Auk 116: 302–304.
- 446 Cosson, R. P., Amiard, J.-C. and Amiard-Triquet, C. 1988. Trace elements in little egrets and  
447 flamingos of Camargue, France. - Ecotoxicol. Environ. Saf. 15: 107–116.
- 448 Czirják, G. Á., Pap, P. L., Vágási, C. I., Giraudeau, M., Mureşan, C., Mirleau, P. and Heeb, P.  
449 2013. Preen gland removal increases plumage bacterial load but not that of feather-  
450 degrading bacteria. - Naturwissenschaften 100: 145–151.
- 451 Dauwe, T., Janssens, E., Pinxten, R. and Eens, M. 2005. The reproductive success and quality  
452 of blue tits (*Paruscaeruleus*) in a heavy metal pollution gradient. - Environ. Pollut.  
453 136: 243–251.
- 454 Dauwe, L. Bervoets, R. Blust, M. Ee, T. 2002. Tissue levels of lead in experimentally  
455 exposed zebra finches (*Taeniopygiaguttata*) with particular attention on the use of  
456 feathers as biomonitors. - Arch. Environ. Contam. Toxicol. 42: 88–92.
- 457 Dotterud, L. K. are, Wilsgaard, T., Vorland, L. H. and Falk, E. S. 2008. The effect of UVB  
458 radiation on skin microbiota in patients with atopic dermatitis and healthy controls. -  
459 Int. J. Circumpolar Health in press.

460 Eeva, T., Ahola, M. and Lehikoinen, E. 2009. Breeding performance of blue tits  
 461 (*Cyanistescaeruleus*) and great tits (*Parus major*) in a heavy metal polluted area. -  
 462 Environ. Pollut. 157: 3126–3131.

463 El-Gazzar, R. M., Finelli, V. N., Boiano, J. and Petering, H. G. 1978. Influence of dietary zinc  
 464 on lead toxicity in rats. - Toxicol. Lett. 1: 227–234.

465 Ezenwa, V. O., Gerardo, N. M., Inouye, D. W., Medina, M. and Xavier, J. B. 2012. Animal  
 466 behaviour and the microbiome. - Science 338: 198–199.

467 Frank, D. N., Robertson, C. E., Hamm, C. M., Kpadeh, Z., Zhang, T., Chen, H., Zhu, W.,  
 468 Sartor, R. B., Boedeker, E. C., Harpaz, N., Pace, N. R. and Li, E. 2011. Disease  
 469 phenotype and genotype are associated with shifts in intestinal-associated microbiota  
 470 in inflammatory bowel diseases: - Inflamm. Bowel Dis. 17: 179–184.

471 Frantz, A., Pottier, M.-A., Karimi, B., Corbel, H., Aubry, E., Haussy, C., Gasparini, J. and  
 472 Castrec-Rouelle, M. 2012. Contrasting levels of heavy metals in the feathers of urban  
 473 pigeons from close habitats suggest limited movements at a restricted scale. - Environ.  
 474 Pollut. 168: 23–28.

475 Giller, K. E., Witter, E. and Mcgrath, S. P. 1998. Toxicity of heavy metals to microorganisms  
 476 and microbial processes in agricultural soils: a review. - Soil Biol. Biochem. 30:  
 477 1389–1414.

478 Griffiths, R., Double, M. C., Orr, K. and Dawson, R. J. G. 1998. A DNA test to sex most  
 479 birds. - Mol. Ecol. 7: 1071–1075.

480 Gulson, B., Mizon, K. J., Korsch, M. J., Howarth, D., Phillips, A. and Hall, J. 1996. Impact on  
 481 blood lead in children and adults following relocation from their source of exposure

482 and contribution of skeletal tissue to blood lead. - Bull. Environ. Contam. Toxicol. 56:  
483 543–550.

484 Gunderson, A. R., Forsyth, M. H. and Swaddle, J. P. 2009. Evidence that plumage bacteria  
485 influence feather colouration and body condition of eastern bluebirds *Sialiasialis*. - J.  
486 Avian Biol. 40: 440–447.

487 Hildebrand, F., Nguyen, T. L., Brinkman, B., Yunta, R. G., Cauwe, B., Vandenabeele, P.,  
488 Liston, A. and Raes, J. 2013. Inflammation-associated enterotypes, host genotype,  
489 cage and inter-individual effects drive gut microbiota variation in common laboratory  
490 mice. - Genome Biol 14: R4.

491 Hill, M. 1997. Intestinal flora and endogenous vitamin synthesis. - Eur. J. Cancer Prev.6: S43-  
492 S45.

493 Hill, G. E. and McGraw, K. J. 2006. Bird colouration. - Harvard University Press.

494 Hoff Brait, C. H. and AntoniosiFilho, N. R. 2011. Use of feathers of feral pigeons (*Columba*  
495 *livia*) as a technique for metal quantification and environmental monitoring. - Environ.  
496 Monit. Assess. 179: 457–467.

497 Hojberg, O., Canibe, N., Poulsen, H. D., Hedemann, M. S. and Jensen, B. B. 2005. Influence  
498 of dietary zinc oxide and copper sulfate on the gastrointestinal ecosystem in newly  
499 weaned piglets. - Appl. Environ. Microbiol. 71: 2267–2277.

500 Hsu, M. J., Selvaraj, K. and Agoramoorthy, G. 2006. Taiwan's industrial heavy metal  
501 pollution threatens terrestrial biota. - Environ. Pollut. 143: 327–334.

502 Jacob, S., Colmas, L., Parthuisot, N. and Heeb, P. 2014. Do feather-degrading bacteria  
503 actually degrade feather colour? No significant effects of plumage microbiome



504 modifications on feather colouration in wild great tits. - *Naturwissenschaften* 101:  
505 929–938.

506 Jacob, S., Parthuisot, N., Vallat, A., Ramon-Portugal, F., Helfenstein, F. and Heeb, P. 2015.  
507 Microbiome affects egg carotenoid investment, nestling development and adult  
508 oxidative costs of reproduction in great tits. - *Funct. Ecol* 8 : 1048-1058.

509 Jacquin, L., Lenouvel, P., Haussy, C., Ducatez, S. and Gasparini, J. 2011. Melanin-based  
510 colouration is related to parasite intensity and cellular immune response in an urban  
511 free living bird: the feral pigeon *Columba livia*. - *J. Avian Biol.* 42: 11–15.

512 Jarup, L. 2003. Hazards of heavy metal contamination. - *Br. Med. Bull.* 68: 167–182.

513 Johnston, R. and Janiga, M. 1995. Feral pigeons. - Oxford University Press.

514 Kekkonen, J., Hanski, I. K., Väisänen, R. A. and Brommer, J. E. 2012. Levels of heavy metals  
515 in house sparrows(*Passer domesticus*) from urban and rural habitats of southern  
516 Finland. - *OrnisFenn.* 89: 91-98.

517 Kilgas, P., Saag, P., Mägi, M., Edenberg, M., Tilgar, V. and Mänd, R. 2012. Variation in  
518 assemblages of feather bacteria in relation to plumage color in female great tits. - *The*  
519 *Condor* 114: 606–611.

520 Kim, E. Y., Goto, R., Tanabe, S., Tanaka, H. and Tatsukawa, R. 1998. Distribution of 14  
521 elements in tissues and organs of oceanic seabirds. - *Arch. Environ. Contam. Toxicol.*  
522 35: 638–645.

523 Kulkarni, S. and Heeb, P. 2007. Social and sexual behaviours aid transmission of bacteria in  
524 birds. - *Behav. Processes* 74: 88–92.

525 Leclaire, S., Nielsen, J. F. and Drea, C. M. 2014a. Bacterial communities in meerkat anal  
526 scent secretions vary with host sex, age, and group membership. - Behav. Ecol. 25:  
527 996–1004.

528 Leclaire, S., Pierret, P., Chatelain, M. and Gasparini, J. 2014b. Feather bacterial load affects  
529 plumage condition, iridescent colour, and investment in preening in pigeons. - Behav.  
530 Ecol. 25: 1192–1198.

531 Leclaire, S., Czirjak, G., Hammouda, A. and Gasparini, J. 2015. Feather bacterial load shapes  
532 the trade-off between preening and immunity in pigeons. BMC Evolutionary Biology  
533 15: 60.

534 Liu, Y., Li, Y., Liu, K. and Shen, J. 2014. Exposing to cadmium stress cause profound toxic  
535 effect on microbiota of the mice intestinal tract. - PLoS ONE 9: e85323.

536 Mardon, J., Saunders, S. M. and Bonadonna, F. 2011. From preen secretions to plumage: the  
537 chemical trajectory of blue petrels' *Halobaenacaerulea* social scent. - J. Avian Biol.  
538 42: 29–38.

539 Martinez-Tabche, L. and Gutierr, I. 2000. Toxic effects of zinc from trout farm sediments on  
540 ATP, protein, and hemoglobin concentrations of *Limnodrilushoffmeisteri*. - J. Toxicol.  
541 Environ. Health A 59: 575–583.

542 Mertz, W. 1981. The essential trace elements. - Science 213: 1332–1338.

543 Moreno-Rueda, G. 2010. Uropygial gland size correlates with feather holes, body condition  
544 and wingbar size in the house sparrow *Passer domesticus*. - J. Avian Biol. 41: 229–  
545 236.

546 Moyer, B. R., Rock, A. N. and Clayton, D. H. 2003. Experimental test of the importance of  
547 preen oil in rock doves (*Columba livia*). - The Auk 120: 490.

548 Mueller, S., Saunier, K., Hanisch, C., Norin, E., Alm, L., Midtvedt, T., Cresci, A., Silvi, S.,  
549 Orpianesi, C., Verdenelli, M. C., Clavel, T., Koebnick, C., Zunft, H.-J. F., Dore, J. and  
550 Blaut, M. 2006. Differences in fecal microbiota in different european study  
551 populations in relation to age, gender, and country: a cross-sectional study. - Appl.  
552 Environ. Microbiol. 72: 1027–1033.

553 Muza, M. M., Burt, E. H. and Ichida, J. M. 2000. Distribution of bacteria on feathers of some  
554 eastern North American birds. - Wilson Bull. 112: 432–435.

555 Nam, D.-H., Lee, D.-P. and Koo, T.-H. 2004. Monitoring for lead pollution using feathers of  
556 feral pigeons (*Columba livia*) from Korea. - Environ. Monit. Assess. 95: 13–22.

557 Oh, S., Kim, S.-H., Ko, Y., Sim, J.-H., Kim, K. S., Lee, S.-H., Park, S. and Kim, Y. J. 2006.  
558 Effect of bacteriocin produced by *Lactococcus* sp. HY 449 on skin-inflammatory  
559 bacteria. - Food Chem. Toxicol. 44: 552–559.

560 Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M., Oksanen, M. and Suggests, M.  
561 2007. The vegan package. Community ecology package 661-637.

562 Olsson, J. C., Westerdahl, A., Conway, P. L. and Kjelleberg, S. 1992. Intestinal colonization  
563 potential of turbot (*Scophthalmus maximus*)-and dab (*Limandalimanda*)-associated  
564 bacteria with inhibitory effects against *Vibrio anguillarum*. - Appl. Environ.  
565 Microbiol. 58: 551–556.

566 Pattee, O. H. 1984. Eggshell thickness and reproduction in American kestrels exposed to  
567 chronic dietary lead. - Arch. Environ. Contam. Toxicol. 13: 29–34.

- 568 Piault, R., Gasparini, J., Bize, P., Paulet, M., McGraw, K. J. and Roulin, A. 2008.  
569 Experimental support for the makeup hypothesis in nestling tawny owls (*Strixaluco*). -  
570 Behav. Ecol. 19: 703–709.
- 571 Pilastro, A., Congiu, L., Tallandini, L. and Turchetto, M. 1993. The use of bird feathers for  
572 the monitoring of cadmium pollution. - Arch. Environ. Contam. Toxicol. 24: 355–358.
- 573 Prasad, A. S. 1998. Zinc and immunity. - Mol. Cell. Biochem. 188: 63–69.
- 574 Prasanthi, R. P. J., Devi, C. B., Basha, D. C., Reddy, N. S. and Reddy, G. R. 2010. Calcium  
575 and zinc supplementation protects lead (Pb)-induced perturbations in antioxidant  
576 enzymes and lipid peroxidation in developing mouse brain. - Int. J. Dev. Neurosci. 28:  
577 161–167.
- 578 Ranjard, L., Brothier, E. and Nazaret, S. 2000. Sequencing bands of ribosomal intergenic  
579 spacer analysis fingerprints for characterization and microscale distribution of soil  
580 bacterium populations responding to mercury spiking. - Appl. Environ. Microbiol. 66:  
581 5334–5339.
- 582 Redig, P. T., Lawler, E. M., Schwartz, S., Dunnette, J. L., Stephenson, B. and Duke, G. E.  
583 1991. Effects of chronic exposure to sublethal concentrations of lead acetate on heme  
584 synthesis and immune function in red-tailed hawks. - Arch. Environ. Contam. Toxicol.  
585 21: 72–77.
- 586 Reid, C., McInnes, K., McLelland, J. M. and Gartrell, B. D. 2012. Anthropogenic lead (Pb)  
587 exposure in populations of a wild parrot (kea *Nestor notabilis*). - N. Z. J. Ecol. 36: 56.

588 Rosenthal, M., Goldberg, D., Aiello, A., Larson, E. and Foxman, B. 2011. Skin microbiota:  
 589 Microbial community structure and its potential association with health and disease. -  
 590 Infect. Genet. Evol. 11: 839–848.

591 Roux, K. E. and Marra, P. P. 2007. The presence and impact of environmental lead in  
 592 passerine birds along an urban to rural land use gradient. - Arch. Environ. Contam.  
 593 Toxicol. 53: 261–268.

594 Ruiz-de-Castañeda, R., Vela, A. I., Lobato, E., Briones, V. and Moreno, J. 2011. Bacterial  
 595 loads on eggshells of the pied flycatcher: environmental and maternal factors. - The  
 596 Condor 113: 200–208.

597 Ruiz-Rodriguez, M., Valdivia, E., Soler, J. J., Martin-Vivaldi, M., Martin-Platero, A. M. and  
 598 Martinez-Bueno, M. 2009. Symbiotic bacteria living in the hoopoe's uropygial gland  
 599 prevent feather degradation. - J. Exp. Biol. 212: 3621–3626.

600 Saag, P., Tilgar, V., Mänd, R., Kilgas, P. and Mägi, M. 2011. Plumage bacterial assemblages  
 601 in a breeding wild passerine: relationships with ecological factors and body condition.  
 602 - Microb. Ecol. 61: 740–749.

603 Scheifler, R., Cœurdassier, M., Morilhat, C., Bernard, N., Faivre, B., Flicoteaux, P.,  
 604 Giraudoux, P., Noël, M., Piotte, P., Rieffel, D., de Vaufleury, A. and Badot, P.-M.  
 605 2006. Lead concentrations in feathers and blood of common blackbirds  
 606 (*Turdusmerula*) and in earthworms inhabiting unpolluted and moderately polluted  
 607 urban areas. - Sci. Total Environ. 371: 197–205.

608 Shawkey, M. D. and Hill, G. E. 2004. Feathers at a fine scale. - The Auk 121: 652.

- 609 Shawkey, M. D., Pillai, S. R. and Hill, G. E. 2003. Chemical warfare? Effects of uropygial oil  
610 on feather-degrading bacteria. - J. Avian Biol. 34: 345–349.
- 611 Shawkey, M. D., Pillai, S. R., Hill, G. E., Siefferman, L. M. and Roberts, S. R. 2007. Bacteria  
612 as an agent for change in structural plumage colour: correlational and experimental  
613 evidence. - Am. Nat. 169: S112–S121.
- 614 Smith, M. O. 2003. Effects of different levels of zinc on the performance and  
615 immunocompetence of broilers under heat stress. - Poult. Sci. 82: 1580–1588.
- 616 Snoeijs, T., Dauwe, T., Pinxten, R., Vandesande, F. and Eens, M. 2004. Heavy metal  
617 exposure affects the humoral immune response in a free-living small songbird, the  
618 great tit (*Parus major*). - Arch. Environ. Contam. Toxicol. 46: 399-404.
- 619 Sugarman, B. 1983. Zinc and Infection. - Clin. Infect. Dis. 5: 137–147.
- 620 Vahjen, W., Pieper, R. and Zentek, J. 2010. Bar-Coded Pyrosequencing of 16S rRNA gene  
621 amplicons reveals changes in ileal porcine bacterial communities due to high dietary  
622 zinc intake. - Appl. Environ. Microbiol. 76: 6689–6691.
- 623 Vega-López, A., Amora-Lazcano, E., López-López, E., Terrón, O. and Proal-Nájera, J. B.  
624 2007. Toxic effects of zinc on anaerobic microbiota from Zimapán Reservoir  
625 (Mexico). - Anaerobe 13: 65–73.
- 626 Waite, J. L., Henry, A. R. and Clayton, D. H. 2012. How effective is preening against mobile  
627 ectoparasites? An experimental test with pigeons and hippoboscids. - Int. J.  
628 Parasitol. 42: 463–467.
- 629 Whitaker, J. M., Cristol, D. A. and Forsyth, M. H. 2005. Prevalence and genetic diversity of  
630 *Bacillus licheniformis* in avian plumage. - J. Field Ornithol. 76: 264–270.

- 631 Wolf, B. O. 2000. The role of the plumage in heat transfer processes of birds. - Integr. Comp.  
632 Biol. 40: 575–584.
- 633 Yorinks, N. and Atkinson, C. T. 2000. Effects of malaria on activity budgets of  
634 experimentally infected juvenile apapane (*Himationesanguinea*). - The Auk 117: 731.

## FIGURE LEGENDS

**Fig. 1** Constrained redundancy analysis (RDA function in R) on bacterial community dissimilarities (estimated by Jaccard distances) between *lead*, *zinc*, *lead and zinc* and *control* groups. Both aviaries of a same treatment are distinguished using either filled or empty circles.

**Fig. 2** Mean  $\pm$  SE plumage bacterial community dissimilarities (estimated by Jaccard distance) in dyads of pigeons according to the spatial distance between them (0 means that the individuals belonged to the same aviary, 1 means that they were in side-by-side aviaries, 2 means that 1 aviary was between them, etc.).

**Fig. 3** Mean  $\pm$  SE plumage bacterial richness (number of different OTUs) according to metal exposure. a and b were significantly different ( $p$ -value $<0.05$ ) while ab was not different from a or from b.

**Fig. 4** Mean  $\pm$  SE plumage bacterial load (number of bacterial colonies per slide) according to metal exposure. Tests were performed for each pair of treatment. a and b tended to be different ( $p$ -value $<0.072$ ) while ab was not different from a or from b.



Table 1. Lead blood and feathers concentrations (mean $\pm$ se, ppb) in lead-exposed and lead non-exposed birds, zinc blood and feathers concentrations (mean $\pm$ se, ppm) in zinc-exposed and zinc non-exposed birds and ANOVAs with lead or zinc concentrations in blood or feathers as dependent variable and zinc or lead exposure as explanatory variable.

		Lead exposed	Lead non-exposed	F	P
Lead	Blood	55.49 $\pm$ 6.54	35.73 $\pm$ 6.77	4.47	0.040
	Feathers	402.31 $\pm$ 35.68	255.81 $\pm$ 49.75	19.61	<0.001
		Zinc exposed	Zinc non-exposed	F	P
Zinc	Blood	4.69 $\pm$ 0.15	4.20 $\pm$ 0.18	5.52	0.022
	Feathers	96.62 $\pm$ 2.15	92.31 $\pm$ 2.25	2.13	0.149

Table 2. PERMANOVAs with 5000 permutations based on Jaccard distance for OTU presence/absence data with the metal exposure as the explicative variable. Bacterial communities' similarities were compared between each pair of metal treatment.

	Control	Zinc-exposure	Lead-exposure
Zinc-exposure	$F_{1,40}=2.74, P=0.002$	-	-
Lead-exposure	$F_{1,42}=3.44, P<0.001$	$F_{1,41}=3.16, P<0.001$	-
Zinc and lead-exposure	$F_{1,40}=4.40, P<0.001$	$F_{1,39}=2.10, P=0.006$	$F_{1,41}=5.13, P<0.001$

Fig. 1




Fig. 2




Fig. 3




Fig. 4

