



HAL
open science

First description of underwater acoustic diversity in three temperate ponds

Camille Desjonquères, Fanny Rybak, Marion Depraetere, Amandine Gasc, Isabelle Le Viol, Sandrine Pavoine, Jérôme Sueur

► **To cite this version:**

Camille Desjonquères, Fanny Rybak, Marion Depraetere, Amandine Gasc, Isabelle Le Viol, et al.. First description of underwater acoustic diversity in three temperate ponds. PeerJ, 2015, 3, pp.e1393. 10.7717/peerj.1393 . hal-01277385

HAL Id: hal-01277385

<https://hal.sorbonne-universite.fr/hal-01277385v1>

Submitted on 22 Feb 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

First description of underwater acoustic diversity in three temperate ponds

Camille Desjonquères^{1,2}, Fanny Rybak², Marion Depraetere¹,
Amandine Gasc¹, Isabelle Le Viol³, Sandrine Pavoine^{4,5} and
Jérôme Sueur¹

¹ Institut de Systématique, Évolution, Biodiversité, ISYEB—UMR 7205—CNRS, MNHN, UPMC, EPHE, Muséum national d'Histoire naturelle, Sorbonne Universités, Rue Buffon, Paris, France

² Université Paris-Sud, Neurosciences Paris-Saclay Institute, UMR 9197, Orsay, France

³ Ecologie et Gestion de la Biodiversité, Centre d'Ecologie et des Sciences de la Conservation (CESCO UMR7204), Sorbonne Universités, MNHN, CNRS, UPMC, Paris, France

⁴ Centre d'Ecologie et des Sciences de la Conservation (CESCO UMR7204), Muséum national d'Histoire naturelle, Paris, France

⁵ Mathematical Ecology Research Group, Department of Zoology, University of Oxford, Oxford, United Kingdom

ABSTRACT

The past decade has produced an increased ecological interest in sonic environments, or soundscapes. However, despite this rise in interest and technological improvements that allow for long-term acoustic surveys in various environments, some habitats' soundscapes remain to be explored. Ponds, and more generally freshwater habitats, are one of these acoustically unexplored environments. Here we undertook the first long term acoustic monitoring of three temperate ponds in France. By aural and visual inspection of a selection of recordings, we identified 48 different sound types, and according to the rarefaction curves we calculated, more sound types are likely present in one of the three ponds. The richness of sound types varied significantly across ponds. Surprisingly, there was no pond-to-pond daily consistency of sound type richness variation; each pond had its own daily patterns of activity. We also explored the possibility of using six acoustic diversity indices to conduct rapid biodiversity assessments in temperate ponds. We found that all indices were sensitive to the background noise as estimated through correlations with the signal-to-noise ratio (SNR). However, we determined that the *AR* index could be a good candidate to measure acoustic diversities using partial correlations with the SNR as a control variable. Yet, research is still required to automatically compute the SNR in order to apply this index on a large data set of recordings. The results showed that these three temperate ponds host a high level of acoustic diversity in which the soundscapes were variable not only between but also within the ponds. The sources producing this diversity of sounds and the drivers of difference in daily song type richness variation both require further investigation. Such research would yield insights into the biodiversity and ecology of temperate ponds.

Submitted 26 August 2015
Accepted 17 October 2015
Published 5 November 2015

Corresponding author
Camille Desjonquères,
cdesjonq@mnhn.fr

Academic editor
Claudio Lazzari

Additional Information and
Declarations can be found on
page 13

DOI 10.7717/peerj.1393

© Copyright
2015 Desjonquères et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Animal Behavior, Aquaculture, Fisheries and Fish Science, Ecology
Keywords Ponds, Sound, Monitoring, Acoustic diversity indices

INTRODUCTION

Over the past 15 years, scientists and land managers have started to draw attention to the importance of ponds in terms of patrimonial, recreational, decorative, agricultural, ecological and environmental interests (Oertli *et al.*, 2009). These water bodies are occupied by a large diversity of organisms harbouring an important number of endemic vertebrate and invertebrate species, some of which are highly threatened (Céréghino *et al.*, 2012). Natural processes such as sedimentation or seasonal changes modify ponds and their environmental parameters, they are thus most of the time bound to be temporary (Wood, Greenwood & Agnew, 2003). Ponds are experiencing an unprecedented and severe degradation due to anthropological causes such as drainage, ancient custom abandon (e.g., forest or agricultural ponds), urbanisation or agriculture intensification (Wood, Greenwood & Agnew, 2003). This degradation can take several forms such as habitat fragmentation, reduction, quality alteration (e.g., pollution or desiccation) or even complete habitat loss. Hence, depending on the area, 40–90% of European ponds have disappeared during the twentieth century (United Kingdom: Wood, Greenwood & Agnew, 2003, Europe: Hull, 1997). Combining a unique diversity and a high level of threat, many ponds can be therefore considered as habitats of high concern for biodiversity conservation, in particular under temperate climates (Verberk *et al.*, 2006).

Plant and animal diversity of ponds has been well studied, however an original facet of biodiversity, the acoustic diversity due to the acoustic signals produced by animals, has been totally neglected. The description and analysis of acoustic diversity was proven to be a valuable approach of diversity assessment combining results in animal behaviour and ecology, in particular to estimate the space and time distribution of species (Towsey, Parsons & Sueur, 2014). The acoustics of freshwater ecosystems have rarely been investigated. Yet they are inhabited by many species belonging to taxa for which terrestrial as well as some aquatic members are known to produce sounds. In ponds, a high diversity of amphibians generates sound underwater during breeding (Duguet & Melki, 2003). Sound production in freshwater arthropods is also quite common but has not been given as much attention. A few crayfish species have been proven to produce sound (Sandeman & Wilkens, 1982; Favaro *et al.*, 2011). Moreover insect stridulation probably makes up most of the acoustic diversity in ponds (Aiken, 1985). A comprehensive review of sound-producing aquatic insects lists 15 families belonging to four orders, namely Trichoptera, Odonata, Coleoptera and the order with the largest number of sound-producing species, Heteroptera (Aiken, 1985). The underwater insect signals cover a wide frequency bandwidth, ranging from 200 Hz for the Coleoptera imago *Acilius sulcatus* (Leston, Pringle & White, 1965) up to 100 kHz for Hydropsychidae larvae (Trichoptera, Silver & Halls, 1980).

Sounds produced by animals can be considered as interacting items belonging to a high-level ecological organization such as a community or a landscape (Farina, 2014). Ecoacoustics, a newly formed discipline, aims at considering animal sound as a material for ecology and biodiversity monitoring (Sueur & Farina, 2015). In this global approach, sound is mainly considered as a tool to infer ecological information. In practice, recordings do not focus on a single singing species but on the overall acoustic output emanating from

a community or a landscape. The analysis of these sounds aims to assess and characterize general features of the structure and the diversity of either an acoustic community, defined as an assemblage of species that share a similar acoustic space (Gasc *et al.*, 2013) or a soundscape, defined as ‘the collection of biological, geophysical and anthropogenic sounds that emanate from a landscape’ (Pijanowski *et al.*, 2011). The global approach is mainly based on the assumption that part of biodiversity can be reflected by acoustic diversity. Numerous acoustic diversity indices have been proposed to measure the acoustic community diversity or soundscape composition (Sueur *et al.*, 2014). All these indices forego species identification and produce relative values that aim at quantifying a feature of the community or the soundscape, like the energy, the complexity or the relative importance of the biophony. The indices have been first tested in terrestrial environments (Sueur *et al.*, 2008; Pieretti, Farina & Morri, 2011; Gage & Axel, 2014; Towsey *et al.*, 2014) and then in marine habitats (Parks, Miksis-Olds & Denes, 2014). These first trials revealed mixed results suggesting the importance of background noise in the reliability of indices (Gasc *et al.*, 2015).

No passive acoustic monitoring study considering all sources of acoustic productions had been conducted in freshwater ponds. We thus explored for the first time the acoustic diversity of three temperate ponds in three different habitats. We tested the following hypothesis: (i) the acoustic diversity differs in richness and composition between the three ponds, (ii) the richness of acoustic diversity varies along day and night revealing ecological cycles, and (iii) acoustic indices can represent the richness of acoustic diversity of each pond detected by human-hearing.

MATERIAL AND METHODS

Study area and recordings

Three ponds were monitored in the Parc Naturel Régional de la Haute Vallée de Chevreuse (PNR), a protected area located 40 km south-west of Paris, France. The ponds were located in three different environments differing by the density of the surrounding vegetation: closed forest habitat (pond 1, 48°34.523’N, 1°53.341’E), semi-closed habitat (pond 2, 48°40.772’N, 1°55.840’E) and open field habitat (pond 3, 48°40.560’N, 1°55.865’E). The main characteristics of the three ponds are summarised in [Table S1](#).

Each pond was monitored with an autonomous recording platform including four units: (1) a hydrophone Reson TC4033 (flat frequency response between 20 Hz and 40 kHz) with a 10 m cable, (2) a charge pre-amplifier Avisoft UltraSoundGate with a frequency high-pass filter at 100 Hz and a gain of +20 dB, (3) a digital audio field recorder SM1 (Wildlife Acoustics, Inc., 2015) with a built-in Texas-Instrument anti-alias filter, (4) a 12 V battery connected to the audio recorder and charged with a solar panel. A four-point linear transect was defined to cross each pond to maximize representation of heterogeneous patches of vegetation. Recordings were achieved on each of the four points of this transect to collect spatial heterogeneity within the ponds. A single recording platform was available for each pond. The hydrophone position was changed every three days according to a sampling rotation with a set up allowing the hydrophone to

move without any intrusion in the ponds. The hydrophone was placed 10 cm below the water surface to reduce heterogeneity of sound propagation that is depth-dependent in shallow water (Forrest, Miller & Zagar, 1993). Rainfall data were collected from a local meteorological station (Météo France, 2015).

Each recorder was programmed to record during one minute every fifteen minutes between the 23rd of June and the 15th of September 2010 (84 days) when the activity and abundance of macro-invertebrate species are known to be maximal (Gascón, Boix & Sala, 2009). The sampling design (3 ponds \times 96 recording time slots \times 84 days) resulted in 24,192 files, among which 7,873 were missing due to technical issues (file corruption, material theft and dysfunction). The final number of files obtained was 16,319. These recordings were sampled at 44.1 kHz with a 16 bits digitization. The files were saved in the lossless compressed format .wac and then transformed into the format .wav with the software WAC to WAV Converter Utility version 1.1 (Wildlife Acoustics, Inc., 2015).

Aural and visual classification of sounds: detection of sound types

Visual identification was conducted on a sub-sample of the initial sample of 16 319 files. Five complete recording days for each sampling point in the three ponds were randomly selected avoiding rainy days. For each selected day, six recording times were defined (00:00, 04:00, 08:00, 12:00, 16:00 and 20:00) resulting in 360 recordings (3 ponds \times 4 sampling points \times 5 days \times 6 recordings per day). These 360 recordings can be considered as samples. 28 recordings, spread across the 3 ponds, had to be withdrawn from this sub-sample due to technical problems with the recorders. Due to the lack of a sound bank for most freshwater species, sounds could not be identified at a species level. Therefore sound types, instead of species-specific sounds, were identified and classified based on aural and visual inspections. Aural inspection was achieved using circumaural headphones and by listening to the files as many times as necessary. This aural inspection was accompanied with the sound visualisation of oscillograms and spectrograms (Window length: 256 samples, frame overlap: 0%, window type: Hanning) with the software Audacity (<http://audacity.sourceforge.net/>). The classification of sound types was based on similarity in amplitude and dominant frequency contours and achieved only by MD to avoid any bias due to the experimenter. This identification of sound types was summarized by two variables for each recordings: the richness of sound types per recording (hereafter referred to as richness) which is the number of different sound types in a recording and the abundance of sound types per recording (hereafter referred to as abundance) which is the total number of sound types detected in a recording.

Signal to noise ratio (SNR)

The signal-to-noise ratio (SNR) of each recording was estimated by computing the ratio between the amplitude of one second of signal (extract lasting one second and containing an identified sound type) and the amplitude of one second of noise (e.g., one second of

recording without any signal) in each file as follows:

$$\text{SNR} = (As/An)^2,$$

with As and An the root mean square (RMS) of signal and noise sections respectively. dB values were obtained by computing:

$$\text{SNR}_{\text{dB}} = 20 * \log_{10} \text{SNR}.$$

The SNR of recordings in which no signal could be found was set to one leading to a SNR_{dB} of zero.

Acoustic analysis

Several acoustic indices have been developed recently to assess the acoustic diversity of a community or a landscape (Sueur *et al.*, 2014). Six acoustic indices were chosen to parametrize the files that were aurally and visually inspected. These indices were (1) the temporal entropy Ht that computes the Shannon evenness of the amplitude envelope (Sueur *et al.*, 2008), (2) the spectral entropy Hf that computes the Shannon evenness of the mean frequency spectrum (Sueur *et al.*, 2008), (3) the envelope energy M that returns the median of the amplitude envelope (Depraetere *et al.*, 2012), (4) the acoustic richness AR which is a ranked index based on the multiplication of Ht and M (Depraetere *et al.*, 2012), (5) the number of major peaks of the mean frequency spectrum NP (Gasc *et al.*, 2013), and (6) the Acoustic Complexity Index ACI which calculates the complexity of the spectrogram, i.e., of the short-term Fourier transform (Pieretti, Farina & Morri, 2011). More details regarding these indices can be found in Sueur *et al.* (2014). All spectral data were obtained with a short-term Fourier transform with a 512 samples non-overlapping Hamming window. To obtain Ht , the absolute amplitude envelope was computed. For NP the parameters used were 1/50 for the amplitude slope threshold and 200 Hz for the frequency threshold. The time step for the ACI was 30 s and the frequency bins were 86 Hz. All acoustic analyses were achieved with the package *seewave* (Sueur, Aubin & Simonis, 2008) of the R statistical environment (R Core Team, 2015).

Statistical analysis

Richness in sound types were compared between ponds with sample-based rarefaction curves (Simberloff, 1972) obtained with the R package *vegan* version 2.2-1 (Oksanen *et al.*, 2013).

A Correspondence Analysis (CA) was computed to characterize the community at each recording point along the transect and the way the communities were ordinated considering their sound type composition. Each recording point was considered as a single observation and the presence/absence data for each sound type was used as a variable.

Differences in SNR between ponds were first assessed with an analysis of variance (ANOVA) but the assumptions of normality and homoscedasticity of the model residuals were not met. A Kruskal–Wallis non-parametric test was therefore used followed by pairwise comparisons using Wilcoxon rank sum test with a Bonferroni adjustment.

To investigate the effect of time in the ponds on sound type richness per recording, we used a Generalized Linear Mixed Model (GLMM; [Baayen, 2008](#)) with a Poisson error structure and log link function ([Mc Cullagh & Nelder, 1989](#)). To examine the daily cyclic effects of time on the richness, we transformed time into a circular variable and included its sine and cosine into the model ([Cox, 2006](#)). Since the effect of time is likely to vary among ponds if they host different species, we included the interaction between pond and the sine and cosine in the model. Transect point and recording day were included as random effects. To keep type I error rate at the nominal level of 5% ([Schielzeth & Forstmeier, 2009](#); [Barr et al., 2013](#)) we included all possible random slopes components (sine and cosine of time within both transect point and recording day and pond within recording day) and also the respective correlations between random slopes and intercepts. As an overall test of the fixed effects, we compared the full model with a null model lacking the fixed effects but comprising the same random effects structure as the full model ([Forstmeier & Schielzeth, 2011](#)) using a likelihood ratio test ([Dobson & Barnett, 2008](#)). We assessed model stability by comparing the estimates derived by a model based on all data with those obtained from models with the levels of random effects excluded one at a time. This revealed the model to be stable. To rule out collinearity we determined Variance Inflation Factors (VIF, [Field, 2005](#)) for a standard linear model excluding random effects and interactions. It revealed a VIF of 1.000 for sine and cosine time and pond which means that there was no collinearity issue.

As data were not normally distributed, correlations between indices and aural analysis were calculated using Spearman's formula. To investigate relationships between acoustic indices and richness and abundance of sounds aurally and visually determined, we performed correlations between acoustic indices (*Ht*, *Hf*, *M*, *AR*, *ACI* and *NP*) and the richness and abundance of sound types. To take into account the effect of the noise on the indices we first estimated the correlation between the acoustic indices (*Ht*, *Hf*, *M*, *AR*, *ACI* and *NP*) and the SNR. Then to control for the correlation between the noise and the indices when investigating the relationship between indices and sound type richness and abundance, we used partial Spearman correlations controlling SNR ([Kim, 2012](#)).

All statistical analyses were run with the R statistical controlling for SNR environment ([R Core Team, 2015](#)) with the packages *ade4* version 1.7-2 ([Dray & Dufour, 2007](#)), *ppcor* version 1.0 ([Kim, 2012](#)), *lme4* version 1.1-7 ([Bates et al., 2014](#)) and *car* version 2.0-25 ([Fox & Weisberg, 2011](#)).

RESULTS

Pond acoustic richness

A total of 2,446 sounds were detected and allocated to 48 sound types (see 3 examples in [Fig. 1](#)). 48 sound types were identified in pond 1, 22 in pond 2, 9 in pond 3. Pond 1 and pond 2 shared 18 sound types, pond 1 and 3 shared 6 sound types and ponds 2 and 3 shared 7 sound types. The rarefaction curve showed a plateau for both pond 2 and 3 but not for pond 1 ([Fig. 2](#)). The plateau is reached at 22 sound types for pond 2 and 9 for pond 3.




Figure 1 Spectrograms and oscillograms of chosen sound productions illustrating the acoustic diversity found in the studied ponds (Fourier window length: 1,024 samples, frame overlap: 50%, window type: Hanning). (A) Sound type 18 recorded in pond 1 on the 9th of July at 12:00 pm. (B) Sound types 24 and 31 recorded in pond 2 on the 3rd of July at 4:00 am. (C) Sound types 30 and 33 recorded in pond 3 on the 14th of July at 4:00 am. Sound type numbers refer to [Supplemental Information](#).

The mean number of sound types found in a recording was 2.2 ± 1.8 (mean \pm sd, $n = 119$) for pond 1, 1 ± 1.1 ($n = 118$) for pond 2 and 0.6 ± 0.8 ($n = 95$) for pond 3. The recording points were more similar within ponds than among ponds as shown by the projection of recording sites along the axes 1 and 2 of a CA (Fig. 3). The two first axes explained 43% of the total variance.

There was an impact of pond and time on the richness (likelihood ratio test comparing the full and null model including only the factor pond, $\chi^2 = 17.269$, $df = 8$, p -value = 0.027). The daily variation was different from one pond to the other (significant interaction between ponds and time, likelihood ratio test comparing the full model and the model without the interaction, $\chi^2 = 10.117$, $df = 4$, p -value = 0.039). Using the predictions of the model, we found that pond 1 had an overall higher number of sound types with its highest sound richness at 11:18, pond 2 had an intermediate sound type richness and its highest sound richness at 20:00 and finally pond 3 had the lowest sound type richness with its highest sound richness at 16:22 (Fig. 4).




Figure 2 Sample based rarefaction curves of sound types per recording for each pond. Each one minute recording is a sample and the sound types are equivalent to species in the rarefaction process. Shaded area around the curves indicates 95% confidence intervals.

Level of SNR

The values of the signal-to-noise ratio had a mean of 5.38 equivalent to 7.31 dB and a median of 1.12 equivalent to 0.5 dB. These values were variable with a standard deviation of 24.51 equivalent to 14 dB, and a median absolute deviation of 1.67 equivalent to 2 dB. 137 out of 332 recordings (41%) had a SNR lower than 1 equivalent to 0 dB. The SNR levels differed significantly between pond 1 and 2 and pond 1 and 3 but not between pond 2 and 3 (pairwise Wilcoxon test, p -values adjusted with the Bonferroni correction: pond 1 – pond 2 < 0.0001, pond 1 – pond 3 < 0.0001, pond 2 – pond 3 = 0.33, $n = 332$). Pond 1 had a significantly higher SNR.

Correlation with the acoustic indices

Correlations between indices and aural analysis revealed that the indices Ht and Hf were negatively correlated with the SNR, the sound type richness and the sound type abundance (Table 1). The indices M , ACI and NP were positively correlated with the SNR and both sound type richness and abundance. The index AR was negatively correlated with SNR but was not correlated either with sound type richness or abundance (Table 1). The richness and abundance of sound types were correlated with the SNR (respectively 0.80 and 0.84, p -values < 0.001). Partial correlations given the SNR revealed that AR was positively




Figure 3 Result of the Correspondence Analysis (CA) with the sound types as variables and the recording points as samples. Each point represents a point of recording in the pond, each ellipse corresponds to 67% of the point dispersion around the centroid for each pond. The axes 1 and 2 explain 26% and 17% of the variance, respectively.

Table 1 Spearman correlations between acoustic indices (*Ht*, *Hf*, *M*, *AR*, *ACI* and *NP*) and richness and abundance of sound types and SNR.

	<i>Ht</i>	<i>Hf</i>	<i>M</i>	<i>AR</i>	<i>NP</i>	<i>ACI</i>
SNR	-0.61***	-0.41***	0.48***	-0.19*	0.42***	0.55***
Richness	-0.5***	-0.3***	0.44***	-0.04	0.34***	0.49***
Abundance	-0.53***	-0.34***	0.47***	-0.06	0.36***	0.5***

Notes.

Stars indicate the significance of the correlation test.
Bonferroni adjusted *p*-value * < 0.05, ** < 0.01, *** < 0.001.

correlated with both the richness and abundance of sound types. All other indices were not significantly correlated with either abundance or richness given the SNR (Table 2).

DISCUSSION

The acoustic production of terrestrial and marine animals has been thoroughly studied for more than 60 years in bioacoustic studies (Fletcher, 2014). The research presented here an important sound diversity that is now reconsidered under the theoretical frameworks of soundscape ecology (Farina, 2014) and ecoacoustics to tackle ecological questions (Sueur & Farina, 2015). So far, the acoustics of animal species inhabiting freshwater habitats has been largely neglected probably because of the lack of flagship species or emblematic




Figure 4 Frequency distribution of sound type richness as a function of time. The size of each point is related to the number of recordings containing the same number of different sound types (total number of recordings $N = 332$). The three dashed lines show the fitted model for each of the three ponds.

Table 2 Spearman partial correlations between acoustic indices (Ht , Hf , M , AR , ACI and NP) and richness and abundance of sound types given the SNR.

	Ht	Hf	M	AR	NP	ACI
Richness	-0.02	0.06	0.11	0.2**	0	0.1
Abundance	-0.04	0	0.15	0.19**	0.02	0.08

Notes.

Stars indicate the significance of the correlation test.
Bonferroni adjusted p -value * < 0.05 , ** $P < 0.01$; ***, $P < 0.001$.

habitats. However, pioneer entomological studies suggested that several aquatic insect species could produce sound (Aiken, 1985) and recent conservation research determined ponds as high concern for biodiversity conservation (Verberk et al., 2006). Here, using a long-term passive acoustic monitoring approach, we (i) identified a total of 48 sound types differing in occurrence between the ponds, (ii) revealed a 24 h cycle of acoustic activity differing among the ponds, and (iii) suggested that the use of AR as an index for automatic monitoring was limited by the need for SNR estimation.

Due to the lack of background research just mentioned above and in particular to the lack of an inventory clearly attributing sound types to species names, we were not able to identify the emitting species of the sound types we inventoried. Only a very small portion of the sound types identified were generated by terrestrial animals (4 sound types were identified as birds) but were rather faint. This probably results from the difference in impedance between air and water making it difficult for sound generated in air to transmit through water. The number of sound types, here named sound richness, is therefore probably an overestimation of the number of singing species as we were not able to assess the intra-specific-diversity and some sound types could also result from plant respiration that generates sonorous air bubbles (*Felisberto et al., 2015*, C Desjonquères, pers. obs., 2014) or from terrestrial animal such as birds.

Even if sound richness could not be directly linked to a number of species, it represents an original facet of biodiversity that can be studied for its own. The three ponds showed different levels and dynamics of sound richness as illustrated with rarefaction curves commonly used to assess sampling effort in biodiversity studies (*Gotelli & Colwell, 2001*).

The number of recordings appeared sufficient to capture the sound richness in the ponds 2 and 3 as the rarefaction curves showed a plateau. Conversely, the rarefaction curve of sound types did not reach a plateau for pond 1 which had the highest sound richness. This suggests that pond 1 embedded a richer and more dynamic acoustic community with a higher diversity of sound types than in the two other ponds.

The distribution of the sound types among the three ponds was different enough to clearly pull apart the three ponds through a correspondence analysis indicating different acoustic communities. This multivariate analysis also revealed a higher heterogeneity of the points in pond 3 than in pond 1 and 2. The Generalized Linear Mixed model confirmed the differences in sound richness among ponds and revealed as well differences along time, the sound richness of each pond evolving in a different way along the 24 h cycle. All together, these results suggest that the three ponds harbour three different communities in terms of richness, composition, and abundance. These differences are in accordance with the three distinct ecological conditions we deliberately chose (open, semi-closed, closed habitats).

The signal to noise ratio (SNR) measures the ratio of the amplitude of the signal of interest over the amplitude of the surrounding noise. It is difficult to have reference values as it depends on several parameters such as surrounding noise level, amplitude of the source, distance from the receiver to the sources, obstacles between the source and the receiver or physical parameters of the matter in which the wave travels (e.g., humidity, temperature, viscosity). Compared to values in a terrestrial habitats which are usually comprised between 15 and 25 dB (*Dabelsteen & Mathevon, 2002*), the SNR was rather low here. A high proportion of recordings had a SNR of 0 dB meaning that they only contained signals which are less or as intense as the background noise. It is now rather difficult to identify whether these low SNR values were due to soft signals and/or loud background noise. The acoustic properties of ponds are unfortunately poorly known (*Aiken, 1982; Forrest, Miller & Zagar, 1993*) compared to marine environments

([Buckingham, 1992](#)). These complex and heterogeneous environments may have very peculiar sound propagation patterns impacting the quality of the recordings, in particular the SNR. Further studies should therefore find solutions to increase the SNR such as removing the background noise with lossless filtering techniques.

Acoustic diversity can be estimated through the identification and count of sound types or species-specific songs. Even if very informative, this approach can be very time demanding when handling large sampling covering hours of audio recordings. Recently, acoustic indices have been developed to get a preliminary estimation of the acoustic diversity without sound or species labelling. We therefore tested six alpha acoustic diversity indices. The correlations between the acoustic indices and the aural analysis showed that the entropy based indices Ht and Hf were negatively correlated with the SNR, the richness and the abundance. This confirms that these metrics are very sensitive to background noise and may function in the reverse way as expected with simulations ([Sueur et al., 2008](#)), as it was already pointed out for bird communities ([Depraetere et al., 2012](#); [Gasc et al., 2015](#)). Conversely the envelope energy M , the number of major peaks of the mean frequency spectrum NP , and the Acoustic Complexity Index ACI were significantly positively correlated with richness and abundance. Although these three indices have been designed in the aim of circumventing the potential bias induced by the presence of noise and the absence of signals ([Depraetere et al., 2012](#); [Gasc et al., 2013](#)), they were here also significantly positively associated with the SNR. Altogether our results show that the SNR could be an important confounding and misleading variable for these five indices and that these indices should be used when recordings have been performed in habitats selected for their low noise level, avoiding flowing water such as streams or waterfalls. The AR index was the single index not correlated with neither richness nor abundance and the only index showing a positive and significant correlation with richness and abundance when taking SNR as a control variable in a partial correlation. The index AR is therefore a good candidate for revealing acoustic diversities within ponds. One of the major drawback of this index is that its ranking property makes independent studies hard to compare. Moreover our results show that it would be necessary to assess automatically the SNR to use AR . To compute the SNR without any manual identification of the signal and noise section within each recording is a technical challenge we could not solve preventing the use of the AR index on the complete set of recordings. Development in signal analysis is therefore still required to be able to monitor automatically pond acoustic diversity.

This preliminary study reveals that ponds we sampled were not silent habitats and that each pond revealed a different acoustic diversity. New biodiversity programs should be developed to describe and understand the sound diversity of ponds. Efforts should be achieved through a species-specific approach based on behavioural and systematic sciences to identify the sound sources and, at the same time, through a community approach based on ecological sciences to allow rapid biodiversity assessment. Combining these two research routes should lead to a better knowledge of this still unknown facet of animal diversity.

ACKNOWLEDGEMENTS

We thank the Parc Naturel Régional de la Haute Vallée de Chevreuse (France), the Office national de la Chasse et de la Faune Sauvage (St Benoist, Auffargis, France), Pierre-Yves Martel, and Arnaud Catherine. We are grateful to Roger Mundry (Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany) for providing R functions to study GLMM stability and to Bruno Naylor (Université Paris 13, Paris, France) for reviewing the English. We are grateful to Mitchell Aide and Klaus Riede for their helpful reviews.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This study was supported by a CNRS INEE PEPS grant, ENS grant (CD), and a CNRS INEE PhD grant (AG). This work benefited from state funding managed by the Agence Nationale de la Recherche as regards of the LabEx ANR-10-LABX-0003-BCDiv, in the framework of the program “Investissement d’avenir” referenced no ANR-11-IDEX-0004-02. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

CNRS INEE PEPS.

ENS.

CNRS INEE.

LabEx: ANR-10-LABX-0003-BCDiv.

Investissement d’avenir: ANR-11-IDEX-0004-02.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Camille Desjonquères analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Fanny Rybak contributed reagents/materials/analysis tools, wrote the paper, reviewed drafts of the paper.
- Marion Depraetere conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, prepared figures and/or tables, reviewed drafts of the paper.
- Amandine Gasc conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, reviewed drafts of the paper.
- Isabelle Le Viol conceived and designed the experiments, contributed reagents/materials/analysis tools, reviewed drafts of the paper.

- Sandrine Pavoine contributed reagents/materials/analysis tools, reviewed drafts of the paper.
- Jérôme Sueur conceived and designed the experiments, performed the experiments, contributed reagents/materials/analysis tools, wrote the paper, reviewed drafts of the paper.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.1393#supplemental-information>.

REFERENCES

- Aiken RB. 1982.** Shallow-water propagation of frequencies in aquatic insect sounds. *Canadian Journal of Zoology* **60**:3459–3461 DOI [10.1139/z82-436](https://doi.org/10.1139/z82-436).
- Aiken RB. 1985.** Sound production by aquatic insects. *Biological Reviews* **60**:163–211 DOI [10.1111/j.1469-185X.1985.tb00714.x](https://doi.org/10.1111/j.1469-185X.1985.tb00714.x).
- Baayen RH. 2008.** *Analyzing linguistic data: a practical introduction to statistics using R*. Cambridge: Cambridge University Press.
- Barr DJ, Levy R, Scheepers C, Tily HJ. 2013.** Random effects structure for confirmatory hypothesis testing: Keep it maximal. *Journal of Memory and Language* **68**:255–278 DOI [10.1016/j.jml.2012.11.001](https://doi.org/10.1016/j.jml.2012.11.001).
- Bates D, Maechler M, Bolker B, Walker S. 2014.** *lme4: linear mixed-effects models using Eigen and S4*. R package version 1.1-7.
- Buckingham MJ. 1992.** Ocean-acoustic propagation models. *Journal d'Acoustique*, 223–287.
- Céréghino R, Oertli B, Bazzanti M, Coccia C, Compin A, Biggs J, Bressi N, Grillas P, Hull A, Kalettka T, Scher O. 2012.** Biological traits of European pond macroinvertebrates. *Hydrobiologia* **689**:51–61 DOI [10.1007/s10750-011-0744-y](https://doi.org/10.1007/s10750-011-0744-y).
- Cox NJ. 2006.** Speaking stata: in praise of trigonometric predictors. *The Stata Journal* **6**(19):561–579.
- Dabelsteen T, Mathevon N. 2002.** Why do songbirds sing intensively at dawn? A test of the acoustic transmission hypothesis. *Acta Ethologica* **4**:65–72 DOI [10.1007/s10211-001-0056-8](https://doi.org/10.1007/s10211-001-0056-8).
- Depraetere M, Pavoine S, Jiguet F, Gasc A, Duvail S, Sueur J. 2012.** Monitoring animal diversity using acoustic indices: implementation in a temperate woodland. *Ecological Indicators* **13**:46–54 DOI [10.1016/j.ecolind.2011.05.006](https://doi.org/10.1016/j.ecolind.2011.05.006).
- Dobson AJ, Barnett A. 2008.** *An introduction to generalized linear models*. Boca Raton: CRC Press.
- Dray S, Dufour AB. 2007.** The ade4 package: implementing the duality diagram for ecologists. *Journal of Statistical Software* **22**:1–20 DOI [10.18637/jss.v022.i04](https://doi.org/10.18637/jss.v022.i04).
- Duguet R, Melki F. 2003.** *Les Amphibiens de France, Belgique et Luxembourg*. Collection Parthénope. Mèze: Editions Biotope.
- Farina A. 2014.** *Soundscape ecology: principles, patterns, methods and applications*. New York: Springer.
- Favaro L, Tirelli T, Gamba M, Pessani D. 2011.** Sound production in the red swamp crayfish *Procambarus clarkii* (Decapoda: Cambaridae). *Zoologischer Anzeiger—A Journal of Comparative Zoology* **250**:143–150 DOI [10.1016/j.jcz.2011.01.002](https://doi.org/10.1016/j.jcz.2011.01.002).
- Felisberto P, Jesus SM, Zabel F, Santos R, Silva J, Gobert S, Beer S, Björk M, Mazzuca S, Procaccini G, Runcie JW, Champenois W, Borges AV. 2015.** Acoustic monitoring of O2

- production of a seagrass meadow. *Journal of Experimental Marine Biology and Ecology* **464**:75–87 DOI [10.1016/j.jembe.2014.12.013](https://doi.org/10.1016/j.jembe.2014.12.013).
- Field AP. 2005. *Discovering statistics using SPSS: and sex, drugs and rock “n” roll, 2005*. Thousand Oaks: Sage Publications Limited.
- Fletcher NH. 2014. Animal bioacoustics. In: *Springer Handbook of Acoustics*. New York: Springer, 785–804.
- Forrest TG, Miller GL, Zagar JR. 1993. Sound propagation in shallow water: implications for acoustic communication by aquatic animals. *Bioacoustics* **4**:259–270 DOI [10.1080/09524622.1993.10510437](https://doi.org/10.1080/09524622.1993.10510437).
- Forstmeier W, Schielzeth H. 2011. Cryptic multiple hypotheses testing in linear models: overestimated effect sizes and the winner’s curse. *Behavioral Ecology and Sociobiology* **65**:47–55 DOI [10.1007/s00265-010-1038-5](https://doi.org/10.1007/s00265-010-1038-5).
- Fox J, Weisberg S. 2011. *An R companion to applied regression*. Second edition. Thousand Oaks: Sage.
- Gage SH, Axel AC. 2014. Visualization of temporal change in soundscape power of a Michigan lake habitat over a 4-year period. *Ecological Informatics* **21**:100–109 DOI [10.1016/j.ecoinf.2013.11.004](https://doi.org/10.1016/j.ecoinf.2013.11.004).
- Gasc A, Pavoine S, Lellouch L, Grandcolas P, Sueur J. 2015. Acoustic indices for biodiversity assessments: analyses of bias based on simulated bird assemblages and recommendations for field surveys. *Biological Conservation* **191**:306–312 DOI [10.1016/j.biocon.2015.06.018](https://doi.org/10.1016/j.biocon.2015.06.018).
- Gasc A, Sueur J, Pavoine S, Pellens R, Grandcolas P. 2013. Biodiversity sampling using a global acoustic approach: contrasting sites with microendemics in New Caledonia. *PLoS ONE* **8**:e65311 DOI [10.1371/journal.pone.0065311](https://doi.org/10.1371/journal.pone.0065311).
- Gascón S, Boix D, Sala J. 2009. Are different biodiversity metrics related to the same factors? A case study from Mediterranean wetlands. *Biological Conservation* **142**:2602–2612 DOI [10.1016/j.biocon.2009.06.008](https://doi.org/10.1016/j.biocon.2009.06.008).
- Gotelli NJ, Colwell RK. 2001. Quantifying biodiversity: procedures and pitfalls in the measurement and comparison of species richness. *Ecology Letters* **4**:379–391 DOI [10.1046/j.1461-0248.2001.00230.x](https://doi.org/10.1046/j.1461-0248.2001.00230.x).
- Hull A. 1997. The pond life project: a model for conservation and sustainability. In: Boothby J, ed. *British pond landscape, proceedings from the UK conference of the Pond Life Project*. Liverpool, 101–109.
- Kim S. 2012. *ppcor: Partial and Semi-partial (Part) correlation*. Package version 1.
- Leston D, Pringle JWS, White DCS. 1965. Muscular activity during preparation for flight in a beetle. *Journal of Experimental Biology* **42**:409–414.
- Météo France. Available at <http://www.meteofrance.com> (accessed 29 October 2015).
- Mc Cullagh P, Nelder JA. 1989. *Generalized linear models*. Boca Raton: CRC Press.
- Oertli B, Céréghino R, Hull A, Miracle R. 2009. Pond conservation: from science to practice. *Hydrobiologia* **634**:1–9 DOI [10.1007/s10750-009-9891-9](https://doi.org/10.1007/s10750-009-9891-9).
- Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O’Hara RB, Simpson GL, Solymos P, Stevens MHH, Wagner H. 2013. Package “vegan”. *Community ecology package*. version 2. Available at <http://cran.ism.ac.jp/web/packages/vegan/vegan.pdf>.
- Parks SE, Miksis-Olds JL, Denes SL. 2014. Assessing marine ecosystem acoustic diversity across ocean basins. *Ecological Informatics* **21**:81–88 DOI [10.1016/j.ecoinf.2013.11.003](https://doi.org/10.1016/j.ecoinf.2013.11.003).

- Pieretti N, Farina A, Morri D. 2011.** A new methodology to infer the singing activity of an avian community: the Acoustic Complexity Index (ACI). *Ecological Indicators* **11**:868–873 DOI [10.1016/j.ecolind.2010.11.005](https://doi.org/10.1016/j.ecolind.2010.11.005).
- Pijanowski BC, Farina A, Gage SH, Dumyahn SL, Krause BL. 2011.** What is soundscape ecology? An introduction and overview of an emerging new science. *Landscape Ecology* **26**:1213–1232 DOI [10.1007/s10980-011-9600-8](https://doi.org/10.1007/s10980-011-9600-8).
- R Core Team. 2015.** *R: a language and environment for statistical computing*. Vienna: R Foundation for Statistical Computing.
- Sandeman D, Wilkens LA. 1982.** Sound production by abdominal stridulation in the Australian Murray River crayfish, *Euastacus armatus*. *Journal of Experimental Biology* **99**:469–472.
- Schielzeth H, Forstmeier W. 2009.** Conclusions beyond support: overconfident estimates in mixed models. *Behavioral Ecology* **20**:416–420 DOI [10.1093/beheco/arn145](https://doi.org/10.1093/beheco/arn145).
- Silver SC, Halls JA. 1980.** Recording the sounds of hydropsychid larvae—a cautionary tale. *Journal of Comparative Physiology* **140**:159–161 DOI [10.1007/BF00606307](https://doi.org/10.1007/BF00606307).
- Simberloff D. 1972.** Properties of the rarefaction diversity measurement. *The American Naturalist* **106**:414–418 DOI [10.1086/282781](https://doi.org/10.1086/282781).
- Sueur J, Aubin T, Simonis C. 2008.** Seewave: a free modular tool for sound analysis and synthesis. *Bioacoustics* **18**:213–226 DOI [10.1080/09524622.2008.9753600](https://doi.org/10.1080/09524622.2008.9753600).
- Sueur J, Farina F. 2015.** Ecoacoustics: the ecological investigation and interpretation of environmental sound. *Biosemiotics* DOI [10.1007/s12304-015-9248-x](https://doi.org/10.1007/s12304-015-9248-x).
- Sueur J, Farina A, Gasc A, Pieretti N, Pavoine S. 2014.** Acoustic indices for biodiversity assessment and landscape investigation. *Acta Acustica united with Acustica* **100**:772–781 DOI [10.3813/AAA.918757](https://doi.org/10.3813/AAA.918757).
- Sueur J, Pavoine S, Hamerlynck O, Duvail S. 2008.** Rapid acoustic survey for biodiversity appraisal. *PLoS ONE* **3**:e4065 DOI [10.1371/journal.pone.0004065](https://doi.org/10.1371/journal.pone.0004065).
- Towsey M, Parsons S, Sueur J. 2014.** Ecology and acoustics at a large scale. *Ecological Informatics* **21**:1–3 DOI [10.1016/j.ecoinf.2014.02.002](https://doi.org/10.1016/j.ecoinf.2014.02.002).
- Towsey M, Wimmer J, Williamson I, Roe P. 2014.** The use of acoustic indices to determine avian species richness in audio-recordings of the environment. *Ecological Informatics* **21**:110–119 DOI [10.1016/j.ecoinf.2013.11.007](https://doi.org/10.1016/j.ecoinf.2013.11.007).
- Verberk WCEP, Van Duinen GA, Brock AMT, Leuven RSEW, Siepel H, Verdonschot PFM, Van der Velde G, Esselink H. 2006.** Importance of landscape heterogeneity for the conservation of aquatic macroinvertebrate diversity in bog landscapes. *Journal for Nature Conservation* **14**:78–90 DOI [10.1016/j.jnc.2005.11.002](https://doi.org/10.1016/j.jnc.2005.11.002).
- Wildlife Acoustics, Inc. 2015.** Bioacoustics software and field recording equipment. Available at <http://www.wildlifeacoustics.com/> (accessed 29 October 2015).
- Wood PJ, Greenwood MT, Agnew MD. 2003.** Pond biodiversity and habitat loss in the UK. *Area* **35**:206–216 DOI [10.1111/1475-4762.00249](https://doi.org/10.1111/1475-4762.00249).