



HAL
open science

Zika virus infection in French Polynesia

Jean-Marie Jouannic, Stephanie Friszer, Isabelle Leparc-Goffart, Catherine Garel, Dominique Eyrolle-Guignot

► **To cite this version:**

Jean-Marie Jouannic, Stephanie Friszer, Isabelle Leparc-Goffart, Catherine Garel, Dominique Eyrolle-Guignot. Zika virus infection in French Polynesia. *The Lancet*, 2016, 387 (10023), pp.1051-1052. 10.1016/S0140-6736(16)00625-5 . hal-01291986

HAL Id: hal-01291986

<https://hal.sorbonne-universite.fr/hal-01291986>

Submitted on 22 Mar 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

French Polynesian experience with maternofetal Zika virus infection

Jouannic JM¹, MD, PhD, Friszer S, MD¹, Leparc-Goffart I², MD, PhD, Garel C³, MD, Eyrolle-Guignot D, MD⁴

¹Service de Médecine Foetale, Hôpital Armand Trousseau, APHP, Paris 6, 26 Avenue du Docteur A. Netter, 75012 Paris, France

²Centre National de Référence des arbovirus, Institut de Recherche Biomédicale des Armées, Service de Biologie, Hôpital d'Instruction des Armées Laveran, 34 Boulevard Laveran, 13013 Marseille, France

³Service de Radiologie Pédiatrique, Hôpital Armand Trousseau, APHP, Paris 6, 26 Avenue du Docteur A. Netter, 75012 Paris, France

⁴Service de Gynécologie-Obstétrique, CHT Mamao, 98713 Papeete

Corresponding author : Pr JM Jouannic, Service de Médecine Foetale, Hôpital Armand Trousseau, APHP, Paris 6, 26 Avenue du Docteur A. Netter, 75012 Paris, France

Tel: +33-14473-5228

Fax: +33-14473-5222

Email: jean-marie.jouannic@aphp.fr

Conflict of interest: none

Contribution:

JMJ, SF, CG, DEG were in charge of the patients

JMJ, SF wrote the manuscript

ILG performed the PCR on amniotic fluids

Key words: zika virus, prenatal diagnosis, fetal, cerebral anomalies, ultrasound

Word count: 499

The World Health Organization issued an alert on February 2nd^{1,2} on the potential fetal consequences of the Zika virus outbreak after the Brazilian authorities reported an abnormal increase in the number of cases of neonates born with microcephaly³. Although no causal link could be clearly established, circumstantial evidence was considered worrisome enough for several countries to discourage pregnant women to travel to Central and South America⁴.

French Polynesia was affected by an epidemic of Zika between September 2013 and March 2014 **with an estimated 28,000 patients affected, representing around 11.5% of the Polynesian population**⁵. During 2014, local clinicians were struck by an unusual rate of cerebral congenital anomalies. **Within this territory, approximately 4,000 births are recorded each year.** All cases of suspected fetal abnormalities detected during routine prenatal ultrasound examinations are referred to Papeete Hospital. When the referral examination confirms the presence of fetal cerebral anomalies, neuroimaging examinations (ultrasound and/or MRI) are transmitted to the reference center of Trousseau Hospital in Paris for evaluation by experts in fetal neuroimagery. In 2014, 13 cases of fetal cerebral anomalies were diagnosed **instead of four and three in the two previous years. In addition, five neonates are currently under investigation for neurological signs of brainstem dysfunction. As part of the prenatal evaluation of the 13 cases of fetal cerebral anomalies**, amniocentesis was offered to all women for fetal karyotype and CMV detection by PCR, and performed in 10 cases. Following the Brazilian alert on possible fetal cerebral damage secondary to maternal Zika infection, we retrospectively tested 6 available stored amniotic fluid samples **using** PCR. Among those, four were found positive for Zika virus. The fetal brain anomalies depicted in

these cases are presented in the table. Microcephaly (Head Circumference<3th centile) was constant but **was observed after 28 weeks of gestation in two cases. Nevertheless, in case 1, ventriculomegaly was present at the second trimester routine scan. In case 3, the first scan was performed at 29⁺² weeks of gestation.** Severe abnormalities of midline structures and the cerebellum as well as abnormal gyration were observed, which may reflect an early embryologic hit of the developing brain. **Thus, a common maternal viral history during the 1st trimester of pregnancy was reported in three out of four cases.**

Only two observations of severe fetal brain damage were recently reported in the Brazilian population⁶. Our population-based experience should raise awareness regarding the fetal risks of Zika materno-fetal infection. It is likely that the true incidence of severe forms of fetal infection has been underestimated in our Polynesian population as we were unable to test all cases retrospectively.

Considering the potential severity of the fetal lesions, congenital Zika virus infection seems available to ultrasound imaging. Fetal neurosonography should be considered from the 2nd trimester in cases of maternal proven Zika virus infection in order to detect subtle cerebral anomalies that may precede the onset of microcephaly. Since we currently do not know the natural history of Zika virus fetal congenital infection, repeated ultrasound examinations should be considered.

References

1. Torjesen I. Zika virus outbreaks prompt warnings to pregnant women. *BMJ*. 2016;352:i500.
2. Zika virus: a new global threat for 2016. *Lancet Lond Engl*. 2016 Jan 9;387(10014):96.
3. Schuler-Faccini L, Ribeiro EM, Feitosa IML, Horovitz DDG, Cavalcanti DP, Pessoa A, et al. Possible Association Between Zika Virus Infection and Microcephaly - Brazil, 2015. *MMWR Morb Mortal Wkly Rep*. 2016;65(3):59–62.
4. Petersen EE, Staples JE, Meaney-Delman D, Fischer M, Ellington SR, Callaghan WM, et al. Interim Guidelines for Pregnant Women During a Zika Virus Outbreak - United States, 2016. *MMWR Morb Mortal Wkly Rep*. 2016;65(2):30–3.
5. **European Center for Disease Prevention and Control. Rapid risk assessment: Zika virus infection outbreak, French Polynesia: EDCCD; 2014 [updated 2014 Feb 14].**
6. Oliveira Melo AS, Malinger G, Ximenes R, Szejnfeld PO, Alves Sampaio S, Bispo de Filippis AM. Zika virus intrauterine infection causes fetal brain abnormality and microcephaly: tip of the iceberg? *Ultrasound Obstet Gynecol Off J Int Soc Ultrasound Obstet Gynecol*. 2016 Jan;47(1):6–7.