



**HAL**  
open science

## Fluctuating Minds: Spontaneous Psychophysical Variability during Mind-Wandering

Rodrigo A. Henríquez, Ana B. B. Chica, Pablo Billeke, Paolo Bartolomeo

► **To cite this version:**

Rodrigo A. Henríquez, Ana B. B. Chica, Pablo Billeke, Paolo Bartolomeo. Fluctuating Minds: Spontaneous Psychophysical Variability during Mind-Wandering. PLoS ONE, 2016, 11 (2), pp.e0147174. 10.1371/journal.pone.0147174 . hal-01294435

**HAL Id: hal-01294435**

**<https://hal.sorbonne-universite.fr/hal-01294435>**

Submitted on 29 Mar 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

RESEARCH ARTICLE

# Fluctuating Minds: Spontaneous Psychophysical Variability during Mind-Wandering

Rodrigo A. Henríquez<sup>1,2,3,4\*</sup>, Ana B. Chica<sup>5</sup>, Pablo Billeke<sup>6</sup>, Paolo Bartolomeo<sup>1,2,3,4,7</sup>

**1** Inserm U 1127, F-75013, Paris, France, **2** Sorbonne Universités, UPMC Univ Paris 06, UMR S 1127, F-75013, Paris, France, **3** CNRS, UMR 7225, F-75013, Paris, France, **4** Institut du Cerveau et de la Moelle épinière, ICM, F-75013, Paris, France, **5** Department of Experimental Psychology, and Brain, Mind, and Behaviour Research Center, University of Granada, Granada, Spain, **6** Research Center for Social Complexity, Faculty of Government, University of Desarrollo, Santiago, Chile, **7** Department of Psychology, Catholic University Milan, Milan, Italy

\* [henriquezch@gmail.com](mailto:henriquezch@gmail.com)



**OPEN ACCESS**

**Citation:** Henríquez RA, Chica AB, Billeke P, Bartolomeo P (2016) Fluctuating Minds: Spontaneous Psychophysical Variability during Mind-Wandering. *PLoS ONE* 11(2): e0147174. doi:10.1371/journal.pone.0147174

**Editor:** Martin Walter, Leibniz Institute for Neurobiology, GERMANY

**Received:** June 9, 2015

**Accepted:** December 30, 2015

**Published:** February 10, 2016

**Copyright:** © 2016 Henríquez et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** This work was supported by a doctoral fellowship from the National Commission for Scientific and Technological Research - CONICYT 72120217, Chile to R.A.H. and A.B.C was supported by a Ramón y Cajal fellowship from the Spanish Ministry of Education and Science, RYC-2011-09320. This research was funded by research projects PSI2011-22416 and PSI2014-58681-P to A.B.C. This research also has received funding from the program "Investissements d'Avenir" ANR-10-IAIHU-06.

## Abstract

Mind-wandering is the occasional distraction we experience while performing a cognitive task. It arises without any external precedent, varies over time, and interferes with the processing of sensory information. Here, we asked whether the transition from the on-task state to mind-wandering is a gradual process or an abrupt event. We developed a new experimental approach, based on the continuous, online assessment of individual psychophysical performance. Probe questions were asked whenever response times (RTs) exceeded 2 standard deviations from the participant's average RT. Results showed that mind-wandering reports were generally preceded by slower RTs, as compared to trials preceding on-task reports. Mind-wandering episodes could be reliably predicted from the response time difference between the last and the second-to-last trials. Thus, mind-wandering reports follow an abrupt increase in behavioral variability, lasting between 2.5 and 10 seconds.

## Introduction

Mind-wandering, the occasional distraction we often experience while performing a cognitive task, is a self-generated condition, because it arises without any external precedent, varies over time spontaneously, and often interferes with the online processing of sensory information [1,2,3]. The very existence of mind-wandering supports the view that perception depends not only on its inputs, but also on the internal variability of the system.

Studies have addressed mind-wandering using different experimental definitions, e.g. task-unrelated thoughts [4,5], stimulus-independent thoughts [6], incidental self-processing [7], inner speech [8] momentary attentional lapses [9], or spontaneous thoughts [10,11]. Patterns of performance related to mind-wandering include variations in response times (RTs) during a sustained attention task [12], and increased response variability on a metronome response task

**Competing Interests:** The authors have declared that no competing interests exist.

[13,14]. Studies on the neural bases of mind wandering have described reduced amplitude of event-related potentials (ERPs) such as P300, MMN, and P2 components [1,15,16], as well as the activation of the brain default network [10,11]. A causal role of the dorsolateral prefrontal cortex has also been proposed [17].

Mind wandering is often explored by asking participants thought sampling questions (TSQs) on their state of mind, while they perform a sustained attention task [18]. However, this method is not optimal to assess whether the transition to mind-wandering is a process that develops gradually, or it is a unique event that triggers a global cognitive change. Moreover, the temporal rate of presentation of the TSQs can affect the likelihood of mind-wandering reports, because people are more likely to report mind-wandering as the time between TSQs increases [14].

In the present study, we aimed at defining some of the psychophysical conditions necessary for mind-wandering to occur. We used a new experimental approach, inspired by the classic sustained attention to response task (SART) [19]. We identified online RT outliers exceeding 2 SD from the participant's mean RT. After each RT outlier, a TSQ was automatically asked. In comparison to traditional methods, this procedure allowed us to fit the behavioral variability associated with mind-wandering to the online statement of its occurrence in a more dynamic and ecological way.

## Materials and Methods

### Participants


Thirty-three healthy undergraduates (3 males) from the University of Granada participated in this study for course credit. A further participant was not considered in the analysis because he did not produce consistent responses, by often changing the response key assignment for targets and nontarget. All participants were right-handed, with their age ranging from 18 to 30 years ( $M = 23.71$ ). Participants had normal or corrected to normal vision. They had no history of neurological disease and were free from psychoactive medication use. All participants provided their written informed consent prior to participating in the study. The local ethics committee from the University of Granada approved the experimental protocol.

### Stimuli

E-Prime<sup>®</sup> software (version 2.1, Psychology Software Tools Inc., <http://www.pstnet.com>) was used to control stimuli presentation and response collection. Stimuli were displayed on a 21" monitor with a refresh rate of 60 Hz, situated at approximately 57 cm from the participants' eyes. Ten randomly chosen upper-case letters were presented in rapid visual serial presentation. All the letters appeared inside an empty central white rectangle on a grey background. Each letter was presented for 100 ms, with an inter-stimulus interval varying randomly between 2,500 and 4,000 ms. There were three blocks of trials, each composed of 329 letters, and lasting for about 30 minutes. Participants were allowed to rest as much as they wanted between blocks.

### Procedure

A keyboard with numeric keypad on the right side was used to provide responses. Participants were instructed to press as fast as possible a colored key (either a red "7" or a green "8", in different participants) on the numeric keypad each time they saw the target (letter F) appearing. This condition occurred in 10% of the trials (Fig 1). In the remaining trials, when letters different from F appeared, they had to press the other colored key. The key assignment to target and



**Fig 1. Schematic depiction of the experimental paradigm.**

doi:10.1371/journal.pone.0147174.g001

nontarget letters was counterbalanced across participants. All participants used their right hand to respond. Mean RTs for nontargets were calculated online based on the last 5 responses. Whenever a RT exceeded 2 SDs from each participant’s RT mean within the last 5 trials (see below), the response was defined as an outlier RT, and a thought sampling question (TSQ) automatically appeared on the test display. Thus, average RTs before the TSQ were calculated on a number of consecutive trials, which could range from a minimum of 5 trials to a maximum of 10 trials (if no RT outlier was detected before trial 10). If no outlier RTs were produced during trials 6 to 9, after trial 10 a control TSQ appeared (“Standard TSQ”).

Each participant received an average of 48 TSQs per block. TSQs were formulated as follows: “Just before this question, your attention was distracted by.” Five possible responses were then proposed: (1) My attention was not distracted. (2) External distraction (e.g., uncomfortable posture, itches, sneezing, coughing). (3) Internal distraction (memories, imagination, thoughts). (4) Loosing vigilance (feeling of falling asleep). (5) Others. Participants had to respond by pressing a button corresponding to the number of the chosen alternative over the numeric keypad (respectively, keys 1, 2, 3, 4, or 5) (Fig 1). Response 1 (no perceived distraction) was considered as evidence that participants were on-task, while response 3 (internal distraction) defined mind-wandering (off-task condition).

## Results

The level of statistical significance was set at  $\alpha = .05$ . Bonferroni correction was used to correct for multiple comparisons. Performance was generally accurate, with 86% correct performance for nontarget stimuli (which constituted 90% of the total stimuli), and 81% correct responses to target stimuli (10% of the total stimuli). Participants chose response 1 (on-task) 55.5% of the time, and response 3 (mind-wandering) 18.6% of the time (Table 1). A t-test performed on the arcsin-transformed response rates demonstrated that on-task responses were more frequent than off-task responses,  $t(32) = 6.87, p < 0.001, d = 1.19, 95\% \text{ CIs } [.18, .34]$ . The

**Table 1. Percentages of response types.**

On-task	55.5%
External distraction	14.5%
Internal distraction (Off-task)	18.6%
Falling asleep	8.8%
Other	2.6%

doi:10.1371/journal.pone.0147174.t001

vast majority (67.3%) of these on-task responses followed standard RTs (within 2 SD around the mean, see Procedure); 23.4% of all on-task responses were given after slow RTs (>2SD above the mean), and 9.4% were given for faster responses (>2SD below the mean).

Responses to TSQs fell into two categories: responses related to outlier RTs (slow or fast), and responses related to standard (non-outlier) RTs. Based on previous evidence that RT variability is maximal in the trials just before mind-wandering episodes [13,20,21], we analyzed the 5 trials preceding on-task or off-task responses to TSQs (remember that 5 trials was the minimal possible interval between two TSQs).


Fig 2 shows the distribution of “slow” RTs (> 2 SD above the mean) across blocks. Each line represents a subject; red Xs represent median percentages per block. Slow RTs were not uniformly distributed among blocks (Block 1, 10%; Block 2, 14.5%, Block 3, 12%; Friedman chi-squared = 12.28,  $df = 2$ ,  $p = 0.002$ ), because they increased in Block 2 compared with Block 1 (Friedman test,  $p = 0.007$ ; Bonferroni correction for 3 comparisons yields an alpha value of 0.017). There was no significant difference between Block 2 and Block 3 ( $p = 0.7$ ) or between Block 1 and Block 3 ( $p = 0.045$ ). However, the percentage of off-task responses did not vary across blocks (Block 1, 13%; Block 2, 16%, Block 3, 15%; Friedman chi-squared = 4.06,  $df = 2$ ,  $p = 0.13$ ). Thus, while there may be some general tendency to an RT increase across blocks, perhaps resulting from fatigue, the rate of mind-wandering reports did not significantly vary along the experimental session.

Fig 3 shows the evolution of RT during the last 5 trials before the TSQ. A 2x5 repeated measures ANOVA with condition (on-task and off-task) and trial position before the TSQ (N-1, N-2, N-3, N-4 and N-5) as within-participant factors revealed a main effect of trial position,  $F(4,128) = 44.00$ ,  $p < .01$ ,  $\eta_p^2 = .57$ , which interacted with condition,  $F(4,128) = 16.49$ ,  $p < .01$ ,  $\eta_p^2 = .34$ .

On-task and off-task conditions in trial N-5 led to similar RTs (Fig 3),  $t(32) = 1.86$ ,  $p > .05$ ,  $d = .32$ , 95% CIs [-1.39, 31.36]. In the following three trials (N-4 to N-2), RTs were on average 23-ms faster in the off-task condition compared with the on-task condition (all  $t_s(32) > 2.84$ , all  $p_s < .01$ ,  $d = .49$ , 95% CIs [14.02, 44.05; 6.04, 36.52; 6.89, 32.98], for all comparisons). However, this pattern reversed in trial N-1 (just before the TSQ), which elicited 66-ms slower RTs on the off-task condition than on the on-task condition,  $t(32) = -3.53$ ,  $p = 0.001$ ,  $d = -.61$ , 95% CIs [-104.94, -28.18] (all these comparisons were Bonferroni corrected for 5 comparisons, with alpha = 0.01).

Although our aim was to explore internal distraction (mind wandering), we also assessed the RT profile of external distraction responses. We performed a 3x5 repeated measures ANOVA with response type (on-task, off-task task, and external distraction) and trial position before the TSQ (N-1, N-2, N-3, N-4 and N-5) as factors. The within-participant factors revealed a main effect of trial position,  $F(4, 128) = 96.22$ ,  $p < .01$ ,  $\eta_p^2 = .75$ , which interacted with the response,  $F(4, 128) = 11.02$ ,  $p < .01$ ,  $\eta_p^2 = .25$ . As shown in Table 2 and in S1 Fig, this interaction stemmed from the fact that on-task and external distraction RTs were similar for trials N-2 to N-5 (all  $t_s(32) < .56$ , all  $p_s > .57$ ,  $d = -.09$ , 95% CIs [-18.31, 29.18; -15.06, 26.44; 6.89, 32.98; -23.88, 16.17]; Bonferroni correction for 5 comparisons, alpha = 0.01); however, similar to off-task responses, external distraction responses differed sharply from on-task responses for trial N-1, which elicited 108-ms slower RTs for external distraction responses than for on-task responses,  $t(32) = -6.09$ ,  $p < .001$ ,  $d = -1.05$ , 95% CIs [-145.47, -72.27]. Thus, off-task and external distraction responses gave rise to a similar RT profile.


We also assessed whether the mean RT for each of the five trials preceding a TSQ could predict the participant’s response (“on-task” or “off-task”), by performing a logistic regression analysis on the mean RT of each preceding trial (from N-1 to N-5). Results showed that increasing RTs significantly predicted reports of mind-wandering. The test of model



**Fig 2. Distribution of slow RTs (> 2SD) across blocks.** Each line represents a subject. Red Xs represent median percentages per block.

doi:10.1371/journal.pone.0147174.g002

coefficients was statistically significant, indicating that the predictors as a set reliably distinguished between trial position N-1 and N-2 (chi square = 34.09,  $p < .01$  with  $df = 2$ ). The Wald criterion demonstrated that trial N-2 and N-1 made a significant contribution to the prediction ( $p < .01$ ). This is confirmed by Nagelkerke R square of .54 on second step that indicated a



**Fig 3. Distribution of mean RTs for the five trials before TSQ.** Continuous line, off-task episodes related to mind-wandering; dashed line, on-task episodes. Error bars represent 1 standard error of the mean.

doi:10.1371/journal.pone.0147174.g003

strong relationship between on- and off-task prediction and the position trials N-2 and N-1. Overall prediction success was 75.8% for the second step compared with less than 57.6% for the first step, which included only trial N-1 (Table 3).

RTs for the other tested trial positions (from N-3 to N-5) failed to predict the participant’s response (all  $ps > .1$ ). Finally, the RT difference between N-1 and N-2 trial significantly predicted participants’ responses to the TSQ (paired Wilcoxon signed rank test,  $Z = -4.51$ ,  $p < .01$ ,  $r = -0.57$ ). Thus, not only was mind-wandering indexed by slow RTs to the last trial before the

**Table 2. Mean RT and standard deviation for trials N-1 to N-5 before the TSQ for each possible response (on-task, off-task, external distraction, falling asleep, and other).**

	N-1	N-2	N-3	N-4	N-5
<b>On-task</b>					
Mean	415.61	398.33	389.16	382.57	381.87
SD	77.55	62.34	70.33	63.25	69.44
<b>External distraction</b>					
Mean	524.48	392.89	383.47	387.65	385.72
SD	114.92	90.12	86.97	77.17	69.39
<b>Off-task</b>					
Mean	482.16	369.29	367.88	362.63	366.89
SD	128.51	80.21	70.89	65.74	76.55
<b>Falling asleep</b>					
Mean	323.79	362.67	371.73	368.24	360.16
SD	76.78	111.58	107.95	92.98	93.83
<b>Other</b>					
Mean	363.56	494.26	391.20	417.65	405.72
SD	154.13	310.48	112.57	223.81	135.61

doi:10.1371/journal.pone.0147174.t002

**Table 3. Logistic regression analysis of off-task and on-task condition for the last five trials before the TSQ.**

	Variables	$\beta$	S.E	Wald	df	Sig.	Exp( $\beta$ )
Step 1.	N-1	.006	.003	5.44	1	.020	1.0
	Constant	-2.79	1.21	5.26	1	.022	.06
Step 2.	N-1	.035	.010	12.30	1	.000	1.03
	N-2	-.045	.013	12.73	1	.000	.95
	Constant	2.06	1.76	1.37	1	.24	7.88

doi:10.1371/journal.pone.0147174.t003

TSQ, but also, and more accurately, by the RT difference between the last and the last-but-one trial.


In order to assess for long-range variations in RTs, we carried out a Fast Fourier Transform (FFT) to explore the variation of RTs through the task for each participant [22]. Then, we compared the resulting power spectrum with random distributions of the same data (2,000 permutations). RTs followed a 1/f distribution. Long-range variations (oscillations slower than one cycle per 6-minute period, Fig 4) were significantly greater than the random distribution. These oscillations represent variations longer than the maximum time interval between two TQS (41s). Finally, we assessed the possible relationship between these slow RT oscillations and the rate of off-task responses. We carried out correlation analyses between the rate of off-task responses and the power of these oscillations in 0.2-min bandwidth windows around the three fastest peaks in the FFT (11.1 min, 7.8 min, 6.0 min, see Fig 4). Results showed that subjects with greater power in the 5-min oscillation tended to produce less off-task responses (Spearman Correlation, 5.9–6.1. min,  $\rho = -0.38$ ,  $p = 0.02$ ), while no correlation emerged for the other oscillations (7.9–7.7 min,  $\rho = 0.08$ ,  $p = 0.6$ ; 11.0–11.2 min,  $\rho = 0.27$ ,  $p = 0.2$ ).

## Discussion

A novel experimental procedure allowed us to identify episodes of mind-wandering based on the on-line assessment of RT fluctuations during a sustained attention task. Our procedure automatically detected outlier RTs and consequently triggered a TSQ during task performance. We explored different sources of mind-wandering based on individual responses to the five trials preceding the TSQ in both on- and off-task conditions. We observed that off-task reports were generally preceded by slower RTs as compared with trials before on-task responses. Note, however, that a substantial proportion of slow RTs (23%) occurred before “on-task” reports. Moreover, RTs tended to increase across blocks, without significant variation of the percentage of off-task responses. Thus, RTs alone are not to be considered as a reliable index of mind wandering. We also observed a clear difference between RTs in N-1 and N-2 trials compared to the remaining trials. This suggests that mind-wandering is not an attentional *state* in itself, characterized by a global slowing of responses. Rather, mind-wandering seems best characterized in a more dynamical way as a *transition* between different attentional states. Assessment of 1/f patterns showed the presence of slow oscillations (over several minutes) in RT performance. However, these oscillations either had no correlation or were negatively correlated with the rate of off-task responses. Thus, mind-wandering states seemed to interrupt these slow oscillations, rather than being part of them. Thus, our results suggest that mind-wandering is a local phenomenon lasting between 2.5 and 10 seconds, presumably driven by specific cognitive processes such as spontaneous fluctuations in the alertness system [23].

Previous studies have adjusted the presentation of evaluations of mind-wandering by using a pre-established rate of TSQs [18]. Our online method allowed us to automatically detect and assess different mind-wandering states continuously, and presumably in a more ecological





**Fig 4. Log-log power spectra of time series of RT variation.** Red continuous line: mean of the power spectrum across participants. Red dotted lines: standard error of mean. The black line represents the statistical threshold (Permutation test and false discovery rate (FDR),  $q = 0.05$ ). The figure also shows the three only significant peaks at 11.1, 7.8, and 6.0 min.

doi:10.1371/journal.pone.0147174.g004

way. Our psychophysical evidence demonstrates how dynamical local changes in the attentional system level can trigger global cognitive changes [24], and supports the hypothesis that changes of the attentional state not only depend on the nature of the stimuli, such as their physical properties, but also on the variability of the system itself [25–27]. Our approach could therefore inspire strategies to prevent inattention when mind-wandering can negatively impact cognitive performance [28].

Despite its clear and straightforward findings, this study has limitations. First, the observed RT profile was not able to clearly distinguish between internal and external distraction. Second, as compared to other methods our samples are presumably more specific, but also reduced in number. Third, it is difficult to estimate the potentially disruptive effects of the TSQs on the sustained attention task. Finally, the dichotomy between on- and off-task conditions might not be subtle enough to capture the richness of phenomenological experience. Thus, it is possible that this first attempt to make an online estimation of mind-wandering did not fully capture the richness of mind-wandering content, because our approach did not make any a priori assumptions about the cognitive definition of this phenomenon. Further experiments

including a more precise evaluation of the cognitive content of mind-wandering should address these issues. To conclude, our new approach allowed us to identify mind-wandering as a dynamic transition between attentional states, which results in abruptly decoupling the attentional system from the external world.

## Supporting Information

### S1 Dataset.

(ZIP)

**S1 Fig. Distribution of mean RTs for the five trials before TSQ.** Continuous black line, off-task episodes related to mind-wandering; continuous grey line, external distraction episodes and black dashed line, on-task episodes. Error bars represent 1 standard error of the mean.

(TIFF)

## Acknowledgments

We are grateful to the participants for their patience and good will and to Dr. Julià Amengual for useful discussion.

## Author Contributions

Conceived and designed the experiments: RAH ABC. Performed the experiments: RAH. Analyzed the data: RAH. Contributed reagents/materials/analysis tools: RAH P. Billeke. Wrote the paper: RAH. Provided critical revisions: ABC P. Bartolomeo.

## References

1. Braboszcz C, Delorme A. Lost in thoughts: Neural markers of low alertness during mind wandering. *Neuroimage*. 2011; 54:3040–7. doi: [10.1016/j.neuroimage.2010.10.008](https://doi.org/10.1016/j.neuroimage.2010.10.008) PMID: [20946963](https://pubmed.ncbi.nlm.nih.gov/20946963/)
2. Schooler JW, Smallwood J, Christoff K, Handy TC, Reichle ED, Sayette M a.. Meta-awareness, perceptual decoupling and the wandering mind. *Trends Cogn Sci*; 2011; 15(7):319–26. doi: [10.1016/j.tics.2011.05.006](https://doi.org/10.1016/j.tics.2011.05.006) PMID: [21684189](https://pubmed.ncbi.nlm.nih.gov/21684189/)
3. Smallwood J, Schooler JW. The restless mind. *Psychol Bull*. 2006; 132(6):946–58. PMID: [17073528](https://pubmed.ncbi.nlm.nih.gov/17073528/)
4. Giambra LM. Task-Unrelated-Thought Frequency as a Function of Age: A Laboratory Study. 1989; 4(2):136–43.
5. Smallwood JM, Baracaia SF, Lowe M, Obonsawin M. Task unrelated thought whilst encoding information. *Conscious Cogn*; 2003 Sep 1; 12(3):452–84. PMID: [12941287](https://pubmed.ncbi.nlm.nih.gov/12941287/)
6. Teasdale JD, Dritschel BH, Taylor MJ, Proctor L, Lloyd C a, Nimmo-Smith I, et al. Stimulus-independent thought depends on central executive resources. *Mem Cognit*. 1995; 23(5):551–9. PMID: [7476241](https://pubmed.ncbi.nlm.nih.gov/7476241/)
7. Gilbert SJ, Frith CD, Burgess PW. Involvement of rostral prefrontal cortex in selection between stimulus-oriented and stimulus-independent thought. *Eur J Neurosci*. 2005; 21:1423–31. PMID: [15813952](https://pubmed.ncbi.nlm.nih.gov/15813952/)
8. Morin A. Inner Speech and Consciousness. In: Banks William P, (Editor), *Encyclopedia of Consciousness* volume 1, pp 389–402 Oxford: Elsevier; 2009. p. 1159–66.
9. Weissman DH, Roberts KC, Visscher KM, Woldorff MG. The neural bases of momentary lapses in attention. *Nat Neurosci*. 2006; 9(7):971–8. PMID: [16767087](https://pubmed.ncbi.nlm.nih.gov/16767087/)
10. Christoff K, Gordon AM, Smallwood J, Smith R, Schooler JW. Experience sampling during fMRI reveals default network and executive system contributions to mind wandering. *Proc Natl Acad Sci U S A*. 2009; 106:8719–24. doi: [10.1073/pnas.0900234106](https://doi.org/10.1073/pnas.0900234106) PMID: [19433790](https://pubmed.ncbi.nlm.nih.gov/19433790/)
11. Mason MF, Norton MI, Van Horn JD, Wegner DM, Grafton ST, Macrae CN. Wandering minds: the default network and stimulus-independent thought. *Science* (80-); 2007 Jan 19; 315(5810):393–5.
12. Smallwood J, Davies JB, Heim D, Finnigan F, Sudberry M, O'Connor R, et al. Subjective experience and the attentional lapse: task engagement and disengagement during sustained attention. *Conscious Cogn*; 2004 Dec 1; 13(4):657–90. PMID: [15522626](https://pubmed.ncbi.nlm.nih.gov/15522626/)

13. Seli P, Cheyne JA, Smilek D. Wandering minds and wavering rhythms: linking mind wandering and behavioral variability. *J Exp Psychol Hum Percept Perform*; 2013; 39(1):1–5. doi: [10.1037/a0030954](https://doi.org/10.1037/a0030954) PMID: [23244046](https://pubmed.ncbi.nlm.nih.gov/23244046/)
14. Seli P, Carriere JS a, Levene M, Smilek D. How few and far between? Examining the effects of probe rate on self-reported mind wandering. *Front Psychol*. 2013; 4(July):1–5.
15. Laufs H, Krakow K, Sterzer P, Eger E, Beyerle A, Salek-Haddadi A, et al. Electroencephalographic signatures of attentional and cognitive default modes in spontaneous brain activity fluctuations at rest. *Proc Natl Acad Sci U S A*; 2003 Sep 16; 100(19):11053–8. PMID: [12958209](https://pubmed.ncbi.nlm.nih.gov/12958209/)
16. Smallwood J, Beach E, Schooler JW, Handy TC. Going AWOL in the brain: mind wandering reduces cortical analysis of external events. *J Cogn Neurosci*. 2008; 20:458–69. PMID: [18004943](https://pubmed.ncbi.nlm.nih.gov/18004943/)
17. Axelrod V, Rees G, Lavidor M, Bar M. Increasing propensity to mind-wander with transcranial direct current stimulation. *Proc Natl Acad Sci*; 2015; 201421435.
18. Smallwood J, McSpadden M, Luus B, Schooler J. Segmenting the stream of consciousness: the psychological correlates of temporal structures in the time series data of a continuous performance task. *Brain Cogn*; 2008 Feb 1; 66(1):50–6. PMID: [17614178](https://pubmed.ncbi.nlm.nih.gov/17614178/)
19. Robertson IH, Manly T, Andrade J, Baddeley BT, Yiend J. “Oops!”: performance correlates of everyday attentional failures in traumatic brain injured and normal subjects. *Neuropsychologia*; 1997 Jun 1; 35(6):747–58. PMID: [9204482](https://pubmed.ncbi.nlm.nih.gov/9204482/)
20. Allan Cheyne J, Solman GJF, Carriere JS a, Smilek D. Anatomy of an error: A bidirectional state model of task engagement/disengagement and attention-related errors. *Cognition*; 2009; 111(1):98–113. doi: [10.1016/j.cognition.2008.12.009](https://doi.org/10.1016/j.cognition.2008.12.009) PMID: [19215913](https://pubmed.ncbi.nlm.nih.gov/19215913/)
21. Bastian M, Sackur J. Mind wandering at the fingertips: Automatic parsing of subjective states based on response time variability. *Front Psychol*. 2013; 4(September):1–11.
22. Castellanos FX, Sonuga-Barke EJS, Scheres A, Di Martino A, Hyde C, Walters JR. Varieties of Attention-Deficit/Hyperactivity Disorder-related intra-individual variability. *Biological Psychiatry*. 2005; 57(11):1416–1423. PMID: [15950016](https://pubmed.ncbi.nlm.nih.gov/15950016/)
23. Sadaghiani S, Scheeringa R, Lehongre K, Morillon B, Giraud A-L, Kleinschmidt A. Intrinsic connectivity networks, alpha oscillations, and tonic alertness: a simultaneous electroencephalography/functional magnetic resonance imaging study. *J Neurosci*. 2010; 30(30):10243–50. doi: [10.1523/JNEUROSCI.1004-10.2010](https://doi.org/10.1523/JNEUROSCI.1004-10.2010) PMID: [20668207](https://pubmed.ncbi.nlm.nih.gov/20668207/)
24. Le Van Quyen M. The brainweb of cross-scale interactions. *New Ideas Psychol*; 2011; 29(2):57–63.
25. Feldman H, Friston K. Attention, uncertainty, and free-energy. *Front Hum Neurosci*; 2010 Jan 1; 4:215. doi: [10.3389/fnhum.2010.00215](https://doi.org/10.3389/fnhum.2010.00215) PMID: [21160551](https://pubmed.ncbi.nlm.nih.gov/21160551/)
26. Friston K, Daunizeau J, Kilner J, Kiebel SJ. Action and behavior: a free-energy formulation. *Biol Cybern*; 2010 Mar 1; 102(3):227–60. doi: [10.1007/s00422-010-0364-z](https://doi.org/10.1007/s00422-010-0364-z) PMID: [20148260](https://pubmed.ncbi.nlm.nih.gov/20148260/)
27. Rastelli F, Tallon-Baudry C, Migliaccio R, Toba MN, Ducorps A, Pradat-Diehl P, et al. Neural dynamics of neglected targets in patients with right hemisphere damage. *Cortex*. 2013; 49(7):1989–96. doi: [10.1016/j.cortex.2013.04.001](https://doi.org/10.1016/j.cortex.2013.04.001) PMID: [23664670](https://pubmed.ncbi.nlm.nih.gov/23664670/)
28. Killingsworth M a, Gilbert DT. A wandering mind is an unhappy mind. *Science*. 2010; 330:932. doi: [10.1126/science.1192439](https://doi.org/10.1126/science.1192439) PMID: [21071660](https://pubmed.ncbi.nlm.nih.gov/21071660/)