Lipid dynamics in early life stages of the icefish *Chionodraco hamatus* in the Dumont d'Urville Sea (East Antarctica)

Carolina Giraldo, Marc Boutoute, Patrick Mayzaud, an Vo Quang, Philippe Koubbi, Eric Tavernier

► To cite this version:

Carolina Giraldo, Marc Boutoute, Patrick Mayzaud, an Vo Quang, Philippe Koubbi, et al.. Lipid dynamics in early life stages of the icefish *Chionodraco hamatus* in the Dumont d'Urville Sea (East Antarctica). *Polar Biology*, 2017, 40 (2), pp.313-320. 10.1007/s00300-016-1956-4 . hal-01317616

HAL Id: hal-01317616

<https://hal.sorbonne-universite.fr/hal-01317616>

Submitted on 18 May 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Lipid dynamics in early life stages of the icefish *Chionodraco hamatus* in the**
2 **Dumont d'Urville Sea (East-Antarctica)**

3
4 Carolina Giraldo ^{1,2*}, Marc Boutoute², Patrick Mayzaud^{1,2} Eric Tavernier^{3,4}, An Vo Quang^{1,2}, and
5 Philippe Koubbi ⁵

6 ¹UPMC Université Paris 06, UMR 7093, Laboratoire d'Océanographie de Villefranche, BP28,
7 06234 Villefranche-sur-Mer, France.

8 ²CNRS, UMR 7093, LOV, BP 28, 06234 Villefranche-sur-Mer, France.

9 ³Université Lille Nord de France, F-59000 Lille, France.

10 ⁴ULCO, LOG, 32 avenue Foch, F-62930 Wimereux, France.

11 ⁵ Unité Biologie des organismes et écosystèmes aquatiques (BOREA, UMR 7208), Sorbonne
12 Universités, Muséum national d'Histoire naturelle, Université Pierre et Marie Curie, Université
13 de Caen Basse-Normandie, CNRS, IRD; CP26, 57 rue Cuvier 75005 Paris, France.

14 *Corresponding author: carolina.giraldo@univ-lille1.fr

15 **Abstract**

16
17 Lipids play a crucial role in polar regions and are of particular importance in early life stages of
18 Antarctic fish. This work presents the significance of lipids and fatty acids (FA) in the early life
19 stages of the icefish *Chionodraco hamatus*. Analysis of lipid classes (Polar lipids, LP;
20 Cholesterol, Chol; and Triacylglycerol, TAG) suggested different energy allocation strategies in
21 preflexion (<25 mm) and postflexion larvae (>25 mm). Structural PL dominated lipid dynamics
22 for preflexion larvae, indicating that small individuals allocate the majority of energy toward
23 somatic growth. Conversely, postflexion larvae appear able to switch between growth
24 (contribution of PL) and energy storage (contribution of TAG) strategies. The condition index

ratio TAG:Chol varied from 0.2 to <2 with no differences between the two larval stages. Further, FA composition of the TAG and PL fractions suggests that both developmental stages share the same carnivorous diet and that *C. hamatus* relies on a few key prey items.

Key-words: Larvae, fatty acids, nutritional condition, diet, Antarctica

Introduction

In the pelagic ecosystem of the Southern Ocean, icefish (Channichthyidae, ‘white-blooded’ fish) form a family within the perciform suborder Notothenioidei, the most predominant suborder of demersal Antarctic fishes (Kock 2005). Early life stages of most species of channichthyids have been described (Efremenko 1987; Kellermann and Schadwinkel 1991) and their distribution and relative abundance reviewed elsewhere (La Mesa and Ashford 2008; Loeb et al. 1993). At hatching, larvae of channichthyids are relatively large (13–17 mm), suggesting greater likelihood of survival compared to smaller larvae of other Antarctic fishes (Kock 2005). In addition, larval channichthyids exhibit relatively fast growth rates and feed primarily on young stages of euphausiids and larval nototheniids (Kellermann 1990; La Mesa et al. 2011). Although most channichthyids are demersal-benthic as adults, larvae and postlarvae are typically pelagic. The most common channichthyid within the pelagic communities of the Ross and Dumont D’Urville Sea are larval and juvenile *Chionodraco* sp. (Granata et al. 2002; North 1988). The genus *Chionodraco* includes *C. hamatus*, *C. myersi*, and *C. rastrospinosus*, with *C. hamatus* and *C. myersi* previously identified from the Dumont d’Urville Sea (Koubbi et al. 2010). As the taxonomic identification of larval stages of *Chionodraco* sp. is still unclear (Kellermann 1990), most authors base larval identification on the co-occurrence of adults in the

same area (Hubold 1990; Moteki and Ishimaru 2008). In the Dumont d'Urville Sea, adult *C. hamatus* are overwhelmingly more common than adult *C. myersi* (Koubbi et al. 2009), and some barcoded larvae have been identified as *C. hamatus* (Dettai, pers. comm.). Therefore, larvae of the genus *Chionodraco* are referred to as *C. hamatus* in this study.

The life history and biology for early life stages of *C. hamatus* remain poorly understood. This species has been reported to spawn during summer in the Ross Sea (La Mesa et al. 2003; Vacchi et al. 1996) and demonstrates nesting behavior (Ferrando et al. 2014). Hatching time is not known, although presumably occurs in summer/autumn because few reproductively active females have been observed in the coastal zone near the Dumont d'Urville station in January (Koubbi, pers. comm.).

In polar regions, lipids and their constituent FA play major roles in growth, movement, buoyancy, and reproduction, and also represent the main energy reserves of many polar species (Sargent et al. 2002). Lipid class dynamics reflect the ability of fish larvae to withstand starvation events and can be used to determine a lipid-based condition index (Fraser 1989; Giraldo et al. 2012; Håkanson 1989). The lipid condition index is based on the principal assumption that larval condition in many marine organisms is dependent upon lipid energy reserves, which typically correspond to triacylglycerol (TAG) content. TAG content is dependent on larval size or body mass and must be standardized by cholesterol (Chol) content; Chol is a membrane lipid that is correlated with larval size but is independent of nutritional condition because it is not catabolized during starvation (Fraser 1989).

The relationship between the storage-class lipids (TAG) and the structural membrane lipids (polar lipids, PL) to total lipids (TL) also reflects the main metabolic strategy of larvae and the tradeoff between energy allocation toward growth and/or lipid storage. Further, FA

composition of the TAG fraction directly reflects FA composition of the diet, and differences in FA composition between different organisms or developmental stages therefore reflect dietary similarities/differences (Dalsgaard et al. 2003). In contrast, FA composition of the PL fraction is under genetic control (i.e. independent of diet) and can be used to determine ontogenetic changes within an organism (Giraldo et al. 2015a).

In this study, the lipid composition and dynamics of *C. hamatus* are analyzed to (1) determine lipid composition and the role of lipids in early life strategies of *C. hamatus*, and (2) provide an overview of the lipid class-specific FA composition of *C. hamatus* that can be used to determine trophic pathways in the East-Antarctic pelagic ecosystem.

Materials and Methods

a. Sampling

Field sampling was conducted in the Dumont d'Urville Sea (East Antarctica) during the austral summer 2010-2011 as part of the French IPEV-ICO²TA program (Integrated Coastal Ocean Observations in Terre Adélie) (Koubbi et al. 2011). Larval fish samples were collected from the RV "*L'Astrolabe*" using an Isaacs-Kidd midwater trawl (IKMT) at four stations along transects oriented from nearshore to across the continental shelf located from the Mertz Glacier Tongue to the Adelie Bank. All samples ($n = 26$) were immediately frozen in liquid nitrogen (-196°C) and stored at -80°C until further analysis. Back to our laboratory, samples were thawed on ice, weighted (degree of precision 0.01 mg) and measured to the nearest 0.1 mm with a digital caliper (standard length, SL) before analysis. Two larval developmental stages were differentiated based on flexion and corresponded to specimens smaller (pre-flexion, $n = 16$) and larger (post-flexion, $n = 10$) than 25mm. Preflexion larvae were defined as having a straight notochord with depleted yolk reserves. The second group corresponded to postflexion larvae and was characterized by an

angled notochord with partially developed anal and dorsal fins. These two distinct larval stages were subsequently separated for further lipid analysis.

b. Lipid analysis

Lipid extraction followed the method of Bligh and Dyer (1959) as modified by Mayzaud et al. (2007). Samples were mechanically homogenized and twice-extracted with a one-phase solvent mixture of methanol:chloroform:water (2:1:0.8 v/v/v). Phases were separated overnight by addition of chloroform and NaCl 0.7 % (w/v) to a final solvent ratio of 2:2:1.8 methanol:chloroform:water (v/v/v). The total extract was vacuum concentrated using a rotary evaporator and stored under nitrogen at -80°C in the laboratory.

TL content was determined gravimetrically. Lipid classes were quantified using a chromatographic separation coupled with FID (Flame Photometry Detection) detection on an Iatroscan MK V TH 10. TL extracts were applied to SIII chromarods using a SAS A4100 autospotter programmed to deliver 1 μl of chloroform extract per rod. Analyses were performed in triplicate. Lipid classes were separated by chromatography using a double development procedure with the following solvent systems: n-hexane:benzene:formic acid 80:20:1 (v/v/v) followed by n-hexane:diethyl ether:formic acid 97:3:1.5 (v/v/v). The FID was calibrated for each compound class using commercial standards. For FA analysis, lipid classes were isolated by thin layer chromatography with the hexane:diethyl ether:acetic acid 170:30:2.5 (v/v/v), and the band of PL was then scraped off from the origin and eluted. Lipid classes were visualized using dichlorofluorescein and identification was achieved by comparison with standard mixtures. Following identification, the TAG fraction was then scraped off and eluted. FA from PL and TAG were subsequently converted into methyl esters with 7% boron trifluoride in methanol

(Morrison & Smith 1964). A control thin layer chromatography was performed to verify the efficiency of the trans methylation procedure.

Gas chromatography (GC) of all FA methyl esters (FAME) was carried out on a 30 m (length) x 0.32 mm (width) internal diameter quartz capillary column coated with Famewax (Restek) in a Perkin-Elmer XL Autolab GC equipped with FID. The column was operated isothermally at 185°C for FAME. Helium was used as carrier gas at 7 psig. Injector and detector were maintained at 250°C. Individual components were identified by comparing retention time data with those obtained from laboratory standards (capelin:menhaden oils, 50:50). In addition, FAME samples were hydrogenated to confirm FA determination. The level of accuracy was $\pm 3\%$ for major components, 1–9% for intermediate components, and up to $\pm 25\%$ for minor components ($<0.5\%$ of total FA). In order to have enough material for GC analysis, two to three preflexion larvae were pooled together according to their TAG:Chol ratio. Postflexion larvae were analyzed individually.

c. Statistics

Linear regression of log-transformed values was used to assess the relationship between TL, WW and SL. Contribution of main lipid classes (i.e. Chol, PL and TAG) to TL was assessed using linear regression on standardized values (μg lipid class per mg WW). Correspondence analysis (CA) (Benzécri et al. 1973) has been widely applied to analyze ecological data (Giraldo et al. 2015b) and was chosen here to describe the total inertia in the multi-dimensional FA data as a sample of fewer dimensions that best visualizes patterns in the data. The FA matrix was transformed to relative frequencies and scaled such that each row (or column) can be viewed as a row (or column) of conditional probability distribution. Distances between profiles were

computed with X^2 metrics. This distance gives symmetry to the 2 sets of data such that each factorial axis of the cloud of variables corresponds to a factorial axis of the cloud of observations. Thus, it was possible to represent both descriptors and observations on the plane defined by the factorial axes. The percent variance explained is given for each analysis. Symbol size is proportional to the cosine², illustrating the quality of representation for each point. Computations were performed in the statistical software package R ver. 3.1.0 (R Team Core 2012). Data were transformed to normalize (Zar 1999).

Results

Size, weight and lipid components

The SL of *C. hamatus* (n = 26) ranged from 19.3 to 35.1 mm with WW from 31 to 249 mg. The regression between WW and SL was described by a log-log function (Eq. 1):

$$(1) \log (WW) = -2.1 + 2.9 * \log (SL), F = 100.4, df = 24, p < 0.001, R^2 = 0.80$$

TL content varied from 1.1 to 2.1% (of WW) and was positive correlated with WW and SL, as described by the following equations (Eq. 2 & Eq. 3):

$$(2) \log (TL) = -2.0 + 1.1 * \log (WW), F = 584.4, df = 24, p < 0.001, R^2 = 0.96$$

$$(3) \log (TL) = -4.3 + 3.2 * \log (SL), F = 76.2, df = 24, p < 0.001, R^2 = 0.75$$

Chol and PL content were used to indicate the mass of an individual. A relatively good linear fit (Fig. 1) shows that these lipid components are indeed indicative of larval weight (Eq. 4 & Eq. 5):

$$(4) \log (Chol) = -1.70 + 1.59 * \log (WW), F = 69.1, df = 24, p < 0.001, R^2 = 0.73$$

$$(5) \log (LP) = 0.36 + 1.39 * \log (WW), F = 205.8, df = 24, p < 0.001, R^2 = 0.89$$

PL were the major constituent in both larval stages ($96.2 \pm 1.7\%$ and $94.1\% \pm 2.5\%$ for pre-flexion and post-flexion, respectively) followed by Chol ($1.96 \pm 0.6\%$ and $2.70 \pm 1.1\%$) and TAG ($1.5 \pm 0.2\%$ and $3.0 \pm 1.5\%$). In some samples free FA and diacylglycerols were identified but represented less than 1% of TL. Wax esters were not present.

Lipid dynamics

The contribution of the main lipid classes (i.e. PL, Chol and TG) to TL content was analyzed by standardizing lipids (μg) by WW (mg). In preflexion larvae, PL and Chol were the dominant constituent influencing changes in TL (*Linear regression* $F = 24470$ and 30.84 , respectively; $df=14$, $p < 0.001$), while TAG levels were low relative to other lipid fractions and did not contribute significantly to TL content (*Linear regression*, $F = 0.55$, $df=14$, $p = 0.46$) (Fig. 2). Similarly, PL dominated TL in postflexion larvae, but all lipid classes contributed significantly to changes in TL (*Linear regression* $F = 2039$, 37.15 , 34.6 for PL, Chol and TAG respectively, $df=8$, $p < 0.001$) (Fig. 3).

Condition index TG:Chol

Although there was a slightly better fit between PL and WW than between Chol and WW, Chol was used to standardize TAG content because PL can be mobilized during starvation and might therefore be less stable. The TAG:Chol ratio varied between 0.18 to 1.74 in preflexion larvae and from 0.22 to 2.20 in postflexion larvae. No significant differences were detected for TAG:Chol ratios (*Wilcoxon test*, $W = 45$, $p > 0.05$) between the two larval stages.

Fatty acid signature

Polar lipids

Overall, saturated (~26%) and polyunsaturated FA (PUFA, ~45%) dominated the FA composition of the PL fraction of each larval stage (Table 1). For all developmental stages, saturated FAs were dominated by palmitic acid (16:0) and, to a lesser extent, by stearic acid (18:0). Monoenoic acids (~23%) were dominated by oleic (18:1n-9), vaccenic (18:1n-7) and palmitoleic (16:1n-7), while PUFA were dominated by EPA (20:5n-3) and DHA (22:6n-3). Correspondence analysis explained 52% of the variance between the FA of preflexion and postflexion larvae. Individuals were distributed around the centroid of the analysis with no clear differences between the two stages (Fig 4).

Triacylglycerols

Monoenoic (~28%) and PUFA (~36%) dominated the TAG fraction of both preflexion and postflexion larvae (Table 1). For all developmental stages, saturated FAs were dominated by palmitic acid (16:0), monoenoic acids were dominated by oleic (18:1n-9), palmitoleic (16:1n-7) and vaccenic acid (18:1n-7), and PUFA were dominated by EPA (20:5n-3) and DHA (22:6n-3). Correspondence analysis explained 73.49% of variance in FA signature of the TAG fraction of preflexion and postflexion larvae. However, all samples fell within the centroid of the analysis, indicating no marked differences in FA composition between the two larval stages.

Discussion

Larval development in many marine organisms is largely dependent upon energy reserves, which typically correspond with TAG content. Starvation and predation represent major causes of mortality in larval fishes, particularly once yolk reserves are depleted and larvae must switch to deriving energy from exogenous sources (Leggett and Deblois 1994). Predation risk is lower in individuals with larger body size, and fish larvae are therefore confronted with a

tradeoff between allocating energy reserves to optimize growth, reduce predation risk and increase the opportunity to consume larger prey items, versus storing TAG to increase the likelihood of survival during starvation events. In this study, the dominance of polar structural lipids (PL) compared to TAG content in preflexion larvae (< 25 mm) of *C. hamatus* indicates that small individuals primarily allocate energy reserves toward somatic growth. Conversely, the important contribution of both TAG and PL to TL dynamics of older individuals (> 25 mm, postflexion larvae) suggests an ability to allocate energy toward both growth and lipid storage reserves because the benefits of allocating energy toward growth decrease with increase in body size. Overall, the results of this study highlight a marked ontogeny of energy allocation that likely decreases predation risk for small larvae and favors lipid storage in postlarval *C. hamatus*.

Ratios of TAG:Chol suggested no significant differences in condition factor of pre- and postflexion larvae, indicating that both stages were in good condition and that prey availability during the summer 2011 was sufficient to meet the basal requirements of *C. hamatus* individuals from this study.

PL are known to be strongly regulated and appear to be directly related to membrane requirements and specific energetic pathways (reviewed by Sargent et al., 2002 and Dalsgaard et al., 2003). Although the lipid class dynamics of preflexion and postflexion *C. hamatus* larvae were different, neither of these developmental stages exhibited any detectable ontogeny in the FA pattern of PL and TAG fractions. High levels of DHA with a mean EPA: DHA ratio of 0.6 ± 0.05 reflects the role of polyunsaturated n-3 FA in the maintenance of biological membranes in *C. hamatus* larvae. In contrast to PL, the FA composition of TAG is related to trophic interactions in marine organisms (Dalsgaard et al. 2003; Mayzaud et al. 2011). In this study, there were no significant differences between the pattern of FA trophic markers for pre- and

post- larval *C. hamatus*, indicating similar diet across larval stages. Specifically, early life stages of *C. hamatus* were characterized by low contribution of *Calanus*-type copepod markers ($\Sigma C20:1$, $C22:1$) and a relatively high value of the ratio $C18:1n-9/C18:1n-7$, a typical carnivory index in marine trophic ecology (Hagen et al. 2000). This finding is consistent with previous work demonstrating that young stages of euphausiids (furcilia to juveniles) and fish larvae constitute the most important food source for channichthyids larvae (Kock 2005).

Early life stages of the pelagic fish, *Pleuragramma antarctica* (Antarctic Silverfish), one of the most important key species of Antarctica (along with the krill *Euphausia superba*) (Corsolini et al. 2002) may be of particular importance in the diet of *C. hamatus*. Early life stages of *C. hamatus* have been previously reported as specialized predators in the Ross Sea, feeding on a relatively narrow niche width, with *P. antarctica* larvae as their overwhelmingly dominant prey (La Mesa et al. 2011). Compared to the carnivory index previously described for *P. antarctica* larvae (1.34 and 1.06) (Mayzaud et al. 2011, Tavernier et al. 2012), the relative high value (1.93) of the ratio $C18:1n-9/C18:1n-7$ reported for *C. hamatus* is in good agreement with previous studies. Moreover, stable nitrogen signatures ($\delta^{15}N$), mainly used to establish trophic relationships, have been reported as 10.5 ‰ for *C. hamatus* larvae (SL= 31-38 mm) (Cherel et al. 2011) and 6.7 ‰ for larval *P. antarctica* (Giraldo et al. 2011). Considering that, on average, a ~3‰ enrichment in $\delta^{15}N$ values accompanies each trophic step, these results also support the hypothesis that larval *P. antarctica* could be the main prey item for *C. hamatus* larvae. The hatching time of *C. hamatus* and *P. antarctica* are thought to coincide. Further, *C. hamatus* and *P. antarctica* co-occur spatially and have a similar geographic distribution in the western Ross Sea (La Mesa et al. 2010) and offshore of the Antarctic Peninsula (White and North 1987). Future studies could use FA as dietary biomarkers to test the hypothesis that survival of larval *C.*

hamatus is closely linked to spatial and temporal dynamics of young stages of euphausiids and *P. antarctica* larvae.

Acknowledgements

This project is a contribution to the Zone Atelier Antarctique du CNRS. Field surveys were funded by the French Polar Institute (IPEV). The authors thank the anonymous reviewers for their valuable comments to improve the manuscript and Shannon MacPhee from Fisheries and Oceans Canada for proofreading and editing.

References

- Benzécri JP, Benzécri F, Birou A, Blumenthal S, De Bœck A (1973) L'analyse de données Vol. 2: L'analyse des correspondances. Dunod, Paris
- Bligh EG, Dyer WJ (1959) A rapid method of total lipid extraction and purification. Can J Biochem Phys 37(8):911-917
- Cherel Y, Koubbi P, Giraldo C et al (2011) Isotopic niches of fishes in coastal, neritic and oceanic waters off Adélie Land, Antarctica. Polar Sci 5:286-297
- Corsolini S, Romeo T, Ademollo N, Greco S, Focardi S (200) POPs in key species of marine Antarctic ecosystem. Microchem J 73(1):187-93
- Dalsgaard J, St John M, Kattner G, Müller-Navarra D, Hagen W (2003) Fatty acid trophic markers in the pelagic marine environment. Adv Mar Biol 46:225-340
- Efremenko V (1987) Species composition and distribution of mesopelagic fish eggs and larvae in the Ross Sea. Biological and oceanographic investigations of the Pacific sector of Antarctica 186-193
- Ferrando S, Castellano L, Gallus L, Ghihiotti L, Masini MA, Pisano E., Vacchi M (2014) A Demonstration of Nesting in Two Antarctic Icefish (Genus *Chionodraco*) Using a Fin Dimorphism Analysis and Ex Situ Videos. PLoS ONE 9(3): e90512. doi:10.1371/journal.pone.0090512
- Fraser AJ (1989) Triacylglycerol Content as a Condition Index for Fish, Bivalve, and Crustacean Larvae. Can J Fish Aquat Sci 46:1868-1873
- Giraldo C, Cherel Y, Vallet C, Mayzaud P, Tavernier E, Moteki M, Hosie G, Koubbi P (2011) Ontogenic changes in the feeding ecology of the early life stages of the Antarctic silverfish (*Pleuragramma antarcticum*) documented by stable isotopes and diet analysis in the Dumont d'Urville Sea (East Antarctica). Polar Sci 5(2):252-63
- Giraldo C, Cherel Y, Vallet C et al (2012) Lipid components as a measure of nutritional condition in fish larvae (*Pleuragramma antarcticum*) in East Antarctica. Marine Biol 160:877-887
- Giraldo C, Mayzaud P, Tavernier E, Boutoute M, Penot F, Koubbi P (2015a) Lipid dynamics and trophic patterns in *Pleuragramma antarctica* life stages. Antarctic Sci 27(5):429-438
- Giraldo C, Stasko A, Choy ES, Rosenberg B, Majewski A, Power M, Swanson H, Loseto L, Reist JD (2015b) Trophic variability of Arctic fishes in the Canadian Beaufort Sea: a fatty acids and stable isotopes approach. Polar Biol 1-6. doi:10.1007/s00300-015-1851-4
- Granata A, Cubeta A, Guglielmo L, et al (2002) Ichthyoplankton abundance and distribution in the Ross Sea during 1987–1996. Polar Biol 25:187-202
- Hagen W, Kattner G, Friedrich C (2002) The lipid compositions of high-Antarctic notothenioid fish species with different life strategies. Polar Biol 23(11):785-91

290 Håkanson JL (1989) Analysis of lipid components for determining the condition of anchovy larvae,
 291 *Engraulis mordax*. Marine Biol 102:143-151
 292 Hubold G (1990) Seasonal Patterns of Ichthyoplankton Distribution and Abundance in the Southern
 293 Weddell Sea. In: Kerry KR, Hempel G (eds) Antarctic Ecosystems. Springer Berlin Heidelberg,
 294 pp 149-158
 295 Kellermann A (1990) Catalogue of early life stages of Antarctic notothenioid fishes. Ber Polarforsch
 296 67:45-136
 297 Kellermann A, Schadwinkel S (1991) Winter aspects of the ichthyoplankton community in Antarctic
 298 Peninsula waters. Polar Biol 11:117-127
 299 Kock K-H (2005) Antarctic icefishes (Channichthyidae): a unique family of fishes. A review, Part I Polar
 300 Biol 28:862-895
 301 Koubbi P, Duhamel G, Hecq JH et al (2009) Ichthyoplankton in the neritic and coastal zone of Antarctica
 302 and Subantarctic islands: A review. J Mar Syst 78:547-556
 303 Koubbi P, Ozouf-Costaz C, Goarant A et al. (2010) Estimating the biodiversity of the East Antarctic shelf
 304 and oceanic zone for ecoregionalisation: example of the ichthyofauna of the CEAMARC
 305 (Collaborative East Antarctic Marine Census) CAML surveys. Polar Sci 4:115-133
 306 Koubbi P, O'Brien C, Loots C, Giraldo C, Smith M, Tavernier E, Vacchi M, Vallet C, Chevallier J,
 307 Moteki M (2011) Spatial distribution and inter-annual variations in the size frequency distribution
 308 and abundances of *Pleuragramma antarcticum* larvae in the Dumont d'Urville Sea from 2004 to
 309 2010. Polar Sci 5(2):225-38
 310 La Mesa M, Ashford J (2008) Age and growth of ocellated icefish, *Chionodraco rastrispinosus* DeWitt
 311 and Hureau, 1979, from the South Shetland Islands. Polar Biol 31:1333-1342
 312 La Mesa M, Caputo V, Rampa R, Vacchi M (2003) Macroscopic and histological analyses of gonads
 313 during the spawning season of *Chionodraco hamatus* (Pisces, Channichthyidae) off Terra Nova
 314 Bay, Ross Sea, Southern Ocean. Polar Biol 26:621-628
 315 La Mesa M, Catalano B, Greco S (2011) Larval feeding of *Chionodraco hamatus* (Pisces,
 316 Channichthyidae) in the Ross Sea and its relation to environmental conditions. Polar Biol 34:127-
 317 137
 318 La Mesa M, Catalano B, Russo A, Greco S, Vacchi M, Azzali M (2010) Influence of environmental
 319 conditions on spatial distribution and abundance of early life stages of Antarctic silverfish,
 320 *Pleuragramma antarcticum* (Nototheniidae), in the Ross Sea. Antarct Sci 22:243-254
 321 Leggett WC, Deblois E (1994) Recruitment in marine fishes: Is it regulated by starvation and predation in
 322 the egg and larval stages? Neth J Sea Res 32:119-134
 323 Loeb VJ, Kellermann AK, Koubbi P, North AW, White MG (1993) Antarctic larval fish assemblages: a
 324 review. B Mar Sci 53:416-449
 325 Mayzaud P, Boutoute M, Perissinotto R, Nichols P (2007) Polar and neutral lipid composition in the
 326 pelagic tunicate *Pyrosoma atlanticum*. Lipids 42(7):647-57
 327 Mayzaud P, Chevallier J, Tavernier E, Moteki M, Koubbi P (2011) Lipid composition of the Antarctic
 328 fish *Pleuragramma antarcticum*. Influence of age class. Polar Sci 5:264-271
 329 Morrison WR, Smith LM (1964) Preparation of fatty acid methyl esters and dimethylacetals from lipids
 330 with boron fluoride-methanol. J Lipid Res 5(4):600-8
 331 Moteki M, Ishimaru T (2008) Development of feeding and swimming functions in larvae of *Chionodraco*
 332 *rastrispinosus* (Channichthyidae). Cybium 32:247-251
 333 Moteki M, Koubbi P, Pruvost P, Tavernier E, Hulley P-A (2011) Spatial distribution of pelagic fish off
 334 Adélie and George V Land, East Antarctica in the austral summer 2008. Polar Sci 5:211-224
 335 North A (1988) Distribution of fish larvae at South Georgia: horizontal, vertical, and temporal
 336 distribution and early life history relevant to monitoring year-class strength and recruitment
 337 SCCAMLR, SeL Sci Pap 4:1987
 338 R Team Core (2012) R: A language and environment for statistical computing
 339 Sargent JR, Tocher DR, Bell JG (2002) The lipids. Fish nutrition 3:181-257

340 Tavernier E, Mayzaud P, Boutoute M, Vallet C, Koubbi P (2012) Lipid characterization of
 341 *Pleuragramma antarcticum* (Nothoteniidae) larvae off East Antarctica (139 E–145.10 E) during
 342 summer. Polar Biol 35(6):829-40
 343 Vacchi M, Williams R, La Mesa M (1996) Reproduction in three species of fish from the Ross Sea and
 344 Mawson Sea. Antarct Sci 8:185-192
 345 Vallet C, Labat J-P, Smith M, Koubbi P (2011) Interannual variations in euphausiid life stage distribution
 346 in the Dumont d'Urville Sea from 2004 to 2008. Polar Sci 5:166-178
 347 White MG, North AW (1987) Postlarval Notothenioides and midwater fish collected during the SIBEX
 348 cruise by British Antarctic Survey, 1985. In Proc Congr Europ Ichthyol, Stockholm Vol
 349 1985:405-411)
 350 Zar JH (1999) Biostatistical Analysis (4th Edition) Prentice Hall, Englewood Cliffs, New Jersey
 351

352 Figure 1: Linear regression on log-transformed membrane lipids (Chol: cholesterol, PL: polar lipids, μ g)
353 and wet weight (WW).

354 Figure 2: Contribution of the main lipid classes (PL: Polar Lipids, TAG: Triacylglycerol, Chol:
355 Cholesterol) to the total lipid (TL) content of preflexion larvae *C. hamatus*.

356 Figure 3: Contribution of the main lipid classes (PL: Polar Lipids, TAG: Triacylglycerol, Chol:
357 Cholesterol) to the total lipid (TL) content of postflexion larvae of *C. hamatus*.

358 Figure 4: Correspondence Analysis of the fatty acid signature of the polar lipid fraction for preflexion
359 larvae (L1-6) and postflexion larvae (P1-10) of *C. hamatus*.

Table 1: Fatty acid composition of the triacylglycerol and polar lipid fraction in pre-flexion and post-flexion larvae. Minor fatty acids (<1%) common for all individuals are not shown. Data is reported as means $\pm$ standard deviation.

	Triacylglycerol		Polar Lipids	
Fatty acid/Stage	Pre - flexion	Post - flexion	Pre - flexion	Post - flexion
C14	2.49 $\pm$ 0.64	2.60 $\pm$ 0.54	0.96 $\pm$ 0.19	0.97 $\pm$ 0.19
C15	0.47 $\pm$ 0.22	0.30 $\pm$ 0.04	0.18 $\pm$ 0.02	0.18 $\pm$ 0.01
ISO17	0.76 $\pm$ 0.12	0.84 $\pm$ 0.11	0.53 $\pm$ 0.08	0.55 $\pm$ 0.07
ANT17	0.20 $\pm$ 0.06	0.21 $\pm$ 0.06	0.14 $\pm$ 0.05	0.10 $\pm$ 0.05
C16	12.88 $\pm$ 0.77	11.82 $\pm$ 0.37	20.39 $\pm$ 0.49	20.21 $\pm$ 1.37
C17	2.27 $\pm$ 0.36	2.88 $\pm$ 0.23	0.26 $\pm$ 0.03	0.25 $\pm$ 0.04
C18	3.11 $\pm$ 0.34	2.79 $\pm$ 0.22	3.70 $\pm$ 0.25	3.72 $\pm$ 0.29
Σ Saturates	23.17 $\pm$ 1.27	22.15 $\pm$ 0.86	26.43 $\pm$ 0.43	26.27 $\pm$ 1.62
C16:1n7	7.32 $\pm$ 0.56	7.02 $\pm$ 0.47	3.91 $\pm$ 0.16	3.95 $\pm$ 0.35
C16:1n5	0.81 $\pm$ 0.15	0.85 $\pm$ 0.15	1.03 $\pm$ 0.11	1.00 $\pm$ 0.10
C18:1n9	13.98 $\pm$ 1.28	14.11 $\pm$ 1.29	7.21 $\pm$ 0.24	7.92 $\pm$ 0.71
C18:1n7	7.75 $\pm$ 1.33	7.28 $\pm$ 0.95	8.08 $\pm$ 0.50	7.83 $\pm$ 1.01
C18:1n5	0.95 $\pm$ 0.03	0.97 $\pm$ 0.19	0.75 $\pm$ 0.05	0.79 $\pm$ 0.08
C20:1n9	1.09 $\pm$ 0.24	0.90 $\pm$ 0.32	1.10 $\pm$ 0.11	1.09 $\pm$ 0.11
C20:1n7	0.36 $\pm$ 0.42	0.21 $\pm$ 0.03	0.16 $\pm$ 0.03	0.17 $\pm$ 0.03
C24:1n9	0.23 $\pm$ 0.10	0.20 $\pm$ 0.09	0.16 $\pm$ 0.03	0.21 $\pm$ 0.05
Σ Monoenes	33.72 $\pm$ 1.78	32.56 $\pm$ 1.61	22.73 $\pm$ 0.28	23.29 $\pm$ 1.57
C18:2n7	0.30 $\pm$ 0.24	0.37 $\pm$ 0.29	0.11 $\pm$ 0.03	0.13 $\pm$ 0.03
C18:2n6	2.28 $\pm$ 0.22	2.42 $\pm$ 0.10	1.15 $\pm$ 0.11	1.34 $\pm$ 0.13
Σ Dienes	2.79 $\pm$ 0.13	3.06 $\pm$ 0.39	1.51 $\pm$ 0.15	1.76 $\pm$ 0.20
C16:3n6	0.54 $\pm$ 0.17	0.64 $\pm$ 0.11	0.32 $\pm$ 0.06	0.38 $\pm$ 0.07
C16:3n3	0.43 $\pm$ 0.04	0.42 $\pm$ 0.02	0.25 $\pm$ 0.05	0.27 $\pm$ 0.05
C18:3n6	1.41 $\pm$ 0.78	0.57 $\pm$ 0.13	0.17 $\pm$ 0.06	0.14 $\pm$ 0.07
C18:3n3	0.98 $\pm$ 0.37	1.23 $\pm$ 0.13	0.40 $\pm$ 0.13	0.37 $\pm$ 0.14
Σ Trienes	3.46 $\pm$ 0.76	2.97 $\pm$ 0.26	1.22 $\pm$ 0.22	1.21 $\pm$ 0.21
C16:4n3	0.37 $\pm$ 0.08	0.36 $\pm$ 0.11	0.36 $\pm$ 0.05	0.37 $\pm$ 0.06
C18:4n3	1.93 $\pm$ 0.66	2.77 $\pm$ 0.35	0.49 $\pm$ 0.14	0.58 $\pm$ 0.09
C20:4n6	0.62 $\pm$ 0.11	0.63 $\pm$ 0.09	1.36 $\pm$ 0.14	1.35 $\pm$ 0.10
Σ Tetraene	3.36 $\pm$ 0.69	4.14 $\pm$ 0.36	2.41 $\pm$ 0.08	2.44 $\pm$ 0.11
C20:5n3	13.6 $\pm$ 0.69	14.85 $\pm$ 0.59	16.77 $\pm$ 0.89	16.26 $\pm$ 1.23
C22:5n3	0.71 $\pm$ 0.06	0.68 $\pm$ 0.08	0.76 $\pm$ 0.07	0.60 $\pm$ 0.04
C22:6n3	18.58 $\pm$ 1.03	19.24 $\pm$ 1.34	28.02 $\pm$ 1.09	28.03 $\pm$ 2.16
Σ PUFA	33.51 $\pm$ 1.90	35.12 $\pm$ 1.74	45.55 $\pm$ 0.64	44.89 $\pm$ 2.97

Figure 1

Figure 2

Figure 3

Figure 4