



HAL
open science

Why the P3b is still a plausible correlate of conscious access? A commentary on Silverstein et al., 2015

Lionel Naccache, Sébastien Marti, Jacobo D. Sitt, Darinka Trübutschek, Lucie Berkovitch

► To cite this version:

Lionel Naccache, Sébastien Marti, Jacobo D. Sitt, Darinka Trübutschek, Lucie Berkovitch. Why the P3b is still a plausible correlate of conscious access? A commentary on Silverstein et al., 2015. *Cortex*, 2016, 10.1016/j.cortex.2016.04.003 . hal-01317650

HAL Id: hal-01317650

<https://hal.sorbonne-universite.fr/hal-01317650>

Submitted on 18 May 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Commentary:
Why the P3b is still a plausible correlate of conscious access? A commentary on Silverstein et al., 2015.

Lionel Naccache¹²³⁴⁵, Sébastien Marti⁶, Jacobo D. Sitt³⁴, Darinka Trübtschek⁶, Lucie Berkovitch⁶⁷

1-AP-HP, Groupe hospitalier Pitié-Salpêtrière, Department of Neurology, Paris, France

2-AP-HP, Groupe hospitalier Pitié-Salpêtrière, Department of Neurophysiology, Paris, France

3-INSERM, U 1127, F-75013, Paris, France

4-Institut du Cerveau et de la Moelle épinière, ICM, PICNIC Lab, F-75013, Paris, France

5-Sorbonne Universités, UPMC Univ Paris 06, Faculté de Médecine Pitié-Salpêtrière, Paris, France

6-Cognitive Neuroimaging Unit, CEA DSV/I2BM, INSERM, Université Paris-Saclay, NeuroSpin Center, 91191 Gif/Yvette, France

7-Ecole des Neurosciences de Paris Ile-de-France, 75006 Paris, France

We read with interest the article by Silverstein and colleagues (Silverstein, Snodgrass, Shevrin and Kushwaha 2015) who questioned the putative specificity of the P3b event-related potentials (ERP) component as a neural signature of conscious access to a visual representation. Prior to this new study, numerous empirical reports revealed that a brain response peaking ~300ms after stimulus onset and maximally distributed over parietal electrodes – the so called P3b – is closely related to subjective visibility (Vogel, Luck and Shapiro 1998; Sergent, Baillet and Dehaene 2005). These experimental findings provided the bases to develop neuronal and computational theories of consciousness such as the global workspace model (Dehaene and Naccache 2001; Dehaene, Changeux, Naccache, Sackur and Sergent 2006; Dehaene and Changeux 2011). Silverstein and colleagues used a ‘passive attentive’ version of a masked visual odd-ball paradigm while recording scalp ERPs. On each trial, subjects were presented with either the masked word ‘LEFT’ (in 80% or 20% of trials) or the masked word ‘RIGHT’ (in 20% or 80% of trials). Word frequency was balanced across subjects, who were asked to attend carefully to the masked sequence. Not only were they instructed that this sequence contained a masked word, but also that: “however implausible it might seem, our prior data suggested that the stimuli would nonetheless be unconsciously perceived and produce brain wave effects—but only if they maintained their attention”. When contrasting ERPs elicited by rare and frequent masked words, Silverstein and colleagues identified a P3b ERP

component followed by a late, and sustained, slow wave (LSW). Given that subjects subjectively reported the absence of conscious perception of words, and that they performed at chance-level in a stimulus detection task performed after the main experiment, Silverstein and colleagues concluded that a P3b can be observed during unconscious perception. If valid, their interpretation would then simply invalidate the P3b as a possible candidate neural signature of conscious access.

This original and provocative study, however, raises both methodological and conceptual concerns which need to be addressed before one can adopt Silverstein and colleagues interpretation.

A set of methodological problems

The P3b is part of a larger complex of positive deflections – the so-called P300. Of particular importance here, the P3a can be functionally distinguished from the P3b: it is known to occur in the absence of conscious perception (Muller-Gass, Macdonald, Schroger, Sculthorpe and Campbell 2007) and even in non-conscious patients (Faugeras, Rohaut, Weiss, Bekinschtein, Galanaud, Puybasset et al. 2012). P3a and P3b are both positive deflections and occur in similar time windows, but they can nevertheless be separated based on their topographies. The P3b is maximally distributed over parietal electrodes, while the P3a is more frontally distributed. The spatial sampling of the EEG signal is therefore critical to separate these ERPs. Surprisingly, the authors only used 3 midline (Fz, Cz, Pz) electrodes referenced to linked ears, as well as 2 electrodes at the right eye to detect eye movement artifacts. As expected for a P3b component, the effect reported by Silverstein and colleagues was maximal over Pz, but we simply don't have access to scalp topographies of the ERP effects reported in this work. We agree that there is no intrinsic relation between the number of electrodes and the quality of a result, but in the context of distinguishing P3b from P3a ERP components this limitation turns into a genuine problem.

In the same vein, one of the most reliable findings in the vast odd-ball literature, is the existence of a N2 and mismatch negativity (MMN) ERP effect occurring before the P3 complex (Tiitinen, May, Reinikainen and Naatanen 1994). The apparent absence of such an effect (a small inverse difference is seen on Figure 2) confirms the necessity of sampling brain activity with a richer spatial resolution in order to describe reliably the observed effects.

Moreover, shortcomings in statistical analyses of the ERPs deserve further discussion. The visual inspection of 'effects' suggest that the effect size reported by Silverstein et al are not quite different from fluctuations within the baseline and from other periods of ERPs (see e.g. Figure 2C). Actually, the authors did not assess significant differences on the entire time course of the ERP but only on predefined time windows. Thus it is impossible to determine whether the reported effects are temporally and spatially precise and specific to the P3b. A better approach would consist in performing non selective sample-by-sample tests, and then identifying temporal clusters during which ERPs significantly differed.

More importantly, although the article by Silverstein et al. opens by asking the fundamental question “How can perceptual awareness be indexed in humans?”, their experimental design is lacking the crucial comparison of unconscious ‘P3b’ with its conscious equivalent. Rather than using exclusively masked trials, the authors could have added unmasked trials, in order to compare the properties (latency, amplitude and effect size, duration, topography) of conscious and unconscious ERP effects. By doing so previous studies could identify specific components of conscious access (Dehaene, Changeux, Naccache, Sackur and Sergent 2006). From a theoretical perspective, we previously mentioned and modeled the possibility for a masked stimulus to “evoke transient workspace activity of variable intensity and duration” (see also Figure 1 in (Dehaene and Naccache 2001)). Such transient and partial activation of the workspace could appear as brief and small patterns of activity distinct from a large and sustained P3b component. Therefore, without this crucial conscious contrast, it becomes almost impossible to precisely qualify the observed ERP effect.

It is noteworthy that according to our theory, conscious access associated with the P3b is also associated with other signatures (Gaillard, Dehaene, Adam, Clemenceau, Hasboun, Baulac et al. 2009) such as: long-range synchrony in theta-alpha-beta band, decrease of alpha power, and late increase of gamma power. None of these neural signatures, complementary to the P3b, are tested here and the nature of the observed ERP effects therefore remains unclear.

Additionally, the interesting use by Silverstein and al. of ‘LEFT’ and ‘RIGHT’ words opened the possibility of complementing the results by lateralized readiness potentials (LRPs) analyses. Such analyses proved to be very useful to explore both unconscious and conscious processing of masked primes (Dehaene, Naccache, Le Clec, Koechlin, Mueller, Dehaene-Lambertz et al. 1998; Eimer and Schlaghecken 1998). Unfortunately, the use of only 3 midline electrodes, and the absence C3/C4 electrodes precluded this interesting complementary approach.

Furthermore, from a Bayesian perspective, we consider the authors should have mentioned and discussed more extensively the large set of empirical evidence that their finding seems to contradict: numerous studies conducted in normal controls as well as in many clinical settings (e.g.: blindsight, visual neglect) support the P3b theory by reporting rich unconscious processing of visual stimuli without any late P3b signature (for a review see (Dehaene and Changeux 2011)). This literature, acting here as a strong prior against Silverstein and colleagues interpretation, needs to be addressed.

Conscious metacognition of unconscious perceptual processes?

Beyond these notable methodological issues, this article also raises a more profound question. The major difference between this study and previous studies rests in the fact that subjects were told from the very beginning of the presence of masked stimuli, and were instructed to pay attention to them very carefully. Therefore, even if we discard the methodological issues we just raised, and consider that these results are correct, it may be the case that the P3b signature observed here between deviant

and standard stimuli would correspond to a metacognitive effect, that is to say to a conscious access to the consequence of unconscious processing of masked primes. For instance, a motor effect induced by the processing of the rare 'LEFT' prime (or 'RIGHT' for other subjects) in motor areas may well lead to conscious access to a subjective confidence information that the prime was deviant or standard. By amplifying subjects' attention to monitor prime processing, this metacognitive interpretation may well explain the striking pattern of results reported here. Interestingly, a growing empirical evidence demonstrates that a large class of unconscious cognitive processes are strongly influenced by the conscious posture and endogenous attentional allocation (Naccache, Blandin and Dehaene 2002). In addition to such an amplification, it might be the case that subject informed of the presence of subliminal stimuli could more easily introspect a form of surprise originating either from perceptual or from motor-related areas ('LEFT', 'RIGHT'). In other words, this study may illustrate conscious access to the downstream effects of an unconsciously perceived stimulus. Interestingly, a recent study using a visual masked priming paradigm reported that the conflict between masked prime and visible target stimuli modulated two ERP components (Desender, Van Opstal, Hughes and Van den Bussche in press): an early N2 component, as well as a late P3 complex. During this experiment, subjects had to perform two tasks on each trial: they first had to respond to the target, and then to introspect on the difficulty of the trial. Nicely, introspection of the prime-target conflict elicited by the unconscious processing of the prime was possible, and correlated only with the P3 component. Similarly, in the study by Silverstein and colleague, one may suppose that the P3b component and the LSW they observed corresponds to a conscious introspection of processes elicited by the unconsciously perceived prime.

As a conclusion, if the results reported in Silverstein et al do correspond to a genuine P3b ERP component (but see our methodological concerns above), they may elegantly illustrate the complex relations prevailing between conscious and unconscious processes, and still not refute the relation prevailing between conscious access and the P3b ERP component.

Funding

This work has been supported by the Fondation pour la Recherche Médicale ('Equipe FRM 2015' grant to L.N. and by the 'Recovery of consciousness after severe brain injury Phase II' grant of the James S. McDonnell Foundation.

References

- Dehaene, S. and J. P. Changeux (2011). "Experimental and theoretical approaches to conscious processing." *Neuron* **70**(2): 200-227.
- Dehaene, S., J. P. Changeux, L. Naccache, J. Sackur and C. Sergent (2006). "Conscious, preconscious, and subliminal processing: a testable taxonomy." *Trends in Cognitive Sciences* **10**(5): 204-211.

- Dehaene, S. and L. Naccache (2001). "Towards a cognitive neuroscience of consciousness: basic evidence and a workspace framework." *Cognition* **79**: 1-37.
- Dehaene, S., L. Naccache, H. G. Le Clec, E. Koechlin, M. Mueller, G. Dehaene-Lambertz, et al. (1998). "Imaging unconscious semantic priming." *Nature* **395**: 597-600.
- Desender, K., F. Van Opstal, G. Hughes and E. Van den Bussche (in press). "The temporal dynamics of metacognition: Dissociating task-related activity from later metacognitive processes." *Neuropsychologia*.
- Eimer, M. and F. Schlaghecken (1998). "Effects of masked stimuli on motor activation: behavioral and electrophysiological evidence." *J Exp Psychol Hum Percept Perform* **24**: 1737-1747.
- Faugeras, F., B. Rohaut, N. Weiss, T. Bekinschtein, D. Galanaud, L. Puybasset, et al. (2012). "Event related potentials elicited by violations of auditory regularities in patients with impaired consciousness." *Neuropsychologia* **50**: 403-418.
- Gaillard, R., S. Dehaene, C. Adam, S. Clemenceau, D. Hasboun, M. Baulac, et al. (2009). "Converging intracranial markers of conscious access." *PLoS Biol* **7**: e61.
- Muller-Gass, A., M. Macdonald, E. Schroger, L. Sculthorpe and K. Campbell (2007). "Evidence for the auditory P3a reflecting an automatic process: elicitation during highly-focused continuous visual attention." *Brain research* **1170**: 71-78.
- Naccache, L., E. Blandin and S. Dehaene (2002). "Unconscious masked priming depends on temporal attention." *Psychol Sci* **13**: 416-424.
- Sergent, C., S. Baillet and S. Dehaene (2005). "Timing of the brain events underlying access to consciousness during the attentional blink." *Nat Neurosci* **8**: 1391-1400.
- Silverstein, B. H., M. Snodgrass, H. Shevrin and R. Kushwaha (2015). "P3b, consciousness, and complex unconscious processing." *Cortex; a journal devoted to the study of the nervous system and behavior* **73**: 216-227.
- Tiitinen, H., P. May, K. Reinikainen and R. Naatanen (1994). "Attentive novelty detection in humans is governed by pre-attentive sensory memory." *Nature* **372**: 90-92.
- Vogel, E. K., S. J. Luck and K. L. Shapiro (1998). "Electrophysiological evidence for a postperceptual locus of suppression during the attentional blink." *J Exp Psychol Hum Percept Perform* **24**: 1656-1674.