



**HAL**  
open science

## Allometric relationships for intertidal macroalgae species of commercial interest

Doriane Stagnol, Marine Macé, Christophe Destombe, Dominique Davoult

### ► To cite this version:

Doriane Stagnol, Marine Macé, Christophe Destombe, Dominique Davoult. Allometric relationships for intertidal macroalgae species of commercial interest. *Journal of Applied Phycology*, 2016, pp.1-5. 10.1007/s10811-016-0860-1 . hal-01318013

**HAL Id: hal-01318013**

<https://hal.sorbonne-universite.fr/hal-01318013v1>

Submitted on 19 May 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 1 **Allometric relationships for intertidal macroalgae species of commercial interest**

2 Doriane Stagnol <sup>a,b</sup>, Marine Macé <sup>a,b</sup>, Christophe Destombe <sup>a,c</sup>, Dominique Davoult <sup>a,b</sup>

3 <sup>a</sup> Sorbonne Universités, UPMC Univ Paris 6, Station Biologique, Place Georges Teissier, F-29682  
4 Roscoff, France

5 <sup>b</sup> CNRS, UMR 7144 AD2M, Place Georges Teissier, F-29682, Roscoff, France

6 <sup>c</sup> CNRS, UMI 3614 EBEA, Place Georges Teissier, F-29682, Roscoff, France

7 \* Corresponding author: doriane.stagnol@gmail.com Tel: +33 2 98 29 23 23 Fax: +33 2 98 29 23 24

8

## 9 **Abstract**

10 The demand for seaweeds has intensified in recent decades and will most certainly continue to  
11 expand. Several methods exist to evaluate the biomass of seaweeds in the field but most of  
12 them are destructive. The objectives of this study were (1) to develop and evaluate allometric  
13 equations for estimating seaweed biomass in the field for some harvested species, and (2) to  
14 provide uniform calculated dry/wet biomass ratios to estimate the relative water content of  
15 these seaweeds. Sampling and measurements of more than 350 seaweeds individuals were  
16 carried out for 8 species of commercial interest. Our models were fitted for both power and  
17 linear equations and were tested for different explanatory variables. While the power equation  
18 was found to be the best for predicting biomass of all species, we found that the best  
19 descriptive biometric variable varies according to seaweed morphology. Species with a bushy  
20 morphology were best described by the volume, while long stringy species were best  
21 described by the length and flat species by the surface. This study attempts to provide  
22 nondestructive tools that could be used by professional seaweed harvesters, their employers as  
23 well as scientists and public regulators, to assess the harvest potential of a field of seaweed in  
24 a nondestructive approach.

25 **Keywords:** biometrics, biomass, allometry, harvesting

## 26 **Acknowledgements**

27 The authors thank all the people who helped carry out the fieldwork. Our study was funded by  
28 the ALGMARBIO project, InitiativeBioBretagne (IBB), FranceAgriMer (National Institute of  
29 Agricultural and Marine Products), the Brittany Regional Council, the Côte d'Armor  
30 Departmental Council, the European Fisheries Fund (EFF), and benefited from the support of  
31 the French Government run by the National Research Agency and with regards to the  
32 investment expenditure programme IDEALG ANR-10-BTBR-04.

### 33 **Introduction**

34 Seaweed diversity and community structure are highly impacted and threatened by physical  
35 and/or anthropic forcing such as climatic changes (Airoldi and Beck 2007; Mangialajo et al.  
36 2008). These continued stressors cause the fragmentation and loss of canopy-forming algae  
37 worldwide (Connell et al. 2008; Airoldi et al. 2008), and even could lead to their extinction  
38 (Estes et al. 1989). Besides producing a valuable crop to the seaweed harvesters, macroalgae  
39 plays an important role in the primary production of nearshore ecosystems (Golléty et al.  
40 2008; Migné et al. 2015). Within this context of increasing pressures, one can wonder about  
41 the effects of the loss of canopy-forming algae on primary production and on carbon and  
42 nitrogen biochemical cycles. Accurate and efficient estimation of biomass in such populations  
43 is central to understand and monitor their net contribution in providing these ecosystem  
44 services.

45 Ecologists, botanists and foresters estimate biomass for a wide range of purposes, such as  
46 assessment of crop value, site productivity, as well as nutrient recycling. Destructive sampling  
47 has generally been used to obtain an accurate measure of biomass at a particular sampling  
48 point, including in seaweed populations (Mathieson and Guo 1992; Vadas, Sr. et al. 2004).  
49 However, these destructive approaches can have short and long-term consequences on the  
50 associated ecosystem, including decrease in invertebrate abundance and richness (Benedetti-  
51 Cecchi et al. 2001; Watt and Scrosati 2013), replacement by grazers or turfs (Perkol-Finkel  
52 and Airoldi 2010), or reduction in algal biomass and primary productivity (Golléty et al.  
53 2008; Tait and Schiel 2011). In order to reduce these effects, nondestructive methods were  
54 developed to answer specific questions in plants (Niklas and Enquist 2002; Sack et al. 2003;  
55 Scrosati et al. 2005; Mccarthy and Enquist 2007; Poorter et al. 2012). Without losing their  
56 scientific rigor, the use of nondestructive sampling methods permits the absence of laboratory  
57 work, simplifying data processing and reducing the total monitoring costs. One of these

58 nondestructive methods is based on fitting so-called allometric equations to convert field  
59 inventory data to biomass estimates (Chave et al. 2005; Jonson and Freudenberger 2011; Paul  
60 et al. 2013). In seaweeds, this method was mainly applied in population dynamics of red and  
61 brown algae (Åberg 1990; Lindgren et al. 1998; Engel et al. 2001) or to estimate growth  
62 during two sampling events (Vaz-Pinto et al. 2014). Allometric equations are particularly  
63 useful to evaluate biomass allocation pattern (i.e. the relative amount of biomass present in  
64 the various organs; Niklas and Enquist 2002), to measure the temporal evolution of the  
65 biomass on a specific field, or to adjust the harvesting pressure according to biomass  
66 estimates at a given time. Biological ratios are often used in the literature to standardize  
67 biological data. Dry/wet biomass ratios, are generally used to estimate the relative water  
68 content in plants and to homogenize the parameters found in the literature (which may be  
69 expressed either in dry or wet biomass). Moreover, this ratio can be used by professional  
70 seaweed harvesters (or their employers) that are required, under French law, to report monthly  
71 the quantities of algae they have harvested, in fresh biomass.

72 Seaweeds are a polyphyletic group that displays a wide diversity of life cycles and  
73 morphologically diverse thalli with variable growth rates. Because seaweeds species are  
74 highly diverse, estimation of their biomass through allometric relationships is a challenging  
75 task. The overall objective of this study was to develop and evaluate allometric equations for  
76 estimating the biomass in the tree main groups of harvested seaweed (three red algal species  
77 (*Chondrus crispus*, *Mastocarpus stellatus*, *Palmaria palmata*), four brown algal species  
78 (*Fucus serratus*, *Fucus vesiculosus*, *Himanthalia elongata*, *Saccharina latissima*) and one  
79 green algal species (*Ulva* sp.). We also provide uniform calculated dry/wet biomass ratios to  
80 estimate the relative water content of seaweeds.

81

82

## 83 **Materials and methods**

84 Samples were collected in Brittany (Northern France) where more than 80% of macroalgae  
85 are harvested in France. We pooled datasets obtained across several years (2004 to 2015), in  
86 order to create sufficiently powered samples that are large enough to allow for meaningful  
87 analysis. An attempt was made to obtain samples representative of the full length range of  
88 each species. All datasets were obtained between March and November, the time when most  
89 of the biomass is extracted due to greater harvestable biomass and legal harvest period.

90 In this study, we measured individuals, as defined by Scrosati (2005). The whole thallus  
91 corresponding to all the fronds that arise from one holdfast was measured for clonal seaweeds  
92 (*Chondrus crispus*, *Mastocarpus stellatus*, *Palmaria palmata*) and the whole thallus  
93 corresponding to the only upright that arises from one holdfast was measured for unitary  
94 seaweeds (*Fucus serratus*, *F. vesiculosus*, *Himantalia elongata* and *Ulva* sp.). For each  
95 individual, the maximal length (L) and the dry biomass (DW), after drying at 60°C for 48h,  
96 were recorded. For some species, the maximal circumference (C), the maximal width (w) and  
97 the fresh biomass (FW) were also recorded, prior to the drying.

### 98 ***Length-biomass relationships***

99 Allometric length-biomass equations were obtained by regressing dry biomass on maximal  
100 length (L), maximal circumference (C), volume (LC<sup>2</sup>), or surface (Lw). We wrote the models  
101 using R to obtain both linear (Eq 1) and power law equation (Eq 2):

$$102 \quad DW = a \times X + b \quad \text{Eq 1}$$

$$103 \quad DW = a \times X^b \quad \text{Eq 2}$$

104 where DW = dry biomass (g), X = variable or combination of variables (L, C, LC<sup>2</sup>, Lw), and a  
105 and b are constants. Then we selected for each species the best model using the Akaike

106 information criterion (AIC) and the determination coefficient ( $R^2$ ). The best statistical model  
107 minimizes the value of AIC and maximizes the value of  $R^2$ . It is important to note that we also  
108 determined the length-biomass relationship of *C. crispus* and *M. stellatus* blended, because in  
109 the field they usually form a mixed canopy that could not be harvested separately. We also  
110 made a seasonal distinction for *H. elongata* by calculating the allometric equation for only  
111 individuals harvested from March to June on one side (i.e. the harvestable individuals truly  
112 harvested) and the allometric equation for all the individuals harvested between March and  
113 October on the other side. After June or July, large individuals are no more harvested because  
114 they are thick and grainy, thus less appealing for human consumption. Essentially, the first  
115 equation (March-June) should be used by professional seaweed harvesters while the second  
116 equation (March-October) could be better suited for scientist interest.

117 All statistical analyses were carried out with the R software package ([http://www.r-](http://www.r-project.org/)  
118 [project.org/](http://www.r-project.org/)).

### 119 ***Mean water content***

120 The mean water content of the algae was determined by weighing before and after drying. In  
121 order to quantify the relationship between fresh biomass and dry biomass, we used  
122 standardized major axis (SMA) regression (also referred to as reduced major axis regression).  
123 This method is more appropriate than least-squares regression for estimating the line of best  
124 fit for the relationship between two variables (Warton et al. 2006). The obtained fitted line  
125 does not change if the roles of “predictor” and “response” variables are switched; in contrast,  
126 ordinary least squares regression yields a different fitted line if the y-axis and x-axis are  
127 switched (Warton et al. 2006).

128

### 129 **Results and discussion**

130 *Development of allometric equations for estimating seaweed biomass*

131 Relationships between mass (expressed as dry biomass) and biometrics were established. We  
132 tested linear and power models for more than 350 individuals from 8 different species. For  
133 each model, we tested several explanatory variables: L, C, and  $LC^2$  for *C. crispus*, *P. palmata*  
134 *F. serratus*, and *F. vesiculosus*; L, w and Lw for *Ulva* sp.; and L for *H. elongata* and  
135 *S. latissima*. The 10 selected length-biomass relationships are shown in Fig. 1 and their  
136 respective parameters are given in Table 1. These inclusive relationships were all expressed as  
137 a power model. The best descriptive biometric variable varied according to the seaweed  
138 morphology. Species with a bushy morphology were best described by the volume ( $LC^2$ ),  
139 while long stringy species were best described by the length (L) and flat species by the  
140 surface (Lw). All the relationships of the seaweed species analysed in this paper were highly  
141 significant ( $0.77 < R^2 < 0.96$ ) and could consequently be reliably applied (Table 1). Besides,  
142 Gevaert et al. (2001) provided an allometric equation for the species *S. latissima* with a  
143 scaling exponent really close ( $b = 1.357$ ) to the one we calculated ( $b = 1.358$ ). Allometric  
144 equations ( $DW = a \times X^b$ ) were not found for any other species studied.

145 Nondestructive methods of seaweed biomass estimation have successfully been applied in the  
146 past. For example Scrosati and DeWreede (1997) have successfully applied nondestructive  
147 methods to estimate stand biomass in a biomass-density study that was based on the fronds  
148 and not on the individuals of one species (*Mazzaella cornucopiae*).

149 The two allometric equations obtained for *H. elongata* showed different allometric parameter  
150 values, with the scaling exponent (b) of harvestable individuals (March-June) being lower (-  
151 57%) than the one calculated with all individuals (March-October). This difference reveals an  
152 ontogenetic shift, partly because in late summer and autumn, individuals of *H. elongata* get

153 thicker which increases their biomass, become not consumable and so are no more harvested  
154 after June-July.

155 With the exception of *H. elongata*, seasonal variations were not completely taken into account  
156 (no sampling in winter), which may potentially cause a difference between the predicted DW  
157 and the observed DW at the individual scale, due to differences in tissue density (Åberg  
158 1990). However, as stated above, most of the seaweed harvesting occurs between March and  
159 November, which corresponds to the period when we made our samples. Also, we do believe  
160 than any potential biases should be reduced at the scale of the quadrat or seaweed field.  
161 Therefore, these tools can be applied to large populations and are relevant to provide accurate  
162 estimates of the standing biomass of a seaweed field, in a rapid and nondestructive way.

163

#### 164 *Development of ratios for estimating water content*

165 Relationships between DW and FW were expressed as a linear relation and were also highly  
166 significant ( $R^2 > 0.90$ ). They showed that mean water content ranged from 71.7% (*M.*  
167 *stellatus*) to 88.5% (*S. latissima*) (Table 2). While DW:FW ratios may vary depending on the  
168 season, our results are quite consistent with those found in the literature: Scrosati (2006)  
169 described a mean water content of 76.1% for *C. crispus*, 79.3% for *F. vesiculosus* and 87.6%  
170 for *S. latissima*; Gevaert et al. (2001) found a mean water content of 89% for *S. latissima*; and  
171 Alveal and Ponce (1997) estimated a mean water content of 72% for *M. stellatus*. Due to  
172 technical, commercial and infrastructural reasons, harvesters dry some harvested algae prior  
173 to weighing them, and then convert the dry biomass into fresh biomass with a ratio that is  
174 specific to each harvester or employer. These ratios are often confidential and may lead to  
175 over- or underestimate the quantities of algae that are actually harvested. Here we attempt to



176 provide uniform and rigorously calculated ratios that could be used by all the professional  
177 seaweed harvesters and their employers.

178 Global environment change coupled to the increased demand for seaweeds are likely to exert  
179 some significant pressure on the standing seaweed biomass. The relationships established in  
180 the study will provide a basis for future studies to estimate more easily and by a  
181 nondestructive way the biomass of seaweed populations.

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201 **References**

- 202 Åberg P (1990) Measuring size and choosing category size for a transition matrix study of the  
203 seaweed *Ascophyllum nodosum*. *Mar Ecol Prog Ser* 63:281–287
- 204 Airoidi L, Balata D, Beck MW (2008) The Gray Zone: Relationships between habitat loss and  
205 marine diversity and their applications in conservation. *J Exp Mar Biol Ecol* 366:8–15
- 206 Airoidi L, Beck MW (2007) Loss, status and trends for coastal marine habitats of Europe.  
207 *Oceanogr Mar Biol* 45:345–405
- 208 Alveal K, Ponce F (1997) Determination of water content in seaweeds. *Gayana Oceanol* 5:1–  
209 11
- 210 Benedetti-Cecchi L, Pannacciulli F, Bulleri F, et al (2001) Predicting the consequences of  
211 anthropogenic disturbance: large-scale effects of loss of canopy algae on rocky shores.  
212 *Mar Ecol Prog Ser* 214:137–150
- 213 Chave J, Andalo C, Brown S, et al (2005) Tree allometry and improved estimation of carbon  
214 stocks and balance in tropical forests. *Oecologia* 145:87–99
- 215 Connell S, Russell B, Turner D, et al (2008) Recovering a lost baseline: missing kelp forests  
216 from a metropolitan coast. *Mar Ecol Prog Ser* 360:63–72
- 217 Engel C, Åberg P, Gaggiotti OE, et al (2001) Population dynamics and stage structure in a  
218 haploid-diploid red seaweed, *Gracilaria gracilis*. *J Ecol* 89:436–450
- 219 Estes JA, Duggins DO, Rathbun GB (1989) The Ecology of Extinctions in Kelp Forest  
220 Communities. *Conserv Biol* 3:252–264
- 221 Gevaert F, Davoult D, Creach A, et al (2001) Carbon and nitrogen content of *Laminaria*  
222 *saccharina* in the eastern English Channel: biometrics and seasonal variations. *J Mar*  
223 *Biol Assoc UK* 81:727–734
- 224 Golléty C, Migné A, Davoult D (2008) Benthic metabolism on a sheltered rocky shore: role of  
225 the canopy in the carbon budget. *J Phycol* 44:1146–1153
- 226 Jonson JH, Freudenberger D (2011) Restore and sequester: estimating biomass in native  
227 Australian woodland ecosystems for their carbon-funded restoration. *Aust J Bot*  
228 59:640–653
- 229 Lindgren A, Bouza N, Åberg P, Sosa PA (1998) Spatial and temporal variation in distribution  
230 of *Gelidium canariensis* (Rhodophyta) from natural populations of the Canary Islands.  
231 *J Appl Phycol* 10:273–278
- 232 Mangialajo L, Chiantore M, Cattaneo-Vietti R (2008) Loss of furoid algae along a gradient of  
233 urbanisation, and structure of benthic assemblages. *Mar Ecol Prog Ser* 358:63–74
- 234 Mathieson AC, Guo Z (1992) Patterns of furoid reproductive biomass allocation. *Br Phycol J*  
235 27:271–292

- 236 Mccarthy MC, Enquist BJ (2007) Consistency between an allometric approach and optimal  
 237 partitioning theory in global patterns of plant biomass allocation. *Funct Ecol* 21:713–  
 238 720
- 239 Migné A, Delebecq G, Davoult D, Spilmont N, Menu D, Gevaert F (2015) Photosynthetic  
 240 activity and productivity of intertidal macroalgae: in situ measurements, from thallus  
 241 to community scale. *Aquat Bot* 123:6–12
- 242 Niklas KJ, Enquist BJ (2002) On the Vegetative Biomass Partitioning of Seed Plant Leaves,  
 243 Stems, and Roots. *Am Nat* 159:482–497
- 244 Paul KI, Roxburgh SH, England JR, Ritson P, Hobbs T, Brooksbank K, Raison RJ, Larmour  
 245 JS, Murphy S, Norris J, Neumann C, Lewis T, Jonson J, Carter JL, McArthur G,  
 246 Barton C, Rose B (2013) Development and testing of allometric equations for  
 247 estimating above-ground biomass of mixed-species environmental plantings. *For Ecol*  
 248 *Manag* 310:483–494
- 249 Perkol-Finkel S, Airoidi L (2010) Loss and Recovery Potential of Marine Habitats: An  
 250 Experimental Study of Factors Maintaining Resilience in Subtidal Algal Forests at the  
 251 Adriatic Sea. *PLoS ONE* 5:e10791.
- 252 Poorter H, Niklas KJ, Reich PB, Oleksyn J, Poot P and Mommer L (2012), Biomass  
 253 allocation to leaves, stems and roots: meta-analyses of interspecific variation and  
 254 environmental control. *New Phytol*, 193: 30–50
- 255 Sack L, Grubb PJ, Marañón T (2003) The functional morphology of juvenile plants tolerant of  
 256 strong summer drought in shaded forest understories in southern Spain. *Plant Ecol*  
 257 168:139–163
- 258 Santelices B, Hormazábal M, Correa J & Flores V (2004). The fate of overgrown germlings in  
 259 coalescing Rhodophyta. *Phycologia*43:346–52.
- 260 Scrosati R, DeWreede RE (1997) Dynamics of the biomass-density relationship and frond  
 261 biomass inequality for *Mazzaella cornucopiae* (Rhodophyta, Gigartinaceae): implications for  
 262 the understanding of frond interactions. *Phycol* 36: 506-516
- 263 Scrosati R (2005) Review of studies on biomass-density relationships (including self-thinning  
 264 lines) in seaweeds: main contributions and persisting misconceptions. *Phycol. Res.* 53:224–33
- 265 Scrosati R (2006) Length–biomass allometry in primary producers: predominantly  
 266 bidimensional seaweeds differ from the “universal” interspecific trend defined by  
 267 microalgae and vascular plants. *Can J Bot* 84:1159–1166
- 268 Tait LW, Schiel DR (2011) Dynamics of productivity in naturally structured macroalgal  
 269 assemblages: importance of canopy structure on light-use efficiency. *Mar Ecol Prog*  
 270 *Ser* 421:97–107
- 271 Vadas, Sr. RL, Wright WA, Beal BF (2004) Biomass and Productivity of Intertidal  
 272 Rockweeds (*Ascophyllum nodosum* LeJolis) in Cobscook Bay. *Northeast Nat* 11:123–  
 273 142


- 274 Vaz-Pinto F, Martínez B, Olabarria C, Arenas F (2014) Neighbourhood competition in  
275 coexisting species: The native *Cystoseira humilis* vs the invasive *Sargassum muticum*.  
276 J Exp Mar Biol Ecol 454:32–41
- 277 Warton DI, Wright IJ, Falster DS, Westoby M (2006) Bivariate line-fitting methods for  
278 allometry. Biol Rev 81:259–291
- 279 Watt CA, Scrosati RA (2013) Bioengineer effects on understory species richness, diversity,  
280 and composition change along an environmental stress gradient: experimental and  
281 mensurative evidence. Estuar Coast Mar Sci 123: 10-18

**Table 1** Length-biomass relationships of macroalgal species collected in Brittany (NW France). Power equation:  $DW = a \times X^b$ .

Species	Date of sampling	n	Mean total length (cm)	Total length range (cm)	Explanatory variable			
						a	b	R <sup>2</sup>
<i>Chondrus crispus</i> & <i>Mastocarpus stellatus</i>	April-May-July-Oct.	66	10.02	3.5-16	LC <sup>2</sup>	0.0034	0.8259	0.93
<i>Chondrus crispus</i>	May	35	8.70	3.5-13	LC <sup>2</sup>	0.0006	1.0318	0.95
<i>Mastocarpus stellatus</i>	April-July	31	11.52	9-16	LC <sup>2</sup>	0.0067	0.7493	0.93
<i>Fucus serratus</i>	April-Oct.	60	36.50	8-70	LC <sup>2</sup>	0.1763	0.5996	0.92
<i>Fucus vesiculosus</i>	Nov.	48	41.44	13-117	LC <sup>2</sup>	0.0188	0.8028	0.87
<i>Himantalia elongata</i>	March-June	65	79.58	8-232	L	0.0319	1.2878	0.77
<i>Himantalia elongata</i>	March-June-Aug.-Oct.	75	98.20	8-281	L	0.0005	2.2323	0.81
<i>Palmaria palmata</i>	July-Oct.	40	29.73	10-65	LC <sup>2</sup>	0.0006	1.4183	0.91
<i>Saccharina latissima</i>	April	30	97.90	22-214	L	0.0155	1.3587	0.95
<i>Ulva</i> sp.	Oct.	37	21.10	2-87	Lw	0.0077	0.8921	0.93

**Table 2** Mean water content of macroalgal species collected in Brittany (NW France).

Species	n	Mean water content	a	b	R <sup>2</sup>
<i>Chondrus crispus</i> & <i>Mastocarpus stellatus</i>	66	74.4%	0.257	-0.034	0.96
<i>Chondrus crispus</i>	35	77.4%	0.226	0.048	0.99
<i>Mastocarpus stellatus</i>	31	71.7%	0.284	-0.139	0.96
<i>Fucus serratus</i>	30	78.4%	0.216	0.694	0.99
<i>Himantalia elongata</i>	37	83.3%	0.167	-1.365	0.90
<i>Palmaria palmata</i>	40	87.3%	0.127	0.777	0.95
<i>Saccharina latissima</i>	30	88.5%	0.116	0.107	0.99



**Fig. 1** Relationships between dry biomass (g) and biometric variables (in cm, cm<sup>2</sup>, or cm<sup>3</sup>). For *Himanthalia elongata*, the plain line represents individuals harvested between March and June (i.e. the harvestable individuals truly harvested; round data points), while the dotted line also includes the older large individuals (cross-shaped data points) harvested in October.

