



HAL
open science

MP-AzeFlu is more effective than fluticasone propionate for the treatment of allergic rhinitis in children

W Berger, J Bousquet, A.T. Fox, J Just, A Muraro, A Nieto, E Valovirta, M Wickman, U Wahn

► To cite this version:

W Berger, J Bousquet, A.T. Fox, J Just, A Muraro, et al.. MP-AzeFlu is more effective than fluticasone propionate for the treatment of allergic rhinitis in children. *Allergy*, 2016, 10.1111/all.12903 . hal-01326275

HAL Id: hal-01326275

<https://hal.sorbonne-universite.fr/hal-01326275>

Submitted on 3 Jun 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

BRIEF COMMUNICATION

MP-AzeFlu is more effective than fluticasone propionate for the treatment of allergic rhinitis in children

W. Berger^{1,2}, J. Bousquet^{3,4,5,6}, A. T. Fox⁷, J. Just^{8,9}, A. Muraro¹⁰, A. Nieto¹¹, E. Valovirta^{12,13}, M. Wickman^{14,15} & U. Wahn¹⁶

¹Division of Basic Clinical Immunology, School of Medicine, University of California, Irvine; ²Allergy & Asthma Associates, Mission Viejo, CA, USA; ³University Hospital, Montpellier; ⁴MACVIA-LR, Contre les Maladies Chronique pour un Vieillessement Actif en Languedoc Roussillon, European Innovation Partnership on Active and Healthy Aging Reference Site, Montpellier; ⁵INSERM, VIMA: Ageing and Chronic Diseases, Epidemiological and Public Health Approaches, U1168 Paris; ⁶UVSQ UMR-S1168, Université Versailles St-Quentin-en-Yvelines, Versailles, France; ⁷King's College London, MRC & Asthma UK Centre in Allergic Mechanisms of Asthma, Division of Asthma, Allergy and Lung Biology, Guy's and St Thomas' NHS Foundation Trust, London, UK; ⁸Allergology Department, Centre de l'Asthme et des Allergies. Hôpital d'Enfants Armand-Trousseau (APHP); ⁹Sorbonne Universités, UPMC Univ Paris 06, UMR_S 1136, Institut Pierre Louis d'Epidémiologie et de Santé Publique, Equipe EPAR, Paris, France; ¹⁰Department of Mother and Child Health, Padua General University Hospital, Padua, Italy; ¹¹Pediatric Allergy and Pneumology Unit, Children's Hospital La Fe, Valencia, Spain; ¹²Department of Lung Disease & Clinical Immunology, University of Turku; ¹³Terveyystalo Allergy Center, Turku, Finland; ¹⁴National Institute of Environmental Medicine, Karolinska Institutet; ¹⁵Department of Pediatrics, Sachs' Children's Hospital, Stockholm, Sweden; ¹⁶Department for Pediatric Pneumology and Immunology, Charité Medical University, Berlin, Germany

To cite this article: Berger W, Bousquet J, Fox AT, Just J, Muraro A, Nieto A, Valovirta E, Wickman M, Wahn U. MP-AzeFlu is more effective than fluticasone propionate for the treatment of allergic rhinitis in children. *Allergy* 2016; DOI: 10.1111/all.12903.

Keywords

children; Dymista; fluticasone propionate; MP29-02.

Correspondence

William E. Berger, Allergy and Asthma Associates of Southern California, Mission Viejo, CA 92691, USA.

Tel.: +1 949 364 2900

Fax: +1 949 365 0117

E-mail: weberger@uci.edu

Accepted for publication 1 April 2016

DOI:10.1111/all.12903

Edited by: Wytske Fokkens

Abstract

The objective was to evaluate the efficacy of MP-AzeFlu (Dymista[®]) vs fluticasone propionate (FP), (both 1 spray/nostril bid), in children with allergic rhinitis (AR). MP-AzeFlu combines azelastine hydrochloride, FP and a novel formulation in a single spray. Children were randomized in a 3 : 1 ratio to MP-AzeFlu or FP in this open-label, 3-month study. Efficacy was assessed in children aged ≥ 6 to <12 years (MP-AzeFlu: $n = 264$; FP: $n = 89$), using a 4-point symptom severity rating scale from 0 to 3 (0 = no symptoms; 3 = severe symptoms). Over the 3-month period, MP-AzeFlu-treated children experienced significantly greater symptom relief than FP-treated children (Diff: -0.14 ; 95% CI: -0.28 , -0.01 ; $P = 0.04$), noted from the first day (particularly the first 7 days) and sustained for 90 days. More MP-AzeFlu children achieved symptom-free or mild symptom severity status, and did so up to 16 days faster than FP. MP-AzeFlu provides significantly greater, more rapid and clinically relevant symptom relief than FP in children with AR.

Intranasal corticosteroids (INS) are recommended for the treatment of children with allergic rhinitis (AR) (1). However, they provide insufficient symptom control for many. Considering that AR is associated with poor asthma control (2), is a predictor of wheezing onset in school-aged children (3) and poorer examination performance at school (4), it is important to get it under control. Unfortunately, AR is undiagnosed and undertreated in children (5).

MP-AzeFlu (Dymista[®], Meda, Solna, Sweden) comprises an intranasal antihistamine [azelastine hydrochloride (AZE)], an INS [fluticasone propionate (FP)] and a novel formulation in a single spray. Its efficacy and safety in adults and

adolescent AR patients are well established (6–10), providing twice the overall nasal and ocular symptom relief as an INS or intranasal H1-antihistamine, and more complete and rapid symptom control (6).

A lower treatment effect has been observed in paediatric allergy trials (11–13), possibly confounded by caregiver assessment (14, 15). The present study was primarily designed to assess the long-term safety of MP-AzeFlu (the results of which will be published in full elsewhere). Efficacy was assessed secondarily using a simple scoring system in an effort to minimize this confounder. The objective was to evaluate the efficacy of MP-AzeFlu compared to FP in children aged ≥ 6 to <12 years, with AR.

Methods

Protocol

This was a prospective, randomized, active-controlled, parallel-group, 3-month, open-label safety trial in children with AR carried out at 42 investigational sites in the USA (March–October 2013). Ethics approval was obtained (Chesapeake Research Review Inc., Columbia, MD, USA), and the study conducted in accordance with Good Clinical Practice.

Participants

Male and female children aged ≥ 4 to <12 years (at screening visit), with a history of AR and who may benefit from treatment with MP-AzeFlu, in the opinion of the investigator (based on medical history and physical examination), were included. These children were free of any disease or concomitant treatment that could have interfered with the interpretation of the study. Those with superficial or moderate nasal erosion, nasal mucosal ulceration or nasal septum perforation, who had nasal surgery within the last year, had significant pulmonary disease (excluding intermittent asthma), a hypersensitivity to AZE and/or FP or had used any investigation drug within 30 days prior to providing informed consent were excluded.

Planned interventions and timing

The study comprised a 2- to 30-day lead-in period, and a 3-month treatment period, with study visits on Days, 1, 15, 30, 60 and 90. On Day 1, eligible children were randomized in a 3 : 1 ratio to either MP-AzeFlu nasal spray (daily dose: AZE 548 μg , FP 200 μg) or FP nasal spray (Roxane; daily dose 200 μg), both administered as 1 spray/nostril bid.

Efficacy variables

Each child (or caregiver) completed a 24-h reflective total symptom score (TSS) evaluation, scored daily in the morning during the lead-in period and prior to the AM dose in the treatment period, in response to the question 'how were your allergy symptoms over the past 24 h?'.
Allergy symptoms were scored from 0 (none/absent: no symptoms present), 1 (mild: symptoms clearly present, but minimal awareness and easily tolerated), 2 (moderate: definite awareness of symptoms that were bothersome but tolerable) to 3 (severe: hard to tolerate and caused interference with activities of daily living and/or sleeping).

Safety variables

Adverse events, nasal examination findings and vital signs were assessed. These data will be published in full separately (16).

Statistical analyses

Because it was a safety study, no inferential efficacy analysis were predefined in the protocol. The development focus of the paediatric programme was in the age range of 6- to

11-year-olds with some exploration in 4- to 5-year-olds. The mean change in TSS from baseline to each clinic visit was analysed *post hoc* for all randomised children aged ≥ 6 to <12 years old who took the drug. These data were analysed with a mixed-model ANCOVA (baseline as covariate; age class, visit and treatment as fixed effects). Missing data were not replaced. Time to response was analysed by Kaplan–Meier estimates and log-rank tests. Response was defined as change in TSS from a baseline of 2 or 3 (i.e. moderate to severe) to a maximum of 1 (i.e. mild at most). For symptomatic children (i.e. baseline value ≥ 2), this is equivalent to a 50% reduction from baseline in AM + PM reflective total nasal symptom score (rTNSS) assessed in other studies (6, 7). Mean % of days with none to mild symptoms (i.e. TSS 0 or 1) was also calculated for both groups.

Results

Child disposition

A total of 405 children were randomized to MP-AzeFlu or FP treatment. A total of 51 children (aged ≤ 5 years) were excluded from the efficacy analysis, as per the statistical plan efficacy was assessed in those aged ≥ 6 to <12 years. One child was excluded from the FP group (did not receive medication), yielding $n = 264$ and $n = 89$ in the MP-AzeFlu and FP treatment groups, respectively. Time-to-response analysis was performed for symptomatic children (i.e. TSS above 2 at baseline; MP-AzeFlu: $n = 124$; FP: $n = 44$). Of these, 15 children (6%) did not complete the study in the MP-AzeFlu group ($n = 4$ AE, $n = 1$ treatment failure, $n = 1$ protocol violation, $n = 2$ noncompliance, $n = 3$ withdrawal, $n = 2$ lost to follow-up and $n = 2$ other) and eight children (9%) failed to complete from the FP group ($n = 3$ AE, $n = 1$ protocol violation, $n = 1$ withdrawal, $n = 2$ lost to follow-up, $n = 1$ other).

Child baseline and demographic information

Similar baseline characteristics were observed in the MP-AzeFlu and FP groups (Table 1). The proportion of children with concomitant asthma was 4.86% and 4.90% in the MP-AzeFlu and FP groups, respectively.

Reflective total symptom score

Children treated with MP-AzeFlu experienced a -0.68 pt reduction in TSS, significantly greater than that afforded by FP (-0.54 pt reduction; Diff: -0.14 ; 95% CI: $-0.28, -0.01$; $P = 0.0410$). The superiority of MP-AzeFlu was noted from the first day of assessment, particularly during the first 7 days of treatment, and sustained for 90 days. More children treated with MP-AzeFlu (eight of 10 children) achieved symptom-free or at most mild symptom severity in the first month of treatment, and did so up to 16 days faster than FP (Figure 1). More children treated with MP-AzeFlu experienced none or mild symptoms during the 3-month study period [73.4% (SD 28.8)] than those treated with FP [66.0% (SD 34.2)].

Table 1 Demographic and baseline characteristics in children aged ≥ 6 to <12 years

Characteristic	All patients		Symptomatic patients (baseline TSS ≥ 2)	
	MP-AzeFlu (<i>n</i> = 264)	FP (<i>n</i> = 89)	MP-AzeFlu (<i>n</i> = 124)	FP (<i>n</i> = 44)
Age, <i>n</i> (%)				
≥ 6 to <9 years	128 (48.5)	44 (49.4)	63 (50.8)	21 (47.7)
≥ 9 to <12 years	136 (51.5)	45 (50.6)	61 (49.2)	23 (52.3)
Gender, <i>n</i> (%)				
Male	158 (59.9)	46 (41.7)	81 (65.3)	20 (45.5)
Race, <i>n</i> (%)				
Black/African American	41 (15.5)	16 (18.0)	20 (16.1)	7 (15.9)
White	200 (75.8)	67 (75.3)	99 (79.8)	35 (79.5)
Other	23 (8.7)	6 (6.7)	5 (4.0)	2 (4.5)
TSS, mean (SD)	1.72 \pm 0.76	1.77 \pm 0.73	2.40 \pm 0.35	2.36 \pm 0.35
Range	0–3	0–3		

TSS, total symptom score; SD, standard deviation.

MP-AzeFlu: novel intranasal formulation of azelastine hydrochloride and fluticasone propionate (FP) in a single spray.

Discussion

MP-AzeFlu provided significantly better AR symptom relief than FP in children aged ≥ 6 to <12 years, the first time the efficacy of INS has been exceeded in this population. Effect size was similar to that seen in adult and adolescent SAR patients (6). In the current study, MP-AzeFlu induced a -0.68 point TSS reduction from baseline, a -0.14 point difference vs FP, corresponding to a -5.44 point reduction and -1.12 point difference, respectively, on the 24-point rTNSS scale used in the adult/adolescent SAR trials (6, 7). Furthermore, approximately eight of 10 children treated with MP-AzeFlu in the current study achieved symptom-free or at most mild symptom severity in the first month of treatment, and did so up to 16 days faster than FP; comparable to the proportion of adult and adolescent patients with perennial AR over the same time period (8), which should positively influence concordance with MP-

AzeFlu therapy. Approximately three quarters of children treated with MP-AzeFlu experienced no or only mild symptoms during the 3-month treatment period. Achieving AR symptom control (and rapidly) is particularly important in children due to the negative impact of uncontrolled disease on school performance (4) and on asthma control (2).

Previously, MP-AzeFlu has demonstrated statistical superiority over placebo in children (aged ≥ 6 to <12 years) with moderate/severe SAR (17, 18), and as the extent of child self-rating increased, so too did the treatment difference between MP-AzeFlu and placebo. However, efficacy was assessed using an endpoint designed for adolescent/adult populations (i.e. by rTNSS), in line with regulatory requirement, and could be rated by either children or caregivers. The present study was designed to minimize assessment effort and bias using a much simplified and child-friendly efficacy assessment tool. Using this simple 4-point




Figure 1 Time to achieve at most mild allergic rhinitis symptom severity in children with moderate-to-severe symptoms at baseline during the first month of treatment with either MP-AzeFlu (*n* = 124) or fluticasone propionate (FP; *n* = 44), both 1 spray/nostril bid, in children aged ≥ 6 to 12 years. Time to response was analysed by Kaplan–Meier estimates and log-rank tests.

rating scale significant superiority of MP-AzeFlu was detected vs an active treatment.

Limitations of the current study include the open-label design and that it was primarily a safety study, with no assessment of quality of life or inferential efficacy analysis predefined in the protocol. However, the applied statistical method used to assess efficacy is straightforward and standard. In addition, the low number of clinic visits and the simple method of efficacy assessment make this trial quite pragmatic in nature and representative of response achievable in real life. Use of the simple 4-point severity scoring system was sensitive enough to detect difference in efficacy between two active treatments. MACVIA ARIA recommends another simple tool, the visual analogue scale, to assess disease control and guide treatment decisions (19). These tools should be strongly considered for inclusion in AR trials in children.

In conclusion, MP-AzeFlu provides significantly greater, more complete and more rapid AR symptom control than FP in children (aged ≥ 6 –12 years) and has been granted

approval for use in this age group by the FDA. Efficacy assessment in AR trials in children might be improved by use of simple tools.

Acknowledgments

We thank Dr Ruth Murray (Ireland) for assistance in editing this manuscript.

Conflict of interest

All authors are members of Meda's Dymista Paediatric Advisory Board.

Authors' contributions

All authors have contributed to acquisition of data and/or interpretation of data, drafting of the article and revising it critically for important intellectual content and have provided final approval for this version to be published.

References

1. Brozek JL, Bousquet J, Baena-Cagnani CE, Bonini S, Canonica GW, Casale TB et al. Allergic Rhinitis and its Impact on Asthma (ARIA) guidelines: 2010 revision. *J Allergy Clin Immunol* 2010;**126**:466–476.
2. de Groot EP, Nijkamp A, Duiverman EJ, Brand PL. Allergic rhinitis is associated with poor asthma control in children with asthma. *Thorax* 2012;**67**:582–587.
3. RoCHAT MK, Illi S, Ege MJ, Lau S, Keil T, Wahn U et al. Allergic rhinitis as a predictor for wheezing onset in school-aged children. *J Allergy Clin Immunol* 2010;**126**:1170–1175.
4. Walker S, Khan-Wasti S, Fletcher M, Cullinan P, Harris J, Sheikh A. Seasonal allergic rhinitis is associated with a detrimental effect on examination performance in United Kingdom teenagers: case-control study. *J Allergy Clin Immunol* 2007;**120**:381–387.
5. Ruokonen M, Kaila M, Haataja R, Korppi M, Paasilta M. Allergic rhinitis in school-aged children with asthma – still under-diagnosed and under-treated? A retrospective study in a children's hospital. *Pediatr Allergy Immunol* 2010;**21**:e149–e154.
6. Meltzer E, Ratner P, Bachert C, Carr W, Berger W, Canonica GW et al. Clinically relevant effect of a new intranasal therapy (MP29-02) in allergic rhinitis assessed by responder analysis. *Int Arch Allergy Immunol* 2013;**161**:369–377.
7. Carr W, Bernstein J, Lieberman P, Meltzer E, Bachert C, Price D et al. A novel intranasal therapy of azelastine with fluticasone for the treatment of allergic rhinitis. *J Allergy Clin Immunol* 2012;**129**:1282–1289.
8. Price D, Shah S, Bhatia S, Bachert C, Berger W, Bousquet J et al. A new therapy (MP29-02) is effective for the long-term treatment of chronic rhinitis. *J Investig Allergol Clin Immunol* 2013;**23**:495–503.
9. Klimek L, Bachert C, Mosges R, Munzel U, Price D, Virchow JC et al. Effectiveness of MP29-02 for the treatment of allergic rhinitis in real-life: results from a noninterventional study. *Allergy Asthma Proc* 2015;**36**:40–47.
10. Berger WE, Shah S, Lieberman P, Hadley J, Price D, Munzel U et al. Long-term, randomized safety study of MP29-02 (a novel intranasal formulation of azelastine hydrochloride and fluticasone propionate in an advanced delivery system) in subjects with chronic rhinitis. *J Allergy Clin Immunol Pract* 2014;**2**:179–185.
11. Meltzer EO, Lee J, Tripathy I, Lim J, Ellsworth A, Philpot E. Efficacy and safety of once-daily fluticasone furoate nasal spray in children with seasonal allergic rhinitis treated for 2 wk. *Pediatr Allergy Immunol* 2009;**20**:279–286.
12. Storms WW, Segall N, Mansfield LE, Amar NJ, Kelley L, Ding Y et al. Efficacy and safety of beclomethasone dipropionate nasal aerosol in pediatric patients with seasonal allergic rhinitis. *Ann Allergy Asthma Immunol* 2013;**111**:408–414.
13. Maspero JF, Walters RD, Wu W, Philpot EE, Naclerio RM, Fokkens WJ. An integrated analysis of the efficacy of fluticasone furoate nasal spray on individual nasal and ocular symptoms of seasonal allergic rhinitis. *Allergy Asthma Proc* 2010;**31**:483–492.
14. Danell CS, Bergstrom A, Wahlgren CF, Hallner E, Bohme M, Kull I. Parents and school children reported symptoms and treatment of allergic disease differently. *J Clin Epidemiol* 2013;**66**:783–789.
15. Dahlen E, Almqvist C, Bergstrom A, Wettermark B, Kull I. Factors associated with concordance between parental-reported use and dispensed asthma drugs in adolescents: findings from the BAMSE birth cohort. *Pharmacoepidemiol Drug Saf* 2014;**23**:942–949.
16. Berger W, Ratner P, Soteres D. Randomized trial of the safety of MP29-02* compared with fluticasone propionate nasal spray in children aged =4 years to <12 years with allergic rhinitis. Presented at the Paediatric Allergy & Asthma Congress, Berlin, Germany 15-17 Oct 2015. Available at http://www.eaaci-paam2015.com/PAAM2015-Abstracts/abstracts/PAAM_2015_PP76.pdf
17. Berger W, Meltzer EO, Amar NJ, Muraro A, Wickman M, Just J et al. MP-AzeFlu for nasal and ocular symptom relief in children with seasonal allergic rhinitis: importance of patient symptom assessment. Accepted by European Academy of Allergy and Clinical Immunology Congress 2016.
18. Berger W, Meltzer EO, Amar NJ, Muraro A, Wickman M, Just J et al. MP-AzeFlu and time to clinically-meaningful response in the treatment of children with seasonal allergic rhinitis: importance of patient symptom assessment. Accepted by European Academy of Allergy and Clinical Immunology Congress 2016.
19. Bousquet J, Schunemann HJ, Fonseca J, Samolinski B, Bachert C, Canonica GW et al. MACVIA-ARIA Sentinel Network for allergic rhinitis (MASK-rhinitis): the new generation guideline implementation. *Allergy* 2015;**70**:1372–1392.