



HAL
open science

Pleural effusion in difficult weaning from mechanical ventilation

M Dres, Daniel Roux, Thong Tien Pham, Muriel Fartoukh, Jd Ricard, A Demoule

► **To cite this version:**

M Dres, Daniel Roux, Thong Tien Pham, Muriel Fartoukh, Jd Ricard, et al.. Pleural effusion in difficult weaning from mechanical ventilation. *Intensive Care Medicine Experimental*, 2015, 3 (Suppl 1), pp.A92. 10.1186/2197-425X-3-S1-A92 . hal-01332342

HAL Id: hal-01332342

<https://hal.sorbonne-universite.fr/hal-01332342>

Submitted on 15 Jun 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

POSTER PRESENTATION

Open Access

Pleural effusion in difficult weaning from mechanical ventilation

M Dres^{1,2*}, D Roux³, T Pham^{1,4}, M Fartoukh^{1,4}, JD Ricard³, A Demoule^{1,2}

From ESICM LIVES 2015

Berlin, Germany. 3-7 October 2015

Introduction

Pleural effusions (PE) are common in intensive care unit (ICU) patients, especially in patients under mechanically ventilation. Since PE can alter gas exchanges, one could hypothesize that PE could play a role in the outcome of weaning from mechanical ventilation.

However, no study has yet reported the incidence and characteristics of PE in the specific context of weaning failure.

Objectives

To describe the incidence of pleural effusion in patients who failed a first spontaneous breathing trial (SBT) and to describe the characteristics of these patients.

Methods

We conducted a prospective observational study in three medical ICU. All mechanically ventilated patients were screened daily. In patients who failed their first SBT, a pleural ultrasonography was performed and the presence of PE was qualitatively quantified according to a 4-steps classification: 0: no PE; 1: small PE; 2: moderate PE and 3: large PE. In addition the main clinical characteristics of patients were collected. For statistical analysis, patients with classes 0 and 1 were regrouped, as were patients with classes 2 and 3.

Results

From November 2014 to March 2015, 336 patients were screened and 56 (17%) failed their first SBT. Among them, 28 patients (50%) had no PE, 18 patients (32%) had small PE (left or right), 8 (14%) patients had moderate PE and 2 (4%) patients had large PE.

Patients with moderate and large PE stayed longer in ICU (19 ± 7 vs. 9 ± 7 days, $p < 0.01$) and had a longer duration of mechanical ventilation (14 ± 9 vs. 24 ± 12 days, $p < 0.01$) as compared with patients without PE and with small PE.

Patients with moderate or large PE and patients without PE or with small PE shared similar characteristics at admission (age: 66 ± 14 and 61 ± 15 and SAPS 2: 55 ± 18 vs. 53 ± 15 , $p > 0.05$).

Shock as the cause of initiation of mechanical ventilation was more frequent in patients with moderate and large PE as compared to patients without and small PE (40% vs. 4%, $p < 0.01$).

Only one patient with a large PE had a pleural evacuation but without clinical improvement.

Conclusions

Significant pleural effusion is often detected in difficult to wean patients (18%), and is associated with shock state as the reason for mechanical ventilation. The interest of pleural drainage in this population deserves further studies.

Authors' details

¹Sorbonne Universités, UPMC Université Paris 06, INSERM, UMRS 1158 Neurophysiologie Respiratoire Expérimentale et Clinique, Paris, France.

²Hopital Pitie Salpetriere, Assistance Publique Hôpitaux de Paris, Respiratory and Critical Care Department, Paris, France. ³Assistance Publique - Hôpitaux de Paris, Louis Mourier Hospital, Critical Care Department, Colombes, France.

⁴Assistance Publique - Hôpitaux de Paris, Tenon Hospital, Critical Care Department, Paris, France.

Published: 1 October 2015

doi:10.1186/2197-425X-3-S1-A92

Cite this article as: Dres et al.: Pleural effusion in difficult weaning from mechanical ventilation. *Intensive Care Medicine Experimental* 2015 **3**(Suppl 1):A92.

¹Sorbonne Universités, UPMC Université Paris 06, INSERM, UMRS 1158 Neurophysiologie Respiratoire Expérimentale et Clinique, Paris, France
Full list of author information is available at the end of the article