



# On some properties of $SU(3)$ fusion coefficients

Robert Coquereaux, Jean-Bernard Zuber

## ► To cite this version:

Robert Coquereaux, Jean-Bernard Zuber. On some properties of  $SU(3)$  fusion coefficients. Nuclear Physics B, 2016, Mathematical Foundations of Quantum Field Theory: A volume dedicated to the Memory of Raymond Stora., 912, pp.119-150. 10.1016/j.nuclphysb.2016.05.029 . hal-01333044

HAL Id: hal-01333044

<https://hal.sorbonne-universite.fr/hal-01333044>

Submitted on 16 Jun 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# On some properties of SU(3) fusion coefficients

Robert Coquereaux<sup>a,\*</sup>, Jean-Bernard Zuber<sup>b</sup>

<sup>a</sup> Aix Marseille Univ, Univ Toulon, CNRS, CPT, Marseille, France

<sup>b</sup> Laboratoire de Physique Théorique et des Hautes Énergies, CNRS UMR 7589, Université Pierre et Marie Curie, Sorbonne Universités, 4 Place Jussieu, 75252 Paris Cedex 05, France

Received 29 April 2016; accepted 31 May 2016

Editor: Hubert Saleur

## Abstract

Three aspects of the SU(3) fusion coefficients are revisited: the generating polynomials of fusion coefficients are written explicitly; some curious identities generalizing the classical Freudenthal–de Vries formula are derived; and the properties of the fusion coefficients under conjugation of one of the factors, previously analyzed in the classical case, are extended to the affine algebra  $\widehat{su}(3)$  at finite level.

© 2016 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP<sup>3</sup>.

## 1. Introduction

The aim of this paper is threefold:

- 1) Write generating functions for SU(3) fusion matrices and coefficients.
- 2) Use them to get general formulae for dimensions of spaces of essential paths on fusion graphs and derive identities generalizing the classical Freudenthal–de Vries formula.
- 3) Compare multiplicities for  $\lambda \otimes \mu$  and  $\lambda \otimes \bar{\mu}$  (provide a proof that was missing in our paper [1]).

Along the way we discuss several other properties of fusion coefficients that do not seem to have been discussed elsewhere (for instance in section 6).

\* Corresponding author.

E-mail address: [robert.coquereaux@gmail.com](mailto:robert.coquereaux@gmail.com) (R. Coquereaux).

<http://dx.doi.org/10.1016/j.nucphysb.2016.05.029>

0550-3213/© 2016 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP<sup>3</sup>.

But the purpose of this paper is also to make a modest homage to the memory of our distinguished colleague and great friend Raymond Stora. In scientific discussions Raymond was a listener second to none, with unsurpassable insight, critical sharpness and good humor. . . . We hope that he would have found entertaining the following mix of algebraic, geometric and group theoretical considerations, but surely, he would have stimulated us with witty comments and inspiring suggestions.

## 2. Notations

When dealing with  $\widehat{\mathfrak{su}}(3)_k$ , the  $\mathfrak{su}(3)$  affine algebra at level  $k$ , the highest weights (h.w.) will be restricted to the Weyl alcove of level  $k$ , i.e., all dominant weights of level  $\|\lambda\| := \lambda_1 + \lambda_2 \leq k$ , thus

$$\mathcal{P}^{(k)} := \{\lambda | \lambda_1 \geq 0, \lambda_2 \geq 0, \lambda_1 + \lambda_2 \leq k\}. \quad (1)$$

Fusion matrices describe the multiplication, denoted  $\lambda \star \mu$ , and sometimes  $\lambda \star_k \mu$  if needed, of the  $(k+1)(k+2)/2!$  simple objects (irreps) in the fusion category  $\mathcal{A}_k(\mathfrak{su}(3))$ . They are denoted  $N_{\lambda}$ , with  $\lambda = (\lambda_1, \lambda_2)$  in the Weyl alcove, and satisfy the following conjugation property:  $N_{\lambda}^T = N_{(\lambda_1, \lambda_2)}^T = N_{(\lambda_2, \lambda_1)} = N_{\tilde{\lambda}}$ , with  $\tilde{\lambda} = (\lambda_2, \lambda_1)$ .

In  $\mathrm{SU}(3)$  or  $\widehat{\mathfrak{su}}(3)_k$ , there is a  $\mathbb{Z}_3$  grading  $\tau$  (“triality”) on irreps, stemming from the fact that this discrete group is the center of  $\mathrm{SU}(3)$ . We set for the two fundamental weights  $\tau(N_{(1,0)}) = 1$ ,  $\tau(N_{(0,1)}) = 2 \bmod 3$  and more generally

$$\tau(\lambda) \equiv \tau(N_{(\lambda_1, \lambda_2)}) := \lambda_1 + 2\lambda_2 \bmod 3. \quad (2)$$

For tensor product or fusion, this triality is conserved (mod 3):  $N_{\lambda\mu}^{\nu} \neq 0$  only if

$$\tau(\lambda) + \tau(\mu) - \tau(\nu) = 0 \bmod 3. \quad (3)$$

## 3. Generating functions of fusion matrices

Generating series giving  $\mathrm{SU}(3)$  fusion coefficients have been discussed and obtained in [2, 3]. What we do here is to provide generating formulae for the fusion matrices themselves. This material is presumably known to many *affinociados* of fusion algebras, but has never appeared in print to the best of our knowledge, and we think it may be helpful to recall its essentials for the convenience of the reader.

In this section we write down formulae for the generating function of fusion matrices. As the finite set  $\mathcal{P}^{(k)}$  is closed under fusion, this generating function turns out to be a polynomial, see below (6). This property will be taken as granted in the following.

For reference and comparison, the same problem in the case of  $\mathrm{SU}(2)$  is well known [4].

For the affine algebra  $\widehat{\mathfrak{su}}(2)_k$ , consider the matrices obtained by the (Chebyshev) recursion formula

$$N_{\lambda+1} = G \cdot N_{\lambda} - N_{\lambda-1} \quad \lambda \in \mathbb{Z}, \quad (4)$$

with  $N_{(0)} = \mathbb{1}$  and  $G = N_{(1)}$ , the latter being taken as the adjacency matrix of the Dynkin diagram  $A_{k+1}$ . Notice that this implies  $N_{(-1)} = 0$ . The obtained sequence is periodic of period  $2h$  with  $h = k + 2$  (use Cayley–Hamilton for  $G$ ); actually the  $N_{\lambda}$  matrices obey a Weyl group symmetry:

$N_{\lambda+2h} = N_{\lambda}$ ,  $N_{\lambda+h} = -N_{h-2-\lambda}$ . This infinite set of matrices<sup>1</sup> can be obtained from the generating function  $\frac{\mathbb{1}}{\mathbb{1}-sG+s^2} = \sum_{\lambda=0}^{\infty} s^{\lambda} N_{\lambda}$ . When  $0 \leq \lambda \leq k$  (Weyl alcove), the  $N_{\lambda}$  matrices have non-negative integer entries and are the fusion matrices of the affine algebra  $\widehat{su}(2)_k$ . Introducing a term  $+s^h P$ , where  $P = N_{(k)}$  is an involution ( $P \cdot P = \mathbb{1}$ ), in the numerator of the previous generating function, has the effect of truncating the series stemming from the denominator to a finite polynomial, so that the generating function of fusion matrices at level  $k$  reads

$$X(s) = \sum_{\lambda=0}^k s^{\lambda} N_{\lambda} = \frac{\mathbb{1} + s^h P}{\mathbb{1} - sG + s^2}. \quad (5)$$

### 3.1. $SU(3)$ : definition of $X$ and recursion formulae

The generators of the  $\widehat{su}(3)_k$  fusion algebra are  $G = N_{(1,0)}$  and  $G^T = N_{(0,1)}$ , while again  $N_{(0,0)} = \mathbb{1}$ . We define the generating polynomial

$$X(s, t) = \sum_{\lambda=(\lambda_1, \lambda_2) \in \mathcal{P}^{(k)}} s^{\lambda_1} t^{\lambda_2} N_{(\lambda_1, \lambda_2)}. \quad (6)$$

It satisfies the conjugation property

$$X(s, t)^T = X(t, s).$$

We then write the recursion formulae stemming from fusion by  $G$  or  $G^T$

$$G \cdot N_{(\lambda_1, \lambda_2)} = N_{(\lambda_1+1, \lambda_2)} + N_{(\lambda_1-1, \lambda_2+1)} + N_{(\lambda_1, \lambda_2-1)}$$

where each of the three terms in the rhs is present only if respectively,  $\lambda_1 \leq k-1$ ,  $\lambda_1 \geq 1$  and  $\lambda_2 \leq k-1$ ,  $\lambda_2 \geq 1$ . We have a similar expression for  $G^T \cdot N_{(\lambda_1, \lambda_2)}$ . We finally translate the latter formulae into identities on  $X$

$$G \cdot X(s, t) = \frac{1}{s}(X(s, t) - \Lambda_2(t)) + \frac{s}{t}(X(s, t) - \Lambda_1(s)) + t(X(s, t) - \Lambda_3(s, t))$$

where

$$\Lambda_1(s) := X(s, 0), \quad \Lambda_2(t) := X(0, t), \quad \Lambda_3(s, t) := \sum_{\lambda_1+\lambda_2=k} s^{\lambda_1} t^{\lambda_2} N_{(\lambda_1, \lambda_2)}, \quad (7)$$

from which we obtain

$$(s^2 + t + st^2 - stG) \cdot X(s, t) = s^2 \Lambda_1(s) + t \Lambda_2(t) + st^2 \Lambda_3(s, t) \quad (8)$$

and likewise, from the fusion with  $G^T$ ,

$$(t^2 + s + s^2t - stG^T) \cdot X(s, t) = s \Lambda_1(s) + t^2 \Lambda_2(t) + s^2t \Lambda_3(s, t). \quad (9)$$

Subtracting equ. (9) multiplied by  $t$  from equ. (8) multiplied by  $s$ , we eliminate the unwanted  $\Lambda_3$  and get

$$(s^3 \mathbb{1} - s^2tG + st^2G^T - t^3 \mathbb{1}) \cdot X(s, t) = (s^3 - st) \Lambda_1(s) + (st - t^3) \Lambda_2(t). \quad (10)$$

<sup>1</sup> In the non-affine case, the multiplicities for the decomposition of tensor product of irreps are encoded by matrices obtained by taking the large  $k$  limit of the  $N_{\lambda}$  matrices.

Thus, if we can determine the boundary generating polynomials  $\Lambda_1(s) = X(s, 0)$  and  $\Lambda_2(t) = X(0, t)$  we find

$$X(s, t) = \frac{(s^3 - st)\Lambda_1(s) + (st - t^3)\Lambda_2(t)}{s^3\mathbb{1} - s^2tG + st^2G^T - t^3\mathbb{1}} \quad (11)$$

Here the inverse of  $s^3\mathbb{1} - s^2tG + st^2G^T - t^3\mathbb{1}$  may be understood in two alternative ways. Either as the actual matrix inverse, which exists for generic values<sup>2</sup> of  $s$  and  $t$ ; calculating that inverse requires the use of a computer but leads, for each value of the level  $k$ , to an explicit expression. Or as a formal series in, say,  $t/s$ , which actually truncates to a finite degree polynomial. Taking for example  $s \neq 0$  in equ. (10), we write  $(s^3\mathbb{1} - s^2tG + st^2G^T - t^3\mathbb{1}) = s^3(\mathbb{1} - \frac{t}{s}G + \frac{t^2}{s^2}G^T - \frac{t^3}{s^3}\mathbb{1})$ . The latter factor may then be inverted as a formal series<sup>3</sup>  $(\mathbb{1} - \frac{t}{s}G + \frac{t^2}{s^2}G^T - \frac{t^3}{s^3}\mathbb{1})^{-1} = \sum_{\ell} (\frac{t}{s})^{\ell} X_{\ell}$  and equ. (11) reads  $X(s, t) = \sum_{\ell} (\frac{t}{s})^{\ell} X_{\ell} \cdot ((1 - \frac{t}{s^2})X(s, 0) + (\frac{t}{s^2} - \frac{t^3}{s^3})X(0, t))$ . By consistency, this expression is such that the summation over  $\ell$  truncates to a finite order so as to make  $X(s, t)$  a polynomial in  $s$  and  $t$  of total degree  $k$ .

The two issues of determining the boundary terms will be addressed in the following subsections. Here, we just observe that because of the commutativity of the fusion algebra, we may then write the rational fraction in (11) without worrying about the order of terms.

### 3.2. Boundary generating polynomials

We may again write recursion formulae for the boundary generating polynomials  $\Lambda_1(s) = X(s, 0)$  and  $\Lambda_2(t) = X(0, t)$

$$G \cdot \Lambda_1(s) = \sum_{0 \leq \lambda_1 \leq k-1} s_1^{\lambda_1} N_{(\lambda_1+1, 0)} + \sum_{1 \leq \lambda_1 \leq k} s_1^{\lambda_1} N_{(\lambda_1-1, 1)} = \frac{1}{s}(\Lambda_1(s) - \mathbb{1}) + L(s)$$

where  $L(s) := \sum_{\lambda_1=1}^k s^{\lambda_1} N_{(\lambda_1-1, 1)}$ , and likewise

$$G^T \cdot \Lambda_1(s) = s(\Lambda_1(s) - N_{(k, 0)}s^k) + \frac{1}{s}L((s).$$

Eliminating  $L(s)$  between these two relations yields

$$(\mathbb{1} - sG + s^2G^T - s^3\mathbb{1}) \cdot \Lambda_1(s) = (\mathbb{1} - s^h N_{(k, 0)}) \quad (12)$$

where for  $SU(3)$ ,  $h = k + 3$ .

Call  $P = N_{(k, 0)}$ . This permutation matrix describes a rotation of angle  $2\pi/3$  around the center of the Weyl alcove, as

$$P \cdot N_{(\lambda_1 \lambda_2)} = N_{k-\lambda_1-\lambda_2, \lambda_1} \quad (13)$$

<sup>2</sup> For example, for  $s \neq 0$  and  $t$  small enough, the matrix is certainly invertible.

<sup>3</sup> Explicitly, we may write  $1/(\mathbb{1} - \mathcal{X}) = \sum_p \mathcal{X}^p$ , here  $\mathcal{X} = \frac{t}{s}G - \frac{t^2}{s^2}G^T + \frac{t^3}{s^3}\mathbb{1}$ , and we expand  $\mathcal{X}^p$  in terms of multinomial coefficients, hence  $\mathcal{X}^p = \sum_{\substack{q, r \geq 0 \\ q+r \leq p}} (-1)^r \frac{p!}{q!r!(p-q-r)!} (\frac{t}{s})^{q+2r+3(p-q-r)} G^q (G^T)^r$ ; identifying the term of degree  $\ell$  in  $t/s$  gives  $X_{\ell} = \sum_{\substack{p, q, r \geq 0 \\ q+2r+3(p-q-r) = \ell}} (-1)^r \frac{p!}{q!r!(p-q-r)!} (\frac{t}{s})^{q+2r+3(p-q-r)} G^q (G^T)^r$ , which may be restricted to  $\ell \leq k$ .

In particular, for all  $\lambda : 0 \leq \lambda \leq k$ , we have  $P \cdot N_{(\lambda, k-\lambda)} = N_{(0, \lambda)}$ , and for  $\lambda = k$  we have  $N_{(0, k)} = P^2$ . As  $P^3 = \mathbb{1}$  we notice that  $\{\mathbb{1}, P, P^2\}$  is isomorphic with the  $\mathbb{Z}_3$  group. Note that this action is *not* related to the  $\mathbb{Z}_3$  grading (“triality”) on irreps described in section 2.

Now, it is clear that the matrix  $(\mathbb{1} - sG + s^2G^T - s^3\mathbb{1})$  may be inverted as a formal series in  $s$ , thus giving the generating function  $\Lambda_1(s) = X(s, 0)$

$$\Lambda_1(s) = X(s, 0) = \frac{\mathbb{1} - s^h P}{\mathbb{1} - sG + s^2G^T - s^3\mathbb{1}}. \quad (14)$$

The effect of the numerator  $\mathbb{1} - s^h P$  is to truncate these generating series to a polynomial of degree  $k$ . In a similar way, we have

$$\Lambda_2(t) = X(0, t) = \frac{\mathbb{1} - t^h P^2}{\mathbb{1} - tG^T + t^2G - t^3\mathbb{1}}, \quad (15)$$

which satisfies  $\Lambda_2(t) = (\Lambda_1(t))^T$  as anticipated, but also, using (13),

$$\Lambda_2(t) = t^k P^2 \Lambda_1\left(\frac{1}{t}\right). \quad (16)$$

But we may also now determine the third boundary matrix, namely  $\Lambda_3(s, t)$  defined in (7). From (13), it follows that

$$\Lambda_3(s, t) = \sum_{\lambda_1 + \lambda_2 = k} s^{\lambda_1} t^{\lambda_2} N_{(\lambda_1, \lambda_2)} = \sum_{0 \leq \lambda_2 \leq k} s^k \left(\frac{t}{s}\right)^{\lambda_2} P \cdot N_{(\lambda_2, 0)} = s^k P \cdot X\left(\frac{t}{s}, 0\right)$$

which, according to what we already know about  $X(s, 0) = \Lambda_1(s)$ , is indeed a polynomial in  $s$  and  $t$  of total degree  $\leq k$ .

### 3.3. The $X$ generating polynomial

We return to equ. (11), into which the expressions (14) and (15) for  $\Lambda_1(s)$  and  $\Lambda_2(t)$  may now be inserted. After some algebra, the result may be recast in the following form

$$\begin{aligned} X(s, t) = & \frac{(1 - st)\mathbb{1}}{(\mathbb{1} - sG + s^2G^T - s^3\mathbb{1})(\mathbb{1} - tG^T + t^2G - t^3\mathbb{1})} \\ & + \frac{s^h(st - s^3)P}{(\mathbb{1} - sG + s^2G^T - s^3\mathbb{1})(s^3\mathbb{1} - s^2tG + st^2G^T - t^3\mathbb{1})} \\ & + \frac{t^h(st - t^3)P^2}{(\mathbb{1} - tG^T + t^2G - t^3\mathbb{1})(t^3\mathbb{1} - st^2G^T + s^2tG - s^3\mathbb{1})}, \end{aligned} \quad (17)$$

which is clearly symmetric under conjugation (interchange  $P$  and  $P^2$ ,  $s$  and  $t$ ,  $G$  and  $G^T$ ).

Taking the level  $k$ , hence  $h$ , arbitrarily large in (17), i.e., dropping the last two terms and taking for  $G$  the (infinite size) adjacency matrix of the set of  $SU(3)$  dominant weights, gives

$$X_\infty(s, t) = \frac{(1 - st)\mathbb{1}}{(\mathbb{1} - sG + s^2G^T - s^3\mathbb{1})(\mathbb{1} - tG^T + t^2G - t^3\mathbb{1})} \quad (18)$$

which is the generating function (now an infinite series !) for the tensor product of  $SU(3)$  irreps.

There are alternative, different looking, expressions for the generating polynomial  $X(s, t)$ . Adding (rather than subtracting as we did in order to obtain equ. (11)) equ. (9) multiplied by  $t$  to equ. (8) multiplied by  $s$  gives

$$\begin{aligned}
 & (s^3 + t^3 + 2st + 2s^2t^2)\mathbb{1} - st(sG + tG^T)) \cdot X(s, t) \\
 & = (s^3 + st)\Lambda_1(s) + (t^3 + st)\Lambda_2(t) + 2s^2t^2\Lambda_3(s, t).
 \end{aligned} \tag{19}$$

There,  $X(s, t)$  appears as the inverse of the matrix  $(s^3 + t^3 + 2st + 2s^2t^2)\mathbb{1} - st(sG + tG^T)$  times the rhs, the solution of which being given by (17). Note that evaluated at  $s = t = 1$ , the former matrix reads  $K^{-1} = (6\mathbb{1} - (G + G^T))$ , generalizing to  $SU(3)$  what would be  $(2\mathbb{1} - G)$  for  $\widehat{su}(2)_k$ , namely the Cartan matrix of the  $A_{k+1}$  algebra, and the last relation reads  $X = K \Lambda$ , with  $X = X(1, 1)$  and where  $\Lambda = 2(\Lambda_1 + \Lambda_2 + \Lambda_3)$  is the boundary matrix evaluated at  $s = t = 1$ .

#### 4. Generalized Freudenthal–de Vries formulae and dimensions of spaces of paths

From a fusion algebra point of view, all Dynkin diagrams are manifestations of  $SU(2)$ , as they can be introduced as graphs encoding the action of the fusion ring of the affine  $\widehat{su}(2)$  on appropriate modules (nimreps). Strictly speaking this is only true for simply laced diagrams but there is a way to accommodate the non-ADE's in the same common framework. From this point of view, the classical Freudenthal–de Vries formulae, that hold for all simple Lie groups, are also a manifestation of the underlying  $SU(2)$  theory. In turn, those formulae are related to the counting of a particular kind of paths on appropriate Dynkin diagrams. Moving from an underlying  $SU(2)$  to an underlying  $SU(3)$  framework leads to several intriguing formulae that we present in this section.

##### 4.1. $SU(2)$ : dimensions of spaces of essential paths on Dynkin diagrams

Call  $A$  the Cartan matrix associated with some chosen Dynkin diagram of rank  $r$  (the number of vertices), and  $\langle \cdot, \cdot \rangle$  the corresponding inner product in the space of roots. Call  $s$  the vector of scaling coefficients, with components defined as  $\langle \alpha, \alpha \rangle / 2$  where  $\alpha$  runs in a basis of simple roots, the squared norm being 2 if  $\alpha$  is a long root. Call also  $\mathbf{s}$  the corresponding diagonal matrix. It is the identity matrix if the Dynkin diagram is simply laced (ADE cases). Call  $K = A^{-1}\mathbf{s}$  the quadratic form matrix (it gives the scalar products between fundamental weights). Call  $g$  the Coxeter number,  $g^\vee$  the dual Coxeter number (they are equal for simply laced cases) and  $k = g - 2$  the  $(SU(2))$  level to be used below. Call  $\rho$  the Weyl vector (by definition its components are all equal to 1 on the basis of fundamental weights). Its squared norm is then  $\langle \rho, \rho \rangle = \sum_{a,b} K_{ab}$  and it is given by the Freudenthal - de Vries formula :  $\langle \rho, \rho \rangle = g^\vee(g + 1)r/12$ .

Call  $F_\lambda = (F_\lambda)_{a,b}$  the  $r \times r$  matrices defined by the  $SU(2)$  Chebyshev recurrence relation (4), with  $G = 2 - A$ . In a conformal field theory context, and for the ADE cases, they are called the “nimrep” matrices, (for *non-negative integer valued matrix representation* of the fusion algebra) and they describe boundary conditions.

Call  $X = \sum_\lambda F_\lambda$ , the path matrix, and  $u$ , with components<sup>4</sup>  $u_a = \sum_b (X^T)_{b,a} = \sum_b (X)_{a,b}$  the height vector (as defined by Dynkin [5], see also ref. [4]). In this general setup it is also useful to define the vector  $v$ , with components<sup>5</sup>  $v_a = \sum_b (X)_{b,a}$ .

Given  $\lambda \in \{0, 1, \dots, g - 2\}$  and two vertices  $a$  and  $b$  in the chosen Dynkin diagram, the vector space of dimension  $(F_\lambda)_{a,b}$  is called the space of essential paths of length  $\lambda - 1$  from the vertex  $a$  to the vertex  $b$ ; for the purpose of the present paper we don't need to explain how these spaces are realized in terms of actual paths – see [6]. With a slight terminological abuse (dimension

<sup>4</sup> These components are also equal to twice the components of the dual Weyl vector on the basis of simple coroots.

<sup>5</sup> These components are also equal to twice the components of the Weyl vector on the basis of simple roots.

versus cardinality) we may say that there are  $X_{a,b}$  essential paths (of any length) from  $a$  to  $b$  and that the component  $v_b$  of  $v$  gives the total number of essential paths (of arbitrary origin and length) reaching  $b$ . Then,  $d_\lambda = \sum_{a,b} (F_\lambda)_{a,b}$  gives the dimension of the space of essential paths of length  $\lambda$ , and  $d_H = \sum_{\lambda \in \{0,1,\dots,g-2\}} d_\lambda$  is the total dimension of the space of essential paths, but the reader should only take these two equalities as mere definitions for the integers  $d_\lambda$  and  $d_H$ . Now, obviously,  $d_H = \sum_a u_a = \sum_a v_a = \sum_{a,b} X_{a,b}$ . It is also of interest to consider the integer  $d_B = \sum_\lambda d_\lambda^2$  that can be interpreted (at least in the ADE cases, since simply laced Dynkin diagrams classify module-categories over  $\mathcal{A}_k(\mathrm{SU}(2))$ , [7,8]) as the dimension of a weak Hopf algebra [9].

A last relation of interest, for us, relates the path matrix  $X$  to the quadratic form matrix  $K$ . It can be established as we did in the  $\mathrm{SU}(3)$  case (see the end of section 3.3). For an arbitrary Dynkin diagram, it reads again:  $X^T = K s^{-1} \Lambda$ , where  $\Lambda$  is the “boundary matrix”  $\Lambda = F_0 + F_{r-1}$ .

If the diagram is simply laced then  $g^\vee = g$ ,  $s$  is the identity, and  $A$ ,  $K$ ,  $X$  are symmetric. For those cases the former relation reads  $X = K \Lambda$ , but  $F_0$  is the identity and a detailed analysis of all cases shows that  $F_{r-1}$  is a permutation matrix, so that  $\sum_{a,b} X_{ab} = 2 \sum_{a,b} K_{a,b}$ . The Freudenthal–de Vries formula then implies<sup>6,7</sup>

$$d_H = \sum_{a,b} X_{a,b} = 2 \sum_{a,b} K_{a,b} = g(g+1)r/6 \quad (\text{ADE cases}). \quad (20)$$

One may notice that the Freudenthal - de Vries formula (giving  $\langle \rho, \rho \rangle$ ) and the expression giving the dimension  $d_H$  of the space of essential paths only differ by a factor 2. In particular, if the diagram is  $A_r$ , which in particular encodes fusion by the fundamental representation in fusion categories of type  $\mathcal{A}_k(\mathrm{SU}(2))$ , then  $g = r + 1 = k + 2$  and

$$d_H = (k+1)(k+2)(k+3)/6 \quad (A_{k+1} \text{ cases}). \quad (21)$$

#### 4.2. Generalization to $\mathrm{SU}(3)$

The purpose of this section is to show how the above results generalize in the case of graphs (McKay graphs) describing fusion by the fundamental irreps in the case  $\mathcal{A}_k(\mathrm{SU}(3))$ . We will show that the following two formulae hold:

$$\sum_{a,b} K_{a,b} = \frac{1}{2} \frac{(k+1)(k+2)}{2} \frac{(k+4)(k+5)}{60} \quad (22)$$

$$d_H = \sum_{a,b} X_{a,b} = \frac{(k+1)(k+2)(k+3)(k+4)(k+5)((k+3)^2 + 5)}{1680} \quad (23)$$

These formulae were already announced in [11] and [12] where one can find tables containing several other “characteristic numbers” describing the geometry of graphs related to fusion categories of type  $\mathrm{SU}(2)$ , i.e., simply laced Dynkin diagrams, and of type  $\mathrm{SU}(3)$ . However, for these two formulae, no proof was given. The one that we shall give below relies on a crucial property (a theorem that we recall below in sect. 5.1.1) that was actually obtained much later [13].

<sup>6</sup> In the general case (non-necessarily ADE) one would obtain  $d_H = 2 \sum_{a,b} A_{a,b}^{-1}$ .

<sup>7</sup> This number also gives the dimension of the vector space underlying the Gelfand–Ponomarev preprojective algebra associated with the corresponding unoriented quiver [10].



The sum  $d_H$  of matrix elements of  $X$ , for  $SU(3)$  at level  $k$ , equal to the sum of all multiplicities for all possible fusion products up to level  $k$ , can again be interpreted as counting essential “paths” although the “length” is no longer a non-negative integer (i.e., an irrep of  $SU(2)$ ) but a weight of  $SU(3)$  (i.e., a pair of non-negative integers) belonging to the Weyl alcove. The notion of path should therefore be generalized in a way that is appropriate, but we do not intend to enter this discussion and shall stay at the level of combinatorics.

#### 4.3. Proof of the relation (23)

To ease the writing, we introduce, for an arbitrary square matrix  $M$ , the notation  $\Sigma M$  that denotes the sum of its matrix elements, thus  $\Sigma M = \text{Tr} MU$ , where  $U$  is the square matrix of same dimensions that has all its coefficients equal to 1.

$X(s, t)$  being the  $SU(3)$  generating functional introduced in section 3.1, we call  $X = X(1, 1)$ . From the definition of  $U$ , we have  $d_H = \Sigma X = \text{Tr}(XU)$ . One important step of the proof is to show that  $\text{Tr}(G - G^T)X(s, t)U = 0$  for arbitrary values of  $s$  and  $t$ . This property is obvious for  $s = t$  as  $X(s, s)$  is symmetric, but in general we have only  $X(s, t)^T = X(t, s)$ .

**Lemma 1.** *One has  $UX(s, t) = UX(t, s)$ . There is no transpose sign here.*

**Proof.** The matrix element  $(i, j)$  of  $UX(s, t)$ , which is obviously independent of  $i$  is the sum of all matrix elements of the column  $j$  of  $X(s, t)$ . The matrix element  $(i, j)$  of  $UX(t, s)$ , is the sum of all matrix elements of the column  $j$  of  $X(t, s) = X(s, t)^T$ . Since  $X(s, t)^T = \sum_{p,q} N_{(p,q)}^T s^p t^q$ , the equality of these two matrix elements results from the Theorem 1 proved in [13] and recalled below.

**Lemma 2.** *One has  $\text{Tr}(G^T X(s, t)U) = \text{Tr}(GX(s, t)U)$ .*

**Proof.** Using the previous lemma, and the fact that  $G$  commutes with all fusion matrices, and therefore also with  $X(s, t)$ , one gets:  $\text{Tr}(G^T X(s, t)U) = \text{Tr}(U^T X(s, t)^T G) = \text{Tr}(UX(t, s)G) = \text{Tr}(UX(s, t)G) = \text{Tr}(UGX(s, t)) = \text{Tr}(GX(s, t)U)$ . In contrast, the trace of  $(G^T - G)X(s, t)$  has no reason to vanish.

The previous lemma implies:

$$\begin{aligned} \text{Tr}((sG - tG^T)X(s, t)U) &= (s - t) \text{Tr}(GX(s, t)U) \quad \text{and} \\ \text{Tr}((sG + tG^T)X(s, t)U) &= (s + t) \text{Tr}(GX(s, t)U) \end{aligned}$$

Multiplying both equations (19) and (11) on both sides by  $U$ , using the previous lemma, calling  $\Sigma X(s, t) = \text{Tr}(X(s, t)U)$ ,  $\Sigma \Lambda_1(s) = \text{Tr}(\Lambda_1(s)U)$ ,  $\Sigma(GX(s, t)) = \text{Tr}(GX(s, t)U)$ , and taking the trace gives:

$$\begin{aligned} (s^3 + t^3 + 2st + 2s^2t^2)\Sigma X(s, t) - st(s + t)\Sigma(GX(s, t)) \\ = (s^3 + st)\Sigma \Lambda_1(s) + (t^3 + st)t^k \Sigma \Lambda_1(1/t) + 2s^2t^2s^k \Sigma \Lambda_1(t/s) \\ ((s^3 - t^3)\Sigma X(s, t) - st(s - t)\Sigma(GX(s, t))) = (s^3 - st)\Sigma \Lambda_1(s) - (t^3 - st)t^k \Sigma \Lambda_1(1/t) \end{aligned} \quad (24)$$

where we have used the fact that in that calculation  $P$  can be replaced by  $\mathbb{1}$  since  $P \cdot U = U \cdot P = U$ , so  $\text{Tr}(P^2 \Lambda_1(1/t)U) = \text{Tr}(\Lambda_1(1/t)U)$  and  $\text{Tr}(P \Lambda_1(t/s)) = \text{Tr}(\Lambda_1(t/s)U)$ . As it is

easy enough to determine separately the value of  $\Sigma \Lambda_1(s)$  (see below), the above (24) is a system of two linear equations for the two unknown  $\Sigma X(s, t)$  and  $\Sigma(GX(s, t))$ . One finds:

$$\begin{aligned}\Sigma X(s, t) &= (\Sigma \Lambda_1(s)(1-s)s + \Sigma \Lambda_1(1/s)s^{(k+1)}t(s-t) + \Sigma \Lambda_1(1/t)(t^{(k+1)})(t-1)) \\ &\quad /((-1+s)(s-t)(-1+t)) \\ \Sigma(GX(s, t)) &= (\Sigma \Lambda_1(s)(1-s^3) + \Sigma \Lambda_1(1/t)t^k(-1+t^3) + \Sigma \Lambda_1(t/s)s^k(s^3-t^3)) \\ &\quad /((-1+s)(s-t)(-1+t)).\end{aligned}\quad (*)$$

It is known, and in any case it is easy to show, that the sum of matrix elements of  $N_{(p,0)}$  is a product of two triangular numbers:  $(k+2-p)(k+1-p)(1+p)(2+p)/4$ . The common value of  $\Sigma \Lambda_1 = \Sigma \Lambda_1(1)$  and  $\Sigma \Lambda_i = \text{Tr}(\Lambda_i[1].U)$  for  $i = 1, 2, 3$  is obtained by summing the previous quantity along an edge:  $\Sigma \Lambda_1 = \text{Tr}(\Lambda_1(1).U) = \sum_{p=0}^k (k+2-p)(k+1-p)(1+p)(2+p)/4 = \binom{k+5}{5}$ . Unfortunately, this last equality is of little help since one cannot solve the previous system (24) while setting  $s = t = 1$ . So, we plug the value of the polynomial  $\Sigma \Lambda_1(s) = \sum_{p=0}^k (k+2-p)(k+1-p)(1+p)(2+p)/4 s^p$  in the above solutions (\*) for  $\Sigma X(s, t)$  and  $\Sigma(GX(s, t))$ , and calculate the first term of their Taylor series around  $s = t = 1$ . Calling  $X = X(1, 1)$  and  $GX = GX(1, 1)$  one finally gets:

$$\begin{aligned}\Sigma X &= ((k+1)(k+2)(k+3)(k+4)(k+5)((k+3)^2+5))/1680 \\ \Sigma(GX) &= (k+6)!/(k-1)!/560\end{aligned}$$

hence the result (23).

As a side result, we also obtain the value of  $\Sigma \Lambda$ . Indeed,

$$\text{Tr}(6K\Lambda U - GK\Lambda.U - G^T K\Lambda U) = \text{Tr}((6\mathbb{1} - G - G^T)K\Lambda U) = \text{Tr}(AK\Lambda U) = \text{Tr}(\Lambda U)$$

Evaluating the sum  $6\Sigma X - \Sigma(GX) - \Sigma(G^T X)$  using the two previous formulae gives immediately

$$\Sigma \Lambda = (k+1)(k+2)(k+3)(k+4)(k+5)/20. \quad (25)$$

#### 4.4. Proof of the relation (22)

Let  $M$  be an element of the matrix algebra generated by the commuting family of the fusion matrices  $F_\lambda$ . By Verlinde formula [4], all these elements (in particular  $X, A, K, \Lambda, \dots$ ) are diagonalized by the (symmetric) modular matrix  $S$ . We call  $\Delta_M$  the diagonal matrix of eigenvalues of  $M$ , i.e.,  $\Delta_M = SM S^{-1}$ .

**Lemma 3.** *The element  $(i, j)$  of  $SUS^{-1}$  vanishes whenever  $i$  and  $j$  do not both label real representations.*

**Proof.** By definition of  $U$ , the element  $(i, j)$  of  $SUS^{-1}$  is equal to  $\sum_x S_{ix} \sum_y S_{yj}^{-1}$  but  $S^{-1} = S^3 = CS$ , where  $C$  is the conjugation matrix, so that this element is  $\sum_x S_{ix} \sum_y S_{yj} S_{\bar{y}j} = (\sum_x S_{ix})(\sum_y S_{j\bar{y}})$ . Because of theorem 3 of ref. [13], the sum  $\sum_x S_{ix}$  vanishes if  $x$  is not of real type, hence the result. In the present case of  $SU(3)$ , real irreps have highest weight  $\{\mu, \mu\}$ , with  $\mu \in \{0, \dots, [k/2]\}$ .

The  $S$ -matrix elements for those real irreps read [14,4]

$$S_{\{\lambda_1, \lambda_2\}, \{\mu, \mu\}} = \text{const.} \left( \sin \frac{2\pi(\mu+1)(\lambda_1 + \lambda_2 + 2)}{k+3} - \sin \frac{2\pi(\mu+1)(\lambda_1 + 1)}{k+3} - \sin \frac{2\pi(\mu+1)(\lambda_2 + 1)}{k+3} \right)$$

where the  $(\lambda, \mu)$ -independent constant is of no concern to us. A straightforward but tedious calculation leads then to the corresponding eigenvalue of  $\Lambda_1$

$$\begin{aligned} \varpi_\mu(\Lambda_1) &= \sum_{0 \leq \lambda_1 \leq k} \frac{S_{\{\lambda_1, 0\}, \{\mu, \mu\}}}{S_{\{0, 0\}, \{\mu, \mu\}}} \\ &= \frac{(k+3)}{4 \sin^2(\pi(\mu+1)/(k+3))} = \frac{2(k+3)}{6 - 2(1 + 2 \cos(2\pi(\mu+1)/(k+3)))} \end{aligned}$$

Since  $P$  acts as a permutation, and  $P^3 = 1$ , we get immediately the corresponding eigenvalue of  $\Lambda = 2(\Lambda_1 + \Lambda_2 + \Lambda_3) = 2(1 + P + P^2)\Lambda_1$ , namely  $\varpi_\mu(\Lambda) = 6 \varpi_\mu(\Lambda_1)$ .

By the same token, the corresponding eigenvalue of  $A = 6 - (G + G^T)$  is

$$\varpi_\mu(A) = 6 - 2 \frac{S_{\{1, 0\}, \{\mu, \mu\}}}{S_{\{0, 0\}, \{\mu, \mu\}}} = 6 - 2(1 + 2 \cos(2\pi(\mu+1)/(k+3)))$$

and we conclude that the “real” eigenvalues of  $K = A^{-1}$  are proportional to those of  $\Lambda$

$$12(k+3)\varpi_\mu(K) = \varpi_\mu(\Lambda) = 12(k+3) \times \frac{1}{6 - 2(1 + 2 \cos(\frac{2\pi(\mu+1)}{k+3}))}.$$

By Lemma 3, in the calculation of  $SA\Lambda US^{-1} = \Delta_A \Delta_\Lambda SUS^{-1}$ , only real irreps contribute and we have found that  $SAS^{-1}S\Lambda S^{-1}SUS^{-1} = 12(k+3) \times SUS^{-1}$ , i.e.,  $\Lambda U = 12(k+3)KU$ . It follows that  $\Sigma K = \frac{1}{12(k+3)} \Sigma \Lambda$ , thus establishing (22).

Note that in the  $SU(2)$  case, i.e., for Dynkin diagrams, we have seen that the proportionality factor  $\Sigma X / \Sigma K$  was just equal to 2. Equations (22) and (23) illustrates a non-trivial effect of the boundary terms in this  $SU(3)$  situation since the proportionality factor is now a  $k$ -dependent polynomial.

#### 4.5. The non-diagonal cases

Notice that formula (22), using  $h = k + 3$  (generalized Coxeter number, sometimes called “altitude”), and  $r$ , the number of irreps (also the number of vertices of the corresponding fusion graph), can be expressed as  $\Sigma K = \Sigma_K^?$ , with as above  $\Sigma K = \sum_{a,b} K_{a,b}$  and

$$\Sigma_K^? = \frac{1}{2} r \frac{(h+1)(h+2)}{60} \quad (26)$$

Written this way, the r.h.s. of equ. (26), thought as a putative generalisation of Freudenthal–De Vries formula, works for the  $\mathcal{A}_k$  series, as it was proved above, and also for<sup>8</sup>  $\mathcal{E}_5$ , a module at level 5 associated with a non-diagonal invariant ( $r = 12$ ,  $h = 5 + 3 = 8$ ,  $\Sigma K = \Sigma_K^? = 9$ ). Unfortunately it does not work for other exceptional cases like  $\mathcal{E}_9$  ( $r = 12$ ,  $h = 9 + 3 = 12$ ,

<sup>8</sup> The notations that we use in this paragraph to denote these exceptional nimreps with self-fusion use the level  $k$  as a subscript, they differ from those used in [15] where the cases  $\mathcal{E}^5$ ,  $\mathcal{E}^9$ ,  $\mathcal{E}^{21}$  are respectively called  $\mathcal{E}^{(8)}$ ,  $\mathcal{E}_1^{(12)}$ ,  $\mathcal{E}^{(24)}$ .

$\Sigma K = 410/23$ , whereas  $\Sigma_K^2 = 91/5$ ), or like  $\mathcal{E}_{21}$  ( $r = 24$ ,  $h = 21 + 3 = 24$ ,  $\Sigma K = 138$ , whereas  $\Sigma_K^2 = 130$ ). It does not work for the  $\mathcal{D}_{k=3s}$  series either: using  $r = ((k+1)(k+2)/2 - 1)/3 + 3$ ,  $h = k + 3$ , one finds, for the first terms  $k = 3, 6, 9, \dots$ ,  $\Sigma K = 11/4, 10507/963, 64265/2024, \dots$  whereas  $\Sigma_K^2 = 14/5, 11, 637/20, \dots$ . Those values are numerically very close but we did not find a universal formula working in all cases. Something should be understood there.

Formulae generalizing equ. (23) and giving  $d_H$  for modules associated with the non-diagonal  $SU(3)$  invariants have been proposed (without proof) in [11,12].

## 5. Multiplicity properties for fusion products: from $\lambda \otimes \mu$ to $\lambda \otimes \bar{\mu}$

### 5.1. Known results

#### 5.1.1. Known results about multiplicities

The multiplicity of the trivial representation in the decomposition into irreducibles of a tensor product (resp. the fusion product) of three irreducible representations of a group (resp. an affine Lie algebra) is invariant under an arbitrary permutation of the three factors. This property, sometimes called Frobenius reciprocity, is well known. Assuming that the notion of conjugation is defined in the case under study, the multiplicity stays also constant if we conjugate simultaneously the three factors.

Invariance of the multiplicity of  $\lambda \otimes \lambda' \otimes \lambda'' \rightarrow \mathbb{1}$  under the above transformations implies, for example, that this integer is also equal to the multiplicity of  $\lambda \otimes \lambda' \rightarrow \bar{\lambda}''$ , of  $\bar{\lambda} \otimes \bar{\lambda}' \rightarrow \lambda''$ , of  $\lambda'' \otimes \lambda \rightarrow \bar{\lambda}'$ , of  $\lambda'' \otimes \lambda' \rightarrow \bar{\lambda}$ , etc.

These properties implies that the total multiplicity in the decomposition into irreducibles of the product  $\lambda \otimes \mu$  of two irreducible representations is trivially invariant if we conjugate *both* of them.

The following theorem was recently proven [13].

**Theorem 1.** *The total multiplicity in the decomposition into irreducibles of the tensor product of two irreducible representations of a simple<sup>9</sup> Lie algebra stays constant if we conjugate only one of them. At a given level, this property also holds for the fusion multiplicities of affine algebras.*

Another sum rule, which is much easier to prove (see for instance [1], section 1.1), using for instance Verlinde formula or its finite group analog, and which holds at least for finite groups, semi-simple Lie groups and affine algebras, states that the sum of squares of multiplicities is the same for  $\lambda \otimes \mu$  and  $\lambda \otimes \bar{\mu}$ . There are no such properties, in general, for higher powers of multiplicities.

Very recently, cf [1], it was further shown that

**Theorem 2.** *In the special case of  $SU(3)$ , the lists of multiplicities, in the tensor products  $\lambda \otimes \mu$  and  $\lambda \otimes \bar{\mu}$ , are identical up to permutations.*

This does not hold in general for other Lie algebras. It was conjectured that the same property should hold for the fusion product of representations of the affine algebra of  $\mathfrak{su}(3)$  at finite levels, but this stronger result, that we shall call property  $\mathfrak{P}$  in the following, was not proven in the quoted paper.

<sup>9</sup> The simplicity requirement can be lifted, see [16].

**Theorem 3** (property  $\mathfrak{P}$ ). *At any finite level  $k$  and for any integrable highest weights  $\lambda$  and  $\mu$  of  $\widehat{\mathfrak{su}}(3)_k$ , the lists of multiplicities  $N_{\lambda\mu}^{\nu}$  and  $N_{\lambda\bar{\mu}}^{\nu'}$  are identical up to permutations.*

The purpose of the present section is to discuss and prove this property.

A trivial corollary of [Theorem 3](#) is

**Corollary 1.** *At any finite level  $k$  and for any integrable highest weights  $\lambda$  and  $\mu$  of  $\widehat{\mathfrak{su}}(3)_k$ , the number of distinct irreps  $\nu$ , resp.  $\nu'$ , appearing in  $\lambda \star_k \mu$ , resp.  $\lambda \star_k \bar{\mu}$  is the same.*

Strangely, whereas for  $\mathrm{SU}(4)$  or  $\widehat{\mathfrak{su}}(4)_k$  [Theorem 3](#) is invalid (for a counter-example, see [\[1\]](#)), [Corollary 1](#) seems to hold.

**Conjecture 1.** *At any finite level  $k$  and for any integrable weights  $\lambda$  and  $\mu$  of  $\widehat{\mathfrak{su}}(4)_k$ , the number of distinct irreps  $\nu$ , resp.  $\nu'$ , appearing in  $\lambda \star_k \mu$ , resp.  $\lambda \star_k \bar{\mu}$  is the same.*

Of that we have no proof, only some evidence from the computation of all fusion products up to level 15.

In contrast, this property does not hold in general for higher  $\mathrm{SU}(N)$  or  $\widehat{\mathfrak{su}}(N)_k$ . For example, in  $\widehat{\mathfrak{su}}(5)_5$ , and for  $\lambda = (1, 1, 1, 0)$ ,  $\mu = (1, 1, 0, 1)$ , the line  $\mu$  of the matrix  $N_{\lambda}$  has 21 non-vanishing entries, whereas the line  $\bar{\mu}$  has 22.

### 5.1.2. A short summary on couplings, intertwiners, thresholds, and pictographs

In the case of groups or Lie algebras, intertwiners, thought of as equivariant linear maps from a tensor product of three representations to the scalars, are “3J-operators” that, when evaluated on a triple of vectors, become 3J symbols (Clebsch–Gordan coefficients). We shall sometimes refer to a particular space of intertwiners, associated to a particular term  $\nu$  in the decomposition into irreducibles of  $\lambda \otimes \mu$  as a “branching”, and denote it by  $\lambda \otimes \mu \rightarrow \nu$  (and indeed, in the case of a group  $G$ , it also corresponds to a branching from  $G \times G$  to its diagonal subgroup  $G$ ). The multiplicity of  $\nu$  in  $\lambda \otimes \mu$  is also the dimension  $d$  of the associated space of intertwiners. From the point of view of representation theory, all the irreps  $\nu$  that appear on the rhs of the decomposition of  $\lambda \otimes \mu$  are equivalent; nevertheless, it is convenient to consider each of them as a different “coupling” of the chosen representations (this is actually what is done in conformal field theory where such a multiplicity  $d$  is interpreted as the number of distinct couplings of the associated primary fields). In the same way combinatorial models typically associate to a space of intertwiners of dimension  $d$ , and therefore to a given triple or irreducible representations, a set of  $d$  combinatorial and graphical objects, that we call generically pictographs: the reader may look at [\[1\]](#) for a discussion of three kinds of pictographs (KT-honeycombs, BZ-triangles, O-blades), in the framework of the Lie group  $\mathrm{SU}(3)$ . Finally, let us mention that the choice of a basis in a space of intertwiners allows one to decompose a coupling along so-called elementary couplings [\[17\]](#) (equivalently, decompose pictographs along fundamental pictographs, see again [\[1\]](#) and sect. 6.2 of the present paper).

As in the first part, we denote an irreducible representation of a Lie algebra, or the integrable irrep of the corresponding affine algebra at level  $k$  (which may not exist if  $k$  is too small) by its associated highest weight, and therefore by the same symbol. Multiplicities  $N_{\lambda\mu}^{\nu}$ , also called Littlewood–Richardson coefficients, or fusion coefficients in the present context of affine Lie algebras, are the matrix elements of the fusion matrices considered in the first section but here we have to make the level explicit in the notation and call  $N_{\lambda\mu}^{(k)\nu}$  the multiplicity at level  $k$  and keep the notation  $N_{\lambda\mu}^{\nu}$  for the classical multiplicity (infinite level). Beginning with  $k = 0$  and letting  $k$  increase, it is usually so that the three irreps  $\lambda, \mu, \nu$  start to exist at possibly distinct

levels; then, even if the three irreps exist at level  $k$ , it may be that  $N_{\lambda\mu}^{(k)\nu}$  is still 0, or has a value smaller than the classical one (i.e., of infinite level). For instance, one has  $\{1, 1\} \otimes \{0, 2\} = \{0, 2\} \oplus \{1, 0\} \oplus \{1, 3\} \oplus \{2, 1\}$ , but at level  $k = 2$ , although the irrep  $\{0, 2\}$  exists, it does not appear in the fusion product, which is  $\{1, 1\} \star_2 \{0, 2\} = \{1, 0\}$ ; the other irreps  $\{2, 1\}$ ,  $\{0, 2\}$  and  $\{1, 3\}$  appear respectively in the fusion product  $\star_k$  at levels  $k = 3, 3, 4$ .

**Definition.** A triple  $(\lambda, \mu; \nu)$  is called *admissible*, or *classically admissible*, if  $N_{\lambda\mu}^\nu \neq 0$ ; it is called *admissible at level  $k$*  if  $N_{\lambda\mu}^{(k)\nu} \neq 0$ .

The following results can be gathered from [17].

The threshold level (or threshold, for short)  $k_0^{\min}$  of a triple (or of a branching  $\lambda \otimes \mu \rightarrow \nu$ ) is the smallest value of  $k$  for which the fusion coefficient  $N_{\lambda\mu}^{(k)\nu}$  is non-zero. It is known that, for any given triple of irreps, the fusion coefficient is an increasing function of the level, and that it becomes equal to its classical value  $N_{\lambda\mu}^\nu$  when  $k$  reaches a value  $k_0^{\max}$ , then it stays constant. The integers  $k_0^{\min}$  and  $k_0^{\max}$  are functions of  $\lambda, \mu$  and  $\nu$ , but for given  $\lambda$  and  $\mu$  one calls  $k^{\min}$  the infimum of the  $k_0^{\min}$  over the set  $\nu$ , and  $k^{\max}$  the supremum of the  $k_0^{\max}$  over the set  $\nu$ .

In terms of couplings (or pictographs), the discussion goes as follows: for fixed  $\lambda, \mu$  and  $\nu$  and a given  $k$ , we have a set of  $N_{\lambda\mu}^{(k)\nu}$  distinct couplings; these couplings still exist when the level becomes  $k + 1$ , but new couplings may appear, since  $N_{\lambda\mu}^{(k+1)\nu} \geq N_{\lambda\mu}^{(k)\nu}$ . The sets  $I(k)$  of couplings are therefore ordered by inclusion:  $I(k) \subset I(k + 1)$ . The threshold  $k_0(i)$  of a coupling  $i$  is, by definition, the smallest value of  $k$  at which it appears: the coupling  $i$  belongs to the set  $I(k_0(i))$  but it does not belong to the set  $I(k_0(i) - 1)$ .

The threshold  $k_0^{\min}$  of a triple (or of a branching) is therefore equal to the value of  $k$  for which the first associated couplings appear (such couplings have a threshold equal to  $k_0^{\min}$ ). Conversely the value  $k_0^{\max}$  of a branching is equal to the value of  $k$  above which no new couplings appear.

We have  $I(k_0^{\min}) \subset I(k_0^{\min} + 1) \subset I(k_0^{\min} + 2) \subset \dots \subset I(k) \subset \dots \subset I(k_0^{\max})$ . The set  $I(k)$  has cardinality  $N_{\lambda\mu}^{(k)\nu}$ , and the largest set  $I(k_0^{\max})$  can be considered as classical since it contains all the couplings.

From now on, we consider the special case of  $SU(3)$ . In all cases, as we saw, for any given triple of irreps, the fusion coefficient is equal to 0 when  $k < k_0^{\min}$ , equal to the constant  $N_{\lambda\mu}^\nu$  when  $k \geq k_0^{\max}$ , and is an increasing function of  $k$  when  $k_0^{\min} \leq k \leq k_0^{\max}$ . In the special case of the affine algebra  $\mathfrak{su}(3)$ , the multiplicity of the branching  $\lambda \otimes \mu \rightarrow \nu$  is a piecewise affine function of  $k$ , first equal to 0, then it starts to increase with  $k$ , taking values respectively equal to 1, 2, 3, ...,  $N_{\lambda\mu}^\nu$  for successive values of  $k = k_0^{\min}, k_0^{\min} + 1, \dots, k_0^{\max}$ . Then it stays constant. At level  $k$  the multiplicity (the number of couplings) is therefore  $k - k_0^{\min} + 1$ : one new coupling appears for every value of  $k$  between  $k_0^{\min}$  and  $k_0^{\max}$ .

Adapting the results of [17] to our own notations<sup>10</sup> (see also [4] and the discussion at the end of our section 5.1.3), we have<sup>11</sup>

<sup>10</sup> When discussing general couplings  $\lambda \otimes \mu \rightarrow \nu$ , it happens that most formulae are more simply expressed in terms of  $\lambda, \mu$ , and  $\bar{\nu}$ , than in terms of  $\lambda, \mu$  and  $\nu$ , because of Frobenius reciprocity. Admittedly, we should have chosen notations where the roles of  $\nu$  and  $\bar{\nu}$  are exchanged. However we shall not do that, in order to be consistent with the choices made in the companion paper [1].

<sup>11</sup> In order to be interpreted as a minimal or maximal threshold, the argument  $(\lambda, \mu; \nu)$  appearing in eqs. (27), (28), should refer to a classically admissible triple, although the rhs of these two equations make sense for arbitrary arguments. We hope that there should be no confusion.

$$\begin{aligned}
k_0^{\min}(\lambda, \mu; \nu) &= \max \left( \lambda_1 + \lambda_2, \frac{\lambda_1 - \lambda_2 + \mu_1 + 2\mu_2 + 2\nu_1 + \nu_2}{3}, \frac{-\lambda_1 + \lambda_2 + 2\mu_1 + \mu_2 + \nu_1 + 2\nu_2}{3}, \right. \\
&\quad \mu_1 + \mu_2, \frac{2\lambda_1 + \lambda_2 - \mu_1 + \mu_2 + \nu_1 + 2\nu_2}{3}, \frac{\lambda_1 + 2\lambda_2 + \mu_1 - \mu_2 + 2\nu_1 + \nu_2}{3}, \\
&\quad \left. \nu_1 + \nu_2, \frac{\lambda_1 + 2\lambda_2 + \mu_1 + 2\mu_2 - \nu_1 + \nu_2}{3}, \frac{2\lambda_1 + \lambda_2 + 2\mu_1 + \mu_2 + \nu_1 - \nu_2}{3} \right) \quad (27)
\end{aligned}$$

$$\begin{aligned}
k_0^{\max}(\lambda, \mu; \nu) &= \min \left( \frac{2\lambda_1 + \lambda_2 + 2\mu_1 + \mu_2 + \nu_1 + 2\nu_2}{3}, \frac{\lambda_1 + 2\lambda_2 + \mu_1 + 2\mu_2 + 2\nu_1 + \nu_2}{3} \right) \quad (28)
\end{aligned}$$

and

$$k^{\min} = \max(\lambda_1 + \lambda_2, \mu_1 + \mu_2) \quad (29)$$

$$k^{\max} = \lambda_1 + \lambda_2 + \mu_1 + \mu_2$$

The threshold  $k_0(i)$  of a given coupling ( $i$ ) can be read from one of its associated pictographs. Using BZ-triangles for example, this level is obtained<sup>12</sup> as follows:  $k_0(i)$  is the maximum of  $\lambda_1 + \lambda_2 + \alpha$ ,  $\mu_1 + \mu_2 + \beta$ ,  $\nu_1 + \lambda_2 + \gamma$ , where  $\alpha, \beta, \gamma$  are the values of the vertices respectively opposite to the three sides  $\lambda, \mu, \nu$ . Using O-blades, they are the values  $(c, e, a)$  of the internal edges opposite to the three sides in Fig. 10. The same integers can also be easily read from the SU(3)-honeycombs, since they are dual to O-blades (see Fig. 17 of [1]). The translation in terms of KT-honeycombs (which are actually GL(3)-honeycombs) or in terms of the hive model can be done by using methods explained in [1]. We shall illustrate the above considerations on one example, at the end of the next section.

### 5.1.3. Formulae for multiplicities

Explicit formulae for SU(3) multiplicities have been known for quite a while, the first known reference going back to [20], in 1965 (see also [1] and references therein). For the affine case, they were obtained in [17]. Assuming that the triplet  $(\lambda, \mu; \nu)$  is admissible, we have

$$\begin{aligned}
N_{\lambda\mu}^{(k)\nu} &= \begin{cases} 0 & \text{if } k < k_0^{\min}(\lambda, \mu; \nu) \\ k - k_0^{\min}(\lambda, \mu; \nu) + 1 & \text{if } k_0^{\min}(\lambda, \mu; \nu) \leq k \leq k_0^{\max}(\lambda, \mu; \nu) \\ k_0^{\max}(\lambda, \mu; \nu) - k_0^{\min}(\lambda, \mu; \nu) + 1 = N_{\lambda\mu}^{\nu} & \text{if } k \geq k_0^{\max}(\lambda, \mu; \nu) \end{cases} \quad (30)
\end{aligned}$$

This expression entails a recursion formula on fusion coefficients under a shift of all h.w. by  $\rho = (1, 1)$ , the Weyl vector, and of  $k$  by 3. Assume that  $\lambda, \mu, \nu$  are h.w. vectors such that  $\lambda - \rho, \mu - \rho, \nu - \rho$  are also integrable h.w. at level  $k - 3$ , i.e.,  $\lambda_i, \mu_i, \nu_i \geq 1, i = 1, 2$ , and  $\lambda_1 + \lambda_2 \leq k + 1$  etc. Then<sup>13</sup>  $k_0^{\min}(\lambda - \rho, \mu - \rho; \nu - \rho) = k_0^{\min}(\lambda, \mu; \nu) - 2$ , whereas  $k_0^{\max}(\lambda - \rho, \mu - \rho; \nu - \rho) = k_0^{\max}(\lambda, \mu; \nu) - 3$ , thus according to (30), if  $k \geq 3$

<sup>12</sup> This property was proven, in terms of BZ-triangles, by [18,19], see also [17].

<sup>13</sup> More generally, for all integers  $u, v$ , one shows using eq. (27) that  $k_0^{\min}(\lambda + (u, v), \mu + (u, v); \nu + (v, u)) = k_0^{\min}(\lambda, \mu; \nu) + (u + v)$ , and using eq. (28), that  $k_0^{\max}(\lambda + (u, v), \mu + (u, v), \nu + (v, u)) - k_0^{\max}(\lambda, \mu, \nu) = (u + 2v)$ , (resp  $(2u + v)$ ), if  $\lambda_1 + \mu_1 + \nu_2 \geq \lambda_2 + \mu_2 + \nu_1$  and  $u \geq v$ , (resp if  $\lambda_1 + \mu_1 + \nu_2 \leq \lambda_2 + \mu_2 + \nu_1$  and  $u \leq v$ ).



$$N_{\lambda-\rho, \mu-\rho}^{(k-3)v-\rho} = \begin{cases} 0 & \text{if } k-3 < k_0^{\min}(\lambda-\rho, \mu-\rho; v-\rho) = k_0^{\min}(\lambda, \mu; v) - 2 \\ (k-3) - (k_0^{\min}(\lambda, \mu; v) - 2) + 1 & \text{if } k_0^{\min}(\lambda, \mu; v) - 2 \leq k-3 \leq k_0^{\max}(\lambda, \mu; v) - 3 \\ (k_0^{\max}(\lambda, \mu; v) - 3) - (k_0^{\min}(\lambda, \mu; v) - 2) + 1 & \text{if } k-3 \geq k_0^{\max}(\lambda, \mu; v) - 3 \end{cases}$$

$$= \begin{cases} 0 & \text{if } k < k_0^{\min}(\lambda, \mu; v) + 1 \\ k - k_0^{\min}(\lambda, \mu; v) = N_{\lambda\mu}^{(k)v} - 1 & \text{if } k_0^{\min}(\lambda, \mu; v) + 1 \leq k \leq k_0^{\max}(\lambda, \mu; v) \\ k_0^{\max}(\lambda, \mu; v) - k_0^{\min}(\lambda, \mu; v) = N_{\lambda\mu}^v - 1 & \text{if } k \geq k_0^{\max}(\lambda, \mu; v) \end{cases}$$

In other words

$$N_{\lambda-\rho, \mu-\rho}^{(k-3)v-\rho} = \begin{cases} N_{\lambda\mu}^{(k)v} - 1 & \text{if } k \geq \max(k_0^{\min}(\lambda, \mu; v), 3) \\ 0 & \text{if } k \leq k_0^{\min}(\lambda, \mu; v) \text{ or } k < 3 \end{cases}. \quad (31)$$

We will make use of this relation in our proof of property  $\mathfrak{P}$ .

#### 5.1.4. A simple algorithm

In order to simply determine the multiplicity of the branching  $\lambda \otimes \mu \rightarrow v$ , or equivalently of  $\lambda \otimes \mu \otimes \bar{v} \rightarrow \mathbb{1}$ , at level  $k$ , one may proceed as follows, see also [21]. We use the following notations, and drop the label  $k$  for classical multiplicities:

$$\mathfrak{m}_k = \text{mult}_k(\lambda, \mu; v) = \text{mult}_k(\lambda \otimes \mu \otimes \bar{v}) = N_{\lambda\mu}^{(k)v}$$

*Classical case.* Call  $S_1 = \lambda_1 + \mu_1 + \bar{v}_1$ ,  $S_2 = \lambda_2 + \mu_2 + \bar{v}_2$ . If  $S_1 - S_2$  is not a multiple of 3, the multiplicity vanishes [17]. Assuming  $S_1 > S_2$ , we define three new irreps  $\lambda' = (\lambda_1 - x, \lambda_2)$ ,  $\mu' = (\mu_1 - x, \mu_2)$ ,  $\bar{v}' = (\bar{v}_1 - x, \bar{v}_2)$  where  $x = (S_1 - S_2)/3$ . Then<sup>14</sup>  $\text{mult}(\lambda \otimes \mu \otimes \bar{v}) = \text{mult}(\lambda' \otimes \mu' \otimes \bar{v}')$  and the new sums  $S'_1$  and  $S'_2$  are both equal to  $S = \min(S_1, S_2)$ . If  $S_1 < S_2$  one shifts the second components by  $x$  instead. Define  $\ell = S - \sum_i \lambda'_i$ ,  $m = S - \sum_i \mu'_i$ ,  $n = S - \sum_i \bar{v}'_i$  and consider the tableau of order 3 with lines  $(\lambda'_1, \mu'_1, \bar{v}'_1)$ ,  $(\lambda'_2, \mu'_2, \bar{v}'_2)$  and  $(\ell, m, n)$ . By construction, this tableau is a semimagic square of order 3 with magic constant  $S$ ; it is not necessarily magic because the sums along the diagonals do not add to  $S$  in general. The classical multiplicity  $\mathfrak{m}$  is equal to  $c + 1$  where  $c$  is the minimum of the entries of the tableau; indeed, by construction,  $c = k_0^{\max} - k_0^{\min}$ .

*Affine case.* Start from  $k = 0$  and let  $k$  increase: the irrep  $v$  appears for the first time, with multiplicity  $\mathfrak{m}_k = 1$ , in the tensor product  $\lambda \otimes \mu$  when  $k$  reaches<sup>15</sup> the threshold  $k_0^{\min} = S + x - c$ . The multiplicity then increases with the level, each time by one unit, so  $\mathfrak{m}_k = k - k_0^{\min} + 1$ , until

<sup>14</sup> Indeed, from eq. (27) and footnote 13, for an arbitrary integer  $x$ , we have  $k_0^{\min}(\lambda - (x, 0), \mu - (x, 0); v - (0, x)) - k_0^{\min}(\lambda, \mu; v) = -x$ ; moreover, if  $S_1 \geq S_2$ , from eq. (28) and for the particular value  $x = (S_1 - S_2)/3$ , we have  $k_0^{\max}(\lambda - (x, 0), \mu - (x, 0); v - (0, x)) - k_0^{\max}(\lambda, \mu; v) = -x$ .

<sup>15</sup> Remember that  $k_{\min}(\lambda, \mu) = \inf_v k_0^{\min}(\lambda, \mu, v)$  and that  $k_{\max}(\lambda, \mu) = \sup_v k_0^{\max}(\lambda, \mu, v)$ .



$k$  reaches the value  $k_0^{max} = S + x = k_0^{min} + (m - 1)$  for which the multiplicity becomes equal to the classical one,  $m$ . It then stays constant when  $k$  increases beyond  $k_0^{max}$ .

Warning. We have  $mult(\lambda \otimes \mu \otimes \bar{\nu}) = mult(\lambda' \otimes \mu' \otimes \bar{\nu}')$  when the level is infinite (classical case), but their thresholds  $k_0^{min}$  are distinct: it is  $S + x - c$  for the first but  $S - c$  for the next since  $x$  vanishes in the latter case.

Example: let us determine  $m = mult((9, 5), (6, 2); (8, 6)) = mult((9, 5) \otimes (6, 2) \otimes (6, 8))$ . We have  $S_1 = 21, S_2 = 15$ , so  $S = 15$  and  $x = (21 - 15)/3 = 2$ . Then<sup>16</sup>  $m = mult((7, 5) \otimes (4, 2) \otimes (4, 8))$  and  $(\ell, m, n) = (3, 9, 3)$ , the associated semimagic square being  $\begin{smallmatrix} 7 & 4 & 4 \\ 5 & 2 & 8 \\ 3 & 9 & 3 \end{smallmatrix}$ . Therefore  $c = 2$  and  $m = 3$ . The irrep  $(8, 6)$  appears with multiplicity 1 in  $(9, 5) \otimes (6, 2)$  when  $k$  reaches the threshold value  $k_0^{min} = S + x - c = 15 + 2 - 2 = 15$ . Therefore  $m_{15} = 1, m_{16} = 2, m_{17} = 3$ , then it stays constant.

### 5.1.5. One example

$$\begin{aligned} (9, 5) \otimes (6, 2) \\ = \{ (6, 1)_{14}, (7, 2)_{14,15}, (5, 3)_{14}, (8, 3)_{14,15,16}, (6, 4)_{14,15}, (4, 5)_{14}, (7, 5)_{14,15,16}, \\ (5, 6)_{14,15}, (3, 7)_{14}, (6, 7)_{14,15,16}, (4, 8)_{14}, (2, 9)_{14}, (5, 9)_{14,15,16}, (3, 10)_{14,15}, \\ (9, 1)_{15,16}, (10, 2)_{15,16,17}, (9, 4)_{15,16,17}, (8, 6)_{15,16,17}, (4, 8)_{15}, (7, 8)_{15,16,17}, \\ (4, 11)_{15,16}, (1, 11)_{14}, (8, 0)_{15}, (2, 12)_{15}, (11, 0)_{16}, (12, 1)_{16,17}, (11, 3)_{16,17,18}, \\ (10, 5)_{16,17,18}, (9, 7)_{16,17,18}, (6, 10)_{16,17}, (3, 13)_{16}, (14, 0)_{17}, (13, 2)_{17,18}, \\ (12, 4)_{17,18,19}, (11, 6)_{17,18,19}, (8, 9)_{17,18}, (5, 12)_{17}, (15, 1)_{18}, (14, 3)_{18,19}, \\ (13, 5)_{18,19,20}, (10, 8)_{18,19}, (7, 11)_{18}, (16, 2)_{19}, (15, 4)_{19,20}, (12, 7)_{19,20}, \\ (9, 10)_{19}, (17, 3)_{20}, (14, 6)_{20,21}, (11, 9)_{20}, (16, 5)_{21}, (13, 8)_{21}, (15, 7)_{22} \} \end{aligned}$$

The subindices refer to the levels at which the branching rule occurs. Let us consider the branching to  $(8, 6)$  that we already analyzed in the previous subsection. The term  $(8, 6)_{15,16,17}$ , for instance, means that the first coupling associated with the branching  $(9, 5) \otimes (6, 2) \rightarrow (8, 6)$  appears at level 15 (so the multiplicity is 1), another at level 16 (the multiplicity is 2), and a last one at level 17 (the multiplicity is 3). The tensor (i.e., classical) multiplicity of this branching is 3. Note that  $k_0^{min}((9, 5), (6, 2), (8, 6)) = 15$ ,  $k_0^{max}((9, 5), (6, 2), (8, 6)) = 17$  but  $k^{min}((9, 5), (6, 2)) = 14$  and  $k^{max}((9, 5), (6, 2)) = 22$ . The three couplings are illustrated by their respective pictographs, using O-blades and KT-honeycombs,<sup>17</sup> on Figs. 1, 2. Notice that the threshold  $k_0^{(i)}$ ,  $i = 1, 2, 3$ , for each of these couplings can be read from the pictures, as explained previously. The associated BZ-triangles can be immediately obtained from the correspondence explicated on Fig. 16 of [1].

Another way of displaying the results is given on Fig. 3. Rather than giving the irreps  $\nu$  obtained in the fusion of  $(9, 5)$  and  $(6, 2)$ , together with the level at which they appear, we display, for each level, the *couplings* that show up at that level. The highest weight of a particular irrep therefore appears several times on the figure (for instance  $(8, 6)$  appears on the three lines labeled by  $k = 15, 16, 17$ ).

<sup>16</sup> The thresholds of  $((9, 5), (6, 2); (8, 6))$  and  $((7, 5), (4, 2); (8, 4))$  are distinct: 15 for the first and 13 for the next.

<sup>17</sup> These are GL(3) honeycombs, one could prefer to draw instead the corresponding SU(3) honeycombs, where the lengths of all edges are non-negative, see [1].




Fig. 1. The three couplings of  $(9, 5) \otimes (6, 2) \rightarrow (8, 6)$  using the O-blade model.




Fig. 2. The three couplings of  $(9, 5) \otimes (6, 2) \rightarrow (8, 6)$  using the KT-honeycomb model.

What we did for the fusion product  $(9, 5) \otimes (6, 2)$  can be repeated for  $(9, 5) \otimes (2, 6)$ . We leave this as an exercise for the reader. Fig. 4, that should be compared with Fig. 3, summarizes the results.

As it was announced, the two lists of multiplicities at the classical level are the same, up to permutations (we find 21, 16 and 3 irreps with respective multiplicities 1, 2, 3, and therefore a total multiplicity equal to 95), something that we already knew from [1], but they are also equal for all values of the level. In the next section, we turn to a proof of this property.

## 5.2. From $\lambda \otimes \mu$ to $\lambda \otimes \bar{\mu}$ : proof of property $\mathfrak{P}$ using Wesslén polygons

### 5.2.1. New inequalities and the convex polygonal domain

Recall from [22,1] the inequalities satisfied by  $v = (v_1, v_2)$ ,  $v_1, v_2 \geq 0$ , in the intertwiner  $\lambda \otimes \mu \rightarrow v$

$$\begin{aligned} \max(2\mu_1 + \mu_2 - \lambda_1 - 2\lambda_2, 2\lambda_1 + \lambda_2 - \mu_1 - 2\mu_2, \lambda_2 - \lambda_1 + \mu_2 - \mu_1) \\ \leq 2v_1 + v_2 \leq 2\lambda_1 + \lambda_2 + 2\mu_1 + \mu_2 \end{aligned} \quad (32)$$

$$\begin{aligned} \max(\lambda_1 + 2\lambda_2 - 2\mu_1 - \mu_2, \mu_1 + 2\mu_2 - 2\lambda_1 - \lambda_2, \lambda_1 - \lambda_2 + \mu_1 - \mu_2) \\ \leq v_1 + 2v_2 \leq \lambda_1 + 2\lambda_2 + \mu_1 + 2\mu_2 \end{aligned} \quad (33)$$

$$\begin{aligned} \max(\mu_1 - \mu_2 - 2\lambda_1 - \lambda_2, \lambda_1 - \lambda_2 - 2\mu_1 - \mu_2) \\ \leq v_1 - v_2 \leq \min(\lambda_1 - \lambda_2 + \mu_1 + 2\mu_2, \lambda_1 + 2\lambda_2 + \mu_1 - \mu_2) \end{aligned} \quad (34)$$

and from [17] the additional “BMW” ones satisfied in the fusion product  $\lambda \star_k \mu \rightarrow v$  at level  $k$

$$v_1 + v_2 \leq k \quad (35)$$

$$2v_1 + v_2 \leq \min(3k - \lambda_1 + \lambda_2 - \mu_1 - 2\mu_2, 3k - \lambda_1 - 2\lambda_2 - \mu_1 + \mu_2) \quad (36)$$

$$v_1 + 2v_2 \leq \min(3k - 2\lambda_1 - \lambda_2 + \mu_1 - \mu_2, 3k + \lambda_1 - \lambda_2 - 2\mu_1 - \mu_2) \quad (37)$$

$$-3k + \lambda_1 + 2\lambda_2 + \mu_1 + 2\mu_2 \leq v_1 - v_2 \leq 3k - 2\lambda_1 - \lambda_2 - 2\mu_1 - \mu_2 \quad (38)$$




Fig. 3. Thresholds for intertwiners  $(9, 5) \otimes (6, 2) \rightarrow (v_1, v_2)$ . Each line (levels  $k^{\min} = 14$  to  $k^{\max} = 22$ ) lists those irreps  $v$  (components given vertically) for which the multiplicity increases at the chosen level, i.e., for which a new coupling appears. The total number of new couplings at a given level is also given on the left. Same irreps for which there is a multiplicity increase are connected by arrows. The number of arrows with the same color gives the  $u_{pj}$  of sect. 5.2.3: at level  $k = 18$ , i.e.,  $p = k - k^{\min} + 1 = 5$  there are 5 irreps that are not hit by any arrowhead ( $u_{50} = 5$ ), 4 hit by a blue arrow ( $u_{51} = 4$ ), and ( $u_{52} = 3$ ) hit by a red arrow, the total multiplicity increasing of  $5 + 4 + 3 = 12$ . (For interpretation of the references to color in this figure, the reader is referred to the web version of this article.)

which are simple consequences of  $k \geq k_0^{\min}(\lambda, \mu; v)$  with the explicit expression given in (27). Using standard symmetries of fusion coefficients, we may always assume that  $\lambda_1 \geq \max(\lambda_2, \mu_1, \mu_2)$ . Then note that the second inequality (37) follows from the first since  $\mu_1 \leq \lambda_1$ . As discussed above, the issue of finite level is relevant only in the range  $k^{\min} \leq k \leq k^{\max}$ , with  $k^{\min}$  and  $k^{\max}$  given in (29). Below  $k^{\min}$ , one of the weights  $\lambda$  or  $\mu$  is not integrable, above  $k^{\max}$  we are in the classical, “ $k = \infty$ ”, regime.




Fig. 4. Thresholds for intertwiners  $(9, 5) \otimes (2, 6) \rightarrow (\nu'_1, \nu'_2)$ .

For a given  $k \geq k^{\min}$  the set of inequalities (32)–(38) defines in the  $\nu$ -plane a convex polygonal domain. For  $k \geq k^{\max}$  this domain stabilizes to its “classical” shape defined by the only Wesslén’s inequalities (32)–(34), with vertices  $V_i$  defined on Fig. 5. In particular the “highest highest weight” is  $H = V_1 = (\lambda_1 + \mu_1, \lambda_2 + \mu_2)$ , while the “lowest highest weight”  $h = V_5$  is given below in (39). For  $k^{\min} \leq k < k^{\max}$ , we refer to the polygonal domain as the truncated Wesslén’s domain.

By the same type of recursive arguments as in sect. 5.1.3 and eq. (31), we may also find the polygons that bound the domain of multiplicity 2. By (30), they are obtained by imposing that  $k = 1 + k_0^{\min}(\lambda, \mu; \nu)$ , and result in inequalities of the form (35)–(38) in which  $k$  is substituted for  $k - 1$ . But the transformation  $k \rightarrow k - 3$ ,  $\lambda \rightarrow \lambda - \rho$ ,  $\mu \rightarrow \mu - \rho$ ,  $\nu \rightarrow \nu - \rho$  in (35)–(38)) has the same effect. We conclude that the set of  $\nu$  of level  $k$  and multiplicity  $N_{\lambda\mu}^{(k)\nu} = 2$  is the




Fig. 5. The oriented sides of a tensor polygon (with the assumption that  $\lambda_1 \geq \max(\lambda_2, \mu_1, \mu_2)$ ). (We have used orthogonal coordinate axes, although it would be more correct to have them at a  $\pi/3$  angle.) The violet, green, blue and red lines represent respectively the inequalities (35), (36), (37) and (38). As  $k$  decreases from  $k^{max}$  they start cutting the classical domain, hence restricting the allowed values of  $v$ . (For interpretation of the references to color in this figure, the reader is referred to the web version of this article.)

translate by  $\rho$  of the set of  $v'$  of level  $k - 3$  and multiplicity  $N_{\lambda-\rho, \mu-\rho}^{(k-3)v'} = 1$ . It would seem that this argument requires the  $\lambda_i, \mu_i, v_i$  to be  $\geq 1$  in order for  $\lambda - \rho, \mu - \rho, v - \rho$  to be integrable weights, and  $k - 3 \geq 0$ . In fact as the recursion relation is used for quadruplets  $(\lambda, \mu, v; k)$  for which  $N_{\lambda, \mu}^{(k)v} = 2$ , these conditions are automatically enforced.

The recursion extends to higher values of the multiplicity, see below sect. 5.2.3.

A useful remark is that the point  $V_5$  (the “lowest highest weight”) always satisfies the BMW inequalities for any value of  $k^{min} \leq k \leq k^{max}$  and thus always belongs to the truncated Wesslén’s domain.

Proof : a tedious case by case verification using

$$V_5 = \begin{cases} (\mu_1 + \mu_2 - \lambda_1 - \lambda_2, \lambda_1 - \mu_2) & \text{if } 0 \leq \lambda_1 - \mu_2 \leq \mu_1 - \lambda_2 \\ (\lambda_1 + \lambda_2 - \mu_1 - \mu_2, \mu_1 - \lambda_2) & \text{if } 0 \leq \mu_1 - \lambda_2 \leq \lambda_1 - \mu_2 \\ (\lambda_1 - \mu_2, \lambda_2 - \mu_1) & \text{if } \mu_1 - \lambda_2 < 0. \end{cases} \quad (39)$$

Intuitively,  $V_5$  “far away” from the colored lines of Fig. 5.

In contrast all the other vertices of the original (“classical”) tensor polygon may be removed in some cases and for some value of  $k$ .

### 5.2.2. The truncated Wesslén’s domain for $k^{min} \leq k < k^{max}$

As  $k$  is incremented from  $k^{min}$  by one unit, new points  $v$  appear until  $k = k^{max}$ , the new points having multiplicity 1, see [17]. The points  $v$  organize themselves in a pattern of polygons included into one another with the multiplicity increasing from 1 on the outside polygon<sup>18</sup> to a  $k$ -dependent maximum inside. At a given point  $v$  inside one of the polygons, the multiplicity increases as  $k$  grows, until  $k$  reaches the value  $k_0^{max}(\lambda, \mu; v)$ .


This is illustrated on Table 1 for our favorite examples of  $(9, 5) \star (6, 2)$  and  $(9, 5) \star (2, 6)$ .

<sup>18</sup> When  $k = k^{min}$  all the inner points have multiplicity 1 too.

Conversely as  $k$  is decremented from  $k^{max}$  by one unit, the inequality (35), and possibly the other inequalities, start to exclude points from Wesslén's domain. It is clear from Fig. 5 that inequalities (36), (37), (38)<sub>L</sub>, (38)<sub>R</sub>, are satisfied by all points of Wesslén's domain if they are











Table 1

Truncated Wesslén domains in  $v$  for the branchings  $(9, 5) \star (6, 2)$  (left column) and  $(9, 5) \star (2, 6)$  (right column), for increasing  $k$ . The solid polygon is the classical tensor polygon, drawn here in oblique axes. The dot-dashed line is the level line  $v_1 + v_2 = k$ . Points of multiplicity 1, 2 and 3 are displayed respectively in blue, orange and green. The  $u_{pj}$  of the last column are defined below in sect. 5.2.3 and can be read from the pattern of arrows in Fig. 3.



(continued on next page)

Table 1 (continued)

$k = 18$			$u_{40} = 5 \quad u_{41} = 4 \quad u_{42} = 3$
$k = 19$			$u_{50} = 4 \quad u_{51} = 3 \quad u_{52} = 2$
$k = 20$			$u_{60} = 3 \quad u_{61} = 2 \quad u_{62} = 1$
$k = 21$			$u_{70} = 2 \quad u_{71} = 1 \quad u_{72} = 0$
$k = 22$			$u_{80} = 1 \quad u_{81} = 0 \quad u_{82} = 0$

satisfied respectively at points  $V_1 = (\lambda_1 + \mu_1, \lambda_2 + \mu_2)$ ,  $V_8 = (\lambda_1 - \mu_1, \lambda_2 + \mu_1 + \mu_2)$  and  $V_2 = (\lambda_1 + \mu_1 + \min(\lambda_2, \mu_2), \lambda_2 + \mu_2 - 2 \min(\lambda_2, \mu_2))$ . Thus

$$(36) \text{ is satisfied if } k \geq \lambda_1 + \mu_1 + \max(\lambda_2, \mu_2) \quad (40)$$

$$(37) \text{ is satisfied if } k \geq \lambda_1 + \lambda_2 + \mu_2 \quad (41)$$

$$(38)_L \text{ is satisfied if } k \geq \lambda_2 + \mu_1 + \mu_2 \quad (42)$$

$$(38)_R \text{ is satisfied if } k \geq \lambda_1 + \mu_1 + \min(\lambda_2, \mu_2). \quad (43)$$

This introduces several successive boundary values of the level, as  $k$  decreases from  $k^{\max}$ .

### 5.2.3. Proof of property $\mathfrak{P}$

As the level  $k$  is decreased from  $k^{\min} + p$  to  $k = k^{\min} + p - 1$ , there are  $u_{pj}$  terms in the decomposition of  $\lambda \star \mu$  that have their multiplicity decreased from  $j + 1$  to  $j$ , for  $j = 0, \dots, p$ . In this section, we first give an explicit expression for  $u_{p0}$ , then use the recursion formula (31) to extend it to all  $u_{pj}$ . We then observe that these expressions of  $u_{pj}$  are invariant under  $\mu \rightarrow \bar{\mu}$ . Since we know from [1] that property  $\mathfrak{P}$  is satisfied for  $k \geq k^{\max}$ , and since decrementing  $k$  preserves it, it follows that it is satisfied for all  $k$ , qed.

**Computing the  $u_{p0}$ .** As level  $k$  is decreased from  $k^{\min} + p$  to  $k^{\min} + p - 1$ , there are  $u_{p0}$  terms that are excluded from the truncated Weyl domain because they violate one of the inequalities. According to [17], they lie on the boundary of the truncated Weyl domain of level  $k$ . Which inequality is relevant for a given  $k$  depends on the relative values of  $k$ ,  $\lambda_1$ ,  $\lambda_2$ ,  $\mu_1$  and  $\mu_2$ . Recall that by convention, we have  $\lambda_1 \geq \max(\lambda_2, \mu_1, \mu_2)$ .

By a tedious analysis of all possibilities that is not reproduced here, we have found the following expression for  $u_{p0}$ ,  $p = k - k^{\min}$

$$u_{p0} = \begin{cases} 0 & \text{if } k > k^{\max} \\ k^{\max} - k + 1 & \text{if } \max(\lambda_1 + \max(\mu_1, \mu_2), k^{\min} + 1) \leq k \leq k^{\max} = \lambda_1 + \lambda_2 + \mu_1 + \mu_2 \\ \lambda_2 + \min(\mu_1, \mu_2) + 1 & \text{if } k^{\min} + 1 \leq \lambda_1 + \max(\mu_1, \mu_2) \text{ and} \\ \max(\lambda_1 + \min(\mu_1, \mu_2), k^{\min} + 1) \leq k \leq \lambda_1 + \max(\mu_1, \mu_2) \\ k - \lambda_1 + \lambda_2 + 1 & \text{if } k^{\min} + 1 \leq \lambda_1 + \min(\mu_1, \mu_2) \text{ and } k^{\min} + 1 \leq k \leq \lambda_1 + \min(\mu_1, \mu_2) \\ \dots & \\ 0 & \text{if } k = k^{\min} \\ 0 & \text{if } k < k^{\min} \end{cases} \quad (44)$$

Note the “continuity” at the boundary values  $\lambda_1 + \max(\mu_1, \mu_2)$  and  $\lambda_1 + \min(\mu_1, \mu_2)$  of  $k$ .

Note also that this formula *does not* yet determine the value of  $u_{p0}$  for the lowest value 0 of  $p$  (viz  $k = k^{\min}$ ), i.e., the number of points of multiplicity 1 at level  $k^{\min}$ . The latter, however, is determined from the “classical” expression given by Mandel’stweig [20] and denoted  $\sigma(\lambda, \mu; 1)$



in [1] for the total number of  $\nu$  of multiplicity 1 (at levels higher than or equal to  $k^{\max}$ ), and from the successive  $u_{p0}$ :

$$u_{00} = \sigma(\lambda, \mu; 1) - \sum_{p=1}^{k^{\max}-k^{\min}} u_{p0}. \quad (45)$$

We observe that, as anticipated, the resulting expression for  $u_{p0}$  is invariant under  $\mu_1 \leftrightarrow \mu_2$ .

**Computing the  $u_{pj}$ .** Computing the variation of the number of points of multiplicity  $j+1 \geq 2$  as  $k$  decreases by one unit may look more difficult. Fortunately, the recursive argument of sect. 5.2.1 comes to the rescue. For  $k \geq 4$ , the number of  $\nu$  (with  $\nu_i \geq 1$ ) in  $\lambda \star \mu$  that are “downgraded” from multiplicity  $j > 1$  to multiplicity  $j-1$  as  $k \rightarrow k-1$  equals the number of  $\nu'$  in  $(\lambda - \rho) \star (\mu - \rho)$  that are downgraded from multiplicity  $j-1$  to multiplicity  $j-2$  as the level goes from  $k-3$  to  $k-4$ . Whence an expression of  $u_{p1}$  in terms of  $u_{p-1,0}$ : indeed, under a shift of  $\lambda$  and  $\mu$  by  $-\rho$  and of  $k$  to  $k-3$ , one has  $k^{\max} \rightarrow k^{\max} - 4$  and  $k^{\min} \rightarrow k^{\min} - 2$ , see eq. (29), thus  $p = k - k^{\min} \rightarrow p-1$ <sup>19</sup>

$$u_{p1}(\lambda, \mu) = u_{p-1,0}(\lambda - \rho, \mu - \rho) = \begin{cases} 0 & \text{if } k-3 > k^{\max} - 4 \\ (k^{\max} - 4) - (k-3) + 1 & \text{if } \lambda_1 + \max(\mu_1, \mu_2) - 2 \leq k-3 \leq k^{\max} - 4 \\ \lambda_2 + \min(\mu_1, \mu_2) - 2 + 1 & \text{if } \lambda_1 + \min(\mu_1, \mu_2) - 2 \leq k-3 \leq \lambda_1 + \max(\mu_1, \mu_2) - 2 \\ k - \lambda_1 + \lambda_2 - 3 + 1 & \text{if } k^{\min} - 2 + 1 \leq k-3 \leq \lambda_1 + \min(\mu_1, \mu_2) - 2 \\ \dots & \\ 0 & \text{if } k-3 = k^{\min} - 2 \\ 0 & \text{if } k-3 < k^{\min} - 2 \end{cases}$$

or in other words

$$u_{p1}(\lambda, \mu) = \begin{cases} 0 & \text{if } k > k^{\max} - 1 \\ k^{\max} - k & \text{if } \lambda_1 + \max(\mu_1, \mu_2) + 1 \leq k \leq k^{\max} - 1 \\ \lambda_2 + \min(\mu_1, \mu_2) - 1 & \text{if } \lambda_1 + \min(\mu_1, \mu_2) + 1 \leq k \leq \lambda_1 + \max(\mu_1, \mu_2) + 1 \\ k - \lambda_1 + \lambda_2 - 2 & \text{if } k^{\min} + 2 \leq k \leq \lambda_1 + \min(\mu_1, \mu_2) + 1 \\ \dots & \text{if } k = k^{\min} + 1 \\ 0 & \text{if } k < k^{\min} + 1 \end{cases} \quad (46)$$

<sup>19</sup> In order not to clutter the following equations with conditions involving  $k^{\min}$  (like in eq. (44)) we read the consecutive lines as “if clauses” from top to bottom, assuming that they should be used only if  $k^{\min}$  obeys appropriate inequalities...

and more generally

$$u_{pj}(\lambda, \mu) = \begin{cases} 0 & \text{if } k > k^{\max} - j \\ k^{\max} - k - j + 1 & \text{if } \lambda_1 + \max(\mu_1, \mu_2) + j \leq k \leq k^{\max} - j \\ \lambda_2 + \min(\mu_1, \mu_2) - 2j + 1 & \text{if } \lambda_1 + \min(\mu_1, \mu_2) + j \leq k \leq \lambda_1 + \max(\mu_1, \mu_2) + j \\ k - \lambda_1 + \lambda_2 - 3j + 1 & \text{if } k^{\min} + j + 1 \leq k \leq \lambda_1 + \min(\mu_1, \mu_2) + j - 1 \\ \dots & \text{if } k = k^{\min} + j \\ 0 & \text{if } k < k^{\min} + j \end{cases} \quad (47)$$

Again the missing value of  $u_{pp}$  may be derived from Mandel'tsveig's formula (see [20,1]) giving  $\sigma(\lambda, \mu; j+1)$ , the “classical” number of  $v$ 's of multiplicity  $j+1$ :

$$u_{pp} = \sigma(\lambda, \mu; j+1) - \sum_{p=j+1}^{k^{\max}-k^{\min}} u_{pj}.$$

We observe that this expression of  $u_{pj}$  is again invariant under  $\mu \leftrightarrow \bar{\mu}$ . As explained at the beginning of this section, this completes the proof of property  $\mathfrak{P}$ .

## 6. Miscellanea

### 6.1. From $\lambda \otimes \mu$ to $\lambda \otimes \bar{\mu}$ : other approaches

Admittedly the previous proof of property  $\mathfrak{P}$ , with its brute force calculation of the number of points at a given level, lacks elegance and simplicity. We thus attempted to explore other approaches... Although unsuccessful, as far as leading to another proof of the above property, some of these investigations lead to other results that may have a separate interest, we gather them here.

#### 6.1.1. Proof of property $\mathfrak{P}$ in some particular cases (elementary approach)

In addition to the “linear sum rule” proved in [13] (equality of the total multiplicities for  $\lambda \otimes \mu$  and  $\lambda \otimes \bar{\mu}$  at each level), we recall from sect. 5.1.1 the existence of a “quadratic sum rule”: the sum of squares of multiplicities are also the same at each level. These two sum rules put constraints on the two lists of multiplicities.

As in section 5.2.3, and for  $j = 0, \dots, p$ , we call  $u_{pj}$  (resp  $v_{pj}$ ) the number of terms in the decomposition of  $\lambda \otimes \mu$  (resp  $\lambda \otimes \bar{\mu}$ ) that have their multiplicity increased from  $j$  to  $j+1$  when the level is increased from  $k^{\min} + p - 1$  to  $k^{\min} + p$ . Keeping track of the increase in multiplicities at each level, the reader will have no difficulty to show that the two sum rules imply:

$$\sum_{j=0}^p u_{pj} = \sum_{j=0}^p v_{pj} \quad \text{and} \quad \sum_{j=0}^p ((j+1)^2 - j^2) u_{pj} = \sum_{j=0}^p ((j+1)^2 - j^2) v_{pj}.$$

The general result obtained in sect. 5.2.3, namely that  $u_{pj} = v_{pj}$  for all  $p$  and all  $j$ , cannot be deduced from the two sum rules recalled above. However this equality can be immediately obtained at levels  $k = k^{\min}$ ,  $k = k^{\min} + 1$ , and  $k \geq k^{\max}$ . Indeed, when  $k = k^{\min} + 1$  i.e.,  $p =$

1, the two previous equations read, equivalently,  $u_{10} - v_{10} = -(u_{11} - v_{11})$  and  $u_{10} - v_{10} = -3(u_{11} - v_{11})$ , which implies  $u_{10} = v_{10}$  and  $u_{11} = v_{11}$ . When  $k = k^{\min}$  (multiplicities are all equal to 1) and when  $k \geq k^{\max}$  (multiplicities have their classical values), the result is obvious since it is just a way to re-express already known results.

The approach described in the present section provides therefore an elementary proof of property  $\mathfrak{P}$  when  $k = k^{\min}$ ,  $k^{\min} + 1$ , and  $k = k^{\max}$ . By the same token, it also gives a proof for all values of  $k$  if the highest weights  $\lambda$  and  $\mu$  are such that  $k^{\max} - k^{\min} = 0, 1$  or  $2$ .

### 6.1.2. Using automorphisms

The group of automorphisms of the affine version of the  $A_2$  Dynkin diagram is  $\mathbb{Z}_3$ . Let  $(\lambda_1, \lambda_2)$  be the components of a weight  $\lambda$  on the basis of fundamental weights. If the level is  $k$  one introduces an affine component  $\lambda_0 = k - (\lambda_1 + \lambda_2)$  and consider the affine weight  $\hat{\lambda}$  with components  $(\lambda_0, \lambda_1, \lambda_2)$ . The generator  $\zeta$  of  $\mathbb{Z}_3$  acts on affine weights as follows:  $\zeta(\lambda_0, \lambda_1, \lambda_2) = (\lambda_2, \lambda_0, \lambda_1)$ . Obviously,  $\zeta^3 = \mathbb{1}$ . Existence of automorphisms  $a, b$  imply  $N_{a\lambda, b\mu}^{(k)abv} = N_{\lambda, \mu}^{(k)v}$ . In the present case,  $a$  and  $b$  can taken as  $\zeta, \zeta^2$  or  $\mathbb{1}$  and it is understood that automorphisms act on the affine extension of the weights  $\lambda, \mu, v$ , although their affine component (their first component) is usually dropped from the notation: in other words  $\zeta(\lambda_1, \lambda_2) = (\lambda_0, \lambda_1)$ ,  $\zeta^2(\lambda_1, \lambda_2) = (\lambda_2, \lambda_0)$ , etc.

We consider a given branching  $\lambda \otimes \mu \rightarrow v$ . The affine weights at level  $k$  are  $\hat{\lambda} = (k - (\lambda_1 + \lambda_2), \lambda_1, \lambda_2)$ ,  $\hat{\mu} = (k - (\mu_1 + \mu_2), \mu_1, \mu_2)$ ,  $\hat{v} = (k - (v_1 + v_2), v_1, v_2)$ . Let us choose the level  $k$  in such a way that  $k - (\mu_1 + \mu_2) = \mu_1$ , so we take  $k = 2\mu_1 + \mu_2$ . Notice that  $\hat{\mu} = (\mu_1, \mu_1, \mu_2)$ , therefore  $\zeta^2(\mu_1, \mu_1, \mu_2) = (\mu_1, \mu_2, \mu_1)$ . Using automorphisms  $a = \mathbb{1}, b = \zeta^2$ , one has  $N_{\lambda, \zeta^2\mu}^{(k)\zeta^2v} = N_{\lambda, \mu}^{(k)v}$ , i.e.,

$$k = 2\mu_1 + \mu_2, \quad N_{(\lambda_1, \lambda_2), (\mu_2, \mu_1)}^{(k)(v_2, 2\mu_1 + \mu_2 - (v_1 + v_2))} = N_{(\lambda_1, \lambda_2), (\mu_1, \mu_2)}^{(k)(v_1, v_2)}. \quad (48)$$

In order for the components of the weights (including the affine component) to stay non-negative, one needs to assume  $2\mu_1 + \mu_2 \geq \lambda_1 + \lambda_2$  and  $2\mu_1 + \mu_2 \geq v_1 + v_2$ .

In the same way, assuming  $2\mu_2 + \mu_1 \geq \lambda_1 + \lambda_2$  and  $2\mu_2 + \mu_1 \geq v_1 + v_2$ , one chooses the level  $k$  in such a way that  $k - (\mu_1 + \mu_2) = \mu_2$ , so  $k = 2\mu_2 + \mu_1$ . With  $\hat{\mu} = (\mu_2, \mu_1, \mu_2)$ , we have  $\zeta(\mu_2, \mu_1, \mu_2) = (\mu_2, \mu_2, \mu_1)$ . Using automorphisms  $a = \mathbb{1}, b = \zeta$ , one gets  $N_{\lambda, \zeta\mu}^{(k)\zeta v} = N_{\lambda, \mu}^{(k)v}$ , explicitly:

$$k = 2\mu_2 + \mu_1, \quad N_{(\lambda_1, \lambda_2), (\mu_2, \mu_1)}^{(k)(2\mu_2 + \mu_1 - (v_1 + v_2), v_1)} = N_{(\lambda_1, \lambda_2), (\mu_1, \mu_2)}^{(k)(v_1, v_2)} \quad (49)$$

It is clear that the two above transformations are inverse of one another.

We now use  $\text{mult}(\lambda, \mu; v) = \text{mult}(\mu, \lambda; v)$  and  $\text{mult}(\mu, \bar{\lambda}; v') = \text{mult}(\bar{\mu}, \lambda; \bar{v}') = \text{mult}(\lambda, \bar{\mu}; \bar{v}')$ .

Assuming  $2\lambda_2 + \lambda_1 \geq \mu_1 + \mu_2$  and  $2\lambda_2 + \lambda_1 \geq v_1 + v_2$ , one obtains in the same way

$$k = 2\lambda_2 + \lambda_1, \quad N_{(\lambda_1, \lambda_2), (\mu_2, \mu_1)}^{(k)(v_1, 2\lambda_2 + \lambda_1 - (v_1 + v_2))} = N_{(\lambda_1, \lambda_2), (\mu_1, \mu_2)}^{(k)(v_1, v_2)} \quad (50)$$

And, assuming  $2\lambda_1 + \lambda_2 \geq \mu_1 + \mu_2$  and  $2\lambda_1 + \lambda_2 \geq v_1 + v_2$

$$k = 2\lambda_1 + \lambda_2, \quad N_{(\lambda_1, \lambda_2), (\mu_2, \mu_1)}^{(k)(2\lambda_1 + \lambda_2 - (v_1 + v_2), v_2)} = N_{(\lambda_1, \lambda_2), (\mu_1, \mu_2)}^{(k)(v_1, v_2)} \quad (51)$$

For illustration, we apply eq. (48) to our former example  $\lambda = (9, 5)$ ,  $\mu = (6, 2)$ . One checks that the chosen automorphism applied to the list of weights  $v$  appearing on the line<sup>20</sup>  $k = 2\mu_1 + \mu_2 =$

<sup>20</sup> this is also equal to  $k^{\min}(\lambda, \mu)$ , but it is an accident.

14 of Fig. 3 gives, up to reordering, the line  $k = 14$  of Fig. 4. In the same way, the equality of multiplicities for the appropriate triples also holds if we use eq. (51) (now,  $k$  has to be chosen as  $2\lambda_2 + \lambda_1 = 19$ ).

Although giving non-trivial results for particular values of the level, we do not see how to generalize this approach to handle the general case.

### 6.1.3. Using a piece-wise linear map in the space of weights

In reference [1] several proofs of property  $\mathfrak{P}$  in the classical case, i.e., for tensor products, were given. One of them was based on the construction of an involutive piece-wise linear map  $T$  from the set of admissible couplings (or of their corresponding pictographs) associated with the various branchings  $\lambda \otimes \mu \rightarrow \nu$  to the set of couplings associated with the branchings  $\lambda \otimes \bar{\mu} \rightarrow \nu'$ . The two weights  $\lambda, \mu$  being fixed, this particular transformation  $T : (\nu, \alpha) \mapsto (\nu', \alpha')$ , with  $\alpha$  denoting some coupling (or some pictograph) for the triple  $\lambda, \mu, \nu$ , cannot be used in the present situation where we deal with fusion product at level  $k$  because it does not respect the thresholds: in order to prove property  $\mathfrak{P}$  for fusion products, one should exhibit a bijective piecewise-linear transformation  $T$  such that the couplings defined by  $\nu, \alpha$  and  $\nu', \alpha'$  have the same threshold.

Taking into account the specificities of  $SU(3)$ , in particular the fact that the multiplicities of a given branching increase by one unit when the level  $k$  runs from  $k_0^{\min}$  to  $k_0^{\max}$ , it is enough to look for an invertible map  $T$  that is compatible with these two values of the level. In other words, if  $T$  is a one-to-one map from the set of admissible triples  $(\lambda, \mu, \nu)$  to the set of admissible triples  $(\lambda, \bar{\mu}, \nu')$  and is such that  $k_0^{\min}(\lambda, \mu, \nu) = k_0^{\min}(\lambda, \bar{\mu}, \nu')$  and  $k_0^{\max}(\lambda, \mu, \nu) = k_0^{\max}(\lambda, \bar{\mu}, \nu')$ , the theorem is proved.

Unfortunately, many particular cases have to be considered, and the discussion leading to a definition of such a piece-wise linear map seems to be as complicated as the one leading to the proof of property  $\mathfrak{P}$  in sect. 5.2.3.

### 6.1.4. Using the matrix polynomial $X(s, t)$ to prove property $\mathfrak{P}$ (a failed attempt)

The property  $\mathfrak{P}$  can be rephrased as follows: for given weights  $\lambda$  and  $\mu$  and level  $k$ , the multisets<sup>21</sup>  $\{N_{\lambda\mu}^{\nu}\}$  and  $\{N_{\lambda\bar{\mu}}^{\nu'}\}$  are identical and we may use the conjugation properties of the fusion coefficients

$$N_{\lambda\bar{\mu}}^{\nu'} = N_{\lambda\mu}^{\bar{\nu}'} = N_{\lambda\bar{\nu}'}^{\mu}$$

to write this statement in equivalent forms. Thus  $\{N_{\lambda\mu}^{\nu}\} = \{N_{\lambda\bar{\nu}'}^{\mu}\}$  means that

*For an arbitrary  $\widehat{su}(3)_k$  fusion matrix  $N_{\lambda}$ , the content of any row  $\mu$  is the same (up to a permutation that in general depends on the choice of the label  $\mu$ ) as that of the column  $\mu$  of the same matrix.*

Equality of the multisets  $\{N_{\lambda\mu}^{\nu}\} = \{N_{\lambda\bar{\mu}}^{\nu'}\}$  implies that our generating function  $X(s, t)$  satisfies the following property that we call  $\mathcal{P}$ :

$\mathcal{P}$ : For any row label  $\mu$  and for any monomial  $s^{\lambda_1} t^{\lambda_2}$  appearing<sup>22</sup> along the row  $\mu$ , the monomial  $s^{\lambda_2} t^{\lambda_1}$  also appears the same total number of times along the same row  $\mu$ .

<sup>21</sup> i.e., sets with multiplicities, or lists, up to order. As we work with a fixed level  $k$  in this section, we drop this index from the notation used for fusion matrices.

<sup>22</sup> i.e., appearing in the decomposition into monomials of any matrix element (polynomials) of the row  $\mu$ .

Can one derive this result directly from properties of the generating functions? The reader will have no difficulty to prove the following lemmas L1–L5:

**L1** The matrix elements of the fusion matrices of type  $N_{(\lambda_1,0)}$  are either 0 or 1.

**L2** The number of non-zero elements, in the row  $\mu$  (arbitrary) of the fusion matrix  $N_{(\lambda_1,0)}$ , is equal to the number of non-zero elements in the same row  $\mu$  of the fusion matrix  $N_{(0,\lambda_1)}$ , or, equivalently, in the column  $\mu$  of the same matrix  $N_{(\lambda_1,0)}$ .

**L3** The matrix elements of the fusion matrices of type  $N_{(\lambda_1,k-\lambda_1)}$  (i.e., those belonging to the third edge of the Weyl alcove) are either 0 or 1.

**L4** For every choice of an irrep  $\mu$ , the number of non-zero elements in the row  $\mu$  of the matrix  $N_{(\lambda_1,k-\lambda_1)}$  is equal to the number of non-zero elements in the same row  $\mu$  of the matrix  $N_{(k-\lambda_1,\lambda_1)}$  or equivalently, in the column  $\mu$  of the same matrix  $N_{(\lambda_1,k-\lambda_1)}$ .

**L5** Every row of the boundary matrix polynomial<sup>23</sup>  $\Lambda(s, t)$  containing some monomial  $s^{\lambda_1} t^{\lambda_2}$  (in a matrix element) also contains the monomial  $s^{\lambda_2} t^{\lambda_1}$  the same number of times. Equivalently,  $\Lambda(s, t)$  satisfies property  $\mathcal{P}$ .

It is also clear that the polynomial  $A(s, t) = [(s^3 + t^3 + 2st + 2s^2t^2)\mathbb{1} - st(sG + tG^T)]$  satisfies the same property  $\mathcal{P}$ . If we could assert that this property is preserved by matrix multiplication and inversion, we would conclude that  $K(s, t) = A^{-1}(s, t)$  and  $X(s, t) = K(s, t) \cdot \Lambda(s, t)$  also satisfy the same property, thus establishing property  $\mathcal{P}$ , which is equivalent to  $\mathfrak{P}$ . Unfortunately, this is not the case in general and we could not prove property  $\mathfrak{P}$  in this way. It is nevertheless interesting to see how this property is translated when read in terms of generating functions: this justifies the inclusion of the above discussion in the present paper.

## 6.2. An application of O-blades: a new relation between fusion coefficients

**Proposition.** *The transformation defined by eq. (52) is involutive, and, for arbitrary levels  $k$ , one has:*

$$N_{\lambda\mu}^{(k)v} = N_{\lambda'\mu'}^{(k)v'}$$

A generic O-blade for  $SU(3)$  is displayed in Fig. 6(a).

It contains nine edges  $m_{12}, m_{23}, m_{13}, n_{12}, n_{23}, n_{13}, l_{12}, l_{23}, l_{13}$ , labeled with non-negative integers.

It has a single inner vertex, and obeys one single constraint: the three pairs of opposite “angles” defined by the lines intersecting at the inner vertex should be equal, the value of an angle being defined as the sum of its sides. The three weights appearing in the branching rule  $\lambda, \mu \rightarrow \nu$  follow each other as indicated in Fig. 6. Their components, in terms of edges, read (“integer conservation at the external vertices”):

$$\begin{aligned} \lambda_1 &= m_{13} + n_{12} & \mu_1 &= n_{13} + l_{12} & \nu_1 &= m_{13} + l_{23} \\ \lambda_2 &= m_{23} + n_{13} & \mu_2 &= n_{23} + l_{13} & \nu_2 &= m_{12} + l_{13} \end{aligned}$$

The reader will have no difficulty to re-express the “Wesslén inequalities” (32)–(34) as positivity constraints for the nine edges (“blades”) of the O-blades. As in [1] (see this reference for a more complete discussion), we call *fundamental pictographs* (here fundamental O-blades) the combinatorial models associated with intertwiners of the type  $f_1 \otimes f_2 \otimes f_3 \rightarrow \mathbb{1}$  where the  $f_i$ ’s

<sup>23</sup>  $\Lambda(s, t)$  is defined as the rhs of eq. (19).




Fig. 6. (a): A generic O-blade for SU(3) and its nine internal edges. (b): Its seven components ( $a, b, c, d, e, f, g$ ) along the left fundamental basis.




Fig. 7. Fundamental O-blades for SU(3).

denote either a fundamental representation or the trivial one. The dimension of the corresponding spaces of intertwiners being equal to 0 or 1, there is only one such pictograph or none. For SU(3), we have six fundamental O-blades obtained by permuting the factors of  $f \otimes f^* \otimes \mathbb{1}$  where  $f$  is the three-dimensional vector representation (of highest weight  $(1, 0)$ ), and we have two other fundamental O-blades respectively associated with the cubes of  $(1, 0)$  and of  $(0, 1)$ ; the last two (resp. the first six) are called “primitive fundamental” (resp. “non-primitive fundamental”) in [1]. The eight fundamental O-blades are given in Fig. 7 – edges carrying a “1” label have been thickened. Obviously, O-blades, thought of as characteristic functions associated with their set of edges, can be added or multiplied by scalars (they may be called virtual whenever some inner edges carry values that do not belong to the set of non-negative integers. One could think that this family of eight fundamental pictographs can be chosen as a basis in the space of characteristic functions, but this is not so because they are not independent: there is also one relation which is displayed in Fig. 8. The two primitive fundamental O-blades appear as Y-shapes (forks) with a horizontal tail to the left or to the right of the inner vertex; the relation displayed in Fig. 8 exhibits two ways of making a star out of its components. Notice that the space of pictographs comes with a distinguished generating family – those that are fundamental – but not with a distinguished basis. Up to permutations, we may consider two interesting particular basis, each contains seven fundamental intertwiners: the six non-primitive ones, and a last one chosen among the two that are primitive.

Choosing the “left” basis (i.e., the last basis element being Y-shaped, with a horizontal tail to the left of the inner vertex), we see in Fig. 9(a) or in Fig. 6(b) how an arbitrary O-blade  $o$  can be defined as a superposition of its seven components and write<sup>24</sup>  $o_L = (a, b, c, d, e, f, g)$ . The associated weights  $\lambda_1 = b + e + g$ ,  $\lambda_2 = a + f$ ,  $\mu_1 = a + d + g$ ,  $\mu_2 = b + c$ ,  $\nu_1 = d + e$ ,  $\nu_2 = c + f + g$ .

Call  $\Psi$  the linear map that leaves invariant the six non-primitive O-blades, and permutes the two that are Y-shaped. Under this map, an arbitrary element  $o$  (see Fig. 9(a)) becomes  $\Psi(o)$  (see Fig. 9(b)), which, using the relation between fundamental O-blades displayed in Fig. 8, can be

<sup>24</sup> One can easily show that the component  $g$  coincides with the parameter  $x = (S_1 - S_2)/3$  of section 5.1.3.




Fig. 8. The SU(3) intertwiner relation in terms of O-blades.




Fig. 9. (a) Decomposition of an O-blade along the left fundamental basis, (b): its image par  $\Psi$ , and (c): its decomposition along the same basis using the SU(3) intertwiner relation.

expanded on the left basis as in Fig. 9(c). We have therefore<sup>25</sup>  $\Psi(o)_L = (a, b + g, c, d + g, e, f + g, -g)$ , with associated weights  $\lambda', \mu', \nu'$  and  $\lambda'_1 = e + b, \lambda'_2 = a + f + g, \mu_1 = a + d, \mu_2 = b + g + c, \nu_1 = f + g + e, \nu_2 = f + c$ . The linear map  $\Psi$ , defined on the space of couplings, induces a transformation  $(\lambda, \mu; \nu) \rightarrow (\lambda', \mu'; \nu')$  on branchings, that we also call  $\Psi$ , with

$$\begin{aligned} \lambda'_1 &= \frac{1}{3}(2\lambda_1 + \lambda_2 - \mu_1 + \mu_2 + \nu_1 - \nu_2) & \lambda'_2 &= \frac{1}{3}(\lambda_1 + 2\lambda_2 + \mu_1 - \mu_2 - \nu_1 + \nu_2) \\ \mu'_1 &= \frac{1}{3}(-\lambda_1 + \lambda_2 + 2\mu_1 + \mu_2 + \nu_1 - \nu_2) & \mu'_2 &= \frac{1}{3}(\lambda_1 - \lambda_2 + \mu_1 + 2\mu_2 - \nu_1 + \nu_2) \\ \nu'_1 &= \frac{1}{3}(\lambda_1 - \lambda_2 + \mu_1 - \mu_2 + 2\nu_1 + \nu_2) & \nu'_2 &= \frac{1}{3}(-\lambda_1 + \lambda_2 - \mu_1 + \mu_2 + \nu_1 + 2\nu_2) \end{aligned} \quad (52)$$

The nine internal edges and the three highest weights associated with  $\Psi(o)$  are displayed in Fig. 10(b). From its geometrical definition (permutation of left and right forks), it is already clear that  $\Psi$  is an involution, but this can also be checked from the equations given previously. In particular  $\Psi$  is injective, both for branchings and for O-blades (or couplings). This implies that if a given branching has classical multiplicity  $m$ , and can therefore be associated with  $m$  distinct O-blades describing its corresponding couplings, its image by  $\Psi$  is automatically admissible and will have the same multiplicity, the O-blades describing the couplings of the latter being obtained as the image by  $\Psi$  of the O-blades of the former.

We now show that an O-blade  $o$  (characterizing a coupling), and its image  $\Psi(o)$  have the same threshold. In terms of components of highest weights and internal edges, we know (see Fig. 6(a)) that  $k_0(o) = \max(\lambda_1 + \lambda_2 + l_{13}, \mu_1 + \mu_2 + m_{13}, \nu_1 + \nu_2 + n_{13})$ , or, equivalently, in terms of the

<sup>25</sup> One should notice that positivity of edges of a (non-virtual) O-blade implies positivity of its components along the first six intertwiners, those that we called non-primitive fundamental, but not along the last one (the left fork).




Fig. 10. Permuting forks: an O-blade  $o$  and its image  $\Psi(o)$ .

(left) fundamental components  $o_L = (a, b, c, d, e, f, g)$  of  $o$ , that  $k_0(o) = \max(a + b + c + e + f + g, a + c + d + e + f + g, a + b + c + d + e + g)$ . Using  $\Psi(o)_L = (a, b + g, c, d + g, e, f + g, -g)$ , one finds immediately  $k(\Psi(o)) = k_0(o)$ . In particular this also implies that branchings exchanged by  $\Psi$  have the same minimal threshold, so that this transformation also holds in the affine case and that it is compatible with the levels.

**Example.**  $\Psi((9, 5), (6, 2); (10, 5)) = ((8, 6), (5, 3); (11, 4))$ . In both cases, multiplicity and threshold are the same:  $\text{mult}((9, 5), (6, 2); (10, 5)) = \text{mult}((8, 6), (5, 3); (11, 4)) = 3$  and  $k_0^{\min} = 16$ .

**Remark.** It is easy to see that the map  $\Psi$  is compatible with the transposition of the first two arguments and with the six Frobenius transformations.

## Acknowledgements

R.C. thanks hospitality of the Instituto Nacional de Matemática Pura e Aplicada (IMPA, Rio de Janeiro), where part of this work was done.

J.-B.Z. gratefully acknowledges support from the Simons Center for Geometry and Physics, Stony Brook University, at which part of the research of this paper was completed. He thanks Pavel Bleher, Vladimir Korepin and Bernard Nienhuis, the organizers of the workshop “Statistical Physics and Combinatorics”, for their invitation.

## References

- [1] R. Coquereaux, J.-B. Zuber, Conjugation properties of tensor product multiplicities, J. Phys. A 47 (2014) 455202, <http://arxiv.org/abs/1405.4887>.
- [2] L. Bégin, C. Cummins, P. Mathieu, Generating functions for tensor products, J. Math. Phys. 41 (11) (2000) 7611–7639, <http://arxiv.org/abs/hep-th/9811113>.
- [3] L. Bégin, C. Cummins, P. Mathieu, Generating-function method for fusion rules, J. Math. Phys. 41 (11) (2000) 7640–7674, <http://arxiv.org/abs/math-ph/0005002>.
- [4] P. Di Francesco, P. Mathieu, D. Sénéchal, *Conformal Field Theory*, Springer, 1997.
- [5] E.B. Dynkin, Semisimple subalgebras of semisimple Lie algebras, Mat. Sb. (N. S.) 30 (72) (1952) 349–462.
- [6] A. Ocneanu, Paths on Coxeter diagrams: from platonic solids and singularities to minimal models and subfactors, in: Rajarama Bhat, et al. (Eds.), Notes Taken by S. Goto, in: Fields Institute Monographs, AMS, 1999.
- [7] A. Cappelli, C. Itzykson, J.-B. Zuber, The ADE classification of minimal and  $A_1^{(1)}$  conformal invariant theories, Commun. Math. Phys. 13 (1987) 1–26.
- [8] V. Ostrik, Module categories, weak Hopf algebras and modular invariants, Transform. Groups 8 (2) (2003) 177–206, <http://arxiv.org/abs/math.QA/0111139>.
- [9] V. Petkova, J.-B. Zuber, The many faces of Ocneanu cells, Nucl. Phys. B 603 (2001) 449–496, <http://arxiv.org/abs/hep-th/0101151>.



- [10] A. Malkin, V. Ostrik, M. Vybornov, Quiver varieties and Lusztig's algebra, *Adv. Math.* 203 (2) (2006) 514–536, <http://arxiv.org/abs/math.RT/0403222>.
- [11] R. Coquereaux, G. Schieber, Quantum symmetries of  $sl(2)$  and  $sl(3)$  graphs, *Proceedings of the Fifth International Conference on Mathematical Methods in Physics (Rio de Janeiro)*, *Proceedings of Science: PoS (IC2006) 001*, [http://pos.sissa.it/archive/conferences/031/001/IC2006\\_001.pdf](http://pos.sissa.it/archive/conferences/031/001/IC2006_001.pdf).
- [12] R. Coquereaux, G. Schieber, Orders and dimensions for  $sl_2$  or  $sl_3$  module-categories and boundary conformal field theories on a torus, *J. Math. Phys.* 48 (2007) 043511, <http://arxiv.org/abs/math-ph/0610073>.
- [13] R. Coquereaux, J.-B. Zuber, On sums of tensor and fusion multiplicities, *J. Phys. A* 44 (2011) 295208, <http://arxiv.org/abs/1103.2943>.
- [14] V. Kac, D. Peterson, Infinite dimensional Lie algebras, theta functions, and modular forms, *Adv. Math.* 53 (1984) 125–264.
- [15] P. Di Francesco, J.-B. Zuber,  $SU(N)$  lattice integrable models associated with graphs, *Nucl. Phys. B* 338 (1990) 602–646;  
P. Di Francesco, J.-B. Zuber,  $SU(N)$  lattice integrable models and modular invariance, in: S. Randjbar-Daemi, E. Sezgin, J.-B. Zuber (Eds.), *Recent Developments in Conformal Field Theories, Trieste Conference 1989*, World Scientific, 1990.
- [16] R. Zeier, Z. Zimborás, On squares of representations of compact Lie algebras, *J. Math. Phys.* 56 (2015) 081702, <http://arxiv.org/abs/1504.07732>.
- [17] L. Bégin, P. Mathieu, M.A. Walton,  $\widehat{su}(3)_k$  fusion coefficients, <http://arxiv.org/abs/hep-th/9206032>.
- [18] A.N. Kirillov, P. Mathieu, D. Sénéchal, M. Walton, Can fusion coefficients be calculated from the depth rule?, *LAVAL-PHY-20/92*, *LETH-PHY-2/92*, <http://arxiv.org/abs/hep-th/9203004>.
- [19] S. Lu, PhD thesis, MIT, 1990, A.N. Kirillov, unpublished.
- [20] V.B. Mandel'tsveig, Irreducible representations of the  $SU_3$  group, *Sov. Phys. JETP* 20 (1965) 1237–1243.
- [21] L. Suciú, The  $SU(3)$  wire model, PhD thesis, The Pennsylvania State University, 1997, AAT 9802757. A. Ocneanu, unpublished.
- [22] M. Wesslén, A geometric description of tensor product decompositions in  $su(3)$ , *J. Math. Phys.* 49 (2008) 073506.