Subduction zone intermediate-depth seismicity: Insights from the structural analysis of Alpine high-pressure ophiolite-hosted pseudotachylyte (Corsica, France)

Rémi Magott, Olivier Fabbri, Marc Fournier

► To cite this version:

Rémi Magott, Olivier Fabbri, Marc Fournier. Subduction zone intermediate-depth seismicity: Insights from the structural analysis of Alpine high-pressure ophiolite-hosted pseudotachylyte (Corsica, France). *Journal of Structural Geology*, 2016, 87, pp.95-114. 10.1016/j.jsg.2016.04.002 . hal-01346160

HAL Id: hal-01346160

<https://hal.sorbonne-universite.fr/hal-01346160>

Submitted on 18 Jul 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Subduction zone intermediate-depth seismicity: Insights from the structural analysis of Alpine high-pressure ophiolite-hosted pseudotachylyte (Corsica, France).

Rémi Magott¹, Olivier Fabbri^{1*} and Marc Fournier²

1 : UMR CNRS 6249 Chrono-environnement, Université Bourgogne Franche-Comté, Besançon, France

Email : Remi.Magott@univ-fcomte.fr; olivier.fabbri@univ-fcomte.fr

2 : Sorbonne Universités, UPMC Univ. Paris 06, UMR 7193, ISTEP, F-75005 Paris, France

Email : marc.fournier@upmc.fr

* Corresponding author: olivier.fabbri@univ-fcomte.fr

Highlights

Alpine Corsica ophiolite pseudotachylyte formed in a subducting oceanic lithosphere

Displacement sense associated with pseudotachylyte fault veins is determined

Corsican ruptures are similar to seismic ruptures in Pacific plate beneath NE Japan

ABSTRACT

Pseudotachylyte in the Cima di Gratera ophiolite, Alpine Corsica, is distributed in the peridotite unit and in the overlying metagabbro unit and was formed under blueschist to eclogite metamorphic facies conditions, corresponding to a 60-90 km depth range. Peridotite pseudotachylyte is clustered in fault zones either beneath the tectonic contact with overlying metagabbros or at short distance from it. Fault zones are either parallel to the contact or make an angle of 55° to it. Displacement sense criteria associated with fault veins indicate top-to-the-west or top-to-the-northwest reverse senses. Cataclasite flanking most veins was formed before or coevally with frictional melting and likely mechanically weakened the peridotite, facilitating subsequent seismic rupture. In the basal part of the metagabbro unit, post-mylonitization pseudotachylyte can be distinguished from pre-mylonitization pseudotachylyte formed earlier. In the equant metagabbro above the mylonitic sole, only one episode of pseudotachylyte formation can be identified. Kinematics associated with metagabbro pseudotachylyte remain unknown. The geometry and kinematics of the pseudotachylyte veins from the peridotite unit and to a lesser extent from the metagabbro unit are similar to modern

seismic ruptures of the upper parts of the Wadati-Benioff zones such as in the Pacific plate beneath NE Japan.

KEYWORDS

Pseudotachylyte, intermediate-depth seismicity, peridotite, metagabbro, subduction, Alpine Corsica

1. Introduction

Subduction zone seismicity consists mainly of shallow 'megathrust-type' earthquakes (focal depths < 60 km), intermediate-depth earthquakes (focal depths between 60 and 300 km), and deep-focus earthquakes (focal depths > 300 km). Events from the first category typically nucleate and propagate at the interface between the two lithospheric plates, while those from the last two categories nucleate in the crust or the mantle of the subducting slab. In subduction zones, the most destructive earthquakes are those from the first category. Their hypocenters can be shallow, their magnitudes can reach or exceed 8, and they can trigger tsunamis. Though less spectacular, intermediate-depth earthquakes still represent a significant threat because of the proximity of their epicenters with major cities and also because of infrequent but high magnitudes. Events illustrating such hazards include the 1939 Chile earthquake (Frohlich, 2006) or the 2001 El Salvador earthquake (Vallée et al., 2003; Martinez-Diaz et al., 2004).

In subduction zones, the hypocenters of intermediate-depth earthquakes tend to be clustered along a so-called Wadati-Benioff seismic zone. In most cases, precise hypocentral locations allow to divide the Wadati-Benioff seismic zone into two sub-zones. The separation between the two sub-zones is comprised between 8 and 30 km and is a function of the age of the subducting plate (Hasegawa et al., 1978a and b, 2009; Yoshii, 1979; Kao and Chen, 1995; Kao and Liu, 1995; Seno and Yamanaka, 1996; Brudzinski et al., 2007). The upper sub-zone appears to be located in the crust and/or in the uppermost part of the underlying mantle, while the lower sub-zone lies entirely in the mantle (Igarashi et al., 2001; Preston et al., 2003; Abers et al., 2013; Nakajima et al., 2013).

Pseudotachylyte exposed in Cape Corse (Alpine Corsica, France) is of tectonic origin and was generated under blueschist to eclogite facies metamorphic conditions (Austrheim and Andersen, 2004; Andersen and Austrheim, 2006; Andersen et al., 2008, 2014; Deseta et al., 2014a and b). As such, it was likely formed in Cretaceous to Paleogene times in the Wadati-Benioff seismic zone of a subducting Tethysian lithosphere. The aim of this contribution is to analyze the geometry of the pseudotachylyte fault veins and the kinematics during their

formation. The results of this analysis then allows a comparison between the fossil Corsican Cretaceous to Paleogene seismic ruptures and present-day Wadati-Benioff seismic zones observed in cold slabs such as the Pacific plate beneath NE Japan.

2. Geological setting and structure of the study area

2.1. General setting

Alpine Corsica is a segment of the Alpine orogen which displays an imbrication of thrust sheets composed of rocks of various origins and variably deformed and metamorphosed. Peridotite, serpentinite, gabbro, basalt, calcareous schist, siliceous schist and marble represent remnants of the lithosphere of the Jurassic Piemonte-Liguria oceanic basin and its pelagic sedimentary cover. Radiometric dating of the oceanic peridotites and gabbros yielded Middle to Late Jurassic ages between 169 and 152 Ma (Ohnenstetter et al., 1981; Rossi et al., 2002; Rampone et al., 2009; Li et al., 2015). Several observations indicate that the spreading ridge of the Piemonte-Liguria basin was of slow to very slow type (Rampone and Piccardo, 2000; Piccardo, 2008; Manatschal and Müntener, 2009).

Crystalline thrust sheets (composed mainly of orthogneisses) and proximal sedimentary deposits are interpreted as fragments of the stretched European continental paleo-margin and its sedimentary cover (Vitale-Brovarone et al., 2011, 2014; Meresse et al., 2012). The imbrication of such a variety of rocks is classically interpreted as the result of an eastward-dipping Cretaceous subduction of the Piemonte-Liguria oceanic basin and a part of the stretched European margin beneath the continental lithosphere of Apulia followed by an Eocene collision between the European and Apulian continental lithospheres (Mattauer and Proust, 1976; Mattauer et al., 1977, 1981; Warburton, 1986). Initiation of the subduction is poorly dated. Paleogeographic reconstructions suggest a Late Cretaceous age (Stampfli et al., 1998), but Late Cretaceous absolute ages of the HP metamorphism suggest that subduction started in Middle Cretaceous times or earlier. The ophiolite-bearing Schistes Lustrés nappe complex lies between the two blocks involved in the collision.

The classical Alpine evolutionary models were modified by taking into account the Apennine orogeny (e.g., Durand-Delga and Rossi, 2002). Several authors suggested that the east-dipping Cretaceous subduction ceased in Paleocene-Early Eocene times and was followed by a west-dipping subduction of a young oceanic lithosphere of a back-arc basin formed further east (Jolivet et al., 1998; Lacombe and Jolivet, 2005; Molli, 2008; Molli and Malavieille, 2010; Agard and Vitale-Brovarone, 2013). Based on new Late Eocene HP metamorphism

ages, Vitale-Brovarone and Herwartz (2013) suggest that the subducting oceanic plate now preserved as ophiolitic thrust sheets and nappes in Alpine Corsica was of 'Apennine' affinity, i.e., was dipping westward since the very beginning of the convergence. These authors however acknowledge that more datings are needed before invalidating the eastward-dipping ('Alpine') subduction. In this paper, the Late Cretaceous to Early Tertiary oceanic subduction will be considered as an Alpine-type east-dipping subduction, before it is replaced by a west-dipping Apennine-type subduction in Middle Eocene times.

A large part of Alpine Corsica rocks suffered from a high pressure-low temperature (HP-LT) blueschist to lawsonite-eclogite facies metamorphism (Ravna et al., 2010; Vitale-Brovarone et al., 2013 and references therein). This HP-LT metamorphism is of Eocene age (55-34 Ma, Brunet et al., 2000; Martin et al., 2011; Maggi et al., 2012; Vitale-Brovarone and Herwartz, 2013) and is interpreted as the result of the subduction of continental or oceanic units at great depths. A retrograde greenschist facies metamorphism is also recorded in some units and is interpreted as the consequence of a late- to post-orogenic extension during late Oligocene to early Miocene times (Jolivet et al., 1990, 1991, 1998; Fournier et al., 1991).

Non-metamorphic tectonic units (so-called upper or superficial nappes) lie at the top of the structural stacking of thrust sheets. These units consist of sedimentary strata mostly of Jurassic to Eocene age and ophiolitic rocks and related deposits. The rocks constituting these superficial nappes were likely formed or deposited on the margin of the Apulian continent (or an intervening island arc or micro-block) and subsequently transported westward during the collision above the metamorphic units.

Structures associated with the ductile non-coaxial deformation of the metamorphic units include a widespread foliation, various folds (including sheath folds) and a pervasive E-W stretching and mineral lineation. The sense of shear associated with the non-coaxial deformation during the prograde metamorphism is top-to-the-west (Mattaueer et al., 1977, 1981; Faure and Malavieille, 1981; Harris, 1985; Warburton, 1986) whereas that associated with the retrograde metamorphism and the late- to post-orogenic extension is mainly top-to-the east (Fournier et al., 1991; Jolivet et al., 1990, 1991). The late- to post-orogenic extension is also attested by numerous normal faults striking around N-S and dipping eastwards or westwards.

2.2. Structure of the study area

The study area is located in the southern part of the Cape Corse peninsula, around the Cima di Gratera peak, and consists of what will be referred to hereafter as the Cima di Gratera ophiolitic nappe. Through an inferred tectonic contact called ϕ_1 , this nappe overlies ductilely deformed units composed of continental basement rocks, meta-ophiolites and meta-sedimentary cover rocks (Pigno-Olivaccio and Morteda-Farinole units, Fig. 1) which recorded a HP-LT metamorphism (Lahondère, 1981, 1996; Vitale-Brovarone et al., 2013). The Cima di Gratera nappe consists of two superimposed units (Fig. 2): a lower serpentinite-peridotite unit (hereafter peridotite unit) and an upper metagabbro unit separated by a brittle/ductile flat-lying contact referred to as ϕ_2 .

According to Vitale-Brovarone et al. (2013), the highest metamorphic conditions recorded by meta-sedimentary rocks and continental units surrounding the Cima di Gratera nappe are temperatures between 414 and 471°C and pressures between 1.9 and 2.6 GPa, corresponding to blueschist to eclogite facies conditions. Comparable P-T conditions (1.3 GPa, $455 \pm 35^\circ\text{C}$) were estimated by Lahondère and Guerrot (1997) in similar units nearby the Cima di Gratera nappe. No accurate P-T conditions could be determined directly from the Cima di Gratera units. Following Deseta et al. (2014a and b) and despite the presence of faults between them, we suppose that the Cima di Gratera units suffered from P-T conditions comparable to those of the surrounding units, that is, blueschist to eclogite facies P-T conditions.

2.2.1. The peridotite unit.

The peridotite unit is composed of massive or foliated serpentinite embedding fresh to variably serpentinitized peridotite lenses. Near the inferred ϕ_1 tectonic contact, the base of the unit consists of strongly foliated serpentinites whose foliation is severely folded or sheared by C-like surfaces. This intense basal deformation is interpreted as a consequence of the emplacement of the nappe over its substratum. Most serpentinites are distributed in the lower part of the peridotite unit and the degree of serpentinitization generally decreases upwards from ϕ_1 . Most peridotite masses are located near the ϕ_2 contact and have thicknesses between 20 m and 300 m. The peridotite is massive and granoblastic. At locality 3 (Fig. 1), it is cut by a gabbro dyke. The peridotite is lherzolitic in composition, and is constituted by olivine, diopside, enstatite and minor plagioclase, Cr-spinel and magnetite (Deseta et al., 2014a). Pyroxenite was found near the contact ϕ_2 (locality 7), but the poor exposure conditions prevent to determine its actual extent and its nature (dyke, cumulate or sill). Recrystallization of diopside and enstatite to clinochlore and tremolite testifies to a greenschist facies metamorphism of the peridotite (Deseta et al., 2014a and b). On the other hand, and unlike the

metagabbro of the overlying unit (see below), no mineralogical evidence for blueschist facies metamorphism could be found in the peridotite. That the peridotite itself does not contain any evidence for blueschist facies metamorphism most likely reflects the fact that its composition does not allow formation of minerals diagnostic of blueschist facies. Indeed (see our observations below and also Deseta et al., 2014a and b), peridotite-hosted pseudotachylyte veins contain omphacitic microlites indicating that the melt cooled and solidified under blueschist facies to eclogite facies conditions.

According to Deseta et al. (2014a), the greenschist facies metamorphism occurred during two events or successions of events, first *before* the formation of pseudotachylyte and then *after* it. The early metamorphic event or succession of events are tentatively related to hydrothermal alterations having occurred during ocean-continent hyperextension, during seafloor spreading at the ridge, or near the trench where the approaching plate bends. The late metamorphic event or succession of events are likely contemporaneous with slab exhumation or nappe emplacement processes.

In several localities, especially in the northern part of the study area, thin (about 15 m) to thick (> 500 m) lenses of strongly foliated meta-sedimentary rocks and metagabbros are found within the serpentinites, generally at short distances from the inferred basal contact φ_1 (Fig. 1). These lenses likely correspond to slices of the underlying units that were incorporated in the peridotite unit during nappe emplacement. The generally strong deformation (intense folding, shear bands offsetting foliation) of the rocks in the lenses and of the surrounding serpentinites is in favor of this interpretation.

2.2.2. *The metagabbro unit.*

The description of the metagabbro unit by Deseta et al. (2014a) is summarized below and is completed by our observations. The metagabbro unit is predominantly composed of an equant metagabbro. Only its basal part consists of a foliated metagabbro. The primary minerals of the equant metagabbro are plagioclase, diopside, minor olivine and rare ilmenite. Alteration of plagioclase into sericite and of olivine into serpentine, magnetite or iddingsite is common. The texture frequently changes from micro-gabbro to coarse-grained gabbro and locally to pegmatitic gabbro. A magmatic foliation is locally observed but could not be mapped because of exposure scarcity. The metagabbro is intruded by dolerite dykes (locality 12, Fig. 1). According to Deseta et al. (2014a), an early greenschist facies metamorphism of the gabbro is responsible for the partial or total replacement of diopside by actinolite, bastite or Mg-hornblende. This early greenschist facies metamorphism is followed by a blueschist facies

metamorphism as attested by the replacement of diopside, actinolite, Mg-hornblende and plagioclase by glaucophane, barrosite, albite and epidote. A late greenschist facies metamorphism is evidenced by epidote, clinochlore and pumpellyite overprinting the blueschist facies minerals. Like for the peridotite unit, the early greenschist metamorphism of the gabbro is tentatively related to hydrothermal alteration having occurred during ocean-continent hyperextension, during seafloor spreading at the ridge, or near the trench through normal faulting of the bending plate (Deseta et al., 2014a and b). The late metamorphic event is considered to be contemporaneous with slab exhumation or nappe emplacement.

2.2.3. The contact between the two units

The contact ϕ_2 between the two units of the Cima di Gratera nappe, already described by Andersen et al. (2014), can be observed at several places (localities 6, 7, 8 and 9, Fig. 1). It consists in a flat-lying sharp fracture surface which undulates gently and which superimposes equant or foliated gabbro over peridotite or serpentinite. Weakly marked striations or corrugations trending N75°E to N120°E are preserved on the surface. The peridotite below the contact surface is intensely fractured and hosts abundant pseudotachylyte fault veins, most of which being parallel or slightly oblique to the surface (see description below). In the localities where the footwall consists of serpentinites, no pseudotachylyte could be found in these rocks.

Where the hanging-wall consists of equant metagabbro, up to 10 cm thick pseudotachylyte veins locally outline the base of the metagabbro unit. Where the hanging-wall consists of foliated metagabbro, the foliation is generally parallel to the contact but can also be oblique to it, forming angles of up to 35°. The thickness of the basal foliated metagabbro is between 20 cm and 30 m. The mylonitic deformation progressively decreases in intensity when going upward, that is away from ϕ_2 . The foliation bears a stretching lineation around N120°E. Polished hand-sample sections and thin sections perpendicular to the foliation and parallel to the lineation did not provide any consistent shear senses. Going upward and away from ϕ_2 , minor shear zones cross-cut the equant metagabbro. Their thickness is about 15 cm but can locally reach 1.5 m. They strike about N-S and dip between 10 and 40° westward. Their foliation bears a weakly marked stretching lineation along N80°E. No consistent shear senses could be retrieved from hand sample sections or thin sections. The spatial as well as the chronological relationships between these minor shear zones and the main basal foliated zone could not be clarified.

222

223 **3. Pseudotachylyte**

224 *3.1. Pseudotachylyte in the peridotite unit*

225 *3.1.1. General observations*

226 A summary of the description of the pseudotachylyte in the peridotite unit by Andersen et al.
227 (2008, 2014), Andersen and Austrheim (2006), Austrheim and Andersen (2004) and Deseta et
228 al. (2014a and b), completed by our observations, is given in the following. Pseudotachylyte
229 veins in the peridotite are characterized by a positive relief and by an orange to yellowish
230 color which contrasts with the rusty color of the host rock (Fig. 3). Two categories of
231 pseudotachylyte veins are distinguished: fault veins and injection veins. In contrast with
232 injection veins, fault veins can be followed along several tens of centimeters up to a few
233 meters, have similar orientations, and their thickness is between a few millimeters and 30 cm.
234 Injection veins are rare, small (length < 10 cm), and often cut by microfaults or cataclastic
235 shear zones, rendering their recognition difficult. Most fault veins (about 90 %) do not occur
236 as isolated occurrences, but are clustered in fault zones in which a large number of veins
237 (commonly several tens) form anastomosed or tangled networks (Figs 3 to 6).

238 As demonstrated by hand sample polished sections or thin sections, the peridotite hosting the
239 pseudotachylyte veins, whatever these are isolated or forming networks, is cataclastic to
240 varying extents. Where cataclasis is important, the granoblastic texture cannot be recognized
241 any more. Where cataclasis is moderate, olivine crystals, yet fractured, can be distinguished.

242 In the peridotite unit, the fresh (i.e., not serpentinized) peridotite always contains
243 pseudotachylyte. Conversely, the massive serpentinite does not contain any pseudotachylyte.
244 In the vicinity of localities 4 and 12 (Fig. 1), some fault or injection veins can be recognized
245 in serpentinite but, in these occurrences, serpentinization affects both the host rock and the
246 veins, and clearly post-dates pseudotachylyte formation.

247 As already reported by Andersen et al. (2014), unlike all pseudotachylyte occurrences
248 described in non-ductilely deformed host rocks and particularly in other ultramafic rocks
249 (Obata and Karato, 1995; Piccardo et al., 2007, 2010; Souquière and Fabbri, 2010; Ueda et
250 al., 2008), the peridotite fault veins form complex anastomosed or tangled networks inside
251 which no general relative chronology between veins can be established. Indeed, cross-cutting
252 relationships between a limited number (up to 3) veins are often observed at the outcrop scale

and also at the thin section scale, but remain insufficient with respect to the large number of veins to allow a complete chronology of a single fault zone to be reconstructed.

Another difference between the Cima di Gratera peridotite pseudotachylyte and most occurrences from elsewhere lies in the scarcity of sharp and planar fault surfaces adjacent to fault veins. Indeed, since the pioneering work of Sibson (1975), it has been recognized that fault veins are always flanked, on one side or on both sides, by sharp, commonly planar, slip surfaces whose length is between one and several meters or tens of meters (e.g., Allen, 2005; Di Toro and Pennacchioni, 2005; Grocott, 1981; McNulty, 1995; Swanson, 1988; Wenk et al., 2000; Zechmeister et al., 2007). In the case of the Cima di Gratera peridotite unit, such meter-scale slip surfaces are rare. This seems to be also the case for the Lanzo peridotite pseudotachylyte (Piccardo et al., 2007, 2010).

3.1.2. Geometry and kinematics of pseudotachylyte-bearing fault zones

As reported by Andersen and Austrheim (2006), two types of pseudotachylyte fault vein-bearing zones can be distinguished in the peridotite unit: flat-lying fault zones in the upper part of the unit (near ϕ_2), and steeply-dipping fault zones in the middle or lower part of the unit (Fig. 2). Isolated fault veins also follow this geometry: flat-lying veins in the upper part of the peridotite unit, steeply dipping ones in the lower part. To determine the sense of displacement (seismic slip) associated with pseudotachylyte fault veins and given the fact that the fault veins or associated cataclastic surfaces lack displacement direction indicators such as striations, field-oriented hand samples were cut along vertical planes striking N-S, NE-SW, E-W and NW-SE. For all samples, the most coherent sense of displacement criteria are observed along NW-SE surfaces and to a lesser extent along E-W surfaces. This indicates that the NW-SE direction is likely the displacement direction.

Flat-lying pseudotachylyte-bearing fault zones

Two main flat-lying fault zones are found in the peridotite unit (Fig. 2): the first one is located immediately beneath the ϕ_2 contact where it can be followed almost continuously (localities 4 to 12, Fig. 1); the second one is located in the median part of the unit (localities 1 and 3, Fig. 1). The thickness of the fault zone beneath ϕ_2 varies between 25 and 250 m, while that at localities 1 and 3 is about 15 m. In the fault zone beneath ϕ_2 , fault veins are parallel or slightly oblique to ϕ_2 and their density unambiguously increases upwards when getting closer to ϕ_2 . The parallelism between veins also tends to increase when getting closer to the contact. Examination of outcrops, polished hand sample sections and thin sections reveals numerous

cross-cutting relationships between fault veins. As stated above, because of the complexity of the relationships, it is not possible to establish a unique and complete chronology of seismic ruptures in the flat-lying fault zones. The only clear relationship observed at all localities consists in early gently to moderately ($5\sim 20^\circ$) dipping fault veins offset by late flat-lying to gently dipping ($0\sim 10^\circ$) fault veins. Early veins often show blurred boundaries. Conversely, late veins have sharp boundaries, and are frequently thinner than early veins. Cataclasite is always associated with the early veins but is seldom observed along the late veins. For the two fault zones (Figs 3 and 4), the sense of displacement associated with late veins is top-to-the-northwest or top-to-the-west. The sense of displacement associated with the early veins remains undetermined. Top-to-the-northwest or top-to-the-west displacement senses are obtained for isolated flat-lying fault veins scattered in the peridotite unit.

Steeply dipping pseudotachylyte-bearing fault zones

In addition to isolated steeply dipping veins, three steeply dipping pseudotachylyte-bearing fault zones are observed (localities 1, 3 and 5, Fig. 1). They consist of anastomosed networks of fault veins hosted by cataclastic peridotite. Their thickness varies between 30 cm and 1 m. The two first zones strike $N20^\circ E$ to $N40^\circ E$ and dip 55° eastwards. The third zone (locality 5) strikes around $N80\text{--}N100^\circ E$ and dips 55° northwards. The two first fault zones display a clear zonation with cataclastic peridotite predominating in the hanging-wall side and fault veins predominating in the footwall side. This asymmetrical zonation is reminiscent of pseudotachylyte described in different geological settings and host rocks (Fabbri et al., 2000). Kinematic indicators suggest that the sense of displacement during formation of the steeply dipping fault veins was a reverse one, that is a top-to-the-northwest one (Figs 5 and 6). Flat-lying or gently-dipping ($0\sim 10^\circ$) fault surfaces, most of them being coated by pseudotachylyte, cross-cut the steeply dipping fault veins of the three steeply-dipping fault zones. Kinematic indicators associated with these late flat-lying faults indicate a top-to-the-west or top-to-the-northwest sense of displacement. The same cross-cutting relationships and kinematics are obtained for the isolated steeply dipping fault veins scattered in the peridotite unit: steeply dipping fault veins are cross-cut by flat-lying veins, but both types are characterized by a top-to-the-west or top-to-the-northwest sense of displacement.

Relative dating of flat-lying and steeply-dipping fault zones.

Both flat-lying and steeply dipping fault zones were the sites of repeated seismic ruptures, as attested by multiple generations of pseudotachylyte veins. Given the complexity of cross-cutting relationships, it is difficult to establish a chronology of seismic ruptures within each

fault zone and also between fault zones. It is particularly impossible to establish a chronology between flat-lying and steeply-dipping fault zones. However, the latest recorded seismic ruptures are those corresponding to flat-lying or gently dipping, thin (< 5 mm) pseudotachylyte veins displacing all pre-existing veins in either type of fault zones. These veins are rarely associated with cataclasite, hence their sharp boundaries. The ubiquity of these most recent veins suggests that activity of the flat-lying fault zones, especially that beneath φ_2 (which contains a lot of such late veins), lasted for a longer time than that of the steeply dipping fault zones.

3.1.3. Pseudotachylyte in the peridotite unit: microscopic observations

The peridotite pseudotachylyte fault veins are of two types: microlitic and annealed (Fig. 7). Unlike the annealed type which is found only in the flat-lying fault zone beneath φ_2 , the microlitic type is found in all vein types, whatever their relative chronology (early as well as late veins).

The microlitic type is characterized by abundant microlites embedded in a brownish amorphous or crypto-crystalline matrix (Fig. 7A, B and C). Microlites have dendritic shapes, with sizes between $1\ \mu\text{m}$ and $120\ \mu\text{m}$. They commonly draw a zonation in veins, with no or very small microlites at the vein margins and large microlites in the median part of the veins. Such zonations likely reflect quenching at the vein margins. Microlites observed in the thickest veins ($> 5\text{cm}$) have sizes up to $1.5\ \text{mm}$ and display spinifex textures. Microlites consist mostly of olivine. Diopside and enstatite are less common. ‘Survivor’ clasts, about 10% in volume, consist mostly of monocrystalline olivine and minor pyroxene. Some polycrystalline clasts consist of olivine and pyroxene. Clasts of pseudotachylyte are also observed, especially in the flat-lying fault veins beneath φ_2 . The largest clasts are frequently elongated and parallel to the vein walls. Flow folds are common in injection veins, especially at their root. Although most pseudotachylyte vein boundaries are sharp, some are diffuse and show a progressive transition from pseudotachylyte to cataclasites (see below). Such diffuse boundaries are found on one side of the vein wall only, the other showing a sharp transition from pseudotachylyte to host rock. Some microlitic veins, especially flat-lying ones located near the boundary between intact peridotite and surrounding serpentinite, are serpentinitized, but the textures, including microlites and clasts, are preserved, showing that serpentinitization post-dates melting.

The annealed pseudotachylyte type (Fig. 7D and E) was observed only in the flat-lying fault zone of locality 4. The annealed type fault veins have thicknesses between 0.8 and 1.5 cm. They are crossed by numerous cooling cracks perpendicular to the vein walls. The matrix consists of entirely recrystallized olivine with a granoblastic annealed texture. The crystal size is homogeneous, except at the margins, with a mean size of *ca.* 10 μm . Crystal junctions are triple and typically define 120° angles. Chilled margins are thin (< 0.5 mm thick). Survivor clasts are rare (< 5% in volume) and consist of monocrystalline or polycrystalline olivine. Injection veins are not entirely recrystallized, their central part showing a microlitic texture. This suggests that the annealed-type veins are originally microlitic type veins that were subsequently recrystallized.

An important characteristic of the peridotite fault veins, especially the early ones, is that they are frequently flanked, on either side, by cataclastic peridotite (Figs 8 and 9). Cataclasite can be found in association with late veins from flat-lying fault zones, but never with late veins from the steeply dipping fault zones. Where cataclasite is present, a progressive evolution from proto-cataclasite to cataclasite and to ultra-cataclasite can be observed. The cataclasite usually remains on the same side of the fault vein, but can also shift to the opposite side, through a progressive decrease of the cataclastic domain on one side and a correlative increase on the other side.

The kinematics determined from the observation of outcrops or of polished hand sample sections (top-to-the-west or top-to-the-northwest displacement senses) are also observed at the thin section scale. Figure 10 shows examples of criteria of displacement senses associated with early and late veins of steeply dipping fault zones and with late veins from flat-lying fault zones.

3.2. Pseudotachylyte in the metagabbro unit

In the metagabbro unit, pseudotachylyte veins are common in the mylonitic sole and abundant just above it (in the equant metagabbro). Their density decreases upwards. At a distance larger than 300 m above ϕ_2 , no more veins can be found. As already noted by Andersen and Austrheim (2006), within the mylonitized sole, some veins were formed *before* mylonitization and others were formed *after* it (Fig. 11).

3.2.1. Pre-mylonitization pseudotachylyte in the foliated sole

In the foliated metagabbro, dark bluish to blackish fault veins are parallel or slightly oblique to the foliation. Their thickness is less than 5 mm. Because of mylonitization-related stretching and pinch-and-swell, they show a poor lateral continuity and cannot be followed over more than a few centimeters. In some instances, they are flanked by injection veins which are offset by millimeter-thick shear zones parallel to the foliation. No clear kinematics is associated with the pre-mylonitization veins.

Most textures typical of pseudotachylyte are obliterated by the penetrative foliation (Fig. 12). Flow structures, chilled margins and microlites are no longer recognizable. Newly formed minerals are aligned in the foliation and consist of glaucophane, albite, epidote and ilmenite. Survivor clasts, mostly plagioclases, are elongated in the foliation and are flanked with pressure shadows. They are smaller (maximum 50 μm , 30 μm on average) than the clasts in the non-mylonitized veins.

3.2.2. Post-mylonitization pseudotachylyte in the foliated sole

Post-mylonitization fault veins are greenish to greyish and are not foliated (Fig. 11). Most fault veins are parallel or nearly parallel to the foliation. The parallelism between most of the fault veins and the foliation likely reflects a mechanical influence of the foliation on the seismic rupture propagation and hence on the resulting attitude of the veins. Post-mylonitization fault veins as well as injection veins are also found to cross-cut pre-mylonitization veins. No clear kinematics can be attributed to the post-mylonitization pseudotachylyte veins.

Post-mylonitization fault veins are usually less than 5 mm thick, but can reach 10 mm. Clasts consist mostly of plagioclase, with minor tremolite or pyroxene. The largest clasts, especially the pyroxene ones, show embayments. The matrix is cryptocrystalline or glassy, and includes microlites of albite and glaucophane. The length of the microlites is between 10 and 20 μm , but decreases to less than 5 μm near the vein boundaries (chilled margins). Flow folds are abundant and are defined by alternating layers of microlites of different sizes. Injection veins are rare. They include less clasts than the fault veins do. The cryptocrystalline or glassy matrix contains microlites which are slightly larger (40 μm) than in the fault veins. The mineralogical nature of clasts or microlites is the same as in the fault veins.

3.2.3. Pseudotachylyte in the equant metagabbro above the foliated sole

The appearance of pseudotachylyte veins in the equant metagabbro is similar to that of post-mylonitization veins in the foliated sole. In particular, cross-cutting relationships between veins are common and indicate polyphase seismic rupturing. A notable difference is that lenses of pseudotachylyte-supported breccias are observed in equant metagabbro but not in the foliated sole. These lenses are up to 15 cm thick and include rounded fragments of equant metagabbro and foliated metagabbro. They are interpreted as local accumulations in so-called dilational or releasing bends located along slipping surfaces (Sibson, 1986). Another difference is that the fault vein attitudes in the equant metagabbro are scattered. Particularly, veins parallel or slightly oblique to ϕ_2 are scarce.

The metagabbro pseudotachylyte matrix is glassy or crypto-crystalline. Using X-ray diffraction synchrotron, Deseta et al. (2014b) produced Laue patterns without diffraction points, showing unambiguously the presence of glass in the matrix. Survivor clasts are well rounded and show the same mineralogical nature as in post-mylonitization veins. Feldspar clasts are numerous and small whereas pyroxene and olivine clasts are scarce and large. Microlites consist mainly of fibrous or acicular, rarely spherulitic, pyroxene. Deseta et al. (2014b) report the presence of blueschist facies microlites, namely Al-rich omphacite, high-Fe anorthite and accessory ilmenite. A few thin sections show different microlite assemblages, suggesting different stages of vein formation. Some veins contain omphacite but no tremolite or actinolite, some others contain tremolite or actinolite but no omphacite. Omphacite-bearing veins likely formed under high-pressure conditions while amphibole-bearing veins formed in shallower conditions (greenschist facies). Unfortunately, such observations are too scarce to allow a reliable sorting of the equant metagabbro veins.

Pseudotachylyte veins in the equant metagabbro and post-mylonitization veins in the foliated sole can be considered as contemporaneous. However, a part of the veins in the equant metagabbro, especially those cut by other veins, could be older and could have been formed coevally with the pre-mylonitization veins. Only absolute dating of the veins could help clarify the relationships between post-mylonitization veins in the foliated metagabbro and veins in the non-foliated metagabbro.

4. Discussion: Analysis of the seismic ruptures fossilized in the study area and comparison with present-day subduction zone seismology

4.1. Significance of the ϕ_2 contact and formation of metagabbro mylonite and pseudotachylyte

The φ_2 contact is interpreted as an ancient deformed Moho interface by Andersen et al. (2008, 2014). Alternatively, φ_2 can be interpreted as an ancient low-angle detachment fault or shear zone dating back to the initial stretching of the continental lithosphere before formation of the Piemonte-Liguria basin (Meresse et al., 2012), following the model of Manatschal and Müntener (2009). It can also be interpreted as an ancient low-angle detachment fault located at or near the accretion ridge of the Piemonte-Liguria basin (Vitale-Brovarone et al., 2014) following the models of Tucholke and Lin (1994) or Cannat et al. (2009). If φ_2 is an ancient detachment (at the ridge or in the ocean-continent transition), it may not coincide with the Moho interface. The supposed detachment could have been reactivated during plate convergence or during subsequent collision and nappe emplacement. Whatever its origin, the detachment should have been localized beneath the bulging sides of a gabbro diapiric pluton emplaced in the uppermost mantle, to account for the gabbro-over-peridotite succession observed in the study area. Lastly, the two scenarios (reactivated Moho or detachment in the uppermost mantle) are not contradictory. Both involve a low-angle shear zone in the upper part of the lithosphere.

The ductile deformation of the metagabbro sole may result from one or several of the following settings: (1) normal shear along a crustal-scale detachment following continental lithosphere breakup (e.g., Meresse et al., 2012; Vitale-Brovarone et al., 2014), (2) normal shear along an axial detachment fault near the spreading ridge of the Piemonte-Liguria basin, (3) reverse shear along the crust-mantle boundary (Moho) of the subducting slab. The lack of kinematic indicators in the metagabbro sole prevents distinguishing stages (1) or (2) (normal sense of shear) from stage (3) (reverse sense of shear).

Whatever the setting, the ductile deformation was achieved at a place where the ambient temperature was higher than the brittle/ductile transition temperature of gabbro. Since the plagioclase modal content in the metagabbro is about 50%, a minimum estimate of the brittle/ductile transition temperature is *ca.* 550°C (e.g., Molli, 1994; Hansen et al., 2013). Given the abundance of pyroxenes (about 35 % modal content), the actual transition temperature of the metagabbro is likely higher. The maximum metamorphic temperatures in the study area are 414-471°C (Vitale-Brovarone et al., 2013), that is less than 550°C. This could rule out the possibility of a ductile deformation in the subducting slab. However, by admitting that the metagabbro underwent some hydrothermal alteration (as can be expected if φ_2 is a reactivated detachment), its brittle/ductile transition temperature would have been lower (*ca.* 300°C, Stünitz, 1993), thus permitting ductile deformation.

Calling upon ductile deformation of the metagabbro sole along a detachment in the ocean-continent transition or at the mid-oceanic ridge (and hence generation of pre-mylonitization pseudotachylyte at the same place) requires that these early structures will then be transported until the subduction zone at intermediate depths where they will be overprinted by (post-mylonitization) pseudotachylyte veins. In particular, it means that (pre-mylonitization) pseudotachylyte veins formed away (200~600 km away according to Guerrera et al., 1993; Stampfli et al., 1998; Rosenbaum et al., 2002; Marroni and Pandolfi, 2007; Turco et al., 2012) from the subduction zone will be overprinted by (post-mylonitization) pseudotachylyte veins formed at depth (> 60 km) in a subduction zone. Though not impossible, such a coincidence does not seem very plausible.

Figure 13 suggests a simpler scenario in which both pseudotachylyte and mylonite are formed in the subducting slab at shallow to intermediate depths. Pre-mylonitization pseudotachylyte is formed at depths shallower than the gabbro brittle/ductile transition isotherm. It can be formed near the trench, along the Moho (Fig. 13A), following a scenario proposed by Singh et al. (2008, 2011) to account for the location of the hypocenters and rupture propagation geometries of the 2004 Sumatra and 2010 Pagai events. A similar possibility of seismic ruptures inside the oceanic crust of the subducting Philippine Sea plate beneath SW Japan was also suggested by Tsuji et al. (2009, 2013). Pre-mylonitization pseudotachylyte can alternatively be formed deeper, before being subsequently mylonitized when passing through the brittle/ductile transition isotherm. Continuing shear along the crust-mantle boundary would result in φ_2 -parallel seismic ruptures in the brittle peridotite and in φ_2 -parallel foliation in the metagabbro sole (Fig. 13B). That post-mylonitization pseudotachylyte cross-cuts the foliated metagabbro suggests oscillations of the gabbro brittle/ductile transition isotherm. Such oscillations are possible. Indeed, numerical simulations of long-term equilibrium state of the subduction interplate show that the brittle/ductile transition is almost parallel to the crust-mantle boundary of the subducting slab (Arcay, 2012). An alternative view is that post-mylonitization pseudotachylyte veins, especially those perpendicular or highly oblique to the foliation, result from seismic ruptures having nucleated in the underlying mantle and having propagated upwards across the ductile metagabbro (Fig. 13D).

4.2. Formation of peridotite pseudotachylyte

4.2.1. Weakening mechanisms facilitating seismic ruptures at intermediate depths

505 Intermediate-depth seismicity as well as deep-focus seismicity are puzzling. Indeed, given the
506 high stresses expected at depths > 60 km, brittle fracturing or frictional sliding along pre-
507 existing fractures require unrealistic rock strengths or over-pressurized pore fluids which
508 could reduce stresses. Yet earthquakes occur. To solve this paradox, three mechanisms have
509 been proposed (Green and Houston, 1995; Hacker et al., 2003; Frohlich, 2006; Houston,
510 2015): dehydration embrittlement, ductile shear instability and transformational faulting.

511 Transformational faulting calls for the formation of anticracks during phase transformation of
512 olivine to denser phases such as β - or γ -spinel (Green and Burnley, 1989; Kirby et al., 1991;
513 Wiens et al., 1993; Schubnel et al., 2013). The applicability of transformational faulting as a
514 possible mechanism accounting for *intermediate-depth* seismicity in the subducting oceanic
515 lithosphere is questionable (e.g., Hacker et al., 2003) because (1) the expected reactions are
516 too slow compared with earthquake timescales, (2) olivine remains stable at the considered
517 depths, and (3) the metamorphism of basalts or gabbros does not involve the polymorphic
518 reactions required in transformational faulting processes.

519 Dehydration embrittlement is based on a pore fluid pressure increase leading to fracture
520 formation or reactivation by decrease of the otherwise high normal stresses. The pore fluid
521 pressure increase would result from fluid accumulations following dehydration reactions of
522 hydrated minerals. Seismological observations and data modeling, laboratory experiments,
523 field observations and thermal and thermodynamic computations have pointed out possible
524 links between intermediate-depth seismicity and dehydration reactions affecting hydrated
525 rocks or minerals of the subducting slab such as basalt, gabbro, chlorite, antigorite, talc or
526 brucite (Raleigh and Paterson, 1965; Rutter and Brodie, 1988; Green and Burnley, 1989;
527 Green et al., 1990; Green and Houston, 1995; Kirby, 1995; Seno and Yamanaka, 1996; Davis,
528 1999; Peacock, 2001; Seno et al., 2001; Dobson et al., 2002; Wang, 2002; Hacker et al., 2003;
529 Preston et al., 2003; Yamazaki and Seno, 2003; Jung et al., 2004; Wang et al., 2004;
530 Brudzinski et al., 2007; Hirose and Bystricky, 2007; Rondenay et al., 2008; Hasegawa et al.,
531 2009; Nakajima et al., 2009; Angiboust et al., 2012; Abers et al., 2013; Nakajima et al., 2013;
532 Houston, 2015). A strong argument justifying the link between intermediate-depth seismicity
533 and dehydration reactions in the subducting slab lies in overlaps between predicted
534 dehydration reaction isotherms and location of hypocenters (e.g., Peacock, 2001 or Hacker et
535 al., 2003). Regarding the *upper* Wadati-Benioff seismic sub-zone, the source of fluids would
536 lie in dehydration reactions transforming basalts or gabbros into blueschist or eclogite-facies
537 rocks (Hacker et al., 2003; Preston et al., 2003; Yamazaki and Seno, 2003; Kita et al., 2006;
538 Nakajima et al., 2009). Regarding the *lower* Wadati-Benioff sub-zone, the source of fluids

should be searched for in dehydration reactions of chlorite or antigorite since these two minerals are thought to be present in the mantle of subducting slabs (Seno and Yamanaka, 1996; Peacock, 2001; Seno et al., 2001; Hacker et al., 2003). Dehydration reactions have been invoked to account for secondary olivine crystallization along natural fault zones in ophiolitic rocks from the Voltri complex in Italy (Hoogerduijn-Strating and Vissers, 1991; Scambelluri et al., 1991). According to Hoogerduijn-Strating and Vissers (1991), fluid overpressures would nearly reach lithostatic values.

Ductile instability, also called thermal runaway or thermal shear instability, postulates that the temperature-dependent viscosity of a highly localized, ideally fine-grained, creeping ductile shear zone is progressively reduced by the heat provided by continuing creep (Ogawa, 1987; Kameyama et al., 1999; Braeck and Podladchikov, 2007; Kelemen and Hirth, 2007). This positive feedback between continuing creep and temperature rising can eventually lead to seismic failure. This mechanism was evoked in the case of the 1994 M_w 8.2 Bolivian deep-focus earthquake (depth = 637 km) by Kanamori et al. (1998), who further suggested that failure could have led to melting along the newly nucleated fault surface. Indeed, these authors calculated that, for a starting shear zone thinner than 1 cm, temperature elevation in the shear zone would exceed 10,000°C, which is far higher than melting temperature of any rock. Source parameter scaling and energy budget of clusters of intermediate-depth (140-160 km) M_w 4-5 earthquakes beneath northern Colombia led Prieto et al. (2013) to suggest that propagation of these ruptures was caused by a thermal runaway mechanism. A similar mechanism was suggested by Wiens and Snider (2001) to account for deep (550-600 km) earthquakes in the Tonga slab. From the analysis of closely associated mylonitic zones and pseudotachylyte across a gabbro metamorphosed under high- to ultrahigh-pressure conditions in Norway, John et al. (2009) suggested that co-seismic melting was contemporaneous with ductile shear and is the result of a self-localizing thermal runaway process along the shear zones. Another example of possible thermal runaway frozen in the geological record is provided by Andersen et al. (2008, 2014) and Deseta et al. (2014a), as is discussed below.

4.2.2. Did dehydration embrittlement facilitate seismic failure in the peridotite unit?

According to Deseta et al. (2014a), the metagabbro pseudotachylyte was formed under eclogite facies P-T conditions (presence of omphacite microlites in the matrix) before being retrogressed under blueschist facies P-T conditions (presence of glaucophane microlites). These authors estimate that the P-T conditions of crystallization of glaucophane and omphacite microlites in the metagabbro pseudotachylyte are between 430 and 550°C and 1.8

GPa and 2.6 GPa, corresponding to blueschist to eclogite facies conditions. These values are close to those obtained in the units around the Cima di Gratera nappe by Vitale-Brovarone et al. (2013) which are 414-471°C and 1.9-2.6 GPa. With a mean rock density of 3000 kg/m³ and assuming a lithostatic equilibrium, the pressure range corresponds to depths between 60 and 90 km. P-T conditions in the peridotite pseudotachylyte cannot be ascertained. Indeed, the microlites (diopside, olivine, enstatite and clinocllore) that crystallized during cooling of the melt do not bring any constraints on the pressure conditions during pseudotachylyte formation. However, if considering the peridotite unit as *attached* to the metagabbro unit during subduction, then the same metamorphic conditions should apply to both units. In other words, it can be assumed that pseudotachylyte in the peridotite unit formed under blueschist to lawsonite-eclogite facies conditions, as suggested by Deseta et al. (2014a).

As mentioned above, antigorite, which is common in the serpentinites or serpentized peridotites of the Cima di Gratera nappe, is a candidate to account for dehydration embrittlement and subsequent seismicity in subducting slabs. The P-T conditions for antigorite dehydration are known from experiments and are between 550 and 720°C for pressures between 1 and 3 GPa (Ulmer and Trommsdorff, 1995; Wunder and Schreyer, 1997; Dobson et al., 2002; Perrillat et al., 2005; Hilairet et al., 2006; Padron-Navarta et al., 2010). The peak temperatures supposedly recorded by the peridotite pseudotachylyte (430-550°C, see above) or in the surrounding units (414-471°C, Vitale-Brovarone et al., 2013) are lower than the temperatures required for dehydration of antigorite-bearing serpentinite. This temperature difference renders dehydration of the mantle unit antigorite unlikely. In addition to these temperature issues, no optical microscope or SEM observations of secondary olivine newly crystallized at the expense of primary antigorite, such as the assemblages described by Hoogerduijn-Strating and Vissers (1991) or by Scambelluri et al. (1991), could be found in the pseudotachylyte veins or in their vicinity, confirming that antigorite dehydration did not occur in the peridotite. Andersen et al. (2014) and Deseta et al. (2014b) confirm that they did not observe secondary anhydrous minerals resulting from the dehydration of serpentine, talc, clinocllore or amphibole in the Cima di Gratera metagabbros or peridotites. For the time being, we consider that dehydration embrittlement was not an operative mechanism during seismic failure in the peridotite unit.

4.2.3. Did self-localizing thermal runaway facilitate seismic failure in the peridotite unit?

Deseta et al. (2014a) report thin-section scale (20 mm to < 100 µm range) ductile (plastic) deformation structures inside or along pseudotachylyte fault veins. These structures include

(1) elongated wallrock clasts in gabbro-hosted pseudotachylyte veins, (2) crystal plastic deformation of the host metagabbro along the boundaries of pseudotachylyte veins, (3) grain boundary alignment in prolate and lozenge-shaped grains suggesting grain boundary migration in some peridotite-hosted pseudotachylyte veins, and (4) plastic ribbons in gabbro- or peridotite-hosted ultracataclasites. Additionally, by analyzing dislocation slip systems in olivine from peridotite wall rock or from clasts in the pseudotachylyte with the help of electron backscatter diffraction, Andersen et al. (2014) and Deseta et al. (2014b) suggested that ductile deformation preceded pseudotachylyte formation. Based on these microscale ductile precursors to seismic faulting found along pseudotachylyte fault veins in the Cima di Gratera peridotites and gabbros, Andersen et al. (2014) and Deseta et al. (2014a and b) suggested that seismic ruptures were facilitated or triggered by a self-localizing thermal runaway process.

The possible activity of this process in the basal foliated metagabbro sole cannot be demonstrated nor discarded. Indeed, the parallelism between (post-mylonitization) pseudotachylyte fault veins and the foliation of the basal metagabbro probably results from the influence of the pre-existing foliation on the propagation of the seismic rupture, as often invoked in other settings (e.g., Grocott, 1981; Swanson, 1988; Allen, 2005; Zechmeister et al., 2007). More generally, pseudotachylyte veins preserved inside mylonitic zones are quite common (e.g., Sibson, 1980; Passchier, 1982), and their formation, although influenced or guided by the pre-existing planar heterogeneity as stated above, does not necessarily depend on a precursory softening shortly before seismic rupturing, as required in the ductile instability mechanism.

Unlike Deseta et al. (2014a and b), we did not observe any ductile shear zones along the peridotite fault veins, despite a large number of thin sections prepared with samples from the peridotite unit. We rather observe a quasi-ubiquitous association of pseudotachylyte veins with cataclastic peridotite. Consequently, ductile instability does not appear as a predominant mechanism associated with seismic ruptures in the peridotite unit of the study area.

4.2.4. Did cataclasis facilitate seismic failure in the peridotite unit?

Peridotite-hosted pseudotachylyte fault veins are almost always flanked by cataclasite (Figs 8, 9 and 10). Cataclasis may predate or postdate frictional melting, as shown by cataclasite zones crossed by pseudotachylyte veins or by fragments of pseudotachylyte included in cataclasites. Similar pseudotachylyte-cataclasite associations were reported from natural occurrences (Maddock, 1992; Magloughlin, 1992; Swanson, 1992; McNulty, 1995; Obata and Karato,

1995; Curewitz and Karson, 1999; Fabbri et al., 2000; Rowe et al., 2005; Di Toro and Pennacchioni, 2004, 2005; Piccardo et al., 2007, 2010) and also from rock friction experiments (Spray, 1995; Del Gaudio et al., 2009; Hirose et al., 2012). Swanson (1992) considered the cataclasite as the result of the propagating seismic rupture front, frictional melting occurring during seismic slip behind the front. Curewitz and Karson (1999) proposed that cataclasite results from slip surface leveling by asperity grinding and abrasion. Since a cataclastic peridotite is obviously mechanically weaker than an intact peridotite, one can expect cataclasis to be a precursory weakening mechanism facilitating ensuing seismic rupture. Unlike dehydration-derived over-pressurized fluids, cataclasis *per se* does not contribute to counterbalance the high stresses expected at depths > 60 km. Though this mechanism does not bring any answer to the enigma of earthquakes at great depths, it however provides a plausible way to mechanically weaken strong rocks.

4.3. Co-seismic displacement kinematics frozen in the peridotite unit compared with present-day Wadati-Benioff zone earthquakes

Figure 14 shows a comparison between co-seismic kinematics frozen in the peridotite unit and the intermediate-depth seismicity of the Pacific plate presently subducting beneath NE Japan. The Pacific plate is taken as representative of a cold slab. It is also a slab for which high-resolution seismological data are available. The choice of the intermediate-depth seismicity is justified by the fact that the peridotite-hosted pseudotachylyte were formed at depths between 60 and 90 km, as suggested by the metamorphic pressure conditions in the overlying metagabbro unit, supposed attached to the underlying peridotite unit. Since the kinematics of gabbro-hosted pseudotachylyte are undetermined, the discussion will be largely based on peridotite-hosted occurrences, with the assumption of an Alpine-type east-dipping subduction (Section 2.1).

Given the large (55°) angle between flat-lying fault zones and the earliest veins of the steeply-dipping fault zones in the peridotite unit, no unique stress tensor can account for simultaneous slip along the two zones, suggesting that when one fault zone was active, the other was not active. The pervasive intermingling between flat-lying and steeply-dipping veins suggests that the two types of fault zones were active alternatively and under oscillating stress conditions. Such changing stress conditions could result from the near-surface interplate seismic cycle and periodic unlocking of the shallow plate interface during large earthquakes as suggested by Astiz et al. (1988).

Several authors showed that in the case of cold slabs, the upper surface of the intermediate-depth Wadati-Benioff zone is characterized by a so-called downdip compression, meaning that P axes of earthquakes are parallel or almost parallel ($< 30^\circ$) to the dipping direction of the subducting slab (Isacks and Molnar, 1971; Apperson and Frohlich, 1987; Green and Houston, 1995; Kao and Liu, 1995; Igarashi et al., 2001; Chen et al., 2004). The reverse kinematics observed in the steeply-dipping fault zones agree with this configuration. Indeed, in slab coordinates, a 55° dipping earthquake fault plane would have a ‘favorably’ oriented P axis at 30 to 45° to the fault plane, that is, at 10° to 25° to the slab upper surface (taken parallel to ϕ_2).

Refining the comparison with cold subducting plates, a series of events having occurred in the early 2000s in the Pacific plate beneath NE Japan gives some insights on the geometry and sense of slip of the Corsican paleo-ruptures (Fig. 14B and C). First, the November 3, 2002, M 6.1 earthquake which occurred along the upper Wadati-Benioff plane beneath NE Japan (Okada and Hasegawa, 2003) can constitute an analog to seismic ruptures along *flat-lying* fault zones in the peridotite unit or in the base of the metagabbro unit, on either side of ϕ_2 . Indeed, the actual fault plane of this event was parallel to the alignment defining the upper Wadati-Benioff plane (i.e., parallel to the crust-mantle boundary) and the sense of slip was reverse. The analogy is somewhat limited by the shallow focal depth (38 km) of this event (Hasegawa et al., 2007), meaning that it is not strictly speaking an intermediate-depth earthquake. Second, the May 26, 2003 M 7.1 earthquake that occurred in the Pacific plate (focal depth 68 km) can be an analog to seismic ruptures along *steeply-dipping* fault zones in the peridotite unit. This event was located near the upper Wadati-Benioff plane, close to the crust-mantle boundary (Okada and Hasegawa, 2003). Its hypocenter was 50 km away from the November 3, 2002 event. Okada and Hasegawa (2003) further showed that aftershocks were distributed along the steeply-dipping nodal plane, straddling both the oceanic crust and the uppermost mantle. Sense of slip was reverse. The angle between the fault plane and the crust-mantle boundary is 50° (Hasegawa et al., 2007), a value quite comparable to the 55° angle between the flat-lying and the steeply dipping fault zones observed in the peridotite unit. The analogy between the Pacific slab upper Wadati-Benioff seismic events and the Corsican configuration is depicted on Fig. 14. Slab-boundary parallel seismic ruptures would be analogous of the November 3, 2002 earthquake off NE Japan and, more generally, could be deeper equivalents of the low-angle thrust fault (“LT”) type events of Igarashi et al. (2001).

In summary, the reverse senses associated with the seismic ruptures frozen in the study area can be compared with earthquakes with reverse-type focal mechanisms or ‘down-dip compression’ events. More particularly, the lack of normal kinematics associated with

pseudotachylyte generation suggests that normal-type events such as those corresponding to reactivation at depth of normal faults formed in the slab before it starts subducting (e.g., Jiao et al., 2000; Barnhart et al., 2014) either did not occur in the Corsican subduction zone or did not leave any imprint.

5. Conclusion

The structural analysis of pseudotachylyte in the Cima di Gratera ophiolitic nappe leads to the following results.

Pseudotachylyte veins in the peridotite unit are either isolated and scattered in the unit or clustered in fault zones. Isolated veins as well as fault zones are horizontal (flat-lying type) or dip about 55° (steeply dipping type). In fault zones, the abundance of fault veins likely reflects a large number of repeating seismic ruptures, among which some may correspond to small magnitude events like aftershocks. The lack of clear cross-cutting relationships suggests that the flat-lying fault zones and their steeply dipping equivalents were active alternatively, as a consequence of oscillating stress states possibly resulting from periodic unlocking of the shallow plate interface during the seismic cycle. The activity of the flat-lying fault zones probably lasted for a longer time than the steeply dipping fault zones. The sense of displacement associated with steeply dipping fault zones and with most of the flat-lying fault zones is top-to-the-west or top-to-the northwest. Cataclasite flanking most of the veins was formed before or coevally with frictional melting and likely mechanically weakened the peridotite, facilitating subsequent seismic rupture.

The base of the metagabbro unit is mylonitic. The origin of the mylonitization remains undetermined. The scenario retained here suggests that the ductile deformation was achieved in the subducting slab below the brittle-ductile transition depth of the gabbro. The isotherm corresponding to the brittle-ductile transition depth of the gabbro can be taken as parallel to the crust-mantle boundary in the subducting slab. Pseudotachylyte veins in the metagabbro are distributed in the lower part of the unit, in the foliated sole as well as in the equant metagabbro above. They are not as well organized as their equivalents in the peridotite unit. Flat-lying veins are abundant near the contact with the underlying peridotite, steeply dipping veins are scattered in the lower part of the unit. The ductile deformation affecting the base of the unit allows to distinguish pre-mylonitization pseudotachylyte formed above the brittle-ductile transition depth of the gabbro from post-mylonitization veins formed below this depth. In the equant metagabbro, it is no longer possible to distinguish more than one episode of pseudotachylyte formation. No information regarding the sense of displacement associated

with seismic ruptures could be retrieved from gabbro pseudotachylyte veins, whatever their positions or attitudes. It is furthermore not possible to establish a relative chronology between pseudotachylyte formed on either side of the contact ϕ_2 between the peridotite and metagabbro units.

Depth constraints provided by the metamorphic conditions recorded by metagabbro pseudotachylyte (1.9-2.6 GPa pressure range, Deseta et al., 2014 a and b) and geometry as well as kinematics data from peridotite pseudotachylyte show similarities with well documented seismic ruptures occurring in the Wadati-Benioff zone of the Pacific plate beneath NE Japan. These similarities allow to propose a scenario of formation of pseudotachylyte which encompasses shallow seismic ruptures along the crust-mantle boundary as suggested by Singh et al. (2008) for the 2004 Sumatra earthquake and deeper ruptures in the Wadati-Benioff zone (60-100 km depth range). In this scenario, seismic ruptures in the subducting mantle would always occur under brittle conditions while those in the lower part of the subducting crust would partly be coeval with ductile deformation. No relative chronology between pseudotachylyte of either side of ϕ_2 (peridotite vs. metagabbro) can be firmly established. A part of the metagabbro post-mylonitization veins could be the result of large ruptures nucleated in the peridotite unit and propagated upward across ϕ_2 and through a metagabbro sole under ductile conditions.

Deciphering intermediate-depth seismicity from ophiolite-hosted pseudotachylyte is a complex task because rocks may have recorded earthquakes elsewhere than in the subducting slab. The rocks may have been deformed, at least partly, at the axial ridge during oceanic accretion or in the continent-ocean transition, during initial crustal thinning. Final ophiolite emplacement, whatever by obduction or collision, and subsequent episodes (e.g., late- to post-orogenic extension) are also responsible for additional deformation. All these deformation episodes may contribute to clutter the final picture. However, like for the continental lithosphere (Swanson, 1992; Obata and Karato, 1995; Allen, 2005; Di Toro and Pennacchioni, 2005; Ueda et al., 2008), pseudotachylyte is a valuable tool to improve our understanding of the mechanics of the seismic ruptures in the oceanic lithosphere, in complement to geophysical studies.

Acknowledgements.

This work was funded by the CNRS-INSU TelluS-ALEAS program. Electronic microscopy was supported by the RENATECH Network. Reviews by G. Di Toro and an anonymous reviewer significantly helped clarifying several points of the paper.

REFERENCES

- Abers, G.A., Nakajima, J., van Keken, P.E., Kita, S., Hacker, B.R., 2013. Thermal-petrological controls on the location of earthquakes within subducting plates. *Earth and Planetary Science Letters* 369-370, 178-187, 10.1016/j.epsl.2013.03.022.
- Agard, P., Vitale-Brovarone, A., 2013. Thermal regime of continental subduction: The record from exhumed HP-LT terranes (New Caledonia, Oman, Corsica). *Tectonophysics* 601, 206-215, 10.1016/j.tecto.2013.05.011.
- Allen, J.L., 2005. A multi-kilometer pseudotachylyte system as an exhumed record of earthquake rupture geometry at hypocentral depths (Colorado, USA). *Tectonophysics* 402, 37-54.
- Andersen, T., Austrheim, H., 2006. Fossil earthquakes recorded by pseudotachylytes in mantle peridotite from the Alpine subduction complex of Corsica. *Earth and Planetary Science Letters* 242, 58-72.
- Andersen, T.B., Mair, K., Austrheim, H., Podladchikov, Y.Y., Vrijmoed, J.C., 2008. Stress release in exhumed intermediate and deep earthquakes determined from ultramafic pseudotachylyte. *Geology* 36, 995-998.
- Andersen, T.B., Austrheim, H., Deseta, N., Silkose, P., Ashwal, L.D., 2014. Large subduction earthquakes along the fossil Moho in Alpine Corsica. *Geology* 42, 395-398.
- Angiboust, S., Agard, P., Yamato, P., Raimbourg, H., 2012. Eclogite breccias in a subducted eclogite: A record of intermediate-depth earthquake? *Geology* 40, 707-710.
- Apperson, K.D., Frohlich, C., 1987. The relationship between Wadati-Benioff zone geometry and *P*, *T* and *B* axes of intermediate and deep focus earthquakes. *Journal of Geophysical Research* 92, 13821-13831.
- Arcay, D., 2012. Dynamics of interplate domain in subduction zones: influence of rheological parameters and subducting plate age. *Solid Earth* 3, 467-488.
- Astiz, L., Lay, T., Kanamori, H., 1988. Large intermediate-depth earthquakes and the subduction process. *Physics of the Earth and Planetary Interiors* 53, 80-166.
- Austrheim, H., Andersen, T.L., 2004. Pseudotachylytes from Corsica: Fossil earthquakes from subduction complex. *Terra Nova* 16, 193-197, 10.1111/j.1365-3121.2004.00551.x.
- Barnhart, W.D., Hayes, G.P., Samsonov, S.V., Fielding, E.J., Seidman, L.E., 2014. Breaking the oceanic lithosphere of a subducting slab: The 2013 Khash, Iran earthquake. *Geophysical Research Letters* 41, 32-36, 10.1002/2013GL058096.
- Braeck, S., Podladchikov, Y.Y., 2007. Spontaneous thermal runaway as an ultimate failure mechanism of materials. *Physical Review Letters* 98, 095504.
- Brudzinski, M.R., Thurber, C.H., Hacker, B.R., Engdahl, E.R., 2007. Global prevalence of double Benioff zones. *Science* 316, 1472-1474, 10.1126/science.1139204.

799 Brunet, C., Monié, P., Jolivet, L., Cadet, J.P., 2000. Migration of compression and extension in the Tyrrhenian Sea, insights
800 from $^{40}\text{Ar}/^{39}\text{Ar}$ ages on micas along a transect from Corsica to Tuscany. *Tectonophysics* 321, 127-155.

801 Cannat, M., Sauter, D., Escartin, J., Lavier, L., Picazo, S., 2009. Oceanic corrugated surfaces and the strength of the axial
802 lithosphere at slow spreading ridges. *Earth and Planetary Science Letters* 288, 174-183.

803 Chen, P.F., Bina, C.R., Okal, E.A., 2004. A global survey of stress orientations in subducting slabs as revealed by
804 intermediate-depth earthquakes. *Geophysical Journal International* 159, 721-733, 10.1111/j.1365-246X.2004.02450.x.

805 Curewitz, D., Karson, J.A., 1999. Ultracataclasis, sintering, and frictional melting in pseudotachylytes from East Greenland.
806 *Journal of Structural Geology* 21, 1693-1713.

807 Davis, J.H., 1999. The role of hydraulic fractures and intermediate-depth earthquakes in generating subduction-zone
808 magmatism. *Nature* 398, 142-145.

809 Del Gaudio, P., Di Toro, G., Han, R., Hirose, T., Nielsen, S., Shimamoto, T., Cavallo, A., 2009. Frictional melting of
810 peridotite and seismic slip. *Journal of Geophysical Research* 114, B06306, 10.1029/2008JB005990.

811 Deseta, N., Andersen, T.B., Ashwal, L.D., 2014a. A weakening mechanism for intermediate-depth seismicity? Detailed
812 petrographic and microtextural observations from blueschist facies pseudotachylytes, Cape Corse, Corsica. *Tectonophysics*
813 610, 138-149.

814 Deseta, N., Ashwal, L.D., Andersen, T.B., 2014b. Initiating intermediate-depth earthquakes: Insights from a HP-LT ophiolite
815 from Corsica. *Lithos* 206-207, 127-146.

816 Di Toro, G., Pennacchioni, G., 2004. Superheated friction-induced melts in zoned pseudotachylytes within the Adamello
817 tonalites (Italian Southern Alps). *Journal of Structural Geology* 26, 1783-1801.

818 Di Toro, G., Pennacchioni, G., 2005. Fault plane processes and mesoscopic structure of a strong-type seismogenic fault in
819 tonalites (Adamello batholith, Southern Alps). *Tectonophysics* 402, 55-80.

820 Dobson, D.P., Meredith, P.G., Boon, S.A., 2002. Simulation of subduction zone seismicity by dehydration of serpentine.
821 *Science* 298, 1407-1410.

822 Durand-Delga, M., Rossi, P., 2002. About the Ligurian-Piemontese Jurassic Ocean on the transect Corsica-Apennines.
823 *Comptes Rendus Géoscience* 334, 227-228.

824 Fabbri, O., Lin, A.M., Tokushige, H., 2000. Coeval formation of cataclasite and pseudotachylyte in a Miocene forearc
825 granodiorite, southern Kyushu, Japan. *Journal of Structural Geology* 22, 1015-1025.

826 Faure, M., Malavieille, J., 1981. Etude structurale d'un cisaillement ductile : le charriage ophiolitique corse dans la région de
827 Bastia. *Bulletin de la Société Géologique de France* 23, 335-343.

828 Fournier, M., Jolivet, L., Goffé, B., Dubois, R., 1991. Alpine Corsica Metamorphic Core Complex. *Tectonics* 10, 1173-1186.

829 Frohlich, C., 2006. Deep earthquakes. Cambridge University Press, Cambridge, 573 p.

830 Green, H.W., Burnley, P.C., 1989. A new self-organizing mechanism for deep-focus earthquakes. *Nature* 341, 733-737.

831 Green, H.W., Young, T.E., Walker, D., Scholz, C.H., 1990. Anticrack-associated faulting at very high pressure in natural
832 olivine. *Nature* 348, 720-722.

833 Green, H.W., Houston, H., 1995. The mechanics of deep earthquakes. *Annual Reviews of Earth and Planetary Sciences* 23,
834 169-213.

835 Grocott, J., 1981. Fracture geometry of pseudotachylite generation zones: a study of shear fractures formed during seismic
836 events. *Journal of Structural Geology* 3, 169-178.

837 Guerrero, F., Martín-Algarra, A., Perrone, V., 1993. Late Oligocene-Miocene syn-/late-orogenic successions in Western and
838 Central Mediterranean Chains from the Betic Cordillera to the Southern Apennines. *Terra Nova* 5, 525-544.

839 Hacker, B.R., Peacock, S.M., Abers, G.A., Holloway S.D., 2003. Subduction factory 2. Are intermediate-depth earthquakes
840 in subducting slabs linked to metamorphic dehydration reactions? *Journal of Geophysical Research* 108,
841 10.1029/2001JB001129.

842 Hansen, L.N., Cheadle, M.J., John, B.E., Swapp, S.M., Dick, H.J.B., Tucholke, B.E., Tivey, M.A., 2013. Mylonitic
843 deformation at the Kane oceanic core complex: Implications for the rheological behavior of oceanic detachment faults.
844 *Geochemistry, Geophysics, Geosystems* 14, doi:10.1002/ggge.20184.

845 Harris, L., 1985. Progressive and polyphase deformation of the Schistes Lustrés in Cap Corse, Alpine Corsica. *Journal of*
846 *Structural Geology* 7, 637-650.

847 Hasegawa, A., Uchida, N., Igarashi, T., Matsuzawa, T., Okada, T., Miura, S., Suwa, Y., 2007. Asperities and Quasi-Static
848 Slips on the Subducting Plate Boundary East of Tohoku, Northeast Japan. In: *The Seismogenic Zone of Subduction Thrust*
849 *Faults*, edited by T.H. Dixon and J.C. Moore, Princeton University Press, 451-475.

850 Hasegawa, A., Umino, N., Takagi, A., 1978a. Double-planed structure of the deep seismic zone in the Northeastern Japan
851 arc. *Tectonophysics* 47, 43-58.

852 Hasegawa, A., Umino, N., Takagi, A., 1978b. Double-planed deep seismic zone and upper-mantle structure in the
853 Northeastern Japan arc. *Geophysical Journal of the Royal Astronomical Society* 54, 281-296.

854 Hasegawa, A., Nakajima, J., Uchida, N., Okada, T., Zhao, D., Matsuzawa, T., Umino, N., 2009. Plate subduction, and
855 generation of earthquakes and magmas in Japan as inferred from seismic observations: An overview. *Gondwana Research* 16,
856 370-400.

857 Hilalret, N., Daniel, I., Reynard, B., 2006. Equation of state of antigorite, stability field of serpentines, and seismicity in
858 subduction zones. *Geophysical Research Letters* 33, L2302, 10.1029/2005GL024728.

859 Hirose, T., Bystricky, M., 2007. Extreme dynamic weakening of faults during dehydration by coseismic shear heating.
860 *Geophysical Research Letters* 34, L14311, 10.1029/2007GL030049.

861 Hirose, T., Mizoguchi, K., Shimamoto, T., 2012. Wear processes in rocks at slow to high slip rates. *Journal of Structural*
862 *Geology* 38, 102-116, 10.1016/j.jsg.2011.12.007.

863 Hoogerduijn-Strating, E.H., Vissers, R.L.M., 1991. Dehydration-induced fracturing of eclogite-facies peridotites:
864 implications for the mechanical behavior of subducting oceanic lithosphere. *Tectonophysics* 200, 187-198.

865 Houston, H., 2015. Deep Earthquakes. In: G. Schubert (editor) *Treatise on Geophysics*, 2nd edition, Vol. 4. Elsevier, Oxford,
866 329-354.

867 Igarashi, T., Matsuzawa, T., Umino, N., Hasegawa, A., 2001. Spatial distribution of focal mechanisms for interplate and
868 intraplate earthquakes associated with the subducting Pacific plate beneath the northeastern Japan arc- A triple-planed deep
869 seismic zone. *Journal of Geophysical Research* 106, 2177-2191.

870 Isacks, B., Molnar, P., 1971. Distribution of stresses in the descending lithosphere from a global survey of focal mechanism
871 solutions of mantle earthquakes. *Reviews of Geophysics and Space Physics* 9, 103-174.

872 Jiao, W., Silver, P.G., Fei, Y., Prewitt, C.T., 2000. Do intermediate- and deep-focus earthquakes occur on preexisting weak
873 zones? An examination of the Tonga subduction zone. *Journal of Geophysical Research* 105, 28125-28138.

874 John, T., Medvedev, S., Rüpke, L.H., Andersen, T.B., Podladchikov, Y.Y., Austrheim, H., 2009. Generation of intermediate-
875 depth earthquakes by self-localizing thermal runaway. *Nature Geoscience* 2, 137-140, 10.1038/NGEO419.

876 Jolivet, L., Daniel, J.M., Fournier, M., 1991. Geometry and kinematics of extension in Alpine Corsica. *Earth and Planetary
877 Science Letters* 104, 278-291.

878 Jolivet, L., Dubois, R., Fournier, M., Goffé, B., Michard, A., Jourdan, C., 1990. Ductile extension in Alpine Corsica. *Geology*
879 18, 1007-1010.

880 Jolivet, L., Faccenna, C., Goffé, B., Mattei, M., Rossetti, F., Brunet, C., Storti, F., Funiciello, C., Cadet, J.P., D'Agostino, N.,
881 Parra, T., 1998. Mid-crustal shear zones in post-orogenic extension: examples from the northern Tyrrhenian Sea case. *Journal
882 of Geophysical Research* 103, 12123-12160.

883 Jung, H., Green, H.W., Dobrzhinetskaya, L.F., 2004. Intermediate-depth earthquake faulting by dehydration embrittlement
884 with negative volume change. *Nature* 428, 545-549.

885 Kameyama, M., Yuen, D.A., Karato, S.I., 1999. Thermal-mechanical effects of low-temperature plasticity (the Peierls
886 mechanism) on the deformation of a viscoelastic shear zone. *Earth and Planetary Science Letters* 168, 159-172.

887 Kanamori, H., Anderson, D.L., Heaton, T.H., 1998. Frictional melting during the rupture of the 1994 Bolivian earthquake.
888 *Science* 279, 839-842.

889 Kao, H., Chen, W.P., 1995. Transition from interplate slip to double seismic zone along the Kuril-Kamchatka arc. *Journal of
890 Geophysical Research* 100, 9881-9903.

891 Kao, H., Liu, L.G., 1995. A hypothesis for the seismogenesis of a double seismic zone, *Geophysical Journal International*
892 123, 71-84.

893 Kelemen, P., Hirth, G., 2007. A periodic shear-heating mechanism for intermediate-depth earthquake in the mantle. *Nature*
894 446, 787-790.

895 Kirby, S., 1995. Interslab earthquake and phase changes in subducting lithosphere. *Reviews of Geophysics* 33 (suppl., Report
896 to International Union of Geodesy and Geophysics), 287-297.

897 Kirby, S., Durham, W., Stern, L., 1991. Mantle phase changes and deep-earthquake faulting in subducting lithosphere.
898 *Science* 252, p. 216-225.

899 Kita, S., Okada, T., Nakajima, J., Matsuzawa, T., Hasegawa, A., 2006. Existence of a seismic belt in the upper plane of the
900 double seismic zone extending in the along-arc direction at depths of 70-100 km beneath NE Japan. *Geophysical Research*
901 *Letters* 33, L24310, 10.1029/2006GL028239.

902 Lacombe, O., Jolivet, L., 2005. Structural and kinematic relationships between Corsica and the Pyrenees-Provence domain at
903 the time of the Pyrenean orogeny. *Tectonics* 24, TC1003, 10.1029/2004TC001673.

904 Lahondère, D., 1996. Les schistes bleus et les éclogites à lawsonite des unités continentales et océaniques de la Corse alpine,
905 Documents du BRGM 240, Orléans, 294 p.

906 Lahondère, D., Guerrot, C., 1997. Datation Sm-Nd du métamorphisme éclogitique en Corse alpine : un argument pour
907 l'existence, au Crétacé supérieur, d'une zone de subduction active localisée le long du bloc corso-sarde. *Géologie de la France*
908 3, 3-11.

909 Lahondère, J.C., 1981. Relations du "socle ancien" de la région de Bastia (Corse) avec les Schistes lustrés environnants.
910 *Comptes Rendus de l'Académie des Sciences*, 293, 169-172.

911 Li, X.H., Faure, M., Rossi, P., Lahondère, D., 2015. Age of Alpine Corsica ophiolites revisited: Insights from in situ zircon
912 U-Pb age and O-Hf isotopes. *Lithos*, 220-223, 179-190.

913 Maddock, R.H., 1992. Effects of lithology, cataclasis and melting on the composition of fault-generated pseudotachylytes in
914 Lewisian gneiss, Scotland. *Tectonophysics* 204, 261-278.

915 Maggi, M., Rossetti, F., Corfu, F., Theye, T., Andersen, T.B., Faccenna, C., 2012. Clinopyroxene–rutile phyllonites from the
916 East Tenda Shear Zone (Alpine Corsica, France): pressure–temperature–time constraints to the Alpine reworking of Variscan
917 Corsica. *Journal of the Geological Society* 169, 723-732.

918 Magloughlin, J., 1992. Microstructural and chemical changes associated with cataclasis and frictional melting at shallow
919 crustal levels: the cataclasite-pseudotachylyte connection. *Tectonophysics* 204, 243-260.

920 Manatschal G., Müntener, O., 2009. A type sequence across an ancient magma-poor ocean–continent transition: the example
921 of the western Alpine Tethys ophiolites. *Tectonophysics* 473, 4-19.

922 Marroni, M., Pandolfi, L. 2007. The architecture of an incipient oceanic basin: a tentative reconstruction of the Jurassic
923 Liguria-Piemonte basin along the Northern Apennines–Alpine Corsica transect. *International Journal of Earth Sciences* 96,
924 1059-1078.

925 Martin, L.A.J., Rubatto, D., Vitale-Brovarone, A., Hermann, J., 2011. Late Eocene lawsonite-eclogite facies metasomatism
926 of a granitic sliver associated to ophiolites in Alpine Corsica. *Lithos* 125, 620-640.

927 Martinez-Diaz, J.J., Alvarez-Gomez, J.A., Benito, B., Hernandez, D., 2004. Triggering of destructive earthquakes in El
928 Salvador. *Geology* 32, 65-68, 10.1130/G20089.1.

929 Mattauer, M., Proust, F., 1976. La Corse alpine: un modèle de genèse du métamorphisme haute pression par subduction de
930 croûte continentale sous du matériel océanique. *Comptes Rendus de l'Académie des Sciences* 282, 1249-1252.

931 Mattauer, M., Proust, F., Etchecopar, A., 1977. Linéation 'a' et mécanisme de cisaillement simple liés au chevauchement de
932 la nappe des schistes lustrés en Corse. *Bulletin de la Société Géologique de France* 29, 841-847.

- 933 Mattauer, M., Faure, M., Malavieille, J., 1981. Transverse lineation and large-scale structures related to Alpine obduction in
934 Corsica. *Journal of Structural Geology* 3, 401-409.
- 935 McNulty, B.A., 1995. Pseudotachylytes generated in the semi-brittle and brittle regimes, Bench Canyon shear zone, central
936 Sierra Nevada. *Journal of Structural Geology* 17, 1507-1521.
- 937 Meresse, F., Lagabrielle, Y., Malavieille, J., Ildefonse, B., 2012. A fossil Ocean-Continent Transition of the Mesozoic Tethys
938 preserved in the Schistes Lustrés nappe of northern Corsica. *Tectonophysics* 579, 4-16.
- 939 Molli, G., 1994. Microstructural features of high temperature shear zones in gabbros of the Northern Apennine ophiolites.
940 *Journal of Structural Geology* 16, 1535-1541.
- 941 Molli, G., 2008. Northern Apennine-Corsica orogenic system: an updated overview. In Siegesmund, S., Fügenschuh, B.,
942 Froitzheim, N. (Eds.), *Tectonic Aspect of the Alpine-Dinaride-Carpathian System*: Geological Society of London, Special
943 Publication 298, 413-442.
- 944 Molli, G., Malavieille, J., 2010. Orogenic process and the Corsica/Apennine geodynamic evolution: Insight from Taiwan.
945 *International Journal of Earth Sciences* 100, 1207-1224, 10.1007/s00531-010-0598-y.
- 946 Nakajima, J., Tsuji, Y., Hasegawa, A., Kita, S., Okada, T., Matsuzawa, T., 2009. Tomographic imaging of hydrated crust and
947 mantle in the subducting Pacific slab beneath Hokkaido, Japan: Evidence for dehydration embrittlement as a cause of
948 intraslab earthquakes. *Gondwana Research* 16, 470-481.
- 949 Nakajima, J., Uchida, N., Shiina, T., Hasegawa, A., Hacker, B.R., Kirby, S.H., 2013. Intermediate-depth earthquakes
950 facilitated by eclogitization-related stresses. *Geology* 41, 659-662, 10.1130/G33796.1.
- 951 Obata, M., Karato, S.I., 1995. Ultramafic pseudotachylytes from Balmuccia peridotite, Ivrea-Verbano zone, northern Italy.
952 *Tectonophysics* 242, 313-328.
- 953 Ogawa, M., 1987. Shear instability in a viscoelastic material as the cause of deep focus earthquakes. *Journal of Geophysical*
954 *Research* 92, 13801-13810.
- 955 Ohnenstetter, M., Ohnenstetter, D., Vidal, P., Cornichet, J., Hermitte, D., Mace, J., 1981. Crystallization and age of zircon
956 from Corsican ophiolitic albitites: consequences for oceanic expansion in Jurassic times. *Earth and Planetary Science Letters*
957 54, 397-408.
- 958 Okada, T., Hasegawa, A., 2003. The M7.1 May 26, 2003 off-shore Miyagi Prefecture Earthquake in northeast Japan: Source
959 process and aftershock distribution of an intra-slab event. *Earth, Planets, Space* 55, 731-739.
- 960 Padron-Navarta, J.A., Hermann, J., Garrido, C.J., Sanchez-Vizcaino, V.L., Gomez-Pugnaire, M.T., 2010. An experimental
961 investigation of antigorite dehydration in natural silica-enriched serpentinites. *Contributions to Mineralogy and Petrology*
962 159, 25-42, 10.1007/s00410-009-0414-5.
- 963 Passchier, C., 1982. Pseudotachylyte and the development of ultramylonite bands in the Saint-Barthelemy Massif, French
964 Pyrenees. *Journal of Structural Geology* 4, 69-79.
- 965 Peacock, S.M., 2001. Are the lower planes of double seismic zones caused by serpentine dehydration in subduction oceanic
966 mantle? *Geology* 29, 299-302.

- 967 Perrillat, J.P., Daniel, I., Koga, K.T., Reynard, B., Cardon, H., Crichton, W.A., 2005. Kinetics of antigorite dehydration: A
968 real-time X-ray diffraction study. *Earth and Planetary Science Letters* 236, 899-913.
- 969 Piccardo, G.B., 2008. The Jurassic Ligurian Tethys, a fossil ultra-slow spreading ocean: The mantle perspective. In: Coltorti,
970 M., Grégoire, M. (Eds.), *Metasomatism in oceanic and continental lithospheric mantle: Geological Society of London Special*
971 *Publication* 294, 11-33.
- 972 Piccardo, G.B., Ranalli, G., Marasco, M., Padovano, M., 2007. Ultramafic pseudotachylytes in the Mt. Moncuni peridotite
973 (Lanzo Massif, western Alps): tectonic evolution and upper mantle seismicity. *Periodico di Mineralogia* 76, 181-197,
974 10.2451/2007PM0024.
- 975 Piccardo, G.B., Ranalli, G., Guarnieri, L., 2010. Seismogenic shear zone in the lithospheric mantle: Ultramafic
976 pseudotachylytes in the Lanzo peridotite (Western Alps, NW Italy). *Journal of Petrology* 51, 81-100,
977 doi:10.1093/petrology/egp067.
- 978 Preston, L.A., Creager, K.C., Crosson, R.S., Brocher, T.M., Trehu, A.M., 2003. Intraslab earthquakes: Dehydration of the
979 Cascadia Slab. *Science* 302, 1197-1200.
- 980 Prieto, G.A., Florez, M., Barrett, S.A., Beroza, G.C., Pedraza, P., Blanco, J.F., Podeva, E., 2013. Seismic evidence for
981 thermal runaway during intermediate-depth earthquake rupture. *Geophysical Research Letters* 40, 6064-6068,
982 10.1002/2013GL058109.
- 983 Raleigh, C.B., Paterson, M.S., 1965. Experimental deformation of serpentinite and its tectonic implications. *Journal of*
984 *Geophysical Research* 70, 3965-3985.
- 985 Rampone E., Hofmann, A.W., Raczek, I., 2009. Isotopic equilibrium between mantle peridotite and melt: Evidence from the
986 Corsica ophiolite. *Contributions to Mineralogy and Petrology* 156, 453-475, 10.1007/s00410-008-0296-y.
- 987 Rampone, E. Piccardo, G.B., 2000. The ophiolite-oceanic lithosphere analogue: new insights from the Northern Apennines
988 (Italy). *Geological Society of America Special Paper* 349, 21-34.
- 989 Ravna, E.J.K., Andersen, T.B., Jolivet, L., De Capitani, C., 2010. Cold subduction and the formation of lawsonite eclogite -
990 constraints from prograde evolution of eclogitized pillow lava from Corsica. *Journal of Metamorphic Geology* 28, 381-395.
- 991 Rondenay, S., Abers, G.A., van Keken, P.E., 2008. Seismic imaging of subduction zone metamorphism. *Geology* 36, 275-
992 278, 10.1130/G24112A.1.
- 993 Rosenbaum, G., Lister, G.S., Duboz, C., 2002. Relative motions of Africa, Iberia and Europe during Alpine orogeny.
994 *Tectonophysics* 359, 117-129.
- 995 Rossi, P., Cocherie, A., Lahondère, D., Fanning, N., 2002. La marge européenne de la Téthys jurassique en Corse : Datation
996 de trondhjémites de Balagne et indices de croûte continentale sous le domaine Balagno-Ligure. *Comptes Rendus Géoscience*
997 334, 313-322.
- 998 Rowe, C.D., Moore, J.C., Meneghini, F., McKeirnan, A.W., 2005. Large-scale pseudotachylytes and fluidized cataclasites
999 from an ancient subduction thrust fault. *Geology* 33, 937-940, 10.1130/G21856.1.
- 1000 Rutter, E. H., Brodie, K. H. (1988). Experimental “syntectonic” dehydration of serpentinite under conditions of controlled
1001 pore water pressure. *Journal of Geophysical Research* 93, 4907-4932.

- 1002 Scambelluri, M., Hoogerduijn-Strating, E.H., Piccardo, G.B., Vissers, R.L.M., Rampone, E., 1991. Alpine olivine- and
1003 clinohumite-bearing assemblages in the Erro-Tobbio peridotite (Voltri Massif, NW Italy). *Journal of Metamorphic Geology*
1004 9, 79-91.
- 1005 Schubnel, A., Brunet, F., Hilairt, N., Gasc, J., Wang, Y., Green, H.W., 2013. Deep-focus earthquake analogs recorded at
1006 high pressure and temperature in the laboratory. *Science* 341, 1377-1380, 10.1126/science.1240206.
- 1007 Seno, T., Yamanaka, Y., 1996. Double seismic zones, compressional deep trench-outer rise events, and superplumes. In:
1008 Subduction Top to Bottom. *Geophysical Monograph* 96, edited by G. E. Bebout et al., pp. 347– 355, AGU, Washington,
1009 D.C., 347-355.
- 1010 Seno, T., Zhao, D., Kobayashi, Y., Nakamura, M., 2001. Dehydration of serpentinized slab mantle: Seismic evidence from
1011 southwest Japan. *Earth, Planets, Space* 53, 861-871.
- 1012 Sibson, R.H., 1975. Generation of pseudotachylytes by ancient seismic faulting. *Royal Astronomical Society Geophysical*
1013 *Journal* 43, 775-794.
- 1014 Sibson, R.H., 1980. Transient discontinuities in ductile shear zones. *Journal of Structural Geology* 2, 165-171.
- 1015 Sibson, R.H., 1986. Brecciation processes in fault zones: Inferences from earthquake rupturing. *Pure and Applied Geophysics*
1016 124, 159-175.
- 1017 Singh, S.C., Carton, H., Tapponnier, P., Hananto, N.D., Chauhan, A.P.S., Hartoyo, D., Bayly, M., Moeljopranoto, S.,
1018 Bunting, T., Christie, P., Lubis, H., Martin, J., 2008. Seismic evidence for broken oceanic crust in the 2004 Sumatra
1019 earthquake epicentral region. *Nature Geoscience* 1, 777-781, 10.1038/ngeo336.
- 1020 Singh, S.C., Hananto, N., Mukti, M., Permana, H., Djajadihardja, Y., Harjono, H., 2011. Seismic images of the megathrust
1021 rupture during the 25th October 2010 Pagai earthquake, SW Sumatra: Frontal rupture and large tsunami. *Geophysical*
1022 *Research Letters* 38, L16313, 10.1029/2011GL048935.
- 1023 Souquière, F., Fabbri, O., 2010. Pseudotachylytes in the Balmuccia peridotite (Ivrea Zone) as markers of the exhumation of
1024 the southern Alpine continental crust. *Terra Nova* 22, 70-77.
- 1025 Spray, J.G., 1995. Pseudotachylyte controversy: Fact or friction? *Geology* 23, 1119-1122.
- 1026 Stampfli, G.M., Mosar, J., Marquer, D., Marchant, R., Baudin, T., Borel, G., 1998. Subduction and obduction processes in
1027 the Alps. *Tectonophysics* 296, 159-204.
- 1028 Stünitz, H., 1993. Transition from fracturing to viscous flow in a naturally deformed metagabbro. In: *Defects and processes*
1029 *in the solid state: Geoscience applications* (Boland, J.N. and Fitz Gerald, J.D. editors). Elsevier, Amsterdam, 121-150.
- 1030 Swanson, M.T., 1988. Pseudotachylyte-bearing strike-slip duplex structures in the Fort Foster Brittle Zone, S. Maine. *Journal*
1031 *of Structural Geology* 10, 813-828.
- 1032 Swanson, M.T., 1992. Fault structure, wear mechanisms and rupture processes in pseudotachylyte generation.
1033 *Tectonophysics* 204, 223-242.
- 1034 Tsuji, T., Kodaira, S., Ashi, J., Park, J.O., 2013. Widely distributed thrust and strike-slip faults within subducting oceanic
1035 crust in the Nankai Trough off the Kii Peninsula, Japan. *Tectonophysics* 600, 52-62.

1036 Tsuji, T., Park, J.O., Moore, G., Kodaira, S., Fukao, Y., Kuramoto, S.I., Bangs, N., 2009. Intraoceanic thrusts in the Nankai
1037 Trough off the Kii Peninsula: Implications for intraplate earthquakes. *Geophysical Research Letters* 36, L06303,
1038 10.1029/2008GL036974.

1039 Tucholke, B.E., Lin, J., 1994. A geological model for the structure of ridge segments in slow spreading ocean crust. *Journal*
1040 *of Geophysical Research* 99, 11937-11958.

1041 Turco, E., Macchiavelli, C., Mazzoli, S., Schettino, A., Pierantoni, P.P., 2012. Kinematic evolution of Alpine Corsica in the
1042 framework of Mediterranean mountain belts. *Tectonophysics* 579, 193-206.

1043 Ulmer, P., Trommsdorff, V., 1995. Serpentine stability to mantle depths and subduction-related magmatism. *Science* 268,
1044 858-861.

1045 Ueda, T., Obata, M., Di Toro, G., Kanagawa, K., Ozawa, K., 2008. Mantle earthquake frozen in mylonitized ultramafic
1046 pseudotachylytes of spinel-lherzolite facies. *Geology* 36, 607-610.

1047 Vallée, M., Bouchon, M., Schwartz, S.Y., 2003. The 13 January 2001 El Salvador earthquake: A multidata analysis. *Journal*
1048 *of Geophysical Research* 108, 2203, 10.1029/2002JB001922.

1049 Vitale-Brovarone, A., Beltrando, M., Malavielle, J., Giuntoli, F., Tondella, E., Groppo, C., Beyssac, O., Compagnoni, R.,
1050 2011. Inherited Ocean-Continent Transition zones in deeply subducted terranes: Insight from Alpine Corsica. *Lithos* 124,
1051 273-290.

1052 Vitale-Brovarone, A., Beyssac, O., Malavielle, J., Molli, G., Beltrando, M., Compagnoni, R., 2013. Stacking and
1053 metamorphism of continuous segments of subducted lithosphere in a high-pressure wedge: The example of Alpine Corsica
1054 (France). *Earth Science Reviews* 116, 35-56, doi:10.1016/j.earscirev.2012.10.003.

1055 Vitale-Brovarone, A., Herwartz, D., 2013. Timing of HP metamorphism in the Schistes Lustrés of Alpine Corsica: New Lu–
1056 Hf garnet and lawsonite ages. *Lithos* 172-173, 175-191.

1057 Vitale-Brovarone, A., Picatto, M., Beyssac, O., Lagabrielle, Y., Castelli, D., 2014. The blueschist-eclogite transition in the
1058 Alpine chain: P-T paths and the role of slow spreading extensional structures in the evolution of HP-LT mountain belts.
1059 *Tectonophysics* 615-616, 96-121.

1060 Wang, K., 2002. Unbending combined with dehydration embrittlement as a cause for double and triple seismic zones.
1061 *Geophysical Research Letters* 29, 1889, 10.1029/2002GL015441.

1062 Wang, K., Cassidy, J.F., Wada, I., Smith, A.J., 2004. Effects of metamorphic crustal densification on earthquake size in
1063 warm slabs. *Geophysical Research Letters* 31, L01605, 10.1029/2003GL018644.

1064 Warburton, J., 1986. The ophiolite-bearing Schistes Lustrés nappe in Alpine Corsica: A model for the emplacement of
1065 ophiolites that have suffered HP/LP metamorphism. In: *Blueschists and Eclogites* (B.W. Evans and E.H. Brown, eds).
1066 *Geological Society of America Memoir* 164, 313-331.

1067 Wenk, H.R., Johnson, L.R., Ratschbacher, L., 2000. Pseudotachylites in the Eastern Peninsular Ranges of California.
1068 *Tectonophysics* 321, 253-277.

1069 Wiens, D.A., McGuire, J.J., Shore, P.J., 1993. Evidence for transformational faulting from a deep double seismic zone in
1070 Tonga. *Nature* 364, 790-793, 10.1038/364790a0.

1071 Wiens, D.A., Snider N.O., 2001. Repeating deep earthquakes: Evidence for fault reactivation at great depth. *Science* 293,
1072 1465-1466.

1073 Wunder, B., Schreyer, W., 1997. Antigorite: High-pressure stability in the system $\text{MgO-SiO}_2\text{-H}_2\text{O}$ (MSH). *Lithos* 41, 213-
1074 227, 10.1016/S0024-4937(97)82013-0.

1075 Yamasaki, T., Seno, T., 2003. Double seismic zone and dehydration embrittlement of the subducting slab. *Journal of*
1076 *Geophysical Research* 108, 1-21.

1077 Yoshii, T., 1979. A detailed cross-section of the deep seismic zone beneath northeastern Honshu, Japan. *Tectonophysics* 55,
1078 349-360.

1079 Zechmeister, M.S., Ferré, E.C., Cosca, M.A., Geissman, J.W., 2007. Slow and fast deformation in the Dora Maira Massif,
1080 Italian Alps: Pseudotachylytes and inferences on exhumation history. *Journal of Structural Geology* 29, 1114-1130.

1081 **Figure captions**

1082 **Figure 1.** Structural map of the study area (modified after Faure and Malavieille, 1981,
1083 Lahondère, 1996 and Meresse et al., 2012) and lower-hemisphere equal-area projections of
1084 poles to pseudotachylyte fault veins in the peridotite unit. Arrows indicate the sense of shear
1085 associated with pseudotachylyte fault veins. The locations of cross-sections A-A' and B-B'
1086 (see Fig. 2) are shown. CdG: Cima di Gratera; PdM: Punta di Muzzelli.

1087 **Figure 2.** Geological cross-sections of the study area (location in Fig. 1).

1088 **Figure 3.** Examples of outcrop-scale top-to-the-west or top-to-the-northwest displacement
1089 sense criteria from the flat-lying fault zones in the peridotite unit. A, B and C: West-dipping
1090 or northwest-dipping Riedel-like pseudotachylyte-coated normal faults (labeled by R)
1091 offsetting earlier fault veins. A and B from locality 6, C from locality 5. D: Southeast-dipping
1092 pseudotachylyte-coated reverse faults offsetting earlier fault veins at locality 3.

1093 **Figure 4.** Polished surface (A) and corresponding sketches (B and C) of a peridotite hand
1094 sample from the upper flat-lying fault zone at locality 2 (Fig. 1) showing two stages of
1095 pseudotachylyte formation. The kinematics associated with the early pseudotachylyte veins is
1096 undetermined, while that associated with the late pseudotachylyte veins is top-to-the-west
1097 (N280°E). Rectangle on (A) corresponds to the thin section scanner image of Fig. 10C.

1098 **Figure 5.** Detailed field view (A) and corresponding sketch (B) of a steeply-dipping reverse
1099 fault zone (locality 2) showing anastomosed fault veins crossing cataclastic peridotite.

1100 **Figure 6.** Polished surface (A) and corresponding sketch (B) of a peridotite hand sample from
1101 the steeply-dipping fault zone of locality 2 showing three stages of pseudotachylyte

1102 formation. Senses of shear of the intermediate and late seismic ruptures are top-to-the-
1103 northwest (N320°E). Rectangle on (A) corresponds to Fig. 10D.

1104 Figure 7. Thin section scanner images, SEM images and photomicrographs of microlitic-type
1105 and annealed-type pseudotachylyte veins in the peridotite. A: Thin section scanner image of a
1106 microlitic-type fault vein showing a zonation parallel to the boundaries with the host rock
1107 (locality 3). B: Chilled margin (C.m) of a microlitic-type fault vein showing a sharp decrease
1108 in microlite size. Microlites consist of olivine and pyroxene (locality 1). C: Pyroxene
1109 microlites in an injection vein (locality 4). D: Annealed-type fault vein from locality 4.
1110 Arrows indicate cooling cracks. Square is for E. E: SEM image of D showing olivine with a
1111 granoblastic annealed texture.

1112 Figure 8. Examples of pseudotachylyte-cataclasite associations in the peridotite-hosted
1113 steeply dipping fault zone at locality 2. A: Photomicrograph. (B) Thin section scanner image.
1114 Ct: cataclasite; Pct: proto-cataclasite; Pst: pseudotachylyte; Uc: ultra-cataclasite.

1115 Figure 9. SEM images of associated cataclasites and pseudotachylytes in peridotite from
1116 locality 6. A: From top left to bottom right, juxtaposition of proto-cataclasite (PCt), cataclasite
1117 (Ct), pseudotachylyte (Pst) and moderately fractured peridotite (wall). B: Detail of the Ct-Pst-
1118 host rock zoned domain of (A). C: Detail of the Ct domain of (A), showing angular clasts. D:
1119 Detailed view of a cataclasite-wall rock contact.

1120 Figure 10. Thin section scanner images and photomicrographs of fault veins from the steeply
1121 dipping fault zone at locality 2 (A and B) and from the flat-lying fault zone at locality 3 (C, D
1122 and E) showing top-to-the-west or top-to-the-northwest displacement senses. A: Parallel
1123 polarized thin section scanner image showing an anastomosed network of steeply dipping
1124 cataclasite zones with a reverse displacement sense. B: Crossed-polar enlarged image from
1125 (A) showing cataclastic zones offsetting olivine crystals in a reverse sense. C: Parallel-polar
1126 thin section scanner image of a pseudotachylyte vein and associated sheared peridotite
1127 suggesting a top-to-the-west sense of displacement. D: Parallel-polar thin section scanner
1128 image showing an early steeply dipping pseudotachylyte vein left-laterally offset by a late
1129 flat-lying pseudotachylyte vein. E: Detail of D showing that the flat-lying vein is younger
1130 than the vein dipping to the right.

1131 Figure 11. Outcrop aspect of pre- and post-mylonitization pseudotachylyte veins in the
1132 foliated metagabbro sole at locality 4 and attitudes of nearby post-mylonitization veins. A:
1133 Foliated metagabbro showing a post-mylonitization fault vein secant on pre-mylonitization

fault veins and on foliation. The foliation is outlined by the dashed red line. Red arrows point at pre-mylonitization fault veins. B: Lower-hemisphere equal-area projection of poles to post-mylonitization fault veins (solid circles) and poles to foliation (red triangles).

Figure 12. Microscopic aspect of pre- and post-mylonitization pseudotachylyte veins in the foliated sole of the metagabbro unit. A: Parallel-polar thin section scanner image of the foliated metagabbro. B: Detail of A showing a mylonitized pseudotachylyte vein. The foliation inside the vein is slightly oblique to the foliation outside, likely because of some obliquity of the vein with respect to the deformation axes. C: Parallel-polar thin section scanner image showing a post-mylonitization fault vein, a post-mylonitization injection vein and a foliated pre-mylonitization vein. D: sketch of C. Pre-myl. Pst: pre-mylonitization vein; Post-myl. Pst: post-mylonitization vein. Dashed lines outline the foliation trace.

Figure 13. Multi-stage scenario of formation of pseudotachylyte and mylonite in the Cima di Gratera nappe. A: General sketch showing the east-dipping subduction of the Piemonte-Liguria oceanic basin beneath an arc or a micro-continent in Cretaceous times. Also depicted are the hypocenters (asterisks) of the Wadati-Benioff seismic zone. B: Formation of pre-mylonitization pseudotachylyte at shallow depth at or near the mantle-crust boundary. C: Ductile deformation of the base of the oceanic crust and coeval formation of pseudotachylyte in the underlying peridotite. D: Formation of post-mylonitization veins in the ductilely deforming metagabbro by seismic ruptures nucleated in the peridotite and having propagated upward across and beyond the foliated metagabbro.

Figure 14. Geometrical and kinematic similarities between the present-day seismic activity of the Wadati-Benioff zone beneath NE Japan (A and B) and the Corsican fossil seismic ruptures (C and D). A: Possible location of the Corsican seismic ruptures (rectangle) in a cold slab thermal model (isotherms after Peacock, 2001 and Hacker et al., 2003). B: Hypocenters and kinematics of the 2002-2003 seismic activity in the uppermost part of the Pacific plate off NE Japan (Hasegawa et al., 2007). The red rectangle delineates the possible equivalent of the seismic fault zones frozen in Corsica. C: Sketch summarizing the geometry and kinematics of the Corsican fossil seismic ruptures in the peridotite unit (approximate width: 5 km). D: Detail of C emphasizing large ruptures propagating upwards across the crust-mantle boundary and beyond the foliated basal metagabbro (approximate width: 1 km).

***Figure**
[Click here to download high resolution image](#)

*Figure

[Click here to download high resolution image](#)

***Figure**
[Click here to download high resolution image](#)

*Figure
[Click here to download high resolution image](#)

*Figure
[Click here to download high resolution image](#)

[Click here to download high resolution image](#)

***Figure**
[Click here to download high resolution image](#)

*Figure

[Click here to download high resolution image](#)

*Figure
[Click here to download high resolution image](#)

***Figure**
[Click here to download high resolution image](#)

*Figure

[Click here to download high resolution image](#)

***Figure**
[Click here to download high resolution image](#)