



HAL
open science

**Near-future pH conditions severely impact calcification,
metabolism and the nervous system in the pteropod
*Heliconoides inflatus***

Aurelie Moya, Ella L Howes, Thomas Lacoue-Labarthe, Sylvain Forêt, Bishoy Hanna, Mónica Medina, Philip L Munday, Jue-Sheng Ong, Jean-Louis Teyssié, Gergely Torda, et al.

► **To cite this version:**

Aurelie Moya, Ella L Howes, Thomas Lacoue-Labarthe, Sylvain Forêt, Bishoy Hanna, et al.. Near-future pH conditions severely impact calcification, metabolism and the nervous system in the pteropod *Heliconoides inflatus*. *Global Change Biology*, 2016, 10.1111/gcb.13350 . hal-01346620

HAL Id: hal-01346620

<https://hal.sorbonne-universite.fr/hal-01346620v1>

Submitted on 19 Jul 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Received Date : 08-Feb-2016
Revised Date : 22-Apr-2016
Accepted Date : 29-Apr-2016
Article type : Primary Research Articles

Near future pH conditions severely impact calcification, metabolism and the nervous system in the pteropod *Heliconoides inflatus*

Running head: Effect of low pH on Mediterranean pteropods

Aurelie Moya^{1,2§}, Ella L Howes^{2,3}, Thomas Lacoue-Labarthe^{4,5,6}, Sylvain Forêt⁷, Bishoy Hanna⁸, Mónica Medina⁸, Philip L Munday¹, Jue-Sheng Ong⁷
Jean-Louis Teysse⁶, Gergely Torda^{1,9}, Sue-Ann Watson¹, David J Miller^{1,10}, Jelle Bijma³,
Jean-Pierre Gattuso^{2,4}

¹ ARC Centre of Excellence for Coral Reef Studies, James Cook University, Townsville, Qld, 4811, Australia

² Sorbonne Universités, UPMC Univ Paris 06, CNRS-INSU, Laboratoire d'Océanographie de Villefranche, 181 chemin du Lazaret, F-06230 Villefranche-sur-mer, France

³ Marine Biogeosciences, Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, Am Handelshafen 12, D-27570 Bremerhaven, Germany

⁴ Institute for Sustainable Development and International Relations, Sciences Po, 27 rue Saint Guillaume, F-75007 Paris, France

⁵ Littoral, Environnement et Sociétés (LIENSs), UMR 7266, CNRS-Université de La Rochelle, 2 rue Olympe de Gouges, 17000 La Rochelle Cedex, France

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/gcb.13350

⁶ Marine Environmental Laboratories, International Atomic Energy Agency, 4 quai Antoine 1er, 98000 Monaco, Principality of Monaco

⁷ Research School of Biology, Australian National University, Canberra, Australian Capital Territory 0200, Australia

⁸ Department of Biology, Penn State University, University Park, PA 16802, USA

⁹ Australian Institute of Marine Science, PMB 3, Townsville, Queensland 4810, Australia

¹⁰ Comparative Genomics Centre and Department of Molecular and Cell Biology, James Cook University, Townsville, Qld, 4811, Australia

[§] Corresponding author: aurelie.moya@jcu.edu.au. Phone: +61 7 4781 3654, fax: +61 7 4725 5043.

Keywords: global change biology, ocean acidification, pteropods, RNA sequencing, gene expression, calcification, respiration, nervous system, GABA_A Receptor.

Type of paper: Primary research article

Abstract (300 words)

Shelled pteropods play key roles in the global carbon cycle and food webs of various ecosystems. Their thin external shell is sensitive to small changes in pH and shell dissolution has already been observed in areas where aragonite saturation state is ~ 1 . A decline in pteropod abundance has the potential to disrupt trophic networks and directly impact commercial fisheries. Therefore it is crucial to understand how pteropods will be affected by global environmental change, particularly ocean acidification. In the present study, physiological and molecular approaches were used to investigate the response of the Mediterranean pteropod, *Heliconoides inflatus*, to pH values projected for 2100 under a

moderate emissions trajectory (RCP6.0). Pteropods were subjected to pH_T 7.9 for 3 d, and gene expression levels, calcification and respiration rates were measured relative to pH_T 8.1 controls. Gross calcification decreased markedly under low pH conditions, while genes potentially involved in calcification were up-regulated, reflecting the inability of pteropods to maintain calcification rates. Gene expression data imply that under low pH conditions both metabolic processes and protein synthesis may be compromised, while genes involved in acid-base regulation were up-regulated. A large number of genes related to nervous system structure and function were also up-regulated in the low pH treatment, including a GABA_A receptor subunit. This observation is particularly interesting because GABA_A receptor disturbances, leading to altered behaviour, have been documented in several other marine animals after exposure to elevated CO_2 . The up-regulation of many genes involved in nervous system function suggests that exposure to low pH could have major effects on pteropod behaviour. This study illustrates the power of combining physiological and molecular approaches. It also reveals the importance of behavioural analyses in studies aimed at understanding the impacts of low pH on marine animals.

INTRODUCTION

Shelled pteropods both play a critical role in the export of calcium carbonate from the sea surface into the deep ocean and contribute approximately 12 % of the global carbonate flux ([Bednaršek *et al.*, 2012a](#)). Pteropods deposit CaCO_3 as external aragonite shells, which makes them more vulnerable to ambient oceanic conditions than organisms with internal aragonite skeletons such as hard corals. As aragonite is the most soluble form of CaCO_3 , calcification by pteropods is predicted to be sensitive to subtle changes in ambient pH. The key role of pteropods in the global carbon cycle is thus likely to be compromised by ocean acidification ([Ries, 2012](#)). Dissolution of shelled organisms is predicted to occur when the

aragonite saturation state (Ω_a) falls below 1. In the case of pteropods, however, shell dissolution has already been observed in the upper layers of the Southern Ocean where Ω_a levels were around 1 (Bednaršek *et al.*, 2012b) as well as in the North Pacific/California current (Bednaršek *et al.*, 2014; Bednaršek & Ohman, 2015).

Because pteropods are also an important food source for a variety of organisms, ranging from zooplankton to whales and including commercial fish species (e.g. salmon) (Foster & Montgomery, 1993), a decline in pteropod abundance has the potential to disrupt trophic networks and directly impact commercial fisheries. In light of their ecological and economic significance, it is crucial to understand how pteropods will be affected by global environmental change, particularly ocean acidification.

Previous studies have reported declines in calcification rates (Comeau *et al.*, 2012b; Comeau *et al.*, 2009; Comeau *et al.*, 2010b), linear extension rates (Comeau *et al.*, 2012a; Comeau *et al.*, 2010a; Comeau *et al.*, 2009; Lischka *et al.*, 2011; Lischka & Riebesell, 2012) and shell integrity (Comeau *et al.*, 2012b; Lischka *et al.*, 2011; Manno *et al.*, 2012; Orr *et al.*, 2005) with decreasing seawater pH. However, the possible effects of ocean acidification on physiological processes in pteropods has received much less attention (Gazeau *et al.*, 2013 for a review). One study showed that respiration rates of four subtropical and tropical pteropods species that naturally migrate into oxygen minimum zones were not affected by elevated carbon dioxide while the one species studied that do not migrate shown reduced oxygen consumption (Maas *et al.*, 2012). Results on polar species were more complex and influenced by other external parameters such as temperature (Comeau *et al.*, 2010b) or phytoplankton abundance (Seibel *et al.*, 2012). A recent study reported that low salinity in consort with lower pH negatively affects the swimming activity of the pteropod *Limacina retroversa* (Manno *et al.*, 2012). Such finding is consistent with a growing body of evidence

that elevated CO₂ and low pH causes behavioral disturbances in fish, molluscs and other marine organisms ([Briffa et al., 2012](#); [Clements & Hunt, 2015](#); [Watson et al., 2014](#)).

Two recent transcriptomic studies have examined the responses of pteropods to elevated CO₂ ([Koh et al., 2015](#); [Maas et al., 2015](#)). However sample sizes and high inter-individual variability severely limited the numbers of differentially expressed genes that could be detected ([Maas et al., 2015](#)) or no attempts were made to link gene expression data with physiological parameters ([Koh et al., 2015](#)).

In the present study, both physiological and molecular approaches were used to investigate the response of the Mediterranean pteropod, *Heliconoides inflatus*, to pH values projected to occur by 2100 under a moderate emissions trajectory (RCP6.0). *Heliconoides inflatus*

(d'Orbigny, 1836) is a shelled species formerly known as *Limacina inflata*. It is one of the most common warm-water cosmopolitan pteropods and is widely distributed in the tropical and subtropical regions of all oceans. It is an epiplanktonic species living primarily in the upper 300 - 600 m of the water column depending on the season ([Bé & Gilmer, 1977](#)). Both adults and juveniles undertake diurnal vertical migration, moving to surface waters at night.

In adult stages, nocturnal migration occurs year round but is more pronounced during periods when they are deeper living (Feb-Mar). Nocturnal migration of juveniles is seasonally dependent ([Rampal, 1975](#)).

Individuals were subjected to pH_T 8.1 (control pH, on the total scale) or pH_T 7.9 for 3 days.

In addition to changes in gene expression caused by the treatment, rates of calcification and respiration were measured in the two experimental conditions. Changes in mRNA levels were assessed by RNA sequencing (Illumina technology), note that proteins levels were not measured in the present study. Experimental and control gene expression levels were compared by mapping the individual reads onto a *de novo* reference transcriptome assembly

(see *Supporting Information*). A global approach based on Gene Ontology enrichment analysis was first implemented in order to identify which molecular pathways were being repressed or activated in response to low pH. This step was followed by a targeted approach focused on genes involved in calcification and respiration to facilitate comparison between the physiological and molecular results.

MATERIAL AND METHODS

Collection of *Heliconoides inflatus*

Heliconoides inflatus individuals (Fig. 1) were collected in the Bay of Villefranche-sur-mer (43°40' N, 7°18' E) between 16th and 29th February 2012. Sampling was undertaken using very slow oblique tows of 2-4 min in duration at up to 100 m depth with a 57 cm diameter WP2 plankton net (200 µm mesh size). Plankton samples were immediately transported to the Laboratoire d'Océanographie de Villefranche. *Heliconoides inflatus* individuals were inspected under binocular microscope to assess their condition and reproductive status (as *H. inflatus* broods its young, care taken to ensure that brooding females were not included in the experiments). Using a wide aperture pipette, individuals were transferred one by one to filtered seawater medium (pore size 0.2 µm).

pH manipulation

Individuals were exposed for 3 d to pH_T 8.1 (control) or pH_T 7.9 (treatment). The desired pH conditions were maintained using a pH-stat system (IKS Karlsbard) that bubbled filtered seawater (pore size 0.2 µm) with either ambient air (control) or CO₂ (treatment) until the expected pH was reached; the pH electrode was calibrated on the total scale using Tris/HCl and 2-aminopyridine/HCl buffer solutions with a salinity of 38.0 ([Dickson *et al.*, 2007](#)).

Three separate experiments were performed to assess (1) respiration rate, (2) ⁴⁵Ca uptake, and

(3) gene expression. In each experiment, 25 individuals of *H. inflatus* were transferred to 2 L sterilized borosilicate glass bottles, together with 1 ml of *Isochrysis galbana* culture to feed the animals. Three replicates were set up for each pH condition for (1) and (3), and four replicates for (2). Bottles were then closed and sealed with parafilm for 72 h of incubation, unless stated otherwise. Total alkalinity (A_T), pH_T , and dissolved inorganic carbon (C_T) were measured at the start and end of the incubations. A_T samples were filtered on GF/F and measured potentiometrically using a Tritando 80, Metrohm titrator and a Metrohm, electrode plus glass electrode. 60 ml samples was also taken at the start and finish of incubations and poisoned with $HgCl_2$ for determination of C_T . Samples were measured using an AIRICA (Marianda, Kiel) with Licor analyser. All other parameters of the carbonate chemistry were calculated using the R package *seacarb* (Lavigne & Gattuso, 2013; R Development Core Team, 2010), constants for K_1 and K_2 were taken from [Lueker *et al.* \(2000\)](#), K_f from Perez & Fraga (1987) and K_s from Dickson (1990) (Table 1).

Experiments on ^{45}Ca uptake were conducted at the Marine Environment Laboratory of the International Atomic Energy Agency (IAEA) in Monaco, while incubations for gene expression and respiration rates measurements were conducted at Laboratoire d'Océanographie de Villefranche.

Experiment 1: respiration rates

Six, 60 ml biological oxygen demand (BOD) bottles with oxygen sensitive microspots (PreSens, Planar, 5 mm in diameter) glued to the inner wall were prefilled with experimental seawater (prepared as described above), 3 control pH and 3 low pH. Animals were incubated for 48 h in seawater adjusted to the experimental treatment levels; after 48 h, 60 actively swimming animals were selected from each treatment and transferred to BOD bottles containing the same seawater (20 individuals in each bottle). Two additional BOD bottles

were prepared with oxygen sensitive microspots and filled with only the treatment seawater to act as blanks. All eight bottles were connected to an oxygen sensor (OXY-4 mini, PreSens), placed into a temperature-controlled bath at 14°C and oxygen concentration was measured semi-continuously (every 15 s) over the following 24 h. At the end of the incubation, animals were removed from the BOD bottles, placed into a petri dish and fixed using 90% ethanol. Individuals were photographed under a binocular microscope, and the maximum shell diameter measured using the imaging software AxioVision (version 6.1.7601); for consistency with the calcification results, wet weights were calculated from the size weight relationship. The average shell diameter of animals used in the respiration experiments was $497.77 \pm 12.93 \mu\text{m}$ and the average wet weight was $11.27 \pm 1.09 \mu\text{g}$, organisms ranged in wet weight from 0.63 to 67.65 μg (see Table S8).

Oxygen consumption rate was calculated by regressing oxygen concentration through time. The initial 2 h of incubation were excluded to avoid any stress effect caused by the transfer from 2 L glass bottles to the BOD bottles. Respiration rates were calculated for the entire pool of individuals in one replicate after correcting for oxygen consumption rates in the blank incubations and expressed as $\mu\text{mol O}_2 \text{ h}^{-1} (\mu\text{g wet weight})^{-1}$.

Experiment 2: ^{45}Ca uptake

Pteropods were photographed under a binocular microscope, and the maximum shell diameter measured using the imaging software AxioVision, version 6.1.7601a; they were then cultured (n= 25 per bottle) in conditions similar to the one described above and the 2 L culture bottles spiked with a $^{45}\text{CaCl}_2$ (Radioisotope Centre Polatum, Poland; $T_{1/2} = 163 \text{ d}$) solution for a final concentration of *ca.* 130 Bq ml^{-1} . At the end of the 72-h incubation, all the individuals from each of the four replicates were filtered (200 μm mesh), pooled and placed

into a scintillation vial. The shells were dissolved with 0.5 N HCl and the soft tissues removed using a stereomicroscope and fine forceps. The soft tissue was rinsed into the vial with Milli-Q water to ensure that all the liquid containing the dissolved shell was collected. The solution was then neutralized with 1 N NaOH and allowed to evaporate on a heated plate before addition of 10 ml of scintillation fluid (Ultima Gold™ XR, Perkin Elmer). ⁴⁵Ca activity was counted using a Packard scintillation counter (Tri-Carb, Packard 1600 TR or Perkin Elmer 2900 TR), and corrected for quenching by employing external standard (0.4 Bq.ml⁻¹) quench correction curves. Counting time was adjusted between 10 min and 24 h to obtain relative errors below 5 %. The incorporation of ⁴⁵Ca in pteropods shell was expressed according to the following equation (see Martin *et al.*, 2011):

$$Q_{Ca} = [(A_{shell} / A_{sw}) \times C_{sw}] \times 10^3$$

where Q_{Ca} is the amount of Ca incorporated per shell (nmol g⁻¹), A_{shell} is the total ⁴⁵Ca activity in each shell (in Bq), A_{sw} is the ⁴⁵Ca activity (in Bq g⁻¹) in seawater during the time of exposure, and C_{sw} is the total Ca concentration in Mediterranean seawater (0.0114 mmol g⁻¹). ⁴⁵Ca uptake was expressed as nmol.μg⁻¹ wet weight. The pteropod weight was calculated from a previously determined size-weight relationship (see paragraph below).

Experiment 3: Gene expression

At the end of the 3-d incubations, the *H. inflatus* individuals from each of the three replicates were collected from the experimental jars by filtering the water over a 200 μm mesh.

Specimens were rinsed with 0.45 μm filtered seawater, immediately snap-frozen in liquid nitrogen and stored at -80 °C until further treatment. Twenty-five individuals were pooled for each of the three replicates.

Size-weight relationship:

A size-weight relationship was used to estimate the individual weight for ^{45}Ca uptake and respiration. The relationship between *H. inflatus* shell size and overall weight was based on 118 individuals representative of the full size range of individuals collected in the Bay of Villefranche-sur-mer in February 2012. Individuals were photographed under a binocular microscope, and the maximum shell diameter measured using the imaging software AxioVision, version 6.1.7601. Individuals were dried with tissue paper prior to weighting with a Mettler Toledo microbalance (precision = 0.1 μg) and the following relationship was derived:

$$W = -a + b \times D^2$$

where W is the wet weight (including the shell), a and b are constants and D is the shell diameter (Fig. S1).

Statistical analyses

For both respiration rates and ^{45}Ca uptake, mean differences between the two treatments were tested using a paired t -test, due to a temporal difference between replicates. Respiration experiments were all performed at the same time so a standard t -test was used. Statistical analyses were performed using R.

RNA extraction and transcriptome sequencing

Total RNA was extracted using RNeasy Plus Micro Kit (Qiagen) according to the manufacturer instructions. RNA quantity and quality were assessed using a NanoDrop ND-1000 spectrophotometer and denaturing gel electrophoresis using standard methods (Sambrook & Russel, 2001). Before being shipped on dry-ice to the Macrogen sequencing facilities in Seoul, South Korea, each RNA sample was precipitated in ethanol and sodium

acetate (2x and 0.1x sample volume, respectively), and stored at -80 °C. Libraries were prepared with Illumina TruSeq Stranded RNA-seq kit by Macrogen (South Korea). The libraries were sequenced using the Illumina HiSeq2000 platform, producing an average of 63 million sequence reads (100 bp paired-end) per sample.

***De novo* assembly and data analysis**

The reads from all the *H. inflatus* samples were trimmed from low quality regions and sequencing adaptors using libngs (<https://github.com/sylvainforet/libngs>) with a minimum quality of 20 and a minimum size of 75 bp. The trimmed reads were then assembled using Trinity (Grabherr *et al.*, 2011). The sequences of the resulting assembly were clustered using cdhit-est (Fu *et al.*, 2012) with a similarity threshold of 90% and a word size of 8 bp. The multi-modal GC profile of this initial assembly is suggestive of contamination (Fig. S2a), probably originating from the *Isochrysis galbana* culture provided to the animals. In order to remove these potential contaminants, the transcripts were blasted against a database containing the NCBI refseq proteins augmented with sequences from the molluscs *Biomphalaria glabrata* (<https://www.vectorbase.org/organisms/biomphalaria-glabrata>) and *Lymnea stagnalis* (Sadamoto *et al.*, 2012). Sequences were then classified as putative *H. inflatus* sequences if they satisfied the following criteria: e-value less than $1e^{-10}$, a minimum bit-score of at least 100, and a delta bit-score of at least 100. Here delta bit-score refers to the difference between the best mollusc hit and the best non-mollusc hit. After the above filtering, the distribution of GC content appeared unimodal (Fig. S2b). Due to the stringent filtering procedure, the sequences in this reduced set are very likely to be bona-fide pteropod sequences. Characteristics of the transcriptome assemblies before and after filtering are presented as Table S1. Protein-coding sequences were predicted using Transdecoder (<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3875132/>) and annotated using Blast2GO

(Conesa *et al.*, 2005). The reads were mapped back to the assembled transcripts using Bowtie2 (Langmead & Salzberg, 2012). The number of fragments mapping to each transcript was computed using RSEM. Only the longest transcript for each assembled trinity locus was used. Differential gene expression was inferred using EdgeR (Robinson *et al.*, 2010) using a FDR threshold of 5%.

GO enrichment analysis was carried out with the Blast2GO software (Conesa *et al.*, 2005) using a Fisher's exact test with a FDR threshold of 5% to reduce false-positive predictions of enriched GO terms. For each gene category of interest, BlastP and HMMER domain searches (e-value cut-off = $1e^{-5}$) were performed on the sets of differentially expressed genes. An additional blast onto the NCBI *nr* database confirmed the identification of each sequence. In addition for the category "calcification", the set of differentially expressed genes was searched against a database for metazoan biomineralization proteins (<https://peerj.com/preprints/1983/>).

RESULTS

Respiration and calcification

There was no difference in respiration rate between individuals kept under control and low pH (Fig. 2b and Table S9; *t*-test, $t = -0.05$, $df = 4$, p -value = 0.97). However, individuals incubated in the low pH treatment exhibited a 37% decrease in gross calcification (^{45}Ca uptake) compared to control treatment (Fig. 2a; paired *t*-test, $t=6.1$, $df=3$, p -value=0.009).

Transcriptomic analyses

Transcript levels differed significantly between the two experimental conditions; relative to the control (pH_T 8.1), 1.8% (400 transcripts) and 0.8% (173 transcripts) of *H. inflatus* transcripts were up- and down-regulated, respectively, at the lower pH condition (adjusted *P*

<0.05). Relatively small changes were observed amongst genes that were up-regulated: in approximately 74% of cases, expression changed between 2 and 10-fold in low pH conditions, whereas changes of ≥ 100 -fold were observed for fewer than 3% of the up-regulated transcripts. In contrast, a much larger proportion of the down-regulated genes showed higher fold changes, 47% of down-regulated transcripts showing more than 100 fold-changes (Fig. 3).

Ion transport, protein synthesis and mitochondrion activity

Gene Ontology (GO) analyses were conducted to infer the overall impacts of the experimental manipulation. Amongst the up-regulated genes, only a single GO category, *Ion transport* (GO:0006811), was significantly enriched (Table 2). This category comprises 26 genes including acid sensing ion channels, potassium, sodium, calcium and proton transporters, phospholipids and amino acid transporters, as well as neuronal transporters (see Table S2).

In contrast, many GO categories were enriched amongst the down-regulated genes (Table 2), a large proportion of these being related to protein synthesis and mitochondrial activity. For example, 75 clusters associated with protein synthesis were significantly down-regulated in the low pH condition (Table 3 and Table S3), with some transcripts levels being essentially undetectable (e.g.: a GTP-binding translation elongation factor, comp474736_c0_seq3). This category comprises 24 clusters encoding subunits of the 40S ribosomal protein, 26 encoding subunits of the 60S ribosomal protein as well as several other genes encoding proteins involved in regulation of transcription, protein translation initiation and elongation (Table 3 and Table S3), suggesting that the entire protein synthesis machinery was affected. Amongst mitochondrion-related genes, 22 components of the electron transport chain (including proteins from 3 of the 4 mitochondrial inner membrane complexes as well as several ATP synthase subunits, see Table 3 and Table S4) were affected, which suggests that oxidative

metabolism is suppressed in the low pH condition. The magnitude of changes in expression was extremely high in this category; with four cytochrome *c* oxidase subunits completely turned-off in the low pH treatment compared to control.

Acid-base regulation

Amongst the differentially expressed genes, 8 were potentially involved in acid-base regulation (Table 3 and Table S5). Two mRNA encoding for Na⁺-K⁺-ATPases, responsible for establishing and maintaining the electrochemical gradients of Na⁺ and K⁺ ions across the plasma membrane, were up-regulated in the low pH condition (average of 4.9-fold). Two mRNA encoding for vacuolar-type H⁺-ATPases, generating proton gradients across membranes of numerous cell types, were also up-regulated more than 5-fold. One mRNA encoding for a carbonic anhydrase (comp456540_c0_seq1) was up-regulated 39 times in the low pH condition compared to control. Carbonic anhydrases are ubiquitous enzymes that catalyze the interconversion of HCO₃⁻ and CO₂ and are involved in a range of physiological processes that include pH homeostasis (see for review Pastorekova *et al.*, 2004; [Supuran, 2008](#)). Three transcripts encoding members of the SLC15A4 family (Solute carrier family 15 member 4) were also up-regulated. SLC15A4 are proton oligopeptide co-transporters that are required for lysosomal pH regulation and V-type H⁺-ATPase integrity (Kobayashi *et al.*, 2014).

Calcification

The set of differentially expressed genes was searched against a database for database for metazoan biomineralization proteins (<https://peerj.com/preprints/1983/>) and 26 candidates were identified. All were up-regulated in response to elevated CO₂ (Table 3 and Table S6). They include 9 transcripts encoding a putative metalloproteinase, an alkaline phosphatase, two chitin synthases, a transcript encoding a collagen protein and four transcripts encoding cartilage matrix proteins. Two transcripts encoding mucin-like proteins were also highly up-

regulated (average of 17.5-fold). We found two perlucin-like transcripts together with a C-type lectin transcript to be up-regulated, the latter being 600-times up-regulated in the low pH condition compared to control. C-type lectins have been proposed to be involved in avian eggshell calcification ([Mann & Siedler, 2004](#)) and perlucin previously found to be involved in Mollusca biomineralization ([Mann *et al.*, 2000](#)). Interestingly, one transcript showing similarities with a dentin sialophosphoprotein was 9-fold up-regulated under elevated CO₂. Sialophosphoprotein has been shown to be involved in tooth and bone formation ([Prasad *et al.*, 2010](#)). Finally, the two transcripts encoding subunits of the vacuolar-type proton ATPase and the transcript encoding a carbonic anhydrase that were up-regulated in this experiment and mentioned in the “*Acid-base regulation*” section could also play a role in pteropod calcification.

Nervous system

Twenty-two percent of the up-regulated transcripts in the low pH condition (88 of the 400 up-regulated clusters) were genes putatively involved in the functioning of the nervous system.

Those 88 transcripts include several neural cell-adhesion molecules, proteins involved in the maintenance and formation of the nervous system, various neuropeptides and neuropeptide receptors, key players in synaptic vesicle and recovery at the synapses, as well as ion channels potentially involved in synapse communication (Table 3 and Table S7).

This set of genes comprised several ligand-gated ion channels and their associated proteins from three types of synapses: GABAergic, cholinergic and glutamatergic. This includes the GABA_A receptor (GABA_AR), which is of particular interest in relation to recent physiological studies showing that elevated CO₂ alters the behavior of fish ([Heuer & Grosell, 2014](#); [Nilsson *et al.*, 2012](#)) and molluscs ([Watson *et al.*, 2014](#)). In the present study, a transcript encoding a subunit of the GABA_AR was up-regulated 16-fold in low pH compared to control conditions. A transcript encoding a glycine receptor subunit was also up-regulated

(GlyR, 7.7-fold up-regulated), an inhibitory ligand-gated ion channel that is known to co-localize with GABA_AR on some hippocampal neurons (Table 3 and Table S7) (Lévi *et al.*, 2004). A surprising number of transcripts involved in cholinergic synapses were also up-regulated. Fourteen transcripts encoding subunits of acetylcholine receptors (nicotinic and muscarinic), together with one voltage-gated potassium channel and three transcripts encoding acetylcholinesterase were up-regulated (Table 3 and Table S7). Acetylcholine receptors occur both on neurons (in ganglia and brain) and on muscles. The diffusion of Na⁺ and K⁺ across the receptor causes depolarization that opens voltage-gated sodium/potassium channels and allows firing of the action potential and potentially muscular contraction. Acetylcholinesterases hydrolyze the neurotransmitter acetylcholine and are, therefore, essential for the termination of synaptic transmission in cholinergic synapses.

Finally, the mRNA of a glutamate receptor responsible for the glutamate-mediated postsynaptic excitation of neural cells, and the mRNA of a glutamate transporter, were also up-regulated at lower pH (4.8 and 4.7 times respectively).

Only two transcripts with potential roles in the nervous system were down-regulated in response to elevated CO₂: a transcript encoding a voltage-dependent calcium channel subunit and a transcript encoding a putative tenascin-R-like protein (see Table 3 and Table S7). It should be noted that 10 transcript encoding putative tenascin-R-like protein were differentially regulated; one being down-regulated 42-fold and the other ones being up-regulated on average 44-fold under low pH. Tenascin-R proteins (TN-R) are extracellular matrix proteins exclusive to the central nervous system in vertebrates ([Anlar & Gunel-Ozcan, 2012](#)). They have versatile roles and can act as adhesive or anti-adhesive molecules towards various neural and non-neural cells but also inhibitors or enhancers of neurite outgrowth ([Pesheva & Probstmeier, 2000](#)).

Lastly, the mRNA of an amiloride-sensitive cation channel 4 (ASIC 4 or ACCN4) was up-regulated about 7-fold under low pH. These channels have been implicated in synaptic transmission, pain perception as well as mechano-reception in mammals, and were found in zebrafish neurons ([Chen *et al.*, 2007](#)).

DISCUSSION

Physiological and molecular tools were used to investigate the response of the Mediterranean pteropod *Heliconoides inflatus* to seawater pH likely to be reached by 2100 under a moderate emissions trajectory (RCP6.0). *Heliconoides inflatus* individuals were subjected to pH_T 8.1 (control) or pH_T 7.9 for 3 d and changes in gene expression, calcification and respiration were measured. Gross calcification strongly decreased in low pH conditions while genes potentially involved in calcification were up-regulated, suggesting that pteropods attempt to maintain calcification rates when faced with unfavorable conditions. Interestingly, a large number of genes related to nervous system structure and function were also up-regulated in the low pH treatment, including a GABA_A receptor subunit. This is particularly interesting given that GABA_A receptor disturbances have been documented in several marine organisms after exposures to elevated CO₂ ([Heuer & Grosell, 2014](#); [Nilsson *et al.*, 2012](#); [Watson *et al.*, 2014](#)).

It is important to recognize that our experiment was an acute exposure of pteropods to low pH conditions for only three days, and physiological and molecular effects could be different under longer exposure, as previously shown in corals ([Moya *et al.*, 2015](#)). While our study may seem short in comparison to similar studies on other organisms, the difficulties in maintaining these planktonic molluscs under laboratory conditions ([Howes *et al.*, 2014](#)) places our study amongst the longest experiments for non-polar pteropods species.

Suppression of metabolism and protein synthesis

During the course of the present experiment, oxygen consumption did not differ between pH_T 7.9 and pH_T 8.1. Previous studies also found no effect of lower pH on the respiration of *Limacina helicina* (Comeau *et al.*, 2010b), *Clio pyramidata*, *Hyalocylis striata*, *Cavolinia longirostris*, *Creseis virgula* (Maas *et al.*, 2015; Maas *et al.*, 2012) and *Creseis acicula* (Comeau *et al.*, 2012b) while it decreased in *L. helicina forma antarctica* (Seibel *et al.*, 2012) and *Diacria quadridentata* (Maas *et al.*, 2012), suggesting that the natural seawater chemistry may influence their resilience to ocean acidification. However in the present study, exposure of *H. inflatus* to pH_T 7.9 for 3 d led to a decreased expression of metabolism-related genes, indicating that metabolism was suppressed under acidified conditions. The apparent contradiction between the transcriptomic and physiological measurements of the present study may be due to the long turnover times typical of respiratory complex proteins. The latter is not unexpected as all four respiratory complexes contain subunits encoded in the mitochondrial genome. Half-lives of human mitochondrial proteins range from 6 to 16 days (Eden *et al.*, 2011). Studies on the effects of elevated CO_2 on pteropods (Maas *et al.*, 2015), sea urchins (O'Donnell *et al.*, 2010; Todgham & Hofmann, 2009) and coral larvae (Moya *et al.*, 2012) have congruently documented the suppression of metabolic gene expression under elevated CO_2 as observed in the present study. It is widely accepted that metabolic depression is an adaptive strategy for survival in short-term energy limitation in aquatic organisms (Seibel & Walsh, 2003) and is accomplished, at least in part, by shutting down processes such as protein synthesis (Guppy & Withers, 1999), and in particular mitochondrial protein synthesis (Kwast & Hand, 1996). In our experiment, the entire protein synthesis machinery, including transcripts of initiation and elongation factors as well as transcripts of ribosomal proteins, was subject to down-regulation at low pH. Depression of metabolic gene expression

upon exposure to acute stress potentially allows the reallocation of transcriptional resources (and energy) to more immediate demands such as pH homeostasis.

Acid-base regulation

pH homeostasis is crucial for a large range of systemic and cellular functions, including calcification and neural function, but the pathways involved, as well as the efficiency of compensation mechanisms of acid-base imbalance differ between taxa and are often species-specific (Clements & Hunt, 2015; Melzner *et al.*, 2009). Although acid-base regulation is often achieved in molluscs by controlling levels of bicarbonate ions (Pörtner, 2008), in the present case, the expression of bicarbonate transporter genes did not differ between low pH and control treatments. In contrast, eight genes involved in acid-base regulation were found to be up-regulated, including carbonic anhydrases and V-type H⁺-ATPases. This observation suggests that pH homeostasis in *H. inflatus* could be achieved by means other than active bicarbonate transport. For example, carbonic anhydrases interconvert CO₂ into HCO₃⁻ while the protons that this generates are removed by V-type H⁺-ATPases, contributing to the accumulation of bicarbonate ions and pH homeostasis at lower pH. Consistent with the strong decrease in calcification rates observed in this study at low pH, prior research on bivalves suggests that bicarbonate buffering can be partly achieved by the dissolution of the CaCO₃ exoskeletons (Lindinger *et al.*, 1984; Michaelidis *et al.*, 2005). However, this strategy is of limited usefulness in the case of pteropods, considering the thin shell of these planktonic molluscs.

Irrespective of the nature of the strategy used by pteropods to achieve acid-base balance, the extent to which compensation is achieved is unknown, and this has important implications for other physiological processes. In fish, for example, the compensatory response during acid-base regulation is claimed to be responsible for the observed disturbances in neural function

and behavior ([Heuer & Grosell, 2014](#)). An additional consideration is that several proteins involved in acid-base regulation are also involved in other physiological processes, such as calcification and neural functioning (e.g. carbonic anhydrases, V-type H⁺-ATPases). At this stage, however, it is unclear in which processes the differentially expressed isoforms identified in this study are involved. Understanding the mechanisms underlying basic processes such as acid-base regulation or calcification is a prerequisite for predicting the effects of future ocean conditions on pteropod populations.

Calcification decreases despite increased expression of calcification genes

The major (37%) decrease in gross calcification observed after exposure of *H. inflatus* to pH_T 7.9 is consistent with data for the Mediterranean pteropod species *Creseis acicula* ([Comeau et al., 2012b](#) reported a 30% decrease). The decline in calcification could be the result of an active reduction in this energetically expensive process at low pH in order to permit reallocation of energy to other processes. However, this hypothesis is difficult to reconcile with the observation that a number of genes involved in calcification were up-regulated at lower pH. Although the detail of where these genes function in the calcification process is unknown, these data imply that the observed decrease in gross calcification reflects the inability of the pteropod to sustain calcification rates under acidic conditions, rather than active down-regulation of the calcification process.

Amongst the calcification-related genes up-regulated in our study are two perlucin transcripts and a C-type lectin which were also differentially expressed in similar experiments on the pteropods *Clio pyramidata* ([Maas et al., 2015](#)) and *Limacina helicina* ([Koh et al., 2015](#)).

While comparisons between these studies are complicated by major differences in experimental design, it is noteworthy that following a 10 h exposure of *C. pyramidata* to a pH level similar to the one used in the present experiment (pH_T 7.8), a perlucin homolog was

one of the few differentially expressed genes, with a 154-fold increase in expression ([Maas et al., 2015](#)). In contrast, [Koh et al. \(2015\)](#) documented an opposite trend for a C-type lectin in *L. helicina* under more extreme (pH 7.5 and 6.5) conditions than those used here. One factor complicating comparison of data reported here with published analyses is that many of the genes involved in calcification are members of large multi-gene families, hence the assignment of orthology is difficult. Large numbers of perlucin and C-type lectins are known to be present in *C. pyramidata* ([Maas et al., 2015](#)), and in the present study, approximately 300 contigs containing C-type lectin domains (Pfam PF00059) were retrieved from the *H. inflatus* transcriptome (data not shown). C-type lectin domain proteins have a diverse range of roles, including cell-cell adhesion, immune response and apoptosis ([Drickamer, 1999](#)). It is therefore likely that different isoforms have different roles in the response of pteropods to elevated CO₂.

The contrast between physiological and molecular results suggests that pteropods unsuccessfully attempt to maintain calcification rates under unfavorable conditions. If shelled pteropods could calcify faster than their shell dissolves, as has been observed for some aragonite-based corals ([Rodolfo-Metalpa et al., 2011](#)), then they could survive under near future ocean conditions ([Ries, 2012](#)). However, when considered in conjunction with the extensive shell dissolution that has previously been documented for Ω_a levels ~ 1 ([Bednaršek et al., 2012b](#)), the data presented here implies a poor prognosis for shelled pteropod populations.

Low pH affects the nervous system

In the present study, 20% of the genes up-regulated under low pH are likely to function in the nervous system. This is particularly interesting in light of recent studies showing that ocean acidification influences the behavior of both vertebrates and invertebrates in multiple ways

(see Table 1 and 2 of [Clements & Hunt, 2015](#) for a review). Some of the previous studies implicate the GABA-A receptor (GABA_AR, up-regulated 16-fold in the present study) in behavioral changes by demonstrating that gabazine, a specific GABA_AR antagonist, almost completely restores the behavioral performance of fishes ([Chivers *et al.*, 2014](#); [Hamilton *et al.*, 2014](#); [Lai *et al.*, 2015](#); [Nilsson *et al.*, 2012](#)) and gastropod molluscs (e.g. [Watson *et al.*, 2014](#)). The GABA_A receptor is a ligand-gated ion channel activated by GABA, the most common inhibitory neurotransmitter in vertebrates and some invertebrates ([Jessen *et al.* 1979](#)). GABA_A has specific conductance for Cl⁻ and HCO₃⁻, two ions likely to be impacted by ocean acidification ([Nilsson *et al.*, 2012](#)). Under normal conditions, when GABA_A opens, letting Cl⁻ and HCO₃⁻ ions flow into the cell, depolarization of the membrane is prevented, reducing neural activity. GABA_AR are pH sensitive proteins; minor changes in external pH are sufficient to affect their function, affecting neuronal excitability ([Wilkins *et al.*, 2005](#)).

When extracellular pH decreases, most animals (depending on their acid-base regulation strategy) excrete Cl⁻ and accumulate HCO₃⁻ in order to prevent cellular acidosis ([Heuer & Grosell, 2014](#)), which renders GABA_AR signaling excitatory ([Lambert & Grover, 1995](#)), thereby affecting behavior and causing dramatic shifts in sensory preferences ([Nilsson *et al.*, 2012](#)). Whether this hypothesis applies to pteropods remains to be determined, and this will depend on elucidating the acid-base balance strategies of the taxon. If pteropods accumulate HCO₃⁻ (through carbonic anhydrase rather than bicarbonate transporters) in order to avoid cellular acidosis, a mechanism similar to that of fishes is likely to be present. If the acid-base balance is not bicarbonate dependent, GABA_AR should not be specifically affected.

However, irrespective of the acid-base strategy used, if compensation is incomplete then more pH sensitive (neural and other) proteins will be affected at low pH. In the present study, the up-regulation of many other genes involved in neuron function, including several other voltage-dependent neural proteins belonging to glutamatergic and cholinergic synapses,

supports the latter hypothesis and suggests that the effect of low pH might be less specific than proposed in fishes, and could potentially affect other cellular functions. Alternatively, if GABA_A receptor function is altered, that could have wide ranging effects leading to responses in other transmitter systems, which in turn could explain the widespread changes seen in neural gene expression.

A limitation of our study is the lack of behavioral observations that could provide context for the interpretation of the observed changes in neural gene expression. To our knowledge, only one study has investigated the effects of low pH (in combination with altered salinity) on the behavior of pteropods ([Manno *et al.*, 2012](#)). It found that the combined stressors negatively affected upward swimming. In part, the lack of behavioral studies on shelled pteropods is due to the difficulty of maintaining them under laboratory conditions ([Howes *et al.*, 2014](#)).

Clearly, more work is needed to uncover the role of GABA_A receptors in the behavior of invertebrates under elevated CO₂, as well as a better understanding of the link between acid-base regulation and neural function in molluscs.

Planktonic molluscs such as pteropods are likely to normally experience large changes in the carbonate chemistry due to diel vertical migrations in the water column ([Maas *et al.*, 2012](#)).

This life history strategy suggests that pteropods should have molecular mechanisms to cope with sudden changes in pH. This diel migration also involves substantial changes in temperature, a parameter that may act synergistically with CO₂, and potentially amplify or attenuate the pteropod response to high CO₂ alone ([Clements & Hunt, 2015](#)). Scrutinizing the combined effect of CO₂ and temperature on *H. inflatus*, as done by [Comeau *et al.*](#) on the arctic species *L. helicina*, will be the next important step towards better understanding the impacts of global changes in climate and ocean chemistry on these key planktonic organisms.

In the present study, strong decrease in calcification was observed at Ω_a levels around 2.

[Comeau *et al.* \(2010b\)](#) have shown that the Arctic pteropod *Limacina helicina* is unable to

precipitate calcium carbonate at aragonite saturation state close to 1 and that gross calcification declines well above this value according to a logarithmic relationship. The projected increase in temperature does not ameliorate the situation. Both net and gross calcification also decline well before an aragonite saturation state of 1 in the Mediterranean pteropod *Creseis acicula* (Comeau *et al.*, 2012b). At ecologically relevant changes in saturation state in the Mediterranean or tropical/sub-tropical open ocean, saturation states are not expected to go below 1. If these organisms have significant changes in calcification at the reported levels, the scientific community needs to pay particular attention to non-polar pteropods in these regions.

This study advances our understanding of the responses of pteropods to ocean acidification and illustrates the utility of combining physiological and molecular approaches. When considered in the context of previous work on fish (Chivers *et al.*, 2014; [Hamilton *et al.*, 2014](#); [Lai *et al.*, 2015](#); [Nilsson *et al.*, 2012](#)) and other molluscs ([Watson *et al.*, 2014](#)), the major impacts of ocean acidification on expression of genes likely to function in the pteropod nervous system lead us to recommend that future studies of the impacts of ocean acidification on animal biology should include behavioral observations together with physiological and molecular measurements.

Comprehensive studies of this kind will clearly be needed in order to accurately predict the impacts of ocean acidification on individual species and the food webs in which they are involved.

Acknowledgement

This research was supported by a Marie Curie International Outgoing Fellowship (grant agreement # PEOF-GA-2008-235142 project title AMICAL) to AM. This work is a

contribution to the ‘European Project on Ocean Acidification’ (EPOCA) and ‘Mediterranean Sea Acidification under a changing climate’ (MedSeA) projects which received funding from the European Community’s Seventh Framework Program (FP7/2007-2013) under grant agreements # 211384 and # 265103. MM and BH were funded by US NSF grants IOS 1146880 and OCE 1442206. The IAEA is grateful to the Government of the Principality of Monaco for the support provided to its Environment Laboratories. The authors gratefully thank Dr. Sofia Fortunato for constructive comments on previous version of the manuscript.

Figure Legends

Figure 1. Individual of *Heliconoides inflatus*.

Figure 2. (a) Calcium uptake and (b) respiration rates measured at control (pH_T 8.1) and low (pH_T 7.9) pH levels. Asterisk indicates significant difference between control and low pH ($t = 6.0$, $df = 3$, p -value = 0.01).

Figure 3. Distribution of the log₂ (fold-change) of the differentially expressed genes in response to the low pH treatment (adjusted $P < 0.05$).

Supporting information captions

Figure S1. Shell diameter-weight relationship of *Heliconoides inflatus* individuals used in the present study. Individuals were photographed under a binocular microscope, and the maximum shell diameter measured using the imaging software AxioVision, version 6.1.7601. Individuals were dried with tissue paper prior to weighting with a Mettler Toledo microbalance (precision = 0.1 μ g).

Figure S2. Distribution of GC content (a) before and (b) after filtering the contaminating data.

Table S1. Characteristics of the transcriptome assemblies before and after filtering the contaminating data.

Table S2. Genes responsive to the low pH treatment belonging to the GO category *Ion transport* (GO:0006811).

Table S3. Protein synthesis genes responsive to low pH treatment.

Table S4. Mitochondrion genes responsive to low pH treatment.

Table S5. Genes potentially involved in acid-base regulation that were affected by the low pH treatment.

Table S6. Genes potentially involved in pteropod calcification that were affected by the low pH treatment.

Table S7. Genes potentially involved in pteropod nervous system that were affected by the low pH treatment.

Table S8. Shell diameters (μm) and weights (μg) of all individuals used in the respiration experiments. Number in brackets next to pH correspond to the replicate.

Table S9. Results of the respiration experiments showing the $\mu\text{mol O}_2$ consumed per hour per μg of wet weight. Number in brackets next to pH correspond to the replicate.

Data accessibility: The data discussed in this publication have been deposited in NCBI's Gene Expression Omnibus ([Edgar *et al.*, 2002](https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE77934)) and are accessible through GEO Series accession number GSE77934 (<https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE77934>).

References

- Anlar B, Gunel-Ozcan A (2012) Tenascin-R: role in the central nervous system. The International Journal of Biochemistry & Cell Biology, **44**, 1385-1389.
- Bé AWH, Gilmer RW (1977) A zoogeographic and taxonomic review of euthecosomatous pteropoda. In: *Oceanic Micropaleontology*. (ed Ramsey ATS) pp Page. London, Academic Press.
- Bednaršek N, Feely RA, Reum JCP, Peterson B, Menkel J, Alin SR, Hales B (2014) *Limacina helicina* shell dissolution as an indicator of declining habitat suitability owing to ocean acidification in the California Current Ecosystem. Proceedings of the Royal Society of London B: Biological Sciences, **281**.
- Bednaršek N, Možina J, Vogt M, O'Brien C, Tarling GA (2012a) The global distribution of pteropods and their contribution to carbonate and carbon biomass in the modern ocean. Earth System Science Data, **4**, 167-186.
- Bednaršek N, Ohman MD (2015) Changes in pteropod distributions and shell dissolution across a frontal system in the California Current System. Marine Ecology Progress Series, **523**, 93-103.
- Bednaršek N, Tarling GA, Bakker DCE *et al.* (2012b) Extensive dissolution of live pteropods in the Southern Ocean. Nature Geoscience, **5**, 881-885.
- Briffa M, de la Haye K, Munday PL (2012) High CO₂ and marine animal behaviour: Potential mechanisms and ecological consequences. Marine Pollution Bulletin, **64**, 1519-1528.
- Chen X, Polleichtner G, Kadurin I, Gründer S (2007) Zebrafish acid-sensing ion channel (ASIC) 4, characterization of homo- and heteromeric channels, and identification of regions important for activation by H⁺. Journal of Biological Chemistry, **282**, 30406-30413.
- Chivers DP, McCormick MI, Nilsson GE *et al.* (2014) Impaired learning of predators and lower prey survival under elevated CO₂: a consequence of neurotransmitter interference. Global Change Biology, **20**, 515-522.
- Clements JC, Hunt HL (2015) Marine animal behaviour in a high CO₂ ocean. Marine Ecology Progress Series, **536**, 259-279.
- Comeau S, Alliouane S, Gattuso J-P (2012a) Effects of ocean acidification on overwintering juvenile Arctic pteropods *Limacina helicina*. Marine ecology Progress Series, **456**, 279-284.
- Comeau S, Gattuso J-P, Jeffree R, Gazeau F (2012b) Effect of carbonate chemistry manipulations on calcification, respiration, and excretion of a Mediterranean pteropod. Biogeosciences discussions, **9**, 6169-6189.
- Comeau S, Gorsky G, Alliouane S, Gattuso J-P (2010a) Larvae of the pteropod *Cavolinia inflexa* exposed to aragonite undersaturation are viable but shell-less. Marine biology, **157**, 2341-2345.
- Comeau S, Gorsky G, Jeffree R, Teyssié J-L, Gattuso J-P (2009) Impact of ocean acidification on a key Arctic pelagic mollusc (*Limacina helicina*). Biogeosciences, **6**, 1877-1882.
- Comeau S, Jeffree R, Teyssié J-L, Gattuso J-P (2010b) Response of the Arctic pteropod *Limacina helicina* to projected future environmental conditions. PLoS One, **5**, e11362.

- Conesa A, Gotz S, Garcia-Gomez JM, Terol J, Talon M, Robles M (2005) [Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research](#). *Bioinformatics*, **21**, 3674-3676.
- Dickson AG (1990) [Standard potential of the reaction: \$\text{AgCl\(s\)} + 1/2\text{H}_2\text{\(g\)} = \text{Ag\(s\)} + \text{HCl\(aq\)}\$, and the standard acidity constant of the ion \$\text{HSO}_4^-\$ in synthetic sea water from 273.15 to 318.15 K](#). *The Journal of Chemical Thermodynamics*, **22**, 113-127.
- Dickson AG, Sabine CL, Christian JR (2007) [Guide to best practices for ocean \$\text{CO}_2\$ measurements](#), PICES Special Publication 3.
- Drickamer K (1999) [C-type lectin-like domains](#). *Current Opinion in Structural Biology*, **9**, 585-590.
- Eden E, Geva-Zatorsky N, Issaeva I *et al.* (2011) [Proteome half-life dynamics in living human cells](#). *Science*, **331**, 764-768.
- Edgar R, Domrachev M, Lash AE (2002) [Gene Expression Omnibus: NCBI gene expression and hybridization array data repository](#). *Nucleic Acids Research*, **30**, 207-210.
- Foster BA, Montgomery JC (1993) [Planktivory in benthic nototheniid fish in McMurdo Sound, Antarctica](#). *Environmental Biology of Fishes*, **36**, 313-318.
- Fu L, Niu B, Zhu Z, Wu S, Li W (2012) [CD-HIT: accelerated for clustering the next-generation sequencing data](#). *Bioinformatics*, **28**, 3150-3152.
- Gazeau F, Parker LM, Comeau S *et al.* (2013) [Impacts of ocean acidification on marine shelled molluscs](#). *Marine Biology*, **160**, 2207-2245.
- Grabherr MG, Haas BJ, Yassour M *et al.* (2011) [Full-length transcriptome assembly from RNA-Seq data without a reference genome](#). *Nature Biotechnology*, **29**, 644-652.
- Guppy M, Withers P (1999) [Metabolic depression in animals: physiological perspectives and biochemical generalizations](#). *Biological reviews of the Cambridge Philosophical Society*, **74**, 1-40.
- Hamilton TJ, Holcombe A, Tresguerres M (2014) [\$\text{CO}_2\$ -induced ocean acidification increases anxiety in Rockfish via alteration of \$\text{GABA}_A\$ receptor functioning](#). *Proceedings of the Royal Society of London B: Biological Sciences*, **281**, 20132509.
- Heuer RM, Grosell M (2014) [Physiological impacts of elevated carbon dioxide and ocean acidification on fish](#). *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology*, **307**, R1061-R1084.
- Howes EL, Bednaršek N, Bűdenbender J *et al.* (2014) [Sink and swim: a status review of thecosome pteropod culture techniques](#). *Journal of Plankton Research*.
- Kobayashi T, Shimabukuro-Demoto S, Yoshida-Sugitani R *et al.* (2014) [The histidine transporter SLC15A4 coordinates mTOR-dependent inflammatory responses and pathogenic antibody production](#). *Immunity*, **41**, 375-388.
- Koh HY, Lee JH, Han SJ, Park H, Shin SC, Lee SG (2015) [A transcriptomic analysis of the response of the arctic pteropod *Limacina helicina* to carbon dioxide-driven seawater acidification](#). *Polar Biology*, **38**, 1727-1740.
- Kwast KE, Hand SC (1996) [Acute depression of mitochondrial protein synthesis during anoxia: contributions of oxygen sensing, matrix acidification, and redox state](#). *Journal of Biological Chemistry*, **271**, 7313-7319.
- Lai F, Jutfelt F, Nilsson GE (2015) [Altered neurotransmitter function in \$\text{CO}_2\$ -exposed stickleback \(*Gasterosteus aculeatus*\): a temperate model species for ocean acidification research](#). *Conservation Physiology*, **3**, cov018.
- Lambert N, Grover L (1995) [The mechanism of biphasic GABA responses](#). *Science*, **269**, 928-929.
- Langmead B, Salzberg SL (2012) [Fast gapped-read alignment with Bowtie 2](#). *Nature Methods*, **9**, 357-359.

- Lavigne H, Gattuso J-P (2013) Seacarb 1.2.3., an R package to calculate parameters of the seawater carbonate system.
- Lévi S, Logan SM, Tovar KR, Craig AM (2004) Gephyrin is critical for glycine receptor clustering but not for the formation of functional GABAergic synapses in hippocampal neurons. *The Journal of Neuroscience*, **24**, 207-217.
- Lindinger MI, Lauren DJ, McDonald DG (1984) Acid-base balance in the sea mussel, *Mytilus edulis*. III. Effects of environmental hypercapnia on intra- and extracellular acid-base balance. *Marine Biology Letters*, **5**, 371-381.
- Lischka S, Büdenbender J, Boxhammer T, Riebesell U (2011) Impact of ocean acidification and elevated temperatures on early juveniles of the polar shelled pteropod *Limacina helicina*: mortality, shell degradation, and shell growth. *Biogeosciences*, **8**, 919-932.
- Lischka S, Riebesell U (2012) Synergistic effects of ocean acidification and warming on overwintering pteropods in the Arctic. *Global Change Biology*, **18**, 3517-3528.
- Lueker TJ, Dickson AG, Keeling CD (2000) Ocean pCO₂ calculated from dissolved inorganic carbon, alkalinity, and equations for K₁ and K₂: validation based on laboratory measurements of CO₂ in gas and seawater at equilibrium. *Marine Chemistry*, **70**, 105-119.
- Maas AE, Lawson GL, Tarrant AM (2015) Transcriptome-wide analysis of the response of the thecosome pteropod *Clio pyramidata* to short-term CO₂ exposure. *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics*, **16**, 1-9.
- Maas AE, Wishner KF, Seibel BA (2012) The metabolic response of pteropods to acidification reflects natural CO₂-exposure in oxygen minimum zones. *Biogeosciences*, **9**, 747-757.
- Mann K, Siedler F (2004) Ostrich (*Struthio camelus*) eggshell matrix contains two different C-type lectin-like proteins. Isolation, amino acid sequence, and posttranslational modifications. *Biochimica et Biophysica Acta*, **1696**, 41-50.
- Mann K, Weiss IM, André S, Gabius H-J, Fritz M (2000) The amino-acid sequence of the abalone (*Haliotis laevigata*) nacre protein perlucin. *European Journal of Biochemistry*, **267**, 5257-5264.
- Manno C, Morata N, Primicerio R (2012) *Limacina retroversa*'s response to combined effects of ocean acidification and sea water freshening. *Estuarine, Coastal and Shelf Science*, **113**, 163-171.
- Martin S, Richier S, Pedrotti M-L *et al.* (2011) Early development and molecular plasticity in the Mediterranean sea urchin *Paracentrotus lividus* exposed to CO₂-driven acidification. *Journal of Experimental Biology*, **214**, 1357-1368.
- Melzner F, Gutowska MA, Langenbuch M *et al.* (2009) Physiological basis for high CO₂ tolerance in marine ectothermic animals: pre-adaptation through lifestyle and ontogeny? *Biogeosciences*, **6**, 2313-2331.
- Michaelidis B, Ouzounis C, Paleras A, Pörtner H-O (2005) Effects of long-term moderate hypercapnia on acid-base balance and growth rate in marine mussels *Mytilus galloprovincialis*. *Marine Ecology Progress Series*, **293**, 109-118.
- Moya A, Huisman L, Ball E *et al.* (2012) Whole transcriptome analysis of the coral *Acropora millepora* reveals complex responses to CO₂-driven acidification during the initiation of calcification. *Molecular Ecology*, **21**, 2440-2454.
- Moya A, Huisman L, Forêt S, Gattuso J-P, Hayward DC, Ball EE, Miller DJ (2015) Rapid acclimation of juvenile corals to CO₂-mediated acidification by up-regulation of HSP and Bcl-2 genes. *Molecular Ecology*, **24**, 438-452.
- Nilsson GE, Dixon DL, Domenici P, McCormick MI, Sorensen C, Watson SA, Munday PL (2012) Near-future carbon dioxide levels alter fish behaviour by interfering with neurotransmitter function. *Nature Climate Change*, **2**, 201-204.

- O'Donnell MJ, Todgham AE, Sewell MA *et al.* (2010) Ocean acidification alters skeletogenesis and gene expression in larval sea urchins. *Marine Ecology Progress Series*, **398**, 157-171.
- Orr JC, Fabry VJ, Aumont O *et al.* (2005) Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature*, **437**, 681-686.
- Pastorekova S, Parkkila S, Pastorek J, Supuran CT (2004) Carbonic anhydrases: current state of the art, therapeutic applications and future prospects. *Journal of Enzyme Inhibition and Medicinal Chemistry*, **19**, 199-229.
- Perez FF, Fraga F (1987) Association constant of fluoride and hydrogen ions in seawater. *Marine Chemistry*, **21**, 161-168.
- Pesheva P, Probstmeier R (2000) The yin and yang of tenascin-R in CNS development and pathology. *Progress in Neurobiology*, **61**, 465-493.
- Pörtner H-O (2008) Ecosystem effects of ocean acidification in times of ocean warming: a physiologist's view. *Marine Ecology Progress Series*, **373**, 203-217.
- Prasad M, Butler WT, Qin C (2010) Dentin sialophosphoprotein (DSPP) in biomineralization. *Connective Tissue Research*, **51**, 404-417.
- R Development Core Team (2010) R: A language and environment for statistical computing. Vienna, Austria, R Foundation for Statistical Computing.
- Rampal J (1975) Les thécosomes (mollusques pélagiques). Systématique et évolution - Écologies et biogéographie Méditerranéennes. Unpublished Thèse de Doctorat ès Sciences Université Aix-Marseille I 485 pp.
- Ries JB (2012) Oceanography: A sea butterfly flaps its wings. *Nature Geoscience*, **5**, 845-846.
- Robinson MD, McCarthy DJ, Smyth GK (2010) edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*, **26**, 139-140.
- Rodolfo-Metalpa R, Houlbreque F, Tambutte E *et al.* (2011) Coral and mollusc resistance to ocean acidification adversely affected by warming. *Nature Climate Change*, **1**, 308-312.
- Sadamoto H, Takahashi H, Okada T, Kenmoku H, Toyota M, Asakawa Y (2012) *De novo* sequencing and transcriptome analysis of the central nervous system of mollusc *Lymnaea stagnalis* by deep RNA sequencing. *PLoS ONE*, **7**, e42546.
- Sambrook J, Russel DW (2001) *Molecular cloning : a laboratory manual*, Cold Spring Harbor, N.Y, Cold Spring Harbor Laboratory.
- Seibel BA, Maas AE, Dierssen HM (2012) Energetic plasticity underlies a variable response to ocean acidification in the pteropod, *Limacina helicina antarctica*. *PLoS ONE*, **7**, e30464.
- Seibel BA, Walsh PJ (2003) Biological impacts of deep-sea carbon dioxide injection inferred from indices of physiological performance. *Journal of Experimental Biology*, **206**, 641-650.
- Supuran CT (2008) Carbonic anhydrases--an overview. *Current Pharmaceutical Design*, **14**, 603-614.
- Todgham AE, Hofmann GE (2009) Transcriptomic response of sea urchin larvae *Strongylocentrotus purpuratus* to CO₂-driven seawater acidification. *Journal of Experimental Biology*, **212**, 2579-2594.
- Watson S-A, Lefevre S, McCormick MI, Domenici P, Nilsson GE, Munday PL (2014) Marine mollusc predator-escape behaviour altered by near-future carbon dioxide levels. *Proceedings of the Royal Society of London B: Biological Sciences*, **281**, 20132377.

Wilkins ME, Hosie AM, Smart TG (2005) Proton modulation of recombinant GABA(A) receptors: influence of GABA concentration and the β subunit TM2–TM3 domain. *The Journal of Physiology*, **567**, 365-377.

Table 1. Carbonate chemistry parameters in the control and low pH treatments. pH_T

and A_T were measured while all other parameters were estimated using the R package

seacarb. C_T , dissolved inorganic carbon; A_T , total alkalinity, Ω_a , saturation state of aragonite.

Three separate experiments were performed to assess (1) respiration rate, (2) ^{45}Ca uptake, and (3) gene expression. Experiments (1) and (3) were performed at Laboratoire d’Océanographie de Villefranche and experiment (2) was performed at IAEA.

Treatment	pH_T	pCO_2 (μatm)	HCO_3^- ($\mu\text{mol kg}^{-1}$)	CO_3^{2-} ($\mu\text{mol kg}^{-1}$)	C_T ($\mu\text{mol kg}^{-1}$)	A_T ($\mu\text{mol kg}^{-1}$)	Ω_a
Control pH (1) and (3)	8.1	410	2091	188	2294 ± 10	2552 ± 4	2.8
Low pH (2)	7.9	617	2208	139	2370 ± 11	2550 ± 9	2.1
Control pH (1) and (3)	8.1	382	2059	196	2269 ± 17	2540 ± 5	2.9
Low pH (2)	7.9	720	2243	123	2393 ± 21	2561 ± 21	1.9

Table 2. Gene Ontology enrichment table. GO enrichment analysis was carried out with the Blast2GO software using a Fisher's exact test with a False Discovery Rate (FDR) threshold of 5% to reduce false-positive predictions of enriched GO terms. Down, categories enriched in the set of down-regulated genes in the low pH treatment; Up, category enriched in the set of up-regulated genes in the low pH treatment; MF, Molecular Function; CC, Cellular Component; BP Biological Process. The last column indicates the number of differentially expressed genes in each category.

Up- or down-regulated	Category	GO-ID	Term	FDR	# genes
DOWN	MF	GO:0003735	Structural constituent of ribosome	8E-34	46
		GO:0019843	rRNA binding	2E-06	8
		GO:0003729	mRNA binding	9E-05	9
		GO:0016491	Oxidoreductase activity	4E-02	21
		GO:0017111	Nucleoside-triphosphatase activity	4E-02	20
	CC	GO:0005739	Mitochondrion	5E-08	30
		GO:0015935	Small ribosomal subunit	9E-07	7
		GO:0005576	Extracellular region	2E-03	29
		GO:0005875	Microtubule associated complex	3E-03	4
		GO:0005730	Nucleolus	4E-03	13
		GO:0022625	Cytosolic large ribosomal subunit	4E-02	3
	BP	GO:0006091	Generation of precursor metabolites and energy	3E-07	16
		GO:0034655	Nucleobase-containing compound catabolic process	4E-07	16
		GO:0044403	Symbiosis encompassing mutualism through parasitism	9E-07	16
		GO:0044765	Single-organism transport	4E-05	35
		GO:0006605	Protein targeting	1E-04	13
		GO:0042254	Ribosome biogenesis	6E-04	10
		GO:0061024	Membrane organization	2E-03	14
UP	BP	GO:0008584	Male gonad development	2E-02	3
		GO:0006811	Ion transport	2E-02	26

Table 3. Summary of the genes responsive to low pH treatment. See supporting information for more detail about specific processes. Down, down-regulated genes in the low pH treatment; Up, up-regulated genes in the low pH treatment.

Process	Category/gene family	Number of genes	Expression
Protein synthesis	40S ribosomal proteins	24	down
	60S ribosomal proteins	26	down
	Other clusters involved in protein synthesis	19	down
Mitochondrion	Complex I	4	down
	Complex II	0	down
	Complex III	3	down
	Complex IV	10	down
	Complex V	6	down
Acid-base regulation	V-type H ⁺ -ATPase	2	up
	Na ⁺ -K ⁺ -ATPase	2	up
	Carbonic anhydrase	1	up
	Solute carrier family 15 member 4	3	up
Calcification	Metalloproteinases	9	up
	Alkaline phosphatase	1	up
	Chitin synthase	2	up
	Cartilage matrix proteins	4	up
	Mucins	2	up
	C-type lectins	3	up
	Dentin sialophosphoprotein	1	up
	Collagen	1	up
	V-type H ⁺ -ATPase	2	up
	Carbonic anhydrase	1	up
Nervous system	GABAergic synapses	2	up
	Cholinergic synapses	18	up
	Glutamaergic synapses	2	up
	Tenascin-R like	10	9 up, 1 down
	Acid-sensing ion channel	1	up
	Neuronal cell adhesion	15	up
	Neuropeptide receptors	3	up
	Neuronal differentiation and survival	6	up
	Postsynaptic scaffolding protein	4	up
	Vesicular and membrane trafficking	9	up
	Synaptic plasticity	1	up
	Receptor	1	up
	Neuro-transmitter transporters	4	up
	Calcium homeostasis	8	up
	Transporter	1	up
	Electrical coupling	1	up
Unknown role	2	up	



