



HAL
open science

The functional divergence between SPA1 and SPA2 in Arabidopsis photomorphogenesis maps primarily to the respective N-terminal kinase-like domain

Song Chen, Lennart Wirthmueller, Johannes Stauber, Niels Lory, Xu Holtkotte, Lisa Leson, Christian Schenkel, Margaret Ahmad, Ute Hoecker

► **To cite this version:**

Song Chen, Lennart Wirthmueller, Johannes Stauber, Niels Lory, Xu Holtkotte, et al.. The functional divergence between SPA1 and SPA2 in Arabidopsis photomorphogenesis maps primarily to the respective N-terminal kinase-like domain. BMC Plant Biology, 2016, 16 (1), pp.165. 10.1186/s12870-016-0854-9 . hal-01357390

HAL Id: hal-01357390

<https://hal.sorbonne-universite.fr/hal-01357390>

Submitted on 29 Aug 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

RESEARCH ARTICLE

Open Access



The functional divergence between SPA1 and SPA2 in Arabidopsis photomorphogenesis maps primarily to the respective N-terminal kinase-like domain

Song Chen^{1,3}, Lennart Wirthmueller^{1,4}, Johannes Stauber¹, Niels Lory¹, Xu Holtkotte¹, Lisa Leson¹, Christian Schenkel¹, Margaret Ahmad² and Ute Hoecker^{1*} 

Abstract

Background: Plants have evolved complex mechanisms to adapt growth and development to the light environment. The COP1/SPA complex is a key repressor of photomorphogenesis in dark-grown Arabidopsis plants and acts as an E3 ubiquitin ligase to ubiquitinate transcription factors involved in the light response. In the light, COP1/SPA activity is inhibited by photoreceptors, thereby allowing accumulation of these transcription factors and a subsequent light response. Previous results have shown that the four members of the SPA family exhibit partially divergent functions. In particular, SPA1 and SPA2 strongly differ in their responsiveness to light, while they have indistinguishable activities in darkness. The much higher light-responsiveness of SPA2 is partially explained by the much stronger light-induced degradation of SPA2 when compared to SPA1. Here, we have conducted SPA1/SPA2 domain swap experiments to identify the protein domain(s) responsible for the functional divergence between SPA1 and SPA2.

Results: We have individually swapped the three domains between SPA1 and SPA2 - the N-terminal kinase-like domain, the coiled-coil domain and the WD-repeat domain - and expressed them in *spa* mutant Arabidopsis plants. The phenotypes of transgenic seedlings show that the respective N-terminal kinase-like domain is primarily responsible for the respective light-responsiveness of SPA1 and SPA2. Furthermore, the most divergent part of the N-terminal domain was sufficient to confer a SPA1- or SPA2-like activity to the respective SPA protein. The stronger light-induced degradation of SPA2 when compared to SPA1 was also primarily conferred by the SPA2 N-terminal domain. At last, the different affinities of SPA1 and SPA2 for cryptochrome 2 are defined by the N-terminal domain of the respective SPA protein. In contrast, both SPA1 and SPA2 similarly interacted with COP1 in light-grown seedlings.

Conclusions: Our results show that the distinct activities and protein stabilities of SPA1 and SPA2 in light-grown seedlings are primarily encoded by their N-terminal kinase-like domains. Similarly, the different affinities of SPA1 and SPA2 for cry2 are explained by their respective N-terminal domain. Hence, after a duplication event during evolution, the N-terminal domains of SPA1 and SPA2 underwent subfunctionalization, possibly to allow optimal adaptation of growth and development to a changing light environment.

Keywords: Photomorphogenesis, SPA1, SPA2, COP1, E3 ubiquitin ligase, Functional divergence, Arabidopsis

* Correspondence: hoeckeru@uni-koeln.de

¹Botanical Institute and Cluster of Excellence on Plant Sciences (CEPLAS), Biocenter, University of Cologne, Zùlpicher Str. 47b, 50674 Cologne, Germany
Full list of author information is available at the end of the article



Background

Plants have evolved complex strategies to adapt to their changing environments. Light is among the most important environmental factors because it serves as the primary source of energy for photosynthesis. Light is also a signaling cue that controls many aspects of plant growth and development, including seed germination, seedling de-etiolation, phototropism, shade avoidance, anthocyanin production and the induction of flowering [1]. Several classes of photoreceptors evolved that constantly monitor light conditions and allow plants to rapidly respond to changing light conditions. These photoreceptors include the red- (R)/far-red- (FR) perceiving phytochromes, blue light (B)-sensing cryptochromes, phototropins and the ZEITLUPE family and the UV-B receptor UVR8 [2–4].

Arabidopsis seedlings grown in the dark display elongated hypocotyls, closed cotyledons and an apical hook. These etiolation phenotypes require the CONSTITUTIVELY PHOTOMORPHOGENIC1/SUPPRESSOR OF PHYA-105 (COP1/SPA) complex. Hence, dark-grown *cop1* mutants and *spa1 spa2 spa3 spa4* quadruple mutants exhibit constitutive de-etiolation, showing the phenotype of light-grown seedlings in darkness [5–7]. The COP1/SPA complex acts as a repressor of photomorphogenesis in the dark, while its repressor function is inhibited by light through multiple mechanisms. B-activated cryptochrome 1 (*cry1*) interacts with members of the SPA family to disrupt the interaction between COP1 and SPA proteins, leading to reduced COP1/SPA function and de-etiolation of the plant [8, 9]. Similarly, phytochromes interrupt the COP1/SPA interaction in red light [10, 11]. For *cry2*, COP1/SPA function is reduced by the enhancement of the *cry2*-COP1 interaction due to *cry2*-SPA interaction [12]. Another mechanism that inactivates COP1/SPA is based on the light-induced translocation of COP1 from the nucleus to the cytosol [13, 14]. A third mechanism involves the light-induced degradation of SPA1 and SPA2 [15, 16].

The COP1/SPA complex is part of the CULLIN4 (CUL4)-based multi-subunit E3 ubiquitin ligase CUL4-DDB1^{COP1/SPA} [17]. The substrates of this E3 ligase include several transcription factors such as ELONGATED HYPOCOTYL 5 (HY5), LONG HYPOCOTYL IN FR 1 (HFR1) and PRODUCTION OF ANTHOCYANIN PIGMENT (PAP) proteins which are responsible for light-induced photomorphogenesis [18–22]. In the dark, these transcription factors are ubiquitinated by the COP1/SPA-based E3 ubiquitin ligase, leading to their degradation in the 26S proteasome. When the COP1/SPA complex is inhibited by photoreceptors in the light, these transcription factors are stabilized to facilitate their functions in activating the light responses.

COP1 contains an N-terminal RING-finger domain, a coiled-coil domain and C-terminal WD-repeats. The

four SPA proteins (SPA1-SPA4) are structurally related to COP1 in that they contain a coiled-coil domain and WD-repeats. However, the N-termini of COP1 and SPAs are distinct, with COP1 having a RING-finger domain and SPAs harboring a kinase-like domain [22, 23]. COP1 and SPAs can form homo- and heterodimers via their respective coiled-coil domains and the COP1/SPA complex forms a tetramer of two COP1 and two SPA proteins. Both COP1 and SPA proteins can interact with most substrates and with DDB1 through their C-terminal WD-repeat domains [17, 24–26]. The function of the SPA kinase-like domain, in contrast, is not well understood. It displays weak sequence similarity to Ser/Thr protein kinases, though many normally invariant amino acids are not conserved in the kinase-like domain of SPA proteins, suggesting that SPA proteins are pseudokinases [27, 28]. Deletion of the large N-terminus of SPA1 including the kinase-like domain retained SPA1 activity in light-grown transgenic *spa1* mutant seedlings, suggesting that the N-terminus of SPA1 is dispensable for SPA1 function [29, 30]. However, functional redundancy with SPA2, SPA3 and SPA4 might mask a need for the N-terminal domain. Indeed, the N-terminus of SPA1 was necessary for full SPA1 activity in flowering time regulation [12]. The N-terminal domain of SPA1 also de-stabilizes SPA1 in light-grown seedlings [29, 30]. Protein-protein interaction studies have demonstrated that the N-terminal domain of SPA1 is essential for binding *cry2* in blue light and for binding phytochromes in red light [10, 12], though other domains of SPA1 might also be involved in binding phytochromes [11].

Analysis of *spa* single, double and triple mutants showed that the four SPA genes have overlapping but also distinct functions in plant development. In particular, SPA1 and SPA2 act redundantly to repress photomorphogenesis in seedlings in the dark, whereas in the light, only SPA1 serves as a repressor to prevent overstimulation by light. SPA2, in contrast, is extremely effectively inactivated by light of even very low fluence rates. As a consequence, SPA2 has little activity in light-grown plants when compared to SPA1, SPA3 and SPA4 [5, 23, 31–33]. The molecular basis of the difference between SPA1 and SPA2 activity in light-grown seedlings was studied previously. Chimeric SPA1/SPA2 promoter/cDNA swap experiments have shown that the distinct functions of SPA1 and SPA2 genes in light-grown plants are due to differences in the respective protein sequences and independent of the SPA promoter sequences. Moreover, we found that the SPA2 protein is more strongly degraded in the light than SPA1 which correlates with the much stronger light-induced repression of SPA2 activity when compared to the activity of SPA1 [15]. Here, we aimed to define the domain(s) in the SPA proteins which are responsible for the distinct

activities and stabilities of SPA1 and SPA2 in light-grown seedlings. To this end, we expressed chimeric SPA proteins containing domain swaps of SPA1 and SPA2 in transgenic *spa* mutant plants.

Results

Sequence divergence between the N-terminal domains of SPA1 and SPA2 is responsible and sufficient for the functional divergence between SPA1 and SPA2 in the light

We previously found that the diverged function of *SPA1* and *SPA2* genes in the light depends on differences in their protein-coding sequences [15]. In order to identify the domain(s) responsible for the SPA1- and SPA2-specific activities of these SPA proteins, we designed three domain swap constructs that encode chimeric SPA proteins with an N-terminal domain, a coiled-coil domain or a WD-repeat domain from SPA1 fused to the remaining domains of SPA2. The chimeric proteins thus contain one domain from SPA1 and two domains from SPA2 (Fig. 1a). They were expressed under the control of the *SPA2* promoter which is constitutively expressed and thus eliminates any transcriptional effects of light on protein function [15, 29]. The chimeric constructs were transformed into the *spa1 spa2 spa3* mutant which exhibits constitutive photomorphogenesis in darkness.

Dark-grown seedlings expressing any of the three chimeric SPA1/SPA2 proteins fully etiolated and thus exhibited full complementation of the *spa1 spa2 spa3* mutant phenotype (Fig. 1b). This was expected because SPA1 and SPA2 do not differ in their functions in darkness [15, 29]. Moreover, this shows that all three chimeric proteins are fully functional and thus not impaired by the domain swap. By contrast, transgenic light-grown seedlings revealed striking differences in phenotype. Expression of the DS_122-HA chimeric protein harboring the N-terminal domain from SPA1 complemented the phenotype of the *spa1 spa2 spa3* mutant to a similar extent as lines expressing SPA1-HA under low fluence rate of FRc, Rc and Bc (Fig. 1b). Hence, swapping the N-terminal domain was sufficient to confer SPA1-like activity to the SPA2 protein. Transgenic seedlings expressing DS_212-HA or DS_221-HA strongly deetiolated under weak FRc, Rc and Bc and thus behaved similarly as the SPA2-HA protein (Fig. 1b). These results indicate that the coiled-coil domain and the WD-repeat domain of SPA1 are not sufficient to confer a SPA1-like activity to an otherwise SPA2 protein. In summary, the distinct functions of SPA1 and SPA2 in light-grown seedlings can be mapped to the respective N-terminal domain, while the coiled-coil domains and the WD-repeat domains are functionally similar and interchangeable between SPA1 and SPA2.

Based on these findings, we generated homozygous transgenic lines expressing the chimeric DS_122-HA protein and further characterized their responses to Rc, FRc and Bc of different fluence rates. With increasing fluence rate of FRc, Rc and Bc, seedlings of *DS_122-HA*-expressing *spa1 spa2 spa3* lines exhibited only a moderate reduction in hypocotyls length and thus behaved similarly to SPA1-HA-expressing seedlings (Fig. 1c). In contrast, seedlings expressing SPA2-HA responded to light with an extreme shortening of their hypocotyls. These transgenic seedlings deetiolated to a similar extent as the *spa1 spa2 spa3* progenitor, indicating that there was no residual SPA2 activity present in the light, as was reported previously [15]. These results confirm that the SPA1 N-terminal domain is sufficient to maintain repressor activity in an otherwise SPA2 protein in light-grown seedlings.

The divergent region of the N-terminal domain is responsible and sufficient for the functional divergence between SPA1 and SPA2 in the light

We subsequently aimed to narrow down the region in the N-terminal domain of SPA1 and SPA2 that is responsible for the divergence of SPA1 and SPA2 function in the light. Based on sequence similarity, the N-termini of SPA1 and SPA2 were divided into two parts and swapped within the SPA2 protein (Fig. 2a). The N-terminal part is very divergent between SPA1 and SPA2, whereas the sequence of the C-terminal part is quite conserved among all four SPA proteins from *Arabidopsis* and also with SPAs from other species including *Physcomitrella*. The latter also shows moderate sequence similarity with Ser/Thr protein kinases [27, 28, 34].

Dark-grown *spa1 spa2 spa3* mutant seedlings expressing chimeric DS_NT12 or DS_NT21 proteins displayed a fully etiolated phenotypes, indicating that the chimeric proteins complemented the *spa1 spa2 spa3* mutant phenotype and thus were fully functional similar to the non-chimeric SPA1 and SPA2 proteins (Fig. 2b). In FRc, the transgenic seedlings displayed distinct phenotypes: the hypocotyl lengths of the *DS_NT12-HA* lines were similar to those of SPA1-HA expressing lines, while the hypocotyl lengths of *DS_NT21-HA* lines were similar to those of SPA2-HA expressing lines (Fig. 2b). Similar results were obtained in Rc and Bc (Fig. 2c). These findings indicate that swapping the more diverged part of the N-terminal domain was sufficient to confer a SPA1-like activity to the SPA2 protein.

The WD-repeat domain can also contribute to the functional divergence between SPA1 and SPA2 in the light

After examining the domain swaps containing one domain from SPA1 and two domains from SPA2, we conducted the reverse experiment by introducing one




Fig. 1 (See legend on next page.)

(See figure on previous page.)

Fig. 1 The N-terminal domain is involved in the functional divergence between SPA1 and SPA2. **a** Schematic representation of the chimeric SPA1/SPA2 domain-swap proteins DS_122-HA, DS_212-HA and DS_221-HA. "CC" represents the coiled-coil domain. All chimeric proteins were expressed in a *spa1 spa2 spa3* mutant under the control of the SPA2 promoter and fused with a C-terminal HA tag. Numbers above the domains indicate the percent identical amino acids between SPA1 and SPA2. NLS indicates the site of a predicted nuclear localization sequence. **b** Phenotype of 4-day-old *spa1 spa2 spa3* mutant seedlings carrying the indicated domain-swap constructs. Representative T2 seedlings are shown. Seedlings expressing SPA1-HA or SPA2-HA served as controls. All proteins were expressed under the control of the SPA2 promoter. Seedlings were grown in darkness, 0.05 $\mu\text{mol m}^{-2} \text{s}^{-1}$ FRc (FRc), 0.01 $\mu\text{mol m}^{-2} \text{s}^{-1}$ Rc (Rc) or 0.05 $\mu\text{mol m}^{-2} \text{s}^{-1}$ Bc (Bc) for 4 days. Numbers indicate independent transgenic lines. **c** Quantification of hypocotyl length of 4-day-old DS_122-HA expressing *spa1 spa2 spa3* T3 homozygous seedlings grown under various fluence rates of FRc, Rc or Bc. Error bars represent the SEM of at least 20 seedlings

domain from SPA2 into an otherwise SPA1 protein (Fig. 3a). These chimeric proteins were also expressed under the control of the SPA2 promoter in a *spa1 spa2 spa3* mutant. All three constructs - DS_112-HA, DS_121-HA and DS_211-HA - complemented the seedling phenotype of the *spa1 spa2 spa3* mutant in the dark, again indicating that the domains from SPA1 and SPA2 are compatible with each other and that the chimeric proteins are fully functional in darkness (Fig. 3b). In the light, the phenotypes of the transgenic seedlings differed. FRc-grown seedlings expressing DS_112-HA or DS_211-HA exhibited short hypocotyls and open cotyledons similar to the *spa1 spa2 spa3* progenitor and SPA2-HA-expressing *spa1 spa2 spa3* seedlings, indicating that these chimeric proteins, like SPA2, retain barely any activity in the light. Similar results were obtained in Rc and - to a lesser extent, in Bc (Fig. 3b). In contrast, seedlings expressing DS_121-HA showed strong etiolation and behaved similar to the SPA1-HA protein (Fig. 3b). Hence, swapping the coiled-coil domain from SPA2 into the SPA1 protein did not alter SPA1 activity. A similar observation was made previously when introducing the coiled-coil domain from SPA2 into SPA1 [30]. Hence, the coiled-coil domains from SPA1 and SPA2 appear equivalent.

Taken together, these results indicate that both the N-terminal domain and the WD-repeat domain of SPA2 can confer SPA2-like function to an otherwise SPA1 protein. Hence, both the N-terminal domain and the WD-repeat domain, but not the coiled-coil domain, can play a role in the distinct functions of SPA1 and SPA2.

The N-terminal domains of SPA1 and SPA2 differentially regulate SPA protein stability

Our previous studies showed that the SPA2 protein is more rapidly degraded in the light when compared to the SPA1 protein [15]. We therefore used domain swap lines to test which domains contribute to the differential protein stability of SPA1 and SPA2. DS_122-HA protein levels changed only moderately upon irradiation of dark-grown seedlings with FRc, Rc or Bc (Fig. 4). Similarly, the SPA1-HA protein was relatively stable after this short exposure to light, while the SPA2-HA protein was fully degraded upon light exposure, as reported previously [16]. Hence, the DS_122-HA protein behaved similarly to the SPA1 protein, indicating

that the N-terminal domain of SPA1 stabilizes the chimeric DS_122-HA protein in the light.

Levels of the DS_212-HA protein harboring the coiled-coil domain of SPA1 in a SPA2 background strongly decreased in response to light (Additional file 1: Figure S1A). This demonstrates that the coiled-coil domains of SPA1 and SPA2 are functionally equivalent with respect to protein stability. The levels of DS_221-HA harboring the WD-repeat domain of SPA1 in a SPA2 background strongly decreased in response to light, suggesting that the WD-repeats of SPA1 and SPA2 also do not severely differ in their effects on protein stability in this configuration (Additional file 1: Figure S1B). In summary, only a swap-in of the SPA1 N-terminal domain can confer higher stability to a SPA2 protein in the light.

When conducting the reverse experiment, i.e. swapping a SPA2 domain into a SPA1 protein, the N-terminal domain of SPA2 de-stabilized the SPA1 protein in light-grown DS_211-HA lines when compared to SPA1-HA lines (Additional file 2: Figure S2A). Swapping the coiled-coil domain did not dramatically alter protein stability in two out of three lines (Additional file 2: Figure S2B), while swapping the WD-repeat domain tended to destabilize the protein (Additional file 2: Figure S2C). Hence, in an otherwise SPA1 protein, introducing the N-terminal or the WD-repeat domain of SPA2 strongly or moderately destabilized the chimeric protein, respectively.

SPA1 and SPA2 have similar affinities for cry1 in blue light

Since SPA2 is much more strongly inactivated by blue light than SPA1 [15, 29] we tested whether SPA2 might interact with cry1 more strongly than SPA1. Figure 5a shows that SPA1 and SPA2 have similar affinities for cry1 in vivo. This suggests that the higher responsiveness of *spa1 spa3 spa4* mutants to blue light when compared to *spa2 spa3 spa4* mutants [29] is not due to a differential cry1 binding strength. This finding is consistent with our previous observation that inactivation of SPA2 is primarily mediated by phyA [16].

The N-termini of SPA1 and SPA2 differentially interact with cry2

We previously showed that SPA2 does not bind cry2 in vivo, while SPA1 does interact with cry2 in vivo,




Fig. 2 The diverged part of the N-terminal domain of SPA contributes to the divergence of SPA1/SPA2 function in the light. **a** The top row shows a schematic representation of amino acid identity between SPA1 and SPA2 domains. Below indicates the chimeric SPA1/SPA2 domain swap proteins DS_NT12-HA and DS_NT21-HA carrying swaps in the N-terminal domains of SPA1/SPA2. All chimeric constructs were expressed in a *spa1 spa2 spa3* mutant background under the control of the *SPA2* promoter and fused with a C-terminal HA tag. NLS indicates the site of a predicted nuclear localization sequence. **b, c** Phenotype of 4-day-old *spa1 spa2 spa3* mutant seedlings expressing *DS_NT12-HA* or *DS_NT21-HA* swap constructs. Representative T2 seedlings are shown. Seedlings were grown in darkness or in 0.05 $\mu\text{mol m}^{-2} \text{s}^{-1}$ FRC (**b**) and in 0.01 $\mu\text{mol m}^{-2} \text{s}^{-1}$ Rc or 0.05 $\mu\text{mol m}^{-2} \text{s}^{-1}$ Bc (**c**) for 4 days. Numbers indicate independent transgenic lines

as was shown previously [12, 16]. We therefore asked whether swapping of the SPA1 N-terminal domain into SPA2 blocked the interaction between the chimeric SPA protein and cry2. Figure 5b shows that the DS_122-HA chimeric protein was able to associate with cry2 in B, while SPA2-HA was not. Hence, the differential affinity of SPA1 and SPA2 for cry2 is due to the distinct sequences in their N-terminal domains.

SPA1 and SPA2 similarly interact with COP1 in light-grown seedlings

We subsequently asked whether light is more effective in disrupting a COP1/SPA2 complex when compared to a COP1/SPA1 complex. SPA1 and SPA2 co-immunoprecipitated the same amount of COP1 in dark- and light-grown seedlings (Fig. 6). Hence, no differential effect of light on COP1/SPA1 and COP1/SPA2 complex abundance was observed.





Fig. 3 WD-repeat domain is involved in the functional divergence of SPA1 and SPA2. **a** Schematic representation of chimeric SPA1/SPA2 domain swap *DS_112-HA*, *DS_121-HA* and *DS_211-HA* constructs. All chimeric constructs were expressed in a *spa1 spa2 spa3* mutant background under the control of the *SPA2* promoter and fused with a C-terminal HA tag. NLS indicates the location of a predicted nuclear localization sequence. **b** Phenotype of 4-day-old *spa1 spa2 spa3* seedlings expressing *DS_112-HA*, *DS_121-HA* and *DS_211-HA* domain swap constructs. Representative T2 seedlings are shown. Seedlings were grown in darkness or in 0.05 $\mu\text{mol m}^{-2} \text{s}^{-1}$ FRc, 0.01 $\mu\text{mol m}^{-2} \text{s}^{-1}$ Rc or 0.05 $\mu\text{mol m}^{-2} \text{s}^{-1}$ Bc for 4 days. Numbers indicate independent transgenic lines

Discussion

The COP1/SPA protein complex is an important negative regulator of photomorphogenesis in dark-grown Arabidopsis plants. Our previous analysis of *SPA* genes revealed that the four *SPAs* have at least partially distinct functions during plant growth and development. In particular, SPA1 and SPA2 proteins strongly differ in their responsiveness to light, while they have indistinguishable activities in darkness. In light-grown seedlings, SPA1 retains considerable repressor activity to prevent overstimulation by light, while SPA2 is almost fully inactivated by even very low fluences of light [5, 15, 29]. Here, we have shown that the stronger light-responsiveness of SPA2 when compared to SPA1 maps primarily to the N-terminal domain which includes the kinase-like domain. This conclusion is based on our finding that swapping the N-terminal domain of SPA1 into a SPA2 protein confers SPA1-like responsiveness to R, FR and B in the chimeric protein. The same result was found when conducting the reverse experiment, i.e. when introducing the N-terminus of SPA2 into an otherwise SPA1 protein. This finding is consistent with the

observation that sequence divergence between SPA1 and SPA2 is highest in the N-terminal domain (73 %) when compared to the WD-repeat domain (33 %) and the coiled-coil domain (71 %). When we further divided the N-terminal domain into a smaller, highly conserved part and a larger, not conserved part, we found that the non-conserved part of SPA1 and SPA2 conferred the SPA1- or SPA2-specific light-responsiveness, respectively. Hence, these specific SPA activities that map to the N-terminal domain evolved through major sequence alterations and not through smaller sequence changes in a highly conserved domain. It remains to be established which of the two *SPA* genes, *SPA1* or *SPA2*, evolved a new activity since the duplication event leading to *SPA1* and *SPA2* [35]. Since *SPA3* and *SPA4* respond to light similarly to *SPA1*, it is likely that the extreme light-induced inhibition of SPA2 is the recent evolutionary innovation.

Though the N-terminal domain is the primary domain responsible for the distinct activities of SPA1 and SPA2 in light-grown seedlings, the WD-repeat domain can also contribute at least to some extent to the functional divergence between SPA1 and SPA2. We found that



swapping the WD-repeat domain from SPA2 into a SPA1 protein also strongly enhanced light-responsiveness of the chimeric protein. However, the reverse experiment, i.e. introducing the WD-repeat from SPA1 into a SPA2 protein did not reduce light-responsiveness. Hence, the SPA-specific effect of the WD-repeat might be context-dependent. However, it is also possible that the enhanced light-induced inactivation of this chimeric DS₁₁₂ protein is due to a partial loss-of-function caused by the domain swap per se. This partial loss-of function might only be detectable in light-grown seedlings, while the activity of the chimeric protein might be sufficient for full suppression of photomorphogenesis in darkness. Indeed, such a phenotype was observed in the non-constitutively photomorphogenic *cop1^{eid6}* mutant [36]. Moreover, it is known that SPA activities are more limiting in light-grown seedlings than in dark-grown seedlings since *spa* single mutants exhibit a mutant phenotype only in the light [34, 37]. In total, we therefore conclude that the functional divergence between SPA1 and SPA2 primarily maps to the divergent part of the N-terminal domain.

We considered at least two, not mutually exclusive mechanisms that might contribute to the distinct light-

responsiveness of SPA1 and SPA2 proteins. First, the N-termini of SPA1 and SPA2 might have distinct affinities for photoreceptors. There is no evidence that differences in the interactions with cryptochromes might be causal for the higher light-responsiveness of SPA2. Our results show that SPA1 and SPA2 interacted similarly with cry1 in B. With respect to cry2, SPA2 has a severely reduced - rather than increased - affinity for cry2 when compared to SPA1 [16]. Similarly, we have shown here that the N-terminal domain swap protein DS₁₂₂ conferring a lower responsiveness to light exhibits a restored interaction with cry2 when compared to SPA2. The relative in vivo affinities of SPA1 and SPA2 for phytochromes are thus far unknown. It is known, however, that the N-terminal domain of SPA1 which is responsible for the distinct functions of SPA1 and SPA2 interacts with phyA in the yeast-two hybrid system [10]. Also, phytochromes are the primary photoreceptors inactivating SPA2 in R, FR as well as B [16]. On the other hand, SPA2 interacted less with phyA in R-treated yeast cells when compared with SPA1 [10]. Hence, further experiments are necessary to resolve whether differential interactions of SPA1 and SPA2 with phytochromes contribute to the distinct light-responsiveness of SPA1 and SPA2.

Even if affinities for the photoreceptors themselves do not account for the functional divergence of SPA1 and SPA2, downstream effects in repressing COP1/SPA1 and COP1/SPA2 function by photoreceptors may differ and be causative for the distinct light-responsiveness of SPA1 and SPA2. One mechanism by which light inactivates COP1/SPA1 and COP1/SPA2 function is the phytochrome-induced degradation of SPA1 and SPA2. Indeed, SPA2 is much more effectively degraded in the light than SPA1 [15, 16]. Moreover, the degree of photomorphogenesis in transgenic lines expressing chimeric SPA1/SPA2 proteins was negatively correlated with the stability of the chimeric proteins in the light, i.e. highly responsive lines expressed a chimeric protein that is unstable in the light, and vice versa. This correlation supports the notion that light-induced SPA protein degradation is intrinsically related to the light-induced inhibition of COP1/SPA activity. Our finding that a swap of the SPA1/SPA2 N-terminal domains strongly alters the stability of the chimeric SPA proteins also agrees with previous observations showing that the N-terminal domain of SPA1 de-stabilizes the SPA1 protein [29, 30]. Light inactivates COP1/SPA function also by dissociation of the COP1/SPA interaction and by nuclear exclusion of COP1 [22]. Hence, one can hypothesize that the light-induced dissociation of a COP1/SPA2 complex might be more sensitive to light than the dissociation of a COP1/SPA1 complex, and this difference might depend on the N-terminal sequence in SPA. Our results do not provide evidence for this idea because both SPA1 and SPA2 were in a complex with COP1 also in light-treated




Fig. 5 DS_122-HA associates with cry1 and cry2 in blue light. **a, b** Co-immunoprecipitation of cry1 (**a**) and cry2 (**b**) by DS_122-HA. 4-day-old dark-grown seedlings (D) were transferred to $50 \mu\text{mol m}^{-2} \text{s}^{-1}$ B for 1 h (B). DS_122-HA was expressed in *spa1 spa2 spa3* and under the control of the SPA2 promoter. Col-0, SPA2::SPA1-HA and SPA1::SPA2-HA lines were used as controls. SPA-HA proteins were immunoprecipitated using α -HA beads. α -HA antibody was used to detect SPA-HA proteins. α -cry1 and α -cry2 antibodies were used to detect cry1 and cry2, respectively. Asterisks likely indicate phosphorylated cry1 and cry2. To obtain similar SPA-HA protein levels from different transgenic lines in B, all seedlings were treated with proteasome inhibitor to reduce protein degradation in B. Also, SPA2-HA was expressed under the control of the stronger SPA1 promoter to counteract the strong B-induced degradation of SPA2. Five times more protein extract was used for the SPA1::SPA2-HA 64 and DS_122-HA 27 co-immunoprecipitations than for the SPA2::SPA1-HA 28 and DS_122-HA 37 co-immunoprecipitations

seedlings. However, since the light-induced disruption of the COP1/SPA complex is much less detectable in co-immunoprecipitation assays than in yeast three-hybrid or FRET-FLIM studies [8–11], other assays may be necessary to compare the light-induced rearranging in the respective COP1/SPA complexes.

Conclusions

In summary, our results have defined the N-terminal domains of SPA1 and SPA2 as a "light-sensitivity" domain that confers the distinct light-responsiveness to the respective SPA protein regarding light-induced photomorphogenesis and light-induced SPA degradation. The exact mechanism in which the N-terminal domain is involved and which role photoreceptors play remain to be resolved.

Methods


Plant material, light sources and growth conditions

Wild-type *Arabidopsis thaliana* Col-0 and the *spa* triple mutants *spa1-7 spa2-1 spa3-1*, *spa2-1 spa3-1 spa4-1* and *spa1-7 spa3-1 spa4-1* were used [29]. Transgenic lines expressing full-length SPA1 or SPA2 proteins under the control of the SPA2 promoter were SPA2::SPA1-HA 28 and SPA2::SPA1-HA 70 or SPA2::SPA2-HA 32 and

SPA2::SPA2-HA 60, respectively, as described previously [15]. LED light sources and growth conditions were as described previously [5].

Generation of transgenic plants expressing domain swap constructs

The DS_122-HA lines express a chimeric protein with amino acids 1–552 from SPA1 and 580–1036 from SPA2. The DS_212-HA lines express a chimeric protein with amino acids 1–571 from SPA2, 545–703 from SPA1 and 711–1036 from SPA2. The DS_221-HA lines express a chimeric protein with amino acids 1–702 from SPA2 and 696–1029 from SPA1. The DS_112-HA lines express a chimeric protein with amino acids 1–703 from SPA1 and 711–1036 from SPA2. The DS_121-HA lines express a chimeric protein with amino acids 1–552 from SPA1, 580–702 from SPA2 and 696–1029 from SPA1. The DS_211-HA lines express a chimeric protein with amino acids 1–571 from SPA2 and 545–1029 from SPA1. Two highly conserved regions were selected for domain swapping. The first motif lies between N-terminal domain and the coiled-coil domain and reads SELLLHFL for SPA1 and SELLQHFL for SPA2. The second motif lies between the coiled-coil domain and the WD-repeats and reads ARYSKFET which is identical in both SPA1 and SPA2 proteins.



To generate these domain-swap constructs, first of all, a 4 kb *Apal Sall* fragment comprising the full-length SPA2 ORF, the C-terminal triple HA tag and the SPA2 3' UTR was sub-cloned into *pBS SK+* resulting the *pBS SK+ cSPA2* construct. Next, in order to generate domain swap constructs expressing one SPA1 domain and two SPA2 domains (DS_122, DS_212, DS_221) sequences encoding N-terminus, coiled-coil or WD-repeats from SPA1 were amplified from *pJHA212-hpt SPA2::SPA1-HA* [15] by PCR using the primer pairs LW4 and LW5, LW6 and LW7 and LW8 and LW9, respectively, which are compatible with the conserved regions used for swapping. The resulting PCR fragments were gel-purified. The mutagenesis PCRs which create the respective domain swap constructs were performed according to the 'Quik Change Mutagenesis Kit' (Agilent Technologies)

using combinations of the template plasmid *pBS SK+ cSPA2* and the gel-purified PCR products serving as primers. In order to generate swapping constructs expressing two SPA1 domains and one SPA2 domain, sequences encoding N-terminus, coiled-coil or WD-repeats from SPA1 were PCR-amplified from the plasmid *pJHA212-hpt SPA2::SPA1-HA* [15] using the primer pairs LW4 and SC_ds_5 primers, LW6 and SC_ds_7 and LW8 and LW9, respectively, which are compatible with the conserved regions used for swapping. The resulting PCR fragments were gel-purified. The mutagenesis PCRs were performed using a combination of the template plasmid which already expresses the chimeric proteins containing one SPA1 domain and two SPA2 domains generated previously and the newly gel-purified PCR fragments serving as primers. For generating the construct expressing the N-terminus and coiled-coil from SPA1 and the WD-repeats from SPA2 (DS_112), the purified SPA1 N-terminus PCR product was combined with the domain swap construct generated previously expressing the coiled-coil domain from SPA1 and N-terminal domain and WD-repeats from SPA2. For generating the construct expressing both coiled-coil and WD-repeats from SPA1 and N-terminal domain from SPA2 (DS_211), the purified SPA1 coiled-coil PCR product was combined with the domain swap construct generated previously expressing the WD-repeats of SPA1 and N-terminal domain and coiled-coil of SPA2. For generating the construct expressing both N-terminal domain and WD-repeats of SPA1 and coiled-coil of SPA2 (DS_121), the purified SPA1 WD-repeats PCR product was combined with the domain swap construct generated previously expressing the N-terminal domain of SPA1, coiled-coil and WD-Repeats of SPA2.

To swap the SPA1 amino acids 1–425 with SPA2 amino acids 1–450 or SPA1 amino acids 426–552 with SPA2 amino acids 451–579 in a SPA2 background, generating the *DS_NT12-HA* and *DS_NT21-HA* lines respectively, motifs between less conserved N-terminal regions and more conserved C-terminal regions in the N-termini of SPA1 and SPA2 were chosen. In SPA1, the selected motif was LSVSSVSRKQSM while in SPA2, the motif was HCSTVACPFTSV. Sequences encoding amino acids 1–425 and 426–552 of the SPA1 N-terminus were amplified from *pJHA212-hpt SPA2::SPA1-HA* [15] by PCR using the primer pairs LW4 and SC-P2 and SC-P3 and LW5, respectively, which are compatible with the motifs selected for swapping. The resulting PCR fragments were gel-purified. The mutagenesis PCRs which generate the swapping constructs were performed using the combination of the template plasmid *pBS SK+ cSPA2* and the gel-purified PCR products serving as primers.

All mutagenesis PCRs were performed using Pfu polymerase to extend and incorporate the mutagenic primers

resulting in nicked circular strands. The products were digested with DpnI to remove methylated, non-mutated parental templates. After digestion, the products were transformed into *DH5 α* and plated on LB agar plates containing 100 $\mu\text{g/ml}$ ampicillin. Plasmid DNAs were purified from selected colonies and sequenced. Confirmed swapping constructs were cloned back into *pJHA212-hpt SPA2::SPA2-HA* [15] via *ApaI* and *Sall* sites to replace the full length *SPA2*. The final constructs were transformed into *spa1-7 spa2-1 spa3-1* mutant plants by floral dipping. T2 or T3 plants were used for analysis. All primers used for cloning are listed in the Additional file 3: Table S1.

Isolation of total proteins and immunoblot analysis

Total proteins from dark-grown and light-treated transgenic seedlings were isolated and used for immunoblot analysis as described previously [16].

Co-immunoprecipitations

Co-immunoprecipitations for examining SPA-cry and SPA-COP1 associations in B were performed using μMACS Anti-HA Starting Kits (Miltenyi Biotec) as described previously and according to the manufacturer's protocol with minor modification [16]. Prior to the light treatment, seedlings were pre-incubated in 100 μM MG132 and 10 μM clasto-lactacystin β -lactone twice, 15 min each, to prevent degradation of SPA proteins. Five times more protein extract was used for the *SPA1::SPA2-HA 64* and *DS_122-HA 27* line than for the *SPA2::SPA1-HA 28* and *DS_122-HA 37* line. Co-immunoprecipitations for examining SPA-COP1 association in FR and R were performed using Anti-HA Affinity Matrix (Roche) according to the manufacturer's protocol. Equal amount of total protein was used as inputs.

Hypocotyl length measurement

Flattened seedlings were imaged by a Nikon D5000 digital camera on MS plates. Measurements were conducted using ImageJ 1.43u (Wayne Rasband, National Institutes of Health).

Additional files

Additional file 1: Figure S1. The coiled-coil and WD-repeat domains of SPA1 do not provide higher stability to the chimeric SPA2 protein in light-grown seedlings. **A, B.** SPA-HA protein levels in 4-day-old T2 *DS_212-HA* (**A**) or *DS_221-HA* (**B**) transgenic *spa1 spa2 spa3* mutant seedlings. Seedlings were grown in darkness (D) for 4 days and subsequently transferred to 0.35 $\mu\text{mol m}^{-2} \text{s}^{-1}$ FR for 30 min. All transgenes were expressed under the control of the *SPA2* promoter. SPA-HA was detected using an α -HA antibody. HSC70 levels served as a loading control. (PDF 245 kb)

Additional file 2: Figure S2. The N-terminal domain and the WD-repeat domain of SPA2 render the chimeric SPA1 proteins more unstable in light-grown seedlings. **A-C.** SPA-HA protein levels in 4-day-old T2 *DS_211-HA* (**A**), *DS_121-HA* (**B**) and *DS_112-HA* (**C**) transgenic *spa1 spa2*

spa3 seedlings. Seedlings were grown in darkness (D) for 4 days and subsequently transferred to 0.35 $\mu\text{mol m}^{-2} \text{s}^{-1}$ FR for 30 min. All transgenes were expressed under the control of the *SPA2* promoter. SPA-HA was detected using an α -HA antibody. HSC70 levels served as a loading control. (PDF 349 kb)

Additional file 3: Table S1. Primer sequences. (PDF 32 kb)

Abbreviations

B, blue; COP1, CONSTITUTIVELY PHOTOMORPHOGENIC1; cry1, cryptochrome 1; cry2, cryptochrome 2; CUL4, cullin 4; DDB1, damaged DNA binding protein 1; eid6, empfindlicher im dunkelroten Licht 6; FLIM, fluorescence lifetime imaging; FR, far-red; FRET, fluorescence resonance energy transfer; HA, human influenza hemagglutinin; HFR1, long hypocotyl in far-red 1; HY5, long hypocotyl 5; ORF, open reading frame.; PAP, production of anthocyanin pigment; phyA, Phytochrome A; R, red; RING, really interesting new gene; SPA, SUPPRESSOR OF PHYA-105; UV-B, ultraviolet B; UVR8, UV resistance locus8

Acknowledgements

We thank Klaus Menrath, his greenhouse staff and many undergraduate students for their expert care of our plants and important contributions.

Funding

This work was funded by the Deutsche Forschungsgemeinschaft (SFB635 and HO2793/3-1) to U.H. and a grant from AFOSR (FA9550-14-0-0409) to M.A. S.C. was a recipient of a Ph.D. fellowship from the International Graduate School of Genetics and Functional Genomics from the state Northrhine-Westphalia, Germany. X.H. received a Ph.D. fellowship from the International Max Planck Research School of Molecular Plant Development which was co-funded by The Max Planck Institute of Plant Breeding Research and the University of Cologne.

Availability of data and materials

All relevant supporting data can be found within the article and its additional files.

Authors' contributions

SC, LW and UH designed this study, SC, LW, JS, NL, XH, LL and CS performed experiments, MA provided important materials, SC and UH wrote the manuscript, all authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable.

Ethics approval and consent to participate

Not applicable.

Author details

¹Botanical Institute and Cluster of Excellence on Plant Sciences (CEPLAS), Biocenter, University of Cologne, Zùlpicher Str. 47b, 50674 Cologne, Germany. ²UMR 8256 (B2A) CNRS - UPMC, IBPS, Université Pierre et Marie Curie, Bat C, 9 quai Saint-Bernard, 75252 Paris Cedex 05, France. ³Present Address: Department of Botany and Plant Biology, University of Geneva, Sciences III, 30 Quai E. Ansermet, 1211 Geneva 4, Switzerland. ⁴Present Address: Department of Plant Biochemistry, Dahlem Center of Plant Sciences, Freie Universität Berlin, Königin-Luise-Str. 12-16, Berlin, Germany.

Received: 18 May 2016 Accepted: 14 July 2016

Published online: 22 July 2016

References

- Kami C, Lorrain S, Hornitschek P, Fankhauser C. Light-regulated plant growth and development. *Curr Top Dev Biol.* 2010;91:29–66.
- Losi A, Gartner W. The evolution of flavin-binding photoreceptors: an ancient chromophore serving trendy blue-light sensors. *Annu Rev Plant Biol.* 2012;63:49–72.
- Casal JJ. Photoreceptor signaling networks in plant responses to shade. *Annu Rev Plant Biol.* 2013;64:403–27.

4. Tilbrook K, Arongaus AB, Binkert M, Heijde M, Yin R, Ulm R. The UVR8 UV-B photoreceptor: perception, signaling and response. *Arabidopsis Book*. 2013;11, e0164.
5. Laubinger S, Fittinghoff K, Hoecker U. The SPA quartet: a family of WD-repeat proteins with a central role in suppression of photomorphogenesis in *Arabidopsis*. *Plant Cell*. 2004;16(9):2293–306.
6. Deng X-W, Caspar T, Quail PH. *cop1*: A regulatory locus involved in light-controlled development and gene expression in *Arabidopsis*. *Genes Dev*. 1991;5:1172–82.
7. Ordóñez-Herrera N, Fackendahl P, Yu X, Schaefer S, Koncz C, Hoecker U. A *cop1 spa* mutant deficient in COP1 and SPA proteins reveals partial co-action of COP1 and SPA during *Arabidopsis* post-embryonic development and photomorphogenesis. *Mol Plant*. 2015;8:479–81.
8. Liu B, Zuo Z, Liu H, Liu X, Lin C. *Arabidopsis* cryptochrome 1 interacts with SPA1 to suppress COP1 activity in response to blue light. *Genes Dev*. 2011;25(10):1029–34.
9. Lian HL, He SB, Zhang YC, Zhu DM, Zhang JY, Jia KP, Sun SX, Li L, Yang HQ. Blue-light-dependent interaction of cryptochrome 1 with SPA1 defines a dynamic signaling mechanism. *Genes Dev*. 2011;25(10):1023–8.
10. Sheerin DJ, Menon C, zur Oven-Krockhaus S, Enderle B, Zhu L, Johnen P, Schleifenbaum F, Stierhof YD, Huq E, Hiltbrunner A. Light-activated phytochrome A and B interact with members of the SPA family to promote photomorphogenesis in *Arabidopsis* by reorganizing the COP1/SPA complex. *Plant Cell*. 2015;27(1):189–201.
11. Lu XD, Zhou CM, Xu PB, Luo Q, Lian HL, Yang HQ. Red light-dependent interaction of phyB with SPA1 promotes COP1–SPA1 dissociation and photomorphogenic development in *Arabidopsis*. *Mol Plant*. 2015;8:467–78.
12. Zuo Z, Liu H, Liu B, Liu X, Lin C. Blue light-dependent interaction of CRY2 with SPA1 regulates COP1 activity and floral initiation in *Arabidopsis*. *Curr Biol*. 2011;21(10):841–7.
13. Von Arnim AG, Deng X-W. Light inactivation of *Arabidopsis* photomorphogenic repressor COP1 involves a cell-specific regulation of its nucleocytoplasmic partitioning. *Cell*. 1994;79:1035–45.
14. Pacin M, Legris M, Casal JJ. Rapid decline in nuclear constitutive photomorphogenesis1 abundance anticipates the stabilization of its target elongated hypocotyl5 in the light. *Plant Physiol*. 2014;164(3):1134–8.
15. Balcerowicz M, Fittinghoff K, Wirthmueller L, Maier A, Fackendahl P, Fiene G, Koncz C, Hoecker U. Light exposure of *Arabidopsis* seedlings causes rapid de-stabilization as well as selective post-translational inactivation of the repressor of photomorphogenesis SPA2. *Plant J*. 2011;65(5):712–23.
16. Chen S, Lory N, Stauber J, Hoecker U. Photoreceptor specificity in the light-induced and COP1-mediated rapid degradation of the repressor of photomorphogenesis SPA2 in *Arabidopsis*. *PLOS Genetics*. 2015;11, e1005516.
17. Chen H, Huang X, Gusmaroli G, Terzaghi W, Lau OS, Yanagawa Y, Zhang Y, Li J, Lee JH, Zhu D, et al. *Arabidopsis* CULLIN4-damaged DNA binding protein 1 interacts with CONSTITUTIVELY PHOTOMORPHOGENIC1-SUPPRESSOR OF PHYA complexes to regulate photomorphogenesis and flowering time. *Plant Cell*. 2010;22(1):108–23.
18. Osterlund MT, Hardtke CS, Wei N, Deng XW. Targeted destabilization of HY5 during light-regulated development of *Arabidopsis*. *Nature*. 2000;405:462–6.
19. Jang IC, Yang JY, Seo HS, Chua NH. HFR1 is targeted by COP1 E3 ligase for post-translational proteolysis during phytochrome A signaling. *Genes Dev*. 2005;19(5):593–602.
20. Yang J, Lin R, Sullivan J, Hoecker U, Liu B, Xu L, Deng XW, Wang H. Light Regulates COP1-Mediated Degradation of HFR1, a Transcription Factor Essential for Light Signaling in *Arabidopsis*. *Plant Cell*. 2005;17(3):804–21.
21. Maier A, Schrader A, Kokkelink L, Falke C, Welter B, Iniesto E, Rubio V, Uhrig JF, Hulskamp M, Hoecker U. Light and the E3 ubiquitin ligase COP1/SPA control the protein stability of the MYB transcription factors PAP1 and PAP2 involved in anthocyanin accumulation in *Arabidopsis*. *Plant J*. 2013;74(4):638–51.
22. Huang X, Ouyang X, Deng XW. Beyond repression of photomorphogenesis: role switching of COP/DET/FUS in light signaling. *Curr Opin Plant Biol*. 2014;21C:96–103.
23. Menon C, Sheerin DJ, Hiltbrunner A. SPA proteins: SPAnning the gap between visible light and gene expression. *Planta* 2016;244:297–312.
24. Zhu D, Maier A, Lee JH, Laubinger S, Saijo Y, Wang H, Qu LJ, Hoecker U, Deng XW. Biochemical characterization of *Arabidopsis* complexes containing CONSTITUTIVELY PHOTOMORPHOGENIC1 and SUPPRESSOR OF PHYA proteins in light control of plant development. *Plant Cell*. 2008;20(9):2307–23.
25. Hoecker U, Quail PH. The phytochrome A-specific signaling intermediate SPA1 interacts directly with COP1, a constitutive repressor of light signaling in *Arabidopsis*. *J Biol Chem*. 2001;276:38173–8.
26. Saijo Y, Sullivan JA, Wang H, Yang J, Shen Y, Rubio V, Ma L, Hoecker U, Deng XW. The COP1-SPA1 interaction defines a critical step in phytochrome A-mediated regulation of HY5 activity. *Genes Dev*. 2003;17(21):2642–7.
27. Hoecker U, Tepperman JM, Quail PH. SPA1, a WD-repeat protein specific to phytochrome A signal transduction. *Science*. 1999;284:496–9.
28. Ranjan A, Dickopf S, Ullrich KK, Rensing SA, Hoecker U. Functional analysis of COP1 and SPA orthologs from *Physcomitrella* and rice during photomorphogenesis of transgenic *Arabidopsis* reveals distinct evolutionary conservation. *BMC Plant Biol*. 2014;14:178.
29. Fittinghoff K, Laubinger S, Nixdorf M, Fackendahl P, Baumgardt RL, Batschauer A, Hoecker U. Functional and expression analysis of *Arabidopsis* SPA genes during seedling photomorphogenesis and adult growth. *Plant J*. 2006;47(4):577–90.
30. Yang J, Wang H. The central coiled-coil domain and carboxyl-terminal WD-repeat domain of *Arabidopsis* SPA1 are responsible for mediating repression of light signaling. *Plant J*. 2006;47(4):564–76.
31. Laubinger S, Marchal V, Gentilhomme J, Wenkel S, Adrian J, Jang S, Kulajta C, Braun H, Coupland G, Hoecker U. *Arabidopsis* SPA proteins regulate photoperiodic flowering and interact with the floral inducer CONSTANS to regulate its stability. *Development*. 2006;133(16):3213–22.
32. Baumgardt RL, Oliverio KA, Casal JJ, Hoecker U. SPA1, a component of phytochrome A signal transduction, regulates the light signaling current. *Planta*. 2002;215(5):745–53.
33. Rolauffs S, Fackendahl P, Sahm J, Fiene G, Hoecker U. *Arabidopsis* COP1 and SPA genes are essential for plant elongation but not for acceleration of flowering time in response to a low red light to far-red light ratio. *Plant Physiol*. 2012;160(4):2015–27.
34. Laubinger S, Hoecker U. The SPA1-like proteins SPA3 and SPA4 repress photomorphogenesis in the light. *Plant J*. 2003;35(3):373–85.
35. Simillion C, Vandepoele K, Van Montagu MC, Zabeau M, Van de Peer Y. The hidden duplication past of *Arabidopsis thaliana*. *Proc Natl Acad Sci U S A*. 2002;99(21):13627–32.
36. Dieterle M, Buche C, Schäfer E, Kretsch T. Characterization of a novel non-constitutive photomorphogenic *cop1* allele. *Plant Physiol*. 2003;133(4):1557–64.
37. Hoecker U, Xu Y, Quail PH. SPA1: A new genetic locus involved in phytochrome A-specific signal transduction. *Plant Cell*. 1998;10:19–33.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

