



HAL
open science

Intravenous immunoglobulin-induced IL-33 is insufficient to mediate basophil expansion in autoimmune patients

Meenu Sharma, Yoland Schoindre, Pushpa Hegde, Chaitrali Saha, Mohan S Maddur, Emmanuel Stephen-Victor, Laurent Gilardin, Maxime Lecerf, Patrick Bruneval, Luc Mouthon, et al.

► To cite this version:

Meenu Sharma, Yoland Schoindre, Pushpa Hegde, Chaitrali Saha, Mohan S Maddur, et al.. Intravenous immunoglobulin-induced IL-33 is insufficient to mediate basophil expansion in autoimmune patients. *Scientific Reports*, 2014, 4 (1), pp.5672. 10.1038/srep05672 . hal-01358914

HAL Id: hal-01358914

<https://hal.sorbonne-universite.fr/hal-01358914v1>

Submitted on 1 Sep 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



OPEN

SUBJECT AREAS:

AUTOIMMUNITY

DENDRITIC CELLS

IMMUNOTHERAPY

Received
25 April 2014Accepted
26 June 2014Published
11 July 2014

Correspondence and requests for materials should be addressed to S.V.K. (srini.kaveri@crc.jussieu.fr) or J.B. (jagadeesh.bayry@crc.jussieu.fr)

Intravenous immunoglobulin-induced IL-33 is insufficient to mediate basophil expansion in autoimmune patients

Meenu Sharma^{1,2}, Yoland Schoindre^{3,4}, Pushpa Hegde¹, Chaitrali Saha^{1,2}, Mohan S. Maddur^{1,5}, Emmanuel Stephen-Victor^{1,5}, Laurent Gilardin^{1,5}, Maxime Lecerf^{1,5,6}, Patrick Bruneval⁷, Luc Mouthon^{8,9}, Olivier Benveniste^{3,4,10}, Srinivasa V. Kaveri^{1,5,6,11} & Jagadeesh Bayry^{1,5,6,11}

¹Institut National de la Santé et de la Recherche Médicale Unité 1138, Paris, F-75006, France, ²Université de Technologie de Compiègne, Compiègne, F-60205, France, ³Département de Médecine Interne et Immunologie Clinique, Centre de référence maladies neuro-musculaires, Hôpital Pitié-Salpêtrière, AP-HP, Paris F-75013, France, ⁴Université Pierre et Marie Curie - Paris 6, F-75013, France, ⁵Centre de Recherche des Cordeliers, Equipe - Immunopathology and therapeutic immunointervention, Université Pierre et Marie Curie - Paris 6, UMR S 1138, 15 rue de l'École de Médecine, Paris, F-75006, France, ⁶Université Paris Descartes, UMR S 1138 Paris, F-75006, France, ⁷Service d'anatomie pathologique, Hôpital Européen Georges Pompidou, Paris, F-75015, France, ⁸Institut Cochin, Institut National de la Santé et de la Recherche Médicale Unité 1016, CNRS UMR 8104, Université Paris Descartes, F-75014, France, ⁹Pôle de Médecine Interne, Centre de Référence pour les maladies systémiques autoimmunes rares; Assistance Publique-Hôpitaux de Paris (AP-HP); Paris, France, ¹⁰Institut National de la Santé et de la Recherche Médicale Unité 974, Paris, F-75013, France, ¹¹International Associated Laboratory IMPACT (Institut National de la Santé et de la Recherche Médicale, France - Indian council of Medical Research, India), National Institute of Immunohaematology, Mumbai, 400012, India.

Intravenous immunoglobulin (IVIg) is used in the therapy of various autoimmune and inflammatory diseases. Recent studies in experimental models propose that anti-inflammatory effects of IVIg are mainly mediated by $\alpha 2,6$ -sialylated Fc fragments. These reports further suggest that $\alpha 2,6$ -sialylated Fc fragments interact with DC-SIGN⁺ cells to release IL-33 that subsequently expands IL-4-producing basophils. However, translational insights on these observations are lacking. Here we show that IVIg therapy in rheumatic patients leads to significant raise in plasma IL-33. However, IL-33 was not contributed by human DC-SIGN⁺ dendritic cells and splenocytes. As IL-33 has been shown to expand basophils, we analyzed the proportion of circulating basophils in these patients following IVIg therapy. In contrast to mice data, IVIg therapy led to basophil expansion only in two patients who also showed increased plasma levels of IL-33. Importantly, the fold-changes in IL-33 and basophils were not correlated and we could hardly detect IL-4 in the plasma following IVIg therapy. Thus, our results indicate that IVIg-induced IL-33 is insufficient to mediate basophil expansion in autoimmune patients. Hence, IL-33 and basophil-mediated anti-inflammatory mechanism proposed for IVIg might not be pertinent in humans.

Intravenous immunoglobulin (IVIg) is a therapeutic preparation of normal pooled immunoglobulin G (IgG) obtained from the plasma of several thousand healthy donors. High-dose IVIg (1–2 g/kg) is widely used in the treatment of various autoimmune and inflammatory diseases including Kawasaki disease, idiopathic thrombocytopenic purpura, Guillain-Barré syndrome, chronic inflammatory demyelinating polyneuropathy, myasthenia gravis, autoimmune blistering diseases, inflammatory myopathies, graft versus host disease and others^{1–4}. The cellular and molecular mechanisms of action of IVIg in these diverse diseases remain incompletely understood. However, available evidence both from experimental and clinical studies provide an indicator that IVIg could benefit these diverse diseases via several mutually non-exclusive mechanisms^{2,5–10}. These mechanisms include inhibition of activation and functions of innate immune cells such as dendritic cells (DCs), monocytes, macrophages and neutrophils; inhibition of pathogenic effector T cells such as Th1 and Th17 cells; expansion of regulatory T cells (Tregs); modulation of B cell responses; and inhibition of complement pathways. In addition, IVIg has been shown to inhibit inflammatory cytokines and to augment anti-inflammatory molecules such as IL-10 and IL-1 receptor antagonist^{11–21}.

IgGs are glycoproteins and contain fragment antigen-binding (Fab) regions that recognize antigens, and fragment crystallizable (Fc) regions that exert effector functions upon binding to Fc γ receptors. The Fc fragments are glycosylated at Asn297 and recent studies in animal models advocate that anti-inflammatory effects of IVIg



are mediated by a small fraction of antibodies that contain terminal α 2,6-sialylated glycans at Asn297. It was proposed that α 2,6-sialylated Fc fragments interact with dendritic cell-specific intercellular adhesion molecule-3-grabbing non-integrin-positive (DC-SIGN⁺) innate cells to release IL-33, which subsequently expands IL-4-producing basophils²². However, translational insights on these observations are lacking. Therefore, we investigated whether high-dose IVIg therapy induces IL-33 production in autoimmune patients, which in turn would mediate basophil expansion and IL-4 responses.

Results

IVIg therapy induces IL-33 in autoimmune patients. Previous work on the role of IL-33 in IVIg-mediated anti-inflammatory effects was performed in K/BxN serum-induced murine arthritis model. It should be noted that IVIg is not recommended for rheumatoid arthritis due to its inefficacy to relieve inflammation⁴. Therefore, K/BxN serum-induced murine arthritis model might not provide factual image of the mechanisms of IVIg in autoimmune patients. Earlier studies have indicated that IVIg therapy benefits patients with inflammatory myopathies^{1,4}. Therefore, by using heparinized blood samples of these patients (cohort 1 patients), we first investigated the repercussion of IVIg therapy on the induction of IL-33. We found that, out of nine patients, six had minimal level of plasma IL-33 prior to IVIg therapy. The pre-IVIg plasma level of IL-33 was in the range of 150.75 ± 79.52 pg/ml ($n = 9$) (Fig. 1a). Following IVIg therapy, with an exception of one patient, all remaining patients had significant raise in plasma IL-33 and was in the range of 492.23 ± 130.30 pg/ml ($n = 9$) (Fig. 1a). However, the increase in IL-33 following IVIg therapy was heterogeneous and was varying from 1.2 to 911-fold.

To confirm these results, we analyzed the plasma samples from another cohort of patients with inflammatory myopathies ($n = 4$) or anti-neutrophil cytoplasmic antibody-associated vasculitis ($n = 3$) (cohort 2 patients). Importantly, these patients also showed significant increase in plasma IL-33 following IVIg therapy (Fig. 1b) thus confirming the results obtained with cohort 1 patients. The pre-IVIg plasma level of IL-33 was 80.43 ± 24.93 pg/ml ($n = 7$) that increased to 291.58 ± 34.40 pg/ml following IVIg therapy. Together, these results indicate that irrespective of pathologies, IVIg therapy in patients leads to increased plasma level of IL-33.

IVIg-induced IL-33 is not associated with an expansion of basophils. Basophils play a crucial role in the induction of Th2 responses^{23,24}. Recent data from K/BxN serum-induced murine arthritis model suggest that IVIg-induced IL-33 promotes basophil expansion²². Therefore, we investigated changes in the circulating basophils following IVIg therapy in cohort 1 patients. Basophils were identified based on the expression of FcεR1 and CD203c (Fig. 2a)²⁵. In contrast to the results from murine model, we found that IVIg therapy leads to basophil expansion only in two patients who also showed increased plasma level of IL-33 (Fig. 2b). In other patients, basophils were either declined or unaltered. The changes in the proportion of basophils in the circulation following IVIg therapy were not statistically significant. Importantly, the fold-changes in IL-33 and basophils were not correlated (Fig. 2c). Also contrary to previous report²², we could hardly detect IL-4 in the plasma of patients following IVIg therapy. Thus, these results demonstrate that IVIg therapy in patients does not lead to an expansion of basophils. Of note, a recent data from murine models of collagen antibody-induced arthritis and K/BxN serum transfer arthritis also reveal that therapeutic effect of IVIg is independent of sialylation and basophils²⁶.

DC-SIGN-positive human innate cells do not produce IL-33 upon IVIg exposure. DC-SIGN⁺ innate cells (or SIGN-R1⁺ cells in the murine spleen) were proposed to produce IL-33 upon interaction




Figure 1 | Consequence of IVIg therapy in autoimmune patients on the plasma level of IL-33. (a) Heparinized blood samples were obtained from nine patients with inflammatory myopathies (Cohort 1 patients) before (Pre-IVIg) and 2-3 days after initiation of IVIg therapy (Post-IVIg). IL-33 (pg/ml) in the plasma was measured by ELISA. Each symbol in the graph represents individual patient. (b) IL-33 in the plasma of four inflammatory myopathies and three anti-neutrophil cytoplasmic antibody-associated vasculitis patients (Cohort 2 patients) before and post-IVIg therapy. The statistical significance as determined by two-tailed Student-t-test is indicated, where *, $P < 0.05$; **, $P < 0.01$.

with α 2,6-sialylated Fc fragments of IVIg²². By generating humanized DC-SIGN-transgenic mice, the authors found that these transgenic mice express DC-SIGN on DCs, macrophages and monocytes in the blood, bone marrow and spleen. Importantly, higher percentage of monocytes in these transgenic mice expressed DC-SIGN²².

We analyzed the expression of DC-SIGN in human myeloid cells. Contrary to humanized DC-SIGN-transgenic mice, circulating human monocytes did not express DC-SIGN whereas its expression on macrophages was restricted to M2 type macrophages wherein up to 28% cells were positive for DC-SIGN. We could observe high expression of DC-SIGN ($\approx 100\%$) only in monocyte-derived DCs (Mo-DCs) (Fig. 3a). In the human spleen, up to 24% splenocytes were positive for DC-SIGN (Fig. 3b).

Therefore, we explored if Mo-DCs secrete IL-33 upon IVIg treatment. In contrast to proposition by Ravetch and colleagues, we could




Figure 2 | Changes in the proportion of circulating basophils of autoimmune patients following IVIg therapy. Heparinized blood samples were obtained from cohort 1 patients with inflammatory myopathies before (Pre-IVIg) and 2–3 days after initiation of IVIg therapy (Post-IVIg). (a) Representative dot-plots showing basophils from cohort 1 patients gated positive for FcεRI and CD203c (b) Modulation of circulating basophils following IVIg therapy (n = 9). Basophils were analyzed in the whole blood by flow cytometry. The statistical significance as determined by two-tailed Student-t-test is indicated, where NS, non-significant. (c) The correlation between fold-changes in IL-33 and basophils following IVIg therapy.

detect secreted IL-33 from IVIg-exposed DC-SIGN⁺ Mo-DCs neither under non-inflammatory nor under inflammatory conditions (Fig. 3c). Similarly, despite the presence of DC-SIGN⁺ cells in the spleen, human splenocytes did not produce detectable levels of IL-33 upon IVIg exposure both under inflammatory and non-inflammatory conditions (Fig. 3c).

Discussion

Our results demonstrate that IVIg therapy induces IL-33 in autoimmune patients thus confirming the previous observation made in mice. However, IL-33 was not contributed either by splenic




Figure 3 | Effect of IVIg on the IL-33 production from DC-SIGN⁺ human innate cells. (a and b) Histograms showing the expression of DC-SIGN by healthy donor's monocyte-derived human dendritic cells (Mo-DCs) and splenocytes. (c) IVIg does not induce IL-33 from DC-SIGN⁺ human innate cells. Mo-DCs or human splenocytes (n = 5 donors) were exposed to IVIg either under non-inflammatory conditions or under inflammatory conditions (TLR-stimuli or inflammatory cytokine cocktail) for 48 hours. IL-33 in the culture supernatants was analyzed by ELISA.

DC-SIGN⁺ cells or myeloid DCs. Also, the amount of IL-33 induced in the patients was not sufficient to expand basophils. It should be noted that the quantity of IL-33 protein induced in the mice following IVIg treatment was not presented in the previous report. In addition, significant amount of data on IVIg was indirect rather than direct demonstration of IVIg-mediated regulation of cytokine network²². Authors showed that IVIg induces about 12-fold increase in IL-33 mRNA level. However, the contribution of this increased IL-33 mRNA towards IL-33 protein is not clear. Considering five liters as total blood volume in adults, our results show that IVIg induces $\approx 2460 \pm 650$ ng of IL-33 (based on the data from cohort 1 patients). However, to demonstrate the role of IL-33 in IVIg-mediated anti-inflammatory effects, Anthony *et al.*, injected 400 ng of IL-33 for four consecutive days²². As mouse weighing 25 g would have ≈ 1.5 ml of blood, based on the IL-33 data from patients, we could infer that the amount of exogenous IL-33 injected into the mice represents at least 540-times excess of IL-33 that otherwise induced by IVIg. This might explain why IVIg failed to induce expansion of basophils in the patients. Although in our study, patients' sample size was small, we included diseases such as inflammatory myopathies and vasculitis that were shown to benefit from IVIg therapy. Further investigations in a larger number of patients should confirm these observations.



Table 1 | Summary of data for autoimmune rheumatic patients

Cohort 1 patients					
Number	Disease	Age (years)	Sex	IVIg	Additional treatments
1	Polymyositis	59	F	CLAIRYG [®] 1 g/kg	Methylprednisolone
2	Anti-SRP associated necrotizing myopathy	27	F	CLAIRYG [®] 1 g/kg	Prednisone, Methotrexate
3	Anti-HMGCR associated necrotizing myopathy	62	F	CLAIRYG [®] 0.5 g/kg	Prednisone, Methotrexate
4	Anti-HMGCR associated necrotizing myopathy	61	F	CLAIRYG [®] 1 g/kg	Prednisone, Methotrexate
5	Dermatomyositis	52	F	CLAIRYG [®] 1 g/kg	Prednisone, Methotrexate
6	Polymyositis associated with mixed connective tissue disease and Sjögren's syndrome	41	F	CLAIRYG [®] 1 g/kg	Prednisone, Methotrexate
7	Anti-SRP associated necrotizing myopathy	40	M	CLAIRYG [®] 1 g/kg	Prednisone, Methotrexate
8	Anti-Mi2 associated unclassified myositis	30	M	CLAIRYG [®] 1 g/kg	Prednisone, Methotrexate
9	Polymyositis and probable associated Sjögren's syndrome	70	F	CLAIRYG [®] 1 g/kg	Prednisone, Methotrexate
Cohort 2 patients					
Number	Disease	Age (years)	Sex	IVIg	Additional treatments
1	Dermatomyositis	22	F	TEGELINE [®] 1g/kg	Prednisone, Mycophenolate mofetil
2	Polymyositis	42	M	TEGELINE [®] 1g/kg	Prednisone, Methotrexate
3	Dermatomyositis	35	M	TEGELINE [®] 1g/kg	Prednisone
4	Polymyositis	46	F	TEGELINE [®] 1g/kg	Prednisone, ciclosporin
5	Microscopic polyangiitis	61	F	TEGELINE [®] 1g/kg	Prednisone
6	Wegener's granulomatosis	62	M	TEGELINE [®] 1g/kg	None
7	Microscopic polyangiitis	61	M	TEGELINE [®] 1g/kg	Prednisone, Mycophenolate mofetil

SRP, Signal Recognition Particle; HMGCR, 3-hydroxy-3-methylglutaryl-CoA reductase.

The role of Fc-sialylation, DC-SIGN and Fc γ receptor IIB (Fc γ RIIB) in the anti-inflammatory effects of IVIg has been debated recently by several groups²⁷. Mice and humans show wide variations in the expression pattern of Fc γ Rs, and the phenotype and anatomical distribution of innate cells. Unlike mice, human innate cells express both activating Fc γ RIIA and inhibitory Fc γ RIIB. Therefore, the proposition that IVIg enhances Fc γ RIIB on effector macrophages of mice without having corresponding data on Fc γ RIIA might provide a biased picture on the mechanisms of IVIg. Gene array analysis could not confirm IVIg-mediated up-regulation of Fc γ RIIB in the patients with Kawasaki disease²⁸. In line with this report, another recent study failed to demonstrate enhanced expression of Fc γ RIIB on monocytes following IVIg therapy in children with immune thrombocytopenia²⁹. Also, Fc γ R polymorphisms did not predict response to IVIg in myasthenia gravis³⁰. Although DC-SIGN promoter -336 A/G (rs4804803) polymorphism was associated with susceptibility of Kawasaki disease, this variant was found to be not associated with the occurrence of IVIg resistance³¹. Of note, treatment response in Kawasaki disease is apparently associated with sialylation levels of endogenous IgG but not therapeutic IVIg³². All these data thus questions the relevance of DC-SIGN-Fc γ RIIB pathway of anti-inflammatory mechanisms of IVIg in humans.

Several recent studies have challenged the concept of α 2,6-sialylated Fc fragments-mediated anti-inflammatory mechanism of IVIg both in experimental models and in humans. IVIg could inhibit human Th17 cell differentiation and expansion independent of antigen presenting cells and hence independent of interaction of DC-SIGN and α 2,6-sialylated Fc fragments¹³⁻¹⁵. Also, F(ab')₂ fragments of IVIg exerted similar effects thus pointing towards dispensability of α 2,6-sialylated Fc fragments in mediating the suppression of Th17 cells. We have demonstrated that DC-SIGN and α 2,6-sialylated Fc fragment interaction is dispensable for the anti-inflammatory activity of IVIg on human DCs³³. F(ab')₂ fragments but not Fc fragments of IVIg were shown to mediate Treg expansion by inducing cyclooxygenase-2-mediated prostaglandin E2 secretion in human myeloid DCs and was dependent in part on DC-SIGN¹⁹. Similarly, sialylation-

enriched F(ab')₂ fragments could inhibit interferon- α production from toll-like receptor (TLR)7 and TLR9 stimulated human plasmacytoid DCs, although sialic acid itself was not required³⁴.

In the previous reports, Ravetch and colleagues enriched sialic acid-containing IgG-Fc by using sialic acid-specific lectin *Sambucus nigra* agglutinin-based affinity fractionation^{22,35-37}. However, by using same fractionation method, Guhr *et al.*, showed that IVIg fractions depleted for the sialylated antibody fraction exert benefits in a murine model of passive-immune thrombocytopenia similar to that of intact IVIg. However, sialic acid-enriched IVIg fraction failed to enhance platelets count in this model³⁸. Similar sialic-acid independent anti-inflammatory mechanisms were also reported in murine herpes simplex virus encephalitis model³⁹. Further, Käsermann and colleagues showed that lectin fractionation of IVIg results in increased sialylation of Fab fragments but not Fc fragments. By using human whole blood stimulation assay either with lipopolysaccharide or phytohaemagglutinin, they further showed that anti-inflammatory effects of IVIg is associated with F(ab')₂ fraction of IVIg⁴⁰. In animal models of immune thrombocytopenia and multiple sclerosis, the beneficial effects of IVIg were independent of Fc or F(ab')₂ -sialylation and Fc γ RIIB⁴¹⁻⁴⁴. Based on these results, it was suggested that genetic background of the mice and dose of IVIg are the critical factors that determine the role of Fc γ RIIB in IVIg-mediated beneficial effects. In line with these observations, two studies have failed to demonstrate the direct interaction between sialylated IgG Fc fragment and DC-SIGN^{45,46}. These data thus point out that α 2,6-sialylated Fc fragment-DC-SIGN-Fc γ RIIB mechanism merely represents one of the several anti-inflammatory mechanisms of IVIg that were reported. Therefore, this anti-inflammatory pathway of IVIg might be operational in certain pathologies and experimental models and might not be considered as a universal mechanism.

It was proposed that in humanized DC-SIGN-transgenic mice, DC-SIGN⁺ innate cells such as monocytes, macrophages and DCs produce IL-33 upon interaction with α 2,6-sialylated Fc fragments of IVIg²². Recent reports show that IL-33 is an important player for the promotion of Th2 responses and activated DCs are one of the sources of this cytokine^{47,48}. We found that unlike monocytes from huma-



nized DC-SIGN-transgenic mice that were highly positive for DC-SIGN, human monocytes hardly express DC-SIGN. Further, human Mo-DCs despite expressing DC-SIGN, failed to produce IL-33 when exposed to IVIg either under non-inflammatory or inflammatory conditions. In wild type mice, it was suggested that α 2,6-sialylated Fc fragments bind to SIGN-R1 expressed on splenic marginal zone macrophages³⁵. Marginal zone macrophages are absent in human spleen and data from humans show that spleen is dispensable for the anti-inflammatory effects of IVIg. In line with this concept, by using a passive model of induced immune thrombocytopenia, it was shown that IVIg is fully functional in splenectomized mice although this report supported the sialic acid and SIGN-R1-dependent mechanisms of IVIg⁴⁹. We found that despite the presence of DC-SIGN⁺ innate cells in the human spleen, IVIg could not induce IL-33 from the splenocytes. All these data indicate that spleen and DC-SIGN⁺ cells are dispensable for IVIg-mediated IL-33 induction in humans. Thus, the source of IL-33 in humans following IVIg therapy remains elusive. As IVIg is known to cause apoptosis of cells, we suggest that secondary necrosis of late stage apoptotic cells could release IL-33^{50–52}. This process might depend on the signals provided by anti-Fas IgG or anti-Siglec IgG in the IVIg preparations rather than the repercussion of interaction of α 2,6-sialylated Fc fragments with DC-SIGN^{53,54}. In addition, IL-33 is also constitutively expressed in the nucleus of endothelial cells and epithelial cells *in vivo*⁵⁵.

Methods

Patients. All experiments were performed in accordance with relevant guidelines and regulations. We obtained heparinized blood samples of nine patients (cohort 1 patients) with inflammatory myopathies (Table 1). Patients were aged 49.1 ± 15.2 years and include two men. Blood samples were obtained before and 2–3 days following initiation of IVIg therapy (CLAIRYG[®], Laboratoire Français du Fractionnement et des Biotechnologies, France). Informed consent was obtained from all the patients. The study was approved by CPP-Ile-de-France VI, Groupe Hospitalier Pitié-Salpêtrière, Paris. In addition, we also analyzed plasma samples of seven rheumatic patients (cohort 2 patients) before and 2–3 days post-IVIg therapy (TEGELINE[®], Laboratoire Français du Fractionnement et des Biotechnologies). The patients were aged 47 ± 5.8 years (four men) and include inflammatory myopathies and anti-neutrophil cytoplasmic antibody-associated vasculitis (Table 1).

Analysis of basophils. Red blood cells (RBCs) from heparinized blood samples of cohort 1 patients were depleted by using HetaSep[™] (Stemcell Technologies Sarl, France) and nucleated cell suspension was obtained. Basophils were analyzed in RBC-depleted cell suspension by flow cytometry (LSR II, BD Biosciences, France) using fluorochrome-conjugated monoclonal antibodies to Fc ϵ RI (Miltenyi Biotec, France) and CD203c (eBioscience, France). Data were analyzed by FACSDiva[™] software (BD Biosciences).

Generation of monocyte-derived DCs. Buffy coats from the healthy donors were purchased from Centre Necker-Cabanel, Etablissement Français du Sang (EFS), Paris, France. Institut National de la Santé et de la Recherche Médicale-EFS ethical committee permission (N[°]12/EFS/079) was obtained for the use of buffy coats of healthy donors. Peripheral blood mononuclear cells (PBMCs) were purified from the buffy coats by density gradient centrifugation using Ficoll-paque PREMIUM (GE healthcare, France). CD14⁺ monocytes were isolated from PBMCs by using CD14 microbeads (Miltenyi Biotec). Purified monocytes were then cultured for 6 days in RPMI-1640 medium plus 10% fetal calf serum (FCS) containing cytokines GM-CSF (1000 IU/10⁶ cells) and IL-4 (500 IU/10⁶ cells) (both from Miltenyi Biotec) to obtain DCs⁵⁶. The purity of DCs was >98%. DC-SIGN expression on Mo-DCs was examined by flow cytometry using fluorochrome-conjugated monoclonal antibodies (BD Biosciences) and data were analyzed by FACSDiva[™] and FlowJo softwares (Tree Star, USA).

Isolation of human splenocytes. The remnant human spleen sections from individuals submitted for pathological diagnosis were obtained from service d'anatomie pathologique, Hôpital Européen Georges Pompidou, Paris, France. Only healthy spleen tissues were used for the research purpose. Since the study did not require additional sampling, an approval from an ethics committee was not required under French law according to the article L.1121-1 of the public health code. The article states that: The research organized and performed on human beings in the development of biological knowledge and medical research are permitted under the conditions laid down in this book and are hereinafter referred to by the term "biomedical research". The article further states that it does not imply under conditions: For research in which all actions are performed and products used in the usual way, without any additional or unusual diagnostic procedure or surveillance.

The spleen sections were collected in RPMI 1640 medium supplemented with 100 IU/ml penicillin, 100 μ g/ml streptomycin, and 10% FCS. Single-cell suspension

of splenocytes was obtained by mechanical disaggregation of spleen tissue pieces by using gentleMACS dissociator (Miltenyi Biotec) followed by filtration through 70- μ m nylon membrane filter (BD Biosciences). Splenocytes were then subjected to Ficoll-Paque PREMIUM density gradient centrifugation to obtain mononuclear cells. DC-SIGN expression on the splenocytes was investigated by flow cytometry using fluorochrome-conjugated monoclonal antibodies and data were analyzed by FACSDiva[™] and FlowJo softwares.

Stimulation of cells. Mo-DCs (0.5×10^6 /ml) were cultured in RPMI 1640-10% FCS containing GM-CSF and IL-4 in a 12-well plate. The cells were then exposed to IVIg (25 mg/ml) for 48 hours to analyze the effect of IVIg on IL-33 production under non-inflammatory conditions. In parallel, Mo-DCs were stimulated with either TLR4 ligand lipopolysaccharide (100 ng/ml/0.5 $\times 10^6$ cells) (Sigma-Aldrich, France) or inflammatory cytokine cocktail (10 ng/ml each of IL-1 β , IL-6 and TNF- α , all from ImmunoTools, Germany)⁵⁷. After four hours, IVIg was added and cultures were maintained for 48 hours to analyze the effect of IVIg on IL-33 production under inflammatory conditions.

Similarly, splenocytes (0.5×10^6 /ml) were cultured in RPMI 1640-10% FCS for 48 hours either alone or with IVIg. In addition, splenocytes were also stimulated with inflammatory cytokine cocktail and IVIg was added to the cultures after four hours. The cultures were maintained for 48 hours.

Quantification of cytokines. IL-33 in the plasma samples of the patients and in cell-free culture supernatants was quantified by ELISA (R&D systems, France). IL-4 in the plasma was also measured by ELISA (R&D systems).

Statistical analysis. Data was analyzed using Prism 5 software (GraphPad software). Two-tailed Student's t-test was used to determine the statistical significance of the data. Values of $P < 0.05$ were considered as statistically significant.

- Dalakas, M. C. Intravenous immunoglobulin in autoimmune neuromuscular diseases. *JAMA* **291**, 2367–2375 (2004).
- Kazatchkine, M. D. & Kaveri, S. V. Immunomodulation of autoimmune and inflammatory diseases with intravenous immune globulin. *N. Engl. J. Med.* **345**, 747–755 (2001).
- Arnson, Y., Shoenfeld, Y. & Amital, H. Intravenous immunoglobulin therapy for autoimmune diseases. *Autoimmunity* **42**, 553–560 (2009).
- Bayry, J., Negi, V. S. & Kaveri, S. V. Intravenous immunoglobulin therapy in rheumatic diseases. *Nat. Rev. Rheumatol.* **7**, 349–359 (2011).
- Schwab, I. & Nimmerjahn F. Intravenous immunoglobulin therapy: how does IgG modulate the immune system? *Nat. Rev. Immunol.* **13**, 176–189 (2013).
- Seite, J. F., Shoenfeld, Y., Youinou, P. & Hillion, S. What is the contents of the magic draft IVIg? *Autoimmun. Rev.* **7**, 435–439 (2008).
- Tha-In, T., Bayry, J., Metselaar, H. J., Kaveri, S. V. & Kwekkeboom, J. Modulation of the cellular immune system by intravenous immunoglobulin. *Trends Immunol.* **29**, 608–615 (2008).
- Ballou, M. The IgG molecule as a biological immune response modifier: Mechanisms of action of intravenous immune serum globulin in autoimmune and inflammatory disorders. *J. Allergy Clin. Immunol.* **127**, 315–323 (2011).
- Jordan, S. C., Toyoda, M. & Vo, A. A. Intravenous immunoglobulin a natural regulator of immunity and inflammation. *Transplantation* **88**, 1–6 (2009).
- Seemple, J. W. *et al.* Intravenous immunoglobulin prevents murine antibody-mediated acute lung injury at the level of neutrophil reactive oxygen species (ROS) production. *PLoS One* **7**, e31357 (2012).
- Bayry, J. *et al.* Intravenous immunoglobulin abrogates dendritic cell differentiation induced by interferon-alpha present in serum from patients with systemic lupus erythematosus. *Arthritis Rheum.* **48**, 3497–3502 (2003).
- Aubin, E., Lemieux, R. & Bazin, R. Indirect inhibition of *in vivo* and *in vitro* T-cell responses by intravenous immunoglobulins due to impaired antigen presentation. *Blood* **115**, 1727–1734 (2010).
- Maddur, M. S. *et al.* Inhibition of differentiation, amplification, and function of human TH17 cells by intravenous immunoglobulin. *J. Allergy Clin. Immunol.* **127**, 823–830 e821–827 (2011).
- Maddur, M. S., Kaveri, S. V. & Bayry, J. Comparison of different IVIg preparations on IL-17 production by human Th17 cells. *Autoimmun. Rev.* **10**, 809–810 (2011).
- Maddur, M. S. *et al.* Inhibitory effect of IVIG on IL-17 production by Th17 cells is independent of anti-IL-17 antibodies in the immunoglobulin preparations. *J. Clin. Immunol.* **33**, S62–S66 (2013).
- Kessel, A. *et al.* Intravenous immunoglobulin therapy affects T regulatory cells by increasing their suppressive function. *J. Immunol.* **179**, 5571–5575 (2007).
- De Groot, A. S. *et al.* Activation of natural regulatory T cells by IgG Fc- derived peptide "Tregitopes". *Blood* **112**, 3303–3311 (2008).
- Massoud, A. H. *et al.* Intravenous immunoglobulin attenuates airway inflammation through induction of forkhead box protein 3-positive regulatory T cells. *J. Allergy Clin. Immunol.* **129**, 1656–1665 e1653 (2012).
- Trinath, J. *et al.* Intravenous immunoglobulin expands regulatory T cells via induction of cyclooxygenase-2-dependent prostaglandin E2 in human dendritic cells. *Blood* **122**, 1419–1427 (2013).
- Seite, J. F., Goutsmedt, C., Youinou, P., Pers, J. O. & Hillion, S. Intravenous immunoglobulin induces a functional silencing program similar to energy in human B cells. *J. Allergy Clin. Immunol.* **133**, 181–188 e181–189 (2014).



21. Massoud, A. H. *et al.* Dendritic cell immunoreceptor: a novel receptor for intravenous immunoglobulin mediates induction of regulatory T cells. *J. Allergy Clin. Immunol.* **133**, 853–863 e855 (2014).
22. Anthony, R. M., Kobayashi, T., Wermeling, F. & Ravetch, J. V. Intravenous gammaglobulin suppresses inflammation through a novel T(H)2 pathway. *Nature* **475**, 110–113 (2011).
23. Voehringer, D. Protective and pathological roles of mast cells and basophils. *Nat. Rev. Immunol.* **13**, 362–375 (2013).
24. Otsuka, A. *et al.* Basophils are required for the induction of Th2 immunity to haptens and peptide antigens. *Nat. Commun.* **4**, 1739, doi:10.1038/ncomms2740 (2013).
25. Sharma, M. *et al.* Circulating human basophils lack the features of professional antigen presenting cells. *Sci. Rep.* **3**, 1188, doi: 10.1038/srep01188 (2013).
26. Campbell, I. K. *et al.* Therapeutic effect of IVIG on inflammatory arthritis in mice is dependent on the Fc portion and independent of sialylation or basophils. *J. Immunol.* **192**, 5031–5038 (2014).
27. von Gunten, S. *et al.* IVIG pluripotency and the concept of Fc-sialylation: challenges to the scientist. *Nat. Rev. Immunol.* **14**, 349 (2014).
28. Abe, J. *et al.* Gene expression profiling of the effect of high-dose intravenous Ig in patients with Kawasaki disease. *J. Immunol.* **174**, 5837–5845 (2005).
29. Shimomura, M. *et al.* Intravenous immunoglobulin does not increase FcγRIIB expression levels on monocytes in children with immune thrombocytopenia. *Clin. Exp. Immunol.* **169**, 33–37 (2012).
30. Barnett, C. *et al.* Fcγ receptor polymorphisms do not predict response to intravenous immunoglobulin in myasthenia gravis. *J. Clin. Neuromuscul. Dis.* **14**, 1–6 (2012).
31. Yu, H. R. *et al.* DC-SIGN (CD209) promoter -336 A/G (rs4804803) polymorphism associated with susceptibility of Kawasaki disease. *Sci. World J.* **2012**, 634835 (2012).
32. Ogata, S. *et al.* Treatment response in kawasaki disease is associated with sialylation levels of endogenous but not therapeutic intravenous immunoglobulin G. *PLoS One* **8**, e81448 (2013).
33. Bayry, J., Bansal, K., Kazatchkine, M. D. & Kaveri, S. V. DC-SIGN and α2,6-sialylated IgG Fc interaction is dispensable for the anti-inflammatory activity of IVIg on human dendritic cells. *Proc. Natl. Acad. Sci. U S A.* **106**, E24 (2009).
34. Wiedeman, A. E. *et al.* Contrasting mechanisms of interferon-α inhibition by intravenous immunoglobulin after induction by immune complexes versus Toll-like receptor agonists. *Arthritis Rheum.* **65**, 2713–2723 (2013).
35. Anthony, R. M., Wermeling, F., Karlsson, M. C. & Ravetch, J. V. Identification of a receptor required for the anti-inflammatory activity of IVIG. *Proc. Natl. Acad. Sci. U S A.* **105**, 19571–19578 (2008).
36. Anthony, R. M. *et al.* Recapitulation of IVIG anti-inflammatory activity with a recombinant IgG Fc. *Science* **320**, 373–376 (2008).
37. Kaneko, Y., Nimmerjahn, F. & Ravetch, J. V. Anti-inflammatory activity of immunoglobulin G resulting from Fc sialylation. *Science* **313**, 670–673 (2006).
38. Guhr, T. *et al.* Enrichment of sialylated IgG by lectin fractionation does not enhance the efficacy of immunoglobulin G in a murine model of immune thrombocytopenia. *PLoS One* **6**, e21246 (2011).
39. Ramakrishna, C., Newo, A. N., Shen, Y. W. & Cantin, E. Passively administered pooled human immunoglobulins exert IL-10 dependent anti-inflammatory effects that protect against fatal HSV encephalitis. *PLoS Pathog.* **7**, e1002071 (2011).
40. Kasermann, F. *et al.* Analysis and functional consequences of increased Fc-sialylation of intravenous immunoglobulin (IVIg) after lectin fractionation. *PLoS One* **7**, e37243 (2012).
41. Leontyev, D. *et al.* Sialylation-independent mechanism involved in the amelioration of murine immune thrombocytopenia using intravenous gammaglobulin. *Transfusion* **52**, 1799–1805 (2012).
42. Leontyev, D., Katsman, Y. & Branch, D. R. Mouse background and IVIG dosage are critical in establishing the role of inhibitory Fcγ receptor for the amelioration of experimental ITP. *Blood* **119**, 5261–5264 (2012).
43. Othy, S. *et al.* Intravenous gammaglobulin inhibits encephalitogenic potential of pathogenic T cells and interferes with their trafficking to the central nervous system, implicating sphingosine-1 phosphate receptor 1-mammalian target of rapamycin axis. *J. Immunol.* **190**, 4535–4541 (2013).
44. Othy, S. *et al.* Sialylation may be dispensable for reciprocal modulation of helper T cells by intravenous immunoglobulin. *Eur. J. Immunol.* (epub ahead of print) doi:10.1002/eji.201444440 (2014).
45. Crispin, M., Yu, X. & Bowden, T. A. Crystal structure of sialylated IgG Fc: implications for the mechanism of intravenous immunoglobulin therapy. *Proc. Natl. Acad. Sci. U S A.* **110**, E3544–3546 (2013).
46. Yu, X., Vasiljevic, S., Mitchell, D. A., Crispin, M. & Scanlan, C. N. Dissecting the molecular mechanism of IVIg therapy: the interaction between serum IgG and DC-SIGN is independent of antibody glycoform or Fc domain. *J. Mol. Biol.* **425**, 1253–1258 (2013).
47. Grotenboer, N. S., Ketelaar, M. E., Koppelman, G. H. & Nawijn, M. C. Decoding asthma: translating genetic variation in IL33 and IL1RL1 into disease pathophysiology. *J. Allergy Clin. Immunol.* **131**, 856–865 (2013).
48. Williams, J. W. *et al.* Transcription factor IRF4 drives dendritic cells to promote Th2 differentiation. *Nat. Commun.* **4**, 2990, doi:10.1038/ncomms3990 (2013).
49. Schwab, L., Biburger, M., Kronke, G., Schett, G. & Nimmerjahn, F. IVIg-mediated amelioration of ITP in mice is dependent on sialic acid and SIGNR1. *Eur. J. Immunol.* **42**, 826–830 (2012).
50. Bonilla, W. V. *et al.* The alarmin interleukin-33 drives protective antiviral CD8+ T cell responses. *Science* **335**, 984–989 (2012).
51. Cayrol, C. & Girard, J. P. The IL-1-like cytokine IL-33 is inactivated after maturation by caspase-1. *Proc. Natl. Acad. Sci. U S A.* **106**, 9021–9026 (2009).
52. Haraldsen, G., Balogh, J., Pollheimer, J., Sponheim, J. & Kuchler, A. M. Interleukin-33 - cytokine of dual function or novel alarmin? *Trends Immunol.* **30**, 227–233 (2009).
53. Prasad, N. K. *et al.* Therapeutic preparations of normal polyspecific IgG (IVIg) induce apoptosis in human lymphocytes and monocytes: a novel mechanism of action of IVIg involving the Fas apoptotic pathway. *J. Immunol.* **161**, 3781–3790 (1998).
54. von Gunten, S. *et al.* Immunologic and functional evidence for anti-Siglec-9 autoantibodies in intravenous immunoglobulin preparations. *Blood* **108**, 4255–4259 (2006).
55. Moussion, C., Ortega, N. & Girard, J. P. The IL-1-like cytokine IL-33 is constitutively expressed in the nucleus of endothelial cells and epithelial cells in vivo: a novel 'alarmin'? *PLoS One* **3**, e3331 (2008).
56. Maddur, M. S. *et al.* Human B cells induce dendritic cell maturation and favour Th2 polarization by inducing OX-40 ligand. *Nat. Commun.* **5**, 4092, doi:10.1038/ncomms5092 (2014).
57. Banchereau, J. *et al.* Immunobiology of dendritic cells. *Annu. Rev. Immunol.* **18**, 767–811 (2000).

Acknowledgments

This study was supported by Institut National de la Santé et de la Recherche Médicale (INSERM), Centre National de la Recherche Scientifique (CNRS), Université Pierre et Marie Curie, Université Paris Descartes and European Community's Seventh Framework Programme [FP7/2007–2013] under grant agreement HEALTH-2010.2.4.5-2 ALLFUN. We also thank Laboratoire Français du Fractionnement et des Biotechnologies, France for the support.

Author contributions

J.B. designed the research, M.S., C.S., P.H., M.S.M., E.S.-V., L.G. & M.L. performed the experiments, M.S., P.H., M.S.M., S.V.K. & J.B. analyzed the data, Y.S., L.M. & O.B. provided blood samples of the patients, P.B. provided the spleen tissues, J.B. wrote the paper and all authors reviewed and approved the manuscript.

Additional information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Sharma, M. *et al.* Intravenous immunoglobulin-induced IL-33 is insufficient to mediate basophil expansion in autoimmune patients. *Sci. Rep.* **4**, 5672; DOI:10.1038/srep05672 (2014).



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>