



HAL
open science

Changes in Sedentary Behaviours and Associations with Physical Activity through Retirement: A 6-Year Longitudinal Study

Mehdi Menai, Léopold Fezeu, Hélène Charreire, Emmanuelle Kesse-Guyot, Mathilde Touvier, Chantal Simon, Christiane Weber, Valentina A. Andreeva, Serge Hercberg, Jean-Michel Oppert

► **To cite this version:**

Mehdi Menai, Léopold Fezeu, Hélène Charreire, Emmanuelle Kesse-Guyot, Mathilde Touvier, et al.. Changes in Sedentary Behaviours and Associations with Physical Activity through Retirement: A 6-Year Longitudinal Study. PLoS ONE, 2014, 9 (9), pp.e106850. 10.1371/journal.pone.0106850 . hal-01367444

HAL Id: hal-01367444

<https://hal.sorbonne-universite.fr/hal-01367444>

Submitted on 16 Sep 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution| 4.0 International License



Changes in Sedentary Behaviours and Associations with Physical Activity through Retirement: A 6-Year Longitudinal Study

Mehdi Menai¹, Léopold Fezeu¹, Hélène Charreire^{1,2}, Emmanuelle Kesse-Guyot¹, Mathilde Touvier¹, Chantal Simon³, Christiane Weber⁴, Valentina A. Andreeva¹, Serge Hercberg^{1,5}, Jean-Michel Oppert^{1,6*}

1 Université Paris 13, Sorbonne Paris Cité - EREN (Equipe de Recherche en Epidémiologie Nutritionnelle), U1153 Inserm, Inra, Cnam, Centre de Recherche en Epidémiologie et Biostatistiques; CRNH IdF, Bobigny, France, **2** Paris-Est Créteil University, Department of Geography, Lab-Urba, Urbanism Institute of Paris, Paris, France, **3** CARMEN, INSERM U1060/Université de Lyon 1/INRA U1235 Lyon, France, **4** Laboratoire Image, Ville et Environnement, Université de Strasbourg, Strasbourg, France, **5** Department of Public Health, Hôpital Avicenne (AP-HP), Bobigny, France, **6** Université Pierre et Marie Curie-Paris 6, Dept of Nutrition Pitié-Salpêtrière Hospital (AP-HP), Centre for Research on Human Nutrition Ile-de-France (CRNH IdF), Institute of Cardiometabolism and Nutrition (ICAN), Paris, France

Abstract

Changes in sedentary behaviours and physical activity according to retirement status need to be better defined. Retirement is a critical life period that may influence a number of health behaviours. We assessed past-year sedentary behaviours (television, computer and reading time during leisure, occupational and domestic sitting time, in h/week) and physical activity (leisure, occupational and domestic, in h/week) over 6 years (2000–2001 and 2007) using the Modifiable Activity Questionnaire in 2,841 participants (mean age: 57.3±5.0 y) of the SU.VI.MAX (Supplementation with Antioxidants and Minerals) cohort. Analyses were performed according to retirement status. Subjects retired in 2001 and 2007 (40%) were those who spent most time in sedentary behaviour and in physical activity during and outside leisure ($p<0.001$). Leisure-time sedentary behaviours increased in all subjects during follow-up ($p<0.001$), but subjects who retired between 2001 and 2007 (31%) were those who reported the greatest changes (+8.4±0.42 h/week for a combined indicator of leisure-time sedentary behaviour). They also had the greatest increase in time spent in leisure-time physical activity (+2.5±0.2 h/week). In subjects not retired 2001 and 2007 (29%), changes in time spent watching television were found positively associated with an increase in occupational physical activity ($p=0.04$) and negatively associated with changes in leisure-time physical activity ($p=0.02$). No consistent association between changes in sedentary behaviours and changes in physical activity was observed in subjects retired in 2001 and 2007. Public health interventions should target retiring age populations not only to encourage physical activity but also to limit sedentary behaviours.

Citation: Menai M, Fezeu L, Charreire H, Kesse-Guyot E, Touvier M, et al. (2014) Changes in Sedentary Behaviours and Associations with Physical Activity through Retirement: A 6-Year Longitudinal Study. PLoS ONE 9(9): e106850. doi:10.1371/journal.pone.0106850

Editor: Hemachandra Reddy, Oregon Health & Science University, United States of America

Received: March 26, 2014; **Accepted:** August 6, 2014; **Published:** September 26, 2014

Copyright: © 2014 Menai et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: The authors confirm that all data underlying the findings are fully available without restriction. All relevant data are within the paper and its Supporting Information files.

Funding: This work was funded by the French National Research Agency (ANR-05-PNRA-010), the French National Cancer Institute (INCA 2011-1-PL-SHS-10), the French Ministry of Health, Mederic, Sodexo, Ipsen, MGEN and Pierre Fabre. Mederic and MGEN are French health insurance organizations, which are complementary to the National Health Insurance System. Ipsen and Pierre Fabre are private pharmaceutical companies. Sodexo, a food catering company, supported the study by organizing meetings between researchers and study participants. They financially supported the overall implementation of the research project. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors received funding from commercial sources (Mederic, Sodexo, Ipsen, MGEN and Pierre Fabre). This does not alter their adherence to PLOS ONE policies on sharing data and materials.

* Email: jean-michel.oppert@psl.aphp.fr

Background

According to the World Health Organization, the number of people 60 years of age or older worldwide will grow from 600 million to 2 billion by 2050. A large range of determinants (sociological, medical and technological) alter the habits and the behaviours of active and retired populations. The transition to retirement is considered as a major life event in terms of financial as well as behavioural modification, including important changes in sedentary and physical activity behaviours [1].

There is growing interest in sedentary behaviour and related health outcomes in adults [2]. Sedentary behaviour refers to any waking behaviour characterised by an energy expenditure ≤ 1.5 METs while in a sitting or reclining posture [3] (a MET or

Metabolic Equivalent Task is the ratio of the working metabolic rate of an activity divided by the resting metabolic rate [4]). This includes sitting and watching television (TV), along with other forms of screen-based entertainment [5]. Recently, sedentary behaviour was shown to be associated with increased risk of type 2 diabetes [6,7], cardiovascular disease [6,7], metabolic syndrome [8] and all-cause mortality [6,9], independent of habitual physical activity levels. The transition to retirement has been associated with an increase in TV viewing time in only three previous reports [10–12], but there is no evidence for the influence on other sedentary behaviours, such as computer time, reading time or overall sitting time.

In addition to the recognised health benefits of physical activity at all ages, recent studies point to particularly favourable effects in aging subjects. In elderly populations, physical activity was recently shown to be inversely associated with risk of mortality [13], dementia [14], some types of cancer [15], and depression [16]. However, most previous studies have focused on leisure-time physical activity [12,17–21] and only few studies have taken into account domestic and occupational physical activity [20,21].

In a recent systematic review, half of the included studies reported a negative association between physical activity and TV time, no relation was found with computer time, and there was no mention of reading time [22]. Incongruity across study results could reflect differences in methodology as well as in socio-demographic indicators. Considering the protective effect on health of physical activity and the potentially harmful effects of sedentary behaviour [23], defining at-risk populations or stages of life where physical activity and specific sedentary behaviours are related could help fine-tune public health policies targeting both exposures simultaneously. At present, the relationships between sedentary behaviour and physical activity in aging populations remain poorly documented, not permitting the assessment of the association before, during and after retirement [22].

The objectives of the present study, which used a longitudinal design in a sample of middle-aged French adults, were 1) to describe the 6-year changes in different types of sedentary behaviours and different domains of physical activity according to retirement status, and 2) to investigate the relationships between changes in sedentary behaviours and changes in physical activity according to retirement status.

Methods

Ethics statement

Subjects provided written informed consent to the study which was conducted according to guidelines laid down in the Declaration of Helsinki and was approved by the Ethics Committee for Studies with Human Subjects at Paris-Cochin Hospital (CCPPRB n° 706 and n° 2364, respectively) and the

Commission Nationale de l'Informatique et des Libertés (CNIL n° 334641 and n° 907094, respectively).

Subjects

Subjects were participants in the “Supplémentation en Vitamines et Minéraux AntioXydants” (SU.VI.MAX) cohort. The design, methods and rationale of the SU.VI.MAX study have been described elsewhere [24]. It was initially designed as a randomised, double-blind, placebo-controlled primary prevention trial to test the efficacy of daily supplementation with antioxidant vitamins and minerals at nutritional doses in reducing the incidence of ischaemic heart disease, cancer and overall mortality [25]. Following a 5-month national multimedia campaign that included TV, radio and newspapers, volunteer subjects, not selected for any specific risk factors, were included in 1994–1995 for a planned follow-up of 8 years (men: 45–60 y, women: 35–60 y). Each subject underwent a yearly visit alternating between a clinical examination and biological sampling. From the full SU.VI.MAX cohort (N = 12,741), a total of 6,850 subjects who had agreed to participate in a post-supplementation observational follow-up were recruited for the SU.VI.MAX 2 study (2007–2009).

Physical activity and sedentary behaviours

Physical activity and sedentary behaviours were assessed in 2001 and 2007 using a French self-administered version [26] of the Modifiable Activity Questionnaire (MAQ) [27]. This instrument assesses past 12-month physical activity in various domains of everyday life. Physical activity assessment using the MAQ has been validated against energy expenditure measurements using the double-labelled water technique, and the test-retest properties of the questionnaire have been shown [28]. The questionnaire has been described in detail elsewhere [27]. Briefly, subjects were asked to report all leisure-time physical activity performed at least 10 times for 10 min per session over the past 12 months. Detailed information was collected concerning the type of leisure-time activity (walking, cycling, swimming, gardening, etc.). The frequency and duration of each activity was reported. Leisure-time activities were classified according to their intensity, based on their estimated metabolic cost, as moderate (3–6 METs) and

Table 1. Characteristics of the study population according to retirement status.

	Not retired in 2001 and 2007	Retirement between 2001 and 2007	Retired in 2001 and 2007	p
N	1126	891	824	
Sex (% men)	38.3	55.6	64.0	<0.001
Age (years)	53.1 (3.2)	57.1 (3.6)	62.3 (3.3)	<0.001
BMI (kg/m ²)	24.1 (3.5)	24.9 (3.5)	25.4 (3.4)	<0.001
<i>Educational level</i>				
Primary school	14.5	17.3	22.9	<0.001
High school	33.4	39.7	46.6	
University or equivalent	52.1	43	30.4	
<i>Smoking status</i>				
Never smoker	50.9	50.8	49.3	0.075
Former smoker	37.6	37.3	42.1	
Current smoker	11.5	11.9	8.6	

Values are mean (SD) or %.

p values are from ANOVA or Chi-square tests.

doi:10.1371/journal.pone.0106850.t001

Table 2. Sedentary behaviours and physical activity at baseline (2001) and changes during 6-year follow-up.

	Baseline				Change			
	Not retired in 2001 and 2007		Retirement between 2001 and 2007		Not retired in 2001 and 2007 [†]		Retirement between 2001 and 2007 ^{††}	
	n	1126	891	824	n	1126	891	824
<i>Sedentary behaviours (h/week)</i>								
Total leisure sedentary behaviour	21.8 (10.7)	23.3 (11.4)	28.9 (12.1)	<0.001 ^{b,c}	4.7 (0.35)	8.4 (0.42)	4.2 (0.41)	<0.001 ^{ab}
Television viewing during leisure	12.8 (8.2)	13.9 (8.0)	17.9 (9.2)	<0.001 ^{ab,c}	1.5 (0.19)	3.0 (0.24)	0.9 (0.27)	<0.001 ^{ab}
Computer use during leisure	2.1 (4.0)	2.3 (4.5)	2.6 (5.0)	0.047 ^c	2.5 (0.17)	4.1 (0.24)	2.8 (0.21)	<0.001 ^{ab}
Reading during leisure	7.1 (5)	7.1 (5.3)	8.5 (5.6)	0.017 ^{b,c}	0.6 (0.19)	1.3 (0.22)	0.4 (0.20)	0.006 ^b
Occupational sitting	16.9 (13.3)	14.2 (13.0)	-	<0.001 ^a	-1.7 (0.31)	-14.2 (0.45)	-	<0.001 ^a
Domestic sitting	1.8 (4.9)	2.3 (8.0)	4.8 (9.9)	<0.001 ^{b,c}	0.3 (0.26)	1.81 (0.42)	-1.1 (0.53)	<0.001 ^{b,c}
<i>Physical activity (h/week)</i>								
Total leisure	3.4 (3.9)	3.6 (4.2)	6.7 (6.6)	<0.001 ^{b,c}	0.7 (0.11)	2.5 (0.18)	-0.6 (0.23)	<0.001 ^{ab,c}
Moderate leisure	2.7 (3.6)	3.0 (3.9)	5.7 (6.3)	<0.001 ^{b,c}	0.3 (0.11)	1.7 (0.17)	-0.7 (0.22)	<0.001 ^{ab,c}
Vigorous leisure	0.6 (1.5)	0.6 (1.4)	0.9 (2.1)	0.045 ^{b,c}	0.1 (0.06)	0.4 (0.06)	-0.1 (0.07)	<0.001 ^{ab}
Walking	0.8 (1.5)	0.9 (1.7)	1.6 (2.2)	<0.001 ^{b,c}	0.3 (0.08)	0.9 (0.09)	0.1 (0.09)	<0.001 ^{ab}
Gardening	0.7 (1.7)	0.9 (1.9)	2.0 (3.7)	<0.001 ^{b,c}	0.2 (0.05)	0.7 (0.09)	-0.1 (0.13)	<0.001 ^{ab}
Swimming	0.1 (0.3)	0.1 (0.3)	0.1 (0.4)	0.50	0.006 (0.01)	0.04 (0.01)	-0.03 (0.01)	0.002 ^b
Biking	0.2 (0.6)	0.2 (0.7)	0.3 (1.1)	0.043 ^{b,c}	0.04 (0.02)	0.1 (0.05)	-0.05 (0.03)	0.01 ^b
Occupational	14.1 (13.5)	14.1 (13.3)	-	0.78	-3.4 (0.56)	-14.1 (0.46)	-	<0.001 ^a
Domestic	7.6 (8.5)	6.6 (7.3)	9.1 (10.0)	<0.001 ^{b,c}	-0.7 (0.33)	3.6 (0.49)	-0.1 (0.67)	<0.001 ^{ab}

Values are mean (SD) for baseline and mean (SE) for changes. p values are from ANOVA with adjustment on sex and age.

^aComparisons between subjects not retired in 2001 and 2007 and subjects who retired between 2001 and 2007 are significantly different (p<0.05).

^bComparisons between subjects who retired between 2001 and 2007 and subjects retired in 2001 and 2007 are significantly different (p<0.05).

^cComparisons between subjects not retired in 2001 and 2007 and subjects retired in 2001 and 2007 are significantly different (p<0.05).

[†]Except for vigorous leisure time physical activity and swimming, all values for changes are significantly different from 0.

^{††}All values for changes are significantly different from 0.

^{†††}Except for reading, domestic physical activity, vigorous leisure physical activity, walking, gardening and biking, all values for changes are significantly different from 0.

doi:10.1371/journal.pone.0106850.t002

Table 3. Relations between changes in time spent in sedentary behaviours (dependent variables) and changes in physical activity (exposure variables) during the 6-year follow-up for subjects working in 2001 and 2007.

Physical activity (h/week)	Television viewing during leisure		Computer use during leisure		Reading during leisure		Occupational sitting		Domestic sitting	
	β	P	β	P	β	P	β	P	β	P
Total leisure	-0.04	0.02	0.01	0.50	-0.01	0.66	-0.04	0.52	-0.11	0.22
Moderate leisure	-0.04	0.02	0.02	0.31	0.03	0.23	-0.07	0.27	-0.12	0.18
Vigorous leisure	-0.04	0.10	0.01	0.69	-0.03	0.39	-0.10	0.38	-0.19	0.31
Occupational	0.02	0.04	0.002	0.82	0.02	0.13	-0.03	0.53	-0.01	0.87
Domestic	0.08	0.02	0.01	0.7	-0.01	0.84	0.20	0.001	0.05	0.046

Beta coefficients are from linear regression analyses with each change in sedentary behaviour as outcome variable. Models were adjusted for age, sex, educational level, smoking status, physical activity at baseline and baseline values of the respective outcome variable.
doi:10.1371/journal.pone.0106850.t003

vigorous (>6 METs) activity. Thus, physical activity variables used in the analyses included moderate and vigorous leisure-time physical activity, selected physical activities (e.g. walking, gardening, etc.), and the sum of the overall time spent in leisure-time physical activity.

The questionnaire included items about the time usually spent (in hours and minutes per day) at home watching TV/video, using a computer or playing video games, or reading for leisure. Time spent in each leisure sedentary occupation and the sum of all time periods spent in these leisure sedentary occupations were used in the analyses.

Assessment of occupational physical activity was based on the number of hours during which an individual participated in physically demanding activities during an average work day, for each job held over the past year. Occupational sedentary behaviour was based on the number of hours sitting during an average work day, for each job held over the past year. The original version of the MAQ only explored physical activity during leisure-time and at work. To also take into account physical activity unrelated to occupation or leisure, subjects were separately asked to report any domestic physical activity which included activities related to housekeeping, use of active transportation to run errands, etc. that had been performed at least 10 times for 10 min each session over the past 12 months. These questions were asked in the same format as in the occupational section of the questionnaire, including sitting.

Sociodemographic covariables

Retirement status was assessed by self-report during the 2001 (baseline for the present study) and 2007 (follow-up) visits. The population was divided into 3 categories, according to their baseline and follow-up retirement status: 1) subjects who were not retired in 2001 and 2007, 2) subjects who retired between 2001 and 2007, and 3) subjects who were retired in 2001 and 2007. Sex, date of birth and educational level were assessed at entry using a self-administered questionnaire. Level of education was coded into three categories according to the highest certification obtained (primary school, high school, university or equivalent). Smoking status (never, former, current) was assessed in September 1998 by a separate questionnaire sent to the entire cohort. Height and weight were measured during the 2001 and 2007 visits. BMI was calculated as body weight (in kilograms) divided by the squared height (kg/m²).

Statistical analyses

For each subject, changes between 2001 and 2007 in indicators of sedentary behaviour and physical activity were computed as the value recorded in 2007 minus the value recorded in 2001. Continuous variables were summarized by calculating the mean \pm standard deviation (SD). We used one-way ANOVA to assess differences in continuous variables across retirement groups, with a Bonferroni correction applied for multiple comparisons. T-tests were used to assess post-hoc differences between retirement groups and to assess changes in the continuous variables during follow-up. We used chi-square tests to assess differences in categorical variables across retirement groups. Associations between changes in sedentary behaviours and changes in physical activity according to retirement status evolution between 2001 and 2007 were assessed using a multivariate generalised linear model. Covariates included sex, age in 2001, educational level, smoking status and occupational physical activity at baseline when appropriate. For all analyses, the significance level was set at 0.05 and all tests were two-sided. All statistical analyses were performed using SAS software (version 9.3, SAS Institute Inc., Cary, NC, USA).

Results

Comparisons between subjects included and not included in the analyses

Among subjects initially included in the SU.VI.MAX study, we focused the present analyses on subjects with available data from the physical activity questionnaires both in 2001 and 2007 ($n = 3,458$ subjects available), aged 45 years or older at entry into the study (in order to have a similar age range in both genders) ($n = 3,006$ subjects). In addition, subjects who reported being confined to bed more than 4 weeks in the past year before completing the questionnaires were excluded ($n = 165$), thus obtaining a final sample of $n = 2,841$ (1,453 men and 1,388 women).

Compared to subjects examined in 2007 (SU.VI.MAX 2 study) but not included in the present analyses, our study population comprised more men (51.1 vs. 36.2%, $p < 0.001$), more subjects with a university education level (42.9 vs. 39.5%, $p < 0.001$), and they had a lower mean BMI (24.3 kg/m^2 vs. 24.5 kg/m^2 , $p = 0.03$), and were slightly older at baseline (57.0 y vs. 55.2 y, $p < 0.001$).

Baseline characteristics of the study population

Twenty nine percent of subjects were not retired in 2001 and 2007, 31.4% were retired between 2001 and 2007, and 39.6% of the sample were retired in 2001 and 2007 (Table 1). Subjects retired in 2001 and 2007 were mostly men, older and less educated than their employed counterparts (Table 1). At baseline, subjects retired in 2001 and 2007 had higher total levels of leisure-time sedentary behaviour, spent more time watching TV and reading, and spent twice the time in overall leisure-time physical activity (Table 2) compared to other two groups. These subjects also engaged in twice as much walking and gardening.

Six-year changes in sedentary behaviours and physical activity

Between 2001 and 2007, total leisure-time sedentary behaviour, TV viewing and computer use increased in all three groups ($p < 0.001$ in each group for the comparison between follow-up and baseline levels) (Table 2). For subjects retired between 2001 and 2007, there was an increase in both leisure-time and domestic physical activity (all $p < 0.01$, data not shown). For subjects retired in 2001 and 2007, time spent in total leisure-time physical activity and moderate-intensity leisure-time physical activity decreased ($p < 0.005$, data not shown).

Between 2001 and 2007, total leisure-time sedentary behaviour, TV viewing and computer use increased to a larger extent in subjects retired between 2001 and 2007 compared to the two other groups (Table 2). Leisure-time physical activities, except for swimming and biking also increased more in subjects retired between 2001 and 2007 compared to the two other groups (Table 2). When comparing subjects not retired in 2001 and 2007 and subjects retired in 2001 and 2007, there was no significant difference for changes in sedentary behaviours during leisure. In contrast, changes in total and moderate-intensity leisure-time physical activities were significantly higher for subjects not retired in 2001 and 2007 compared to subjects retired in 2001 and 2007 (Table 2).

The associations of retirement with sedentary behaviours and physical activity shown in Table 2 are consistent with models adjusted for age, sex, educational level, smoking status, and initial sedentary and physical activity behaviour levels (data shown in Table S1).

Associations between changes in sedentary behaviour and changes in physical activity

In subjects not retired in 2001 and 2007, changes in time spent TV viewing were positively related to changes in occupational physical activity ($p = 0.04$) (Table 3). In the same group, changes in time spent in total and moderate-intensity leisure-time physical activity was negatively associated with changes in time spent watching TV ($p = 0.02$ and $p = 0.02$ respectively). No association was found for subjects retired between 2001 and 2007 or for subjects retired in 2001 and 2007 (data not shown).

Discussion

In this study in French subjects, we studied the 6-year changes in different types of sedentary behaviour and different domains of physical activity according to retirement status, and we investigated the relationships between changes in sedentary behaviours and changes in physical activity according to retirement status. All subjects increased their sedentary behaviour during follow-up, but those retired between 2001 and 2007 showed the greatest changes. They also had the greatest increase in time spent in leisure-time physical activity. Subjects retired in 2001 and 2007 were those who spent the most time in sedentary behaviour, leisure-time physical activity and domestic leisure-time physical activity. Finally, leisure-time physical activity was inversely associated with time spent TV viewing in subjects not retired in 2001 and 2007. These findings extend those of our previous report documenting changes in sedentary behaviour and physical activity over three years in subjects from the same population [12].

In subjects retired between 2001 and 2007, it was noticeable that the mean increase in total sedentary behaviour was about three times higher (+8.4 h/week) than the mean increase in leisure-time physical activity (+2.5 h/week). The greatest change in sedentary behaviour during leisure-time was for computer use compared to reading and TV viewing. This behavioural change appears in line with the rapidly expanding use of computers and the Internet across different population groups including aging subjects, during these same years (2001–2007) [29]. It underscores the importance of assessing different dimensions of sedentary behaviours and not only TV viewing, known to increase through retirement [11,12].

In addition to sedentary behaviours, this study is one of the few that performed a detailed assessment of changes in different types of leisure-time physical activities in subjects transitioning to retirement [10]. In our subjects, the increase in moderate-intensity activity (such as walking and gardening) accounted on average for about two thirds of the increase in total leisure-time physical activity. This is in agreement with results from previous studies where transition to retirement was associated with an increased sports participation [11], with increased total leisure-time physical activity score [10,18] and with an increased proportion of subjects meeting physical activity recommendations [17].

There are several potential explanations for the increase in leisure-time physical activity with retirement. First, retirement leads to a decrease in time constraints possibly corresponding to more free-time available. Second, individuals likely become increasingly health-conscious in the context of aging, which could increase motivation to engage in more recreational physical activity. Finally, increased leisure-time physical activity after retirement may provide a new daily routine and new opportunities for social interactions [30].

For subjects not retired in 2001 and 2007 the changes in time spent watching TV during leisure-time were differentially associated with changes in time spent being active at work (positive

association) and time spent being active during leisure (negative association). The positive association with changes in active occupational time may intuitively be interpreted as compensatory behaviour and the need for rest outside work. The negative association with changes in active leisure-time might reflect choices made in terms of budget time in this working population. Although there is no previous study reporting such associations in a longitudinal design, negative cross-sectional associations were shown between TV time and leisure-time physical activity [31,32]. The lack of significant results for subjects retired between 2001 and 2007 and those retired in 2001 and 2007 can be interpreted as subjects having more availability after retirement and consequently not to make choices between leisure-time sedentary behaviour and physical activity.

Some limitations of this study need to be mentioned. First, measurements of physical activity and sedentary behaviours were derived from self-reporting, which can cause misclassification bias mostly because of over-reporting of physical activity [33]. However, there is no reason to expect that retirement status would have repercussions on that misclassification. In addition, the applicability of the MAQ to older subjects may be discussed. Regarding reliability, in the first description of the MAQ by Kriska et al. [27], 1–3 week test-retest correlations for past-year leisure-time physical activity were found to be of about the same magnitude in the older compared to the younger adult subjects (37–59 y, $\rho = 0.88$ and 21–36 y, $\rho = 0.92$, respectively). Second, we had information about the employment status (retired or not) at the time of data collection (2001 and 2007) but the actual date of retirement amongst these subjects and the reasons for retirement were not available. A possible selection bias could occur if some participants had retired earlier because of poor health, which might have negatively impacted their physical activity. To prevent, at least in part this limitation, subjects who reported being confined to bed more than 4 weeks in the past year before completing the questionnaires were excluded from the analyses. In addition, smoking habits were self-reported two years before the first wave of physical activity assessment, leading to a potential misclassification bias regarding subjects who had quit smoking between the two assessment periods. Third, despite the retirement age in our population (58–64 y) being consistent with the retirement age in France at the time of the study, our subjects were volunteers participating in a nutritional intervention study.

References

- Ekerdt DJ (2010). Frontiers of research on work and retirement. *J Gerontol B Psychol Sci Soc Sci* 65B(1): 69–80. doi:10.1093/geronb/gbp109.
- Thorp AA, Owen N, Neuhaus M, Dunstan DW (2011) Sedentary behaviors and subsequent health outcomes in adults: a systematic review of longitudinal studies, 1996–2011. *Am J Prev Med* 41(2): 207–15. doi:10.1016/j.amepre.2011.05.004.
- Sedentary Behaviour Research Network (2012) Letter to the editor: standardized use of the terms “sedentary” and “sedentary behaviours”. *Appl Physiol Nutr Metab* 37(3): 540–542. doi:10.1139/h2012-024.
- Ainsworth BE, Haskell WL, Whitt MC, Irwin ML, Swartz AM, et al. (2000) Compendium of physical activities: an update of activity codes and MET intensities. *Medicine and science in sports and exercise* 32(9 Suppl): S498–504.
- Pate RR, O'Neill JR, Lobelo F (2008) The evolving definition of “sedentary”. *Exerc Sport Sci Rev* 36(4): 173–8. doi:10.1097/JES.0b013e3181877d1a.
- Grontved A, Hu FB (2011) Television viewing and risk of type 2 diabetes, cardiovascular disease, and all-cause mortality: a meta-analysis. *JAMA* 305(23): 2448–55. doi:10.1001/jama.2011.812.
- Pinto Pereira SM, Ki M, Power C (2012) Sedentary behaviour and biomarkers for cardiovascular disease and diabetes in mid-life: the role of television-viewing and sitting at work. *PLoS one* 7(2): e31132. doi:10.1371/journal.pone.0031132.
- Edwardson CL, Gorely T, Davies MJ, Gray IJ, Khunti K, et al. (2012) Association of sedentary behaviour with metabolic syndrome: a meta-analysis. *PLoS one* 7(4): e34916. doi:10.1371/journal.pone.0034916.
- Koster A, Caserotti P, Patel KV, Matthews CE, Berrigan D, et al. (2012) Association of sedentary time with mortality independent of moderate to

vigorous physical activity. *PLoS one* 7(6): e37696. doi:10.1371/journal.pone.0037696.

Conclusions

Therefore, they generally had a higher educational levels and occupational status, along with a healthier lifestyle than the general population [25]. For these reasons extrapolation of these findings must be done cautiously.

In subjects transitioning to retirement, both sedentary behaviour and physical activity during leisure-time increased substantially over the 6-year follow-up period. However, the mean increase in sedentary behaviour during leisure-time was about three times higher than the mean increase in leisure-time physical activity, with the greatest change seen in computer use compared to reading and TV viewing. This may be of serious concern considering the growing evidence of the deleterious health effects of sedentary behaviours. This emphasizes the need to test interventions and to reinforce public health measures aimed at decreasing sedentary behaviours and increasing regular physical activity during the critical retirement period.

Supporting Information

Table S1 Relations between changes in time spent in sedentary behaviours or changes in physical activity with retirement status during the 6-year follow-up (linear regression analysis).
(DOC)

Acknowledgments

This work is also part of the ACTI-Cités (Activité physique, transport actif et environnement de vie urbain) project.

The authors thank Younes Esseddik, Paul Flanzly, Mohand Ait Oufella, Yasmina Chelghoum, and Than Duong Van (computer scientists), Florence Charpentier (dietitian), Nathalie Arnault, Véronique Gourlet, Fabien Szabo, Laurent Bourhis, and Stephen Besseau (statisticians), and Rachida Mehroug (logistics assistant) for their technical contribution to the SU.VI.MAX study. We also thank the participants in this cohort.

Author Contributions

Conceived and designed the experiments: SH EKG. Performed the experiments: SH EKG. Analyzed the data: MM. Contributed to the writing of the manuscript: MM LF HC EKG MT CS CW SH JMO VAA.

- Barnett I, van Sluijs E, Ogilvie D, Wareham NJ (2013) Changes in household, transport and recreational physical activity and television viewing time across the transition to retirement: longitudinal evidence from the EPIC-Norfolk cohort. *J Epidemiol Community Health* doi:10.1136/jech-2013-203225.
- Evenston KR, Rosamond WD, Cai J, Diez-Roux AV, Brancati FL (2002) Influence of retirement on leisure-time physical activity: the atherosclerosis risk in communities study. *Am J Epidemiol* 155(8): 692–9.
- Touvier M, Bertrais S, Charreire H, Vergnaud AC, Hercberg S, et al. (2010) Changes in leisure-time physical activity and sedentary behaviour at retirement: a prospective study in middle-aged French subjects. *Int J Behav Nutr Phys Act* 7: 14. doi:10.1186/1479-5868-7-14.
- Samitz G, Egger M, Zwahlen M (2011) Domains of physical activity and all-cause mortality: a systematic review and dose-response meta-analysis of cohort studies. *Int J Epidemiol* 40(5): 1382–400. doi:10.1093/ije/dyr112.
- Barnes DE, Whitmer RA, Yaffe K (2007) Physical activity and dementia: The need for prevention trials. *Exerc Sport Sci Rev* 35(1): 24–9. doi:10.1097/JES.0b013e31802d6bc2.
- Lemane D, Cassileth B, Gubili J (2013) The role of physical activity in cancer prevention, treatment, recovery, and survivorship. *Oncology (Williston Park, N.Y.)* 27(6): 580–5.
- Azevedo Da Silva M, Singh-Manoux A, Brunner EJ, Kaffashian S, Shipley MJ, et al. (2012) Bidirectional association between physical activity and symptoms of

- anxiety and depression: the Whitehall II study. *Eur J Epidemiol* 27(7): 537–46. doi:10.1007/s10654-012-9692-8.
17. Berger U, Der G, Mutric N, Hannah M (2005) The impact of retirement on physical activity. *Ageing Soc* 25(2): 181–95.
 18. Brown WJ, Heesch KC, Miller YD (2009) Life events and changing physical activity patterns in women at different life stages. *Ann Behav Med* 37(3): 294–305. doi:10.1007/s12160-009-9099-2.
 19. Glamsr FD, Hayslip BJ (1985) The impact of retirement on participation in leisure activities. *Ther Recreation J* 19(3): 28–38.
 20. Mein GK, Shipley MJ, Hillsdon M, Ellison GT, Marmot MG (2005) Work, retirement and physical activity: cross-sectional analyses from the Whitehall II study. *Eur J Public Health* 15(3): 317–22. doi:10.1093/eurpub/cki087.
 21. Slingerland AS, van Lenthe FJ, Jukema JW, Kamphuis CB, Looman C, et al. (2007) Aging, retirement, and changes in physical activity: prospective cohort findings from the GLOBE study. *Am J Epidemiol* 165(12): 1356–63. doi:10.1093/aje/kwm053.
 22. Rhodes RE, Mark RS, Temmel CP (2012) Adult sedentary behavior: a systematic review. *Am J Prev Med* 42(3): e3–28. doi:10.1016/j.amepre.2011.10.020.
 23. Hamilton MT, Hamilton DG, Zderic TW (2007) Role of low energy expenditure and sitting in obesity, metabolic syndrome, type 2 diabetes, and cardiovascular disease. *Diabetes* 56(11): 2655–67. doi:10.2337/db07-0882.
 24. Hercberg S, Preziosi P, Briancon S, Galan P, Triol I, et al. (1998) A primary prevention trial using nutritional doses of antioxidant vitamins and minerals in cardiovascular diseases and cancers in a general population: the SU.VI.MAX study—design, methods, and participant characteristics. *Supplementation en Vitamines et Mineraux Antioxydants. Control Clin Trials* 19(4): 336–51.
 25. Hercberg S, Galan P, Preziosi P, Bertrais S, Mennen L, et al. (2004) The SU.VI.MAX Study: a randomized, placebo-controlled trial of the health effects of antioxidant vitamins and minerals. *Arch Intern Med* 164(21): 2335–42. doi:10.1001/archinte.164.21.2335.
 26. Vuillemin A, Oppert JM, Guillemin F, Essermeant L, Fontvieille AM, et al. (2000) Self-administered questionnaire compared with interview to assess past-year physical activity. *Med Sci Sports Exerc* 32(6): 1119–24.
 27. Kriska AM, Knowler WC, LaPorte RE, Drash AL, Wing RR, et al. (1990) Development of questionnaire to examine relationship of physical activity and diabetes in Pima Indians. *Diabetes Care* 13(4): 401–11.
 28. Pereira MA, FitzerGerald SJ, Gregg EW, Joswiak ML, Ryan WJ, et al. (1997) A collection of Physical Activity Questionnaires for health-related research. *Med Sci Sports Exerc* 29(6 Suppl): S1–205.
 29. Bigot R, Crouette P, Daudey E (2013) La diffusion des technologies de l'information et de la communication dans la société française. Centre de Recherche pour l'ÉtuDe et l'Observation des Conditions de vie (CREDOC) <http://www.credoc.fr/pdf/Rapp/R297.pdf>.
 30. Barnett I, Guell C, Ogilvie D (2012) The experience of physical activity and the transition to retirement: a systematic review and integrative synthesis of qualitative and quantitative evidence. *Int J Behav Nutr Phys Act* 9: 97. doi:10.1186/1479-5868-9-97.
 31. Kaleta D, Jegier A (2007) Predictors of inactivity in the working-age population. *Int J Occup Med Environ Health* 20(2): 175–82. doi:10.2478/v10001-007-0019-z.
 32. Sugiyama T, Salmon J, Dunstan DW, Bauman AE, Owen N (2007) Neighborhood walkability and TV viewing time among Australian adults. *Am J Prev Med* 33(6): 444–9. doi:10.1016/j.amepre.2007.07.035.
 33. Shephard RJ (2003) Limits to the measurement of habitual physical activity by questionnaires. *Br J Sports Med* 37(3): 197–206; discussion.