



Rheological implications of extensional detachments: Mediterranean and numerical insights

Loic Labrousse, Benjamin Huet, Laetitia Le Pourhiet, Laurent Jolivet,
Evgenii Burov

► To cite this version:

Loic Labrousse, Benjamin Huet, Laetitia Le Pourhiet, Laurent Jolivet, Evgenii Burov. Rheological implications of extensional detachments: Mediterranean and numerical insights. *Earth-Science Reviews*, 2016, 161, pp.233-258. 10.1016/j.earscirev.2016.09.003 . hal-01371910

HAL Id: hal-01371910

<https://hal.sorbonne-universite.fr/hal-01371910>

Submitted on 26 Sep 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rheological implications of extensional detachments : Mediterranean and numerical insights.

Loïc Labrousse (1,2), Benjamin Huet (3), Laetitia Le Pourhiet (1,2), Laurent Jolivet (4) and Evgenii Burov (1,2).

1- Sorbonne Universités, UPMC Univ Paris 06, UMR 7193, Institut des Sciences de la Terre Paris (iSTeP), F-75005 Paris, France

2- CNRS, UMR 7193, Institut des Sciences de la Terre Paris (iSTeP), F-75005 Paris, France

3- Department of Geodynamics and Sedimentology U.Vienna Althanstrasse 14/2A273 A-1090 VIENNA

4- ISTO, UMR 6113 - CNRS/U. Orléans, 1A, rue de la Férollerie F-45071 ORLEANS

Abstract

The Mediterranean realm shaped by extensive back-arc extension after multiple collisions between Europe and isolated continental blocks is the second densest occurrence of Metamorphic Core Complexes (MCC) after the North American Cordillera. The present review aims at determining the factors controlling detachment system development during post-orogenic extension in the continental lithosphere. 26 different detachment zones over 23 different localities are systematically described and show that MCC structures can be positioned in a three end-member classification. Beside the high-temperature lower plate end-member, most of the time considered as the archetypical MCC type, two cold end-members can be identified: a first one preserving high pressure metasedimentary units in the vicinity of the detachment and represented by numerous cases in the Mediterranean and a second theoretical one with strain localized at the bottom of a resistant upper crust. These two end-members show a strain

pattern guided by inheritance more than thermal state. The relevance of this three end-member classification is then tested via numerical modelling of extension of layered continental lithosphere. The compilation of our results and recently published similar models show that over a 100 of different simulations the most critical parameter for developing a MCC is the intra-crustal strength contrast (S_{\max}/S_{\min} , where S_{\max} and S_{\min} stand respectively for the maximum – at the brittle-ductile transition – and minimum – at the Moho – crustal strength). This contrast has to be risen only by 1 order of magnitude to change extension mode. Below a critical value of 1000, narrow and wide rifts develop. MCCs with a perennial detachment are produced for values slightly higher than 1000, while higher intra-crustal strength contrasts systematically promote double-domes MCCs, with strain progressively relocalizing within the lower plate. Inherited layering is as efficient as temperature for raising this intra-crustal strength contrast while strain rate is actually a second order parameter. Metamorphic reactions are also to be considered as a first order process in localizing detachments.

1. Introduction

Even if the seminal comprehensive descriptions of Metamorphic Core Complexes (MCCs) in the American Cordillera mentioned lower plates constituted of gneiss and intruded by granites (Snake Range, Miller et al. (1983), Whipple Mountains, Davis et al. (1986), the actual definition of MCCs: « *Cordilleran metamorphic core complexes appear to be bodies from the middle crust that have been dragged out from beneath fracturing and extending upper crustal rocks, and exposed beneath shallow-dipping (normal slip) faults of large areal extent* » (Lister and Davis, 1989) refers to rocks exhumed from the middle crust whatever their thermal history. The fundamental property of this middle crust resides in its ability to flow laterally toward the forming dome (Block and Royden, 2010), to accommodate stretching of the upper plate and preserve a relatively flat Moho (Figure 1). Even though thermal reequilibration can induce weakening of the lower crust (Buck, 1991) and increases its ability to flow below a relatively strong upper crust, a similar strength profile can also be inherited from pre-extension evolution of the continental crust and promote development of the original structure leading to the formation of MCCs: the detachment (Davis, pers comm.). In order to unravel the rheological meaning of detachments, we propose here a review of extensional shear zones described as post-orogenic detachments in the Mediterranean realm. These examples share a common tectonic setting and often affect comparable lithotectonic successions. A three end-members typology is proposed with high-temperature MCCs (HT end-member) as one end-member, and two cold MCC end-members with a weak middle crust due in most cases to the interlayering of high-pressure metasedimentary nappes during syn-orogenic wedge building (HP end-member). Even if not met in natural cases, a second theoretical cold end-member can be defined by the presence of a strong upper crust also due to nappe stacking or even obduction (HS end-member). In

both cases, the inherited tectonostratigraphy is responsible for a sensible exaggeration of the intra-crustal strength contrast.

Fully coupled thermo-mechanical modeling experiments allow testing these three end-member typologies and determining the critical intra-crustal strength contrast for the perennial development of a detachment zone and building of a dome at the expense of the lower plate. The comparison of natural cases classification and model-based categories demonstrates the mechanical feasibility of all natural combinations and possibly explains how long-lasting multiple detachment systems such as Menderes, Turkey or North Cycladic Detachment System, Cyclades, Greece, evolve.

2. Mediterranean Extensional Detachments

Metamorphic Core Complexes are described in the Mediterranean within three large extended domains (Carpathians and Pannonian basin, Western Mediterranean from Alboran to Tyrrhenian, and Aegean domain *sensu lato*) subject to slab retreat during Alpine orogeny (Jolivet et al., 2009a). Most of them are now located in continental domains with Moho depths between 25 and 20 km, but not systematically in flat Moho areas (Figure 2). Betic Cordilleras and Menderes are, for instance, on top of dipping Moho with about 10 km vertical drop across them (Karabulut et al., 2013 for instance for Menderes). Metamorphic rocks exhumed within Mediterranean core complexes show a wide scatter in pressure-temperature conditions (Figure 3 and Table 1). Dating peak metamorphic conditions and detachment mylonites or relative dating of the detachment activity allow deciphering between rocks mostly exhumed during pre-extensional thickening stages from rocks exhumed during extensional dynamics leading to the MCC

formation. Furthermore, the presence in the footwall of detachments of lower grade rocks structurally below the High Temperature and Low Pressure (HT-LP) or High Pressure and Low Temperature (HP-LT) rocks is also a criterion to attribute the first stages of the exhumation of the latter to syn-orogenic nappe stacking rather than lithospheric extension. Nappe stacking is indeed the sole process prompt to thrust high-grade metamorphic units on top of a preserved lower grade unit. For instance, the equilibration of the Ichendirene HP-granulite bodies (BB1, Figure 3) included in the lower grade Beni Bousera kinzigites (BB2, Figure 3) in Morrocco relates to stages earlier than the exhumation of the massif itself (Michard et al., 2006).

Even when removing HP-LT peaks inherited from pre-extensional stages and preserved during dome formation, the scatter of PT conditions sampled by detachments footwalls remains high. Noteworthy, some core complexes such as the one described in the Catena Costiera, Calabria, Italy (CC on Figure3, Rossetti et al. (2004), or in the Northern Cyclades (Tn, Tinos, Jolivet et al. (2010b) and references therein) show « cold » footwalls, with conditions theoretically incompatible with the development of detachment systems, according to numerical models (Tirel et al., 2008). The rheological meaning of such « cold MCCs » (Huet et al., 2010), is the purpose of our study.

2.1 Tectonostratigraphic positions

The systematic description of tectonostratigraphic positions of Mediterranean detachments allows highlighting the preferred decoupling levels within the continental crust during its extension (Table 1 and 2). Among relevant characteristics, some were considered necessary for defining a detachment zone in our compilation: the presence of

mylonites along the extensional fault system, in order to compare structures initiated below the ductile-brittle transition only, and a clear extensional setting, in order to compare post-orogenic extensional detachments only, and avoid exhumation shear zone (syn-orogenic detachments in Jolivet et al., 2003). Some characteristics differ from a detachment system to another: the lithology and PT conditions recorded by their footwall (patterns and color code on Figure 4), the lithology of their hangingwall, as well as the presence of plutons, with mantle or crustal origin. The presence of migmatites in the footwall has also been systematically noted. The initiation context of the exhumation process has also been checked : syn-orogenic for rocks partly exhumed during convergence, post-orogenic for rocks exhumed only during post-orogenic extension. The systematic documentation of those characteristics (Figure 4 and Table 2) shows a distribution of natural cases with three theoretical end-members, between which they can be positioned in a triangular diagram (Figure 5). The position of each case was deduced from the relative weighing of high temperature criterion (migmatites or high temperature lower plate or syn-kinematic intrusives), low temperature criterion (preservation of HP-LT paragenesis in the footwall) or high strength upper plate criterion listed in table 2. As will be discussed further, mediterranean natural cases plot within and along edges of the proposed ternary plot. While the HT and HP end-embers are represented by numerous natural cases, no case is found on the HS end-member. None of the documented examples indeed has a strong upper plate as single criterion.

2.1.1 Detachments on top of High-Temperature lower plates

Some of the Mediterranean detachments cap high-temperature lower plates, with HT-LP metamorphic rocks showing syn-extensional pervasive amphibolite facies metamorphism, possibly associated to migmatization, and/or syn-kinematic granite intrusion.

The Kabylia Detachment, Greater Kabylia, Algeria (KaD, table 1C), is located between Permian phyllites in its hanging-wall and amphibolite facies micaschists and orthogneisses in its footwall (Saadallah and Caby, 1996). Migmatitic textures, syn-tectonic aplitic dikes and late peraluminous granite intrusions indicate a temperature above the computed solidus (i. e. 700°C) during the exhumation of paragneiss and micaschists forming most of the lower plate. Rheological contrast between upper and lower plate is mostly due to contrasted thermal histories. The Edough massif (Ed, table 1D) in Algeria shows a similar structure (Caby et al., 2001) on top of various high grade metamorphic rocks reaching anatexis and intruded by syn-kinematic leucogranites. Fragments of sublithospheric mantle were incorporated within crustal stack and now constitute rheologically constrained inclusions (Bosch et al., 2014). Even if exhumation ages in the Edough massif are younger than in Greater Kabylia (Bruguier et al., 2009), both extensional domes actually built from the same crustal stack during Alboran slab retreat or later slab related dynamics.

On Elba island, Italy, even though the cataclastic low angle normal Zuccale fault zone accommodated substantial extension (Collettini and Holdsworth, 2004), the proper detachment zone (Capanne Detachment Zone, Table 1H), exhuming the lower Tuscan unit greenschists, is located at the top of the syntectonic Monte Capanne granodiorite (Daniel and Jolivet, 1995). Mylonites are described in its thermometamorphic aureole and localization of strain is related to its intrusion.

The Lubenik Line (LL, table 1L), in the Central Western Slovakian Carpathians and the Zaouia Fault Zone, in the Beni Bousera massif, Morocco, even though both considered as structures inherited from orogenic stages (Janák et al., 2001; Michard et al., 2006) also belong to the High Temperature detachments class. In the Lubenik Line footwall, the Veporic unit is intruded by the syn-kinematic Rochovce granite and mylonites localized strain at its top (Janák et al., 2001). In the Beni Bousera massif, amphibolite facies metamorphism in the Fillali schists and associated kinzigites indeed indicate temperatures higher than 800°C for their peak conditions, and leucogranite intrusions evidence partial melting of the lower plate (Michard et al., 2006).

HT lower plate detachment class is therefore represented by detachment zones that exhume HT metamorphic rocks, which reached peak P-T conditions just before exhumation or lower plate intruded by plutons, responsible for local thermal weakening and strain localization. This mechanism is discussed in section 4.2.3.

2.1.2 Detachments within interlayered high-pressure metamorphic nappe

Most of the "Cold MCC" (Huet et al., 2010) exhume HP-LT metamorphic rocks in their footwall (Figure 3). Even though most of these HP-LT rocks were actually partly exhumed prior to the proper detachment tectonics related to extension, their preservation from thermal reequilibration during ultimate decompression implies that no significant thermal reequilibration occurred during MCC formation.

Detachment faults described in the Betics, Spain (Mecina-Filabres Shear Zone, MSFZ, table 1A), on Syros, Greece (Vari detachment, VaD, table 1P), in southern Italy (Catena Costiera and Aspromonte, CatCo and Asp, table 1E & F), in Peloponnese, Greece and Crete (Cretan detachments 1 and 2, CrD1 and CrD2, table 1U & V) and also in the Rechnitz window, Austrian Alps (ReW, table 1K) belong to this category.

The Mecina-Filabres Shear Zone develops as mylonites and gouges during extensional reactivation of a former thrust (Platt et al., 1984). Eclogites are preserved within the Nevado-Filabrides lower plate, and most of the syn-tectonic metamorphism is actually greenschist facies (Augier et al., 2005b). No drastic lithological contrast is observed between the upper plate Alpujarrides phyllites, quartzites and graphitic micaschists and the exhumed Bedar-Macael graphitic schists.

The Vari detachment, Syros, exhumes the Cycladic blueschist metasedimentary unit dominated by schists and marble preserving eclogite to blueschist paragenesis, (Trotet et al., 2001), beneath the Vari and "Upper unit" metabasites gneiss and serpentinites which experienced earlier HT reequilibration (Jolivet and Brun, 2008). Even if mainly syn-orogenic, this detachment may have been active in the late stages of post-orogenic extension (Jolivet et al., 2010a).

In Italy, the Catena Costiera detachment is described as a mylonites and cataclasites complex, accounting for the post-orogenic exhumation of the blueschist-bearing metaclastics and marbles associated to the Lower Ophiolitic Unit (LOU, Jolivet and Brun, 2008; Rossetti et al., 2004) beneath the seemingly Upper

Ophiolite Unit, composed of an ophiolitic melange and carbonates (Bonardi et al., 2001). The Aspromonte detachment is a mylonite zone located on top of the Alpine blueschist Aspromonte unit, considered as a syn-to immediate post-orogenic structure (Heymes et al., 2010). Aspromonte and Stilo units on both sides of the detachment are both composed of para and orthogneiss.

Cretan detachments, described on top of the blueschist facies Phyllites-Quartzites nappe in northern Peloponnese (Jolivet et al., 2009b) and in Crete (Kiliass et al., 1994) are represented by extensional mylonites and high-strain phyllites along the reactivated thrust at the base of the Gavrovo-Tripolitza nappe in Crete, or the underlying Tyros Beds in Peloponnese.

In the Rechnitz window group, the detachment zone is composed of serpentinites and quartz-micas mylonites (Cao et al., 2013) at the base of the Austro-Alpine greenschist facies Wechsel unit, and at the top of the blueschist-bearing Rechnitz Penninic unit. A limited HT imprint is recorded during the inferred post-orogenic reactivation of the Austro-Alpine basal thrust (Hoinkes et al., 1999).

All the detachment detailed above are therefore located within a tectonostratigraphic sequence with no significant systematic strength contrast, upper and lower plate being most of the time similar in lithologies, except the HP paragenesis partly preserved in lower plates. Most of the time, extensional strain localized on a former thrust. The reason why such thrusts are prone to extensional reactivation will be discussed further.

2.1.3 Detachments at the base of a strong upper crustal unit

Two close classes of detachment emerge from this review: some localize at the base of a strong upper plate and on top of a HP preserving cover unit and some at the base of a strong upper plate on top of a HT lower plate.

In Nigde massif, Turkey (NiD, table 1W) the amphibolite facies Gumusler formation migmatites (Whitney and Dilek, 1997) are exhumed beneath a massive ophiolite mostly composed of greenschist-facies gabbros (Goncuoglu et al., 1991). Mylonites and cataclasites developed prior to the Uçkapili leucogranite intrusion (Gautier et al., 2002) during an early extension stage affecting the Central Anatolian continental block.

214 In southern and eastern Rhodope (Bulgaria-Greece), the Kerdylion detachment and the Tokachka and
215 Kechros detachment system (KerD and ToKeD, table 1M & N) developed at the base of high-grade rock
216 units: the Vertiskos Gneiss complex (Brun and Sokoutis, 2007) and the Kimi complex garnet-bearing
217 gneiss (Bonev and Beccaletto, 2007) respectively. They exhum the Southern Rhodope Core Complex, the
218 Kesebir-Kardamos and Kechros complexes, all of them being represented by migmatites (Bonev et al.,
219 2006) and amphibolite facies orthogneiss intruded by syn-kinematic granitoids (Vrondou and Symvolon
220 granodiorites in Southern Rhodope, and the Papikjon granitoid in Eastern Rhodope). The base of upper
221 plates is underlined by resistant lithologies such as the Therma and Volvi gabbros and basalts in Southern
222 Rhodope or meta-ophiolitic lenses in Eastern Rhodope (Burg et al., 1996).

223 The Motajica Detachment, Croatia-Bosnia Hercegovina (MoD, table 1, Ustaszewski et al. (2010) is located
224 at the base of the Tisza-Dacia unit ophiolite and at the top of an amphibolite facies meta-accretionary
225 wedge complex intruded by the Motajica granite a few Myr prior to extension initiation (Pamić et al.,
226 2012).

227 In Kazdag, Turkey, extension localized along the northern Alakeçi and southern Selale detachments
228 (AlSeD, table 1S). Mylonites, metaserpentinites and breccias separate the upper plate Cetmi ophiolitic
229 melange from an amphibolite facies basement unit (Beccaletto and Steiner, 2005; Okay and Satir, 2000)
230 topped with migmatites. The Cetmi, Karakaya and Denizgoren ophiolitic units all rest at the base of the
231 upper plate, suggesting a possible reactivated obduction thrust as a precursor for the extensional
232 detachments.

233 These detachment therefore superimpose resistant lithologies on top of HT metamorphic rocks, some
234 other superimpose the same type of resistant units on top of weaker HP-bearing cover units as reviewed
235 in the following.

236 In the Alpi Apuane, northern Italy, the Calcare Cavernoso mylonites (CalCa, table 1G, (Carmignani and
237 Kligfield, 1990) are found between the upper Tuscan nappes made out of massive carbonates at the base
238 and the subjacent blueschist-facies phyllites and metasandstones of the Massa and Autochthonous units.
239 Localization of strain is due to intense cataclasis of anhydrite and dolostone along a reactivated thrust.
240 The Liguride ophiolite on top of the whole lithotectonic pile has been preserved from most of the
241 extensional tectonics.

In Corsica, the Tenda Shear Zone and the Balagne-Nebbio unit basal detachment constitute a multi-level detachment system (Jolivet et al., 1990; Marroni and Pandolfi, 2003) related to post-orogenic extension. In the Cap Corse antiform, the non-metamorphic ophiolitic Balagne nappe directly rests on the Schistes Lustrés calcschists that preserved eclogite to blueschist facies assemblages. On the eastern edge of the Tenda massif, ophiolitic units and basement slices associated to the Schistes Lustrés are separated from the blueschist facies metagranitoids by a thick and complex mylonite zone developed to the expense of the latest.

All these examples therefore lie in intermediate position between the HT end-member or the HP preserving end-member and a theoretical third end-member, which would be represented by a detachment localized at the base of an anomalously strong upper plate.

Some other Mediterranean detachments also lie in intermediate position in the three end-member classification proposed here (Figure 5).

2.1.4 Detachments with intermediate characteristics

The Malaguides-Alpujarrides detachment (MAD, table 1A, Lonergan and Platt, 1995) in the Betics and the Naxos-Paros detachment (NaPaD, table 1Q, Gautier et al., 1993) in Central Cyclades plot along the HT and HP preserving members joint (Figure 5). Both indeed localized at the top of a HP bearing unit, and developed during a thermal reequilibration stage. This thermal reequilibration is more intense in the Naxos-Paros lower plate where prealpine basement underwent pervasive anatexis during exhumation (Vanderhaeghe, 2004), than in the Alpujarrides units, where high-pressure amphibolite facies overprint induced local migmatization in the top Herradura unit and pervasive resetting of radiochronometers (Azañón and Crespo-Blanc, 2000).

Other complex detachment systems such as the North Cycladic Detachment System in the Aegean (NCDS, table 10, Jolivet et al., 2010a) or the Menderes detachments in Turkey also plot within the theoretical triangular classification, but show a singular evolution through time as detailed below.

2.2 Evolution of detachments through time

Three features identified in this review allow discussing how detachment systems evolve through time. When present, the nature of the collapse basin associated to extension along the detachment has been systematically noticed (Figure 6) and evidence how topography locally evolved during extension. The Cyclades and Menderes exhibit complex cases of multi-level detachment systems that evolved through time. Eventually relationship between ductile strain and brittle strain gives indices of how detachment systems behave when crossing the brittle-ductile transition.

2.2.1 Evolution of basins

Most of the syn-tectonic basins described in the hanging-walls of detachment systems have marine affinities or evolved through time from continental to shallow marine, suggesting a topography close to sea level, and most of the time subsiding during extension (Figure 6 and table 1, line 2.1). Accommodation is most of the time moderate, with shallow marine coarse facies, but turbidites are described in the Kazdag massif, suggesting a deep depositional environment (Bonev and Beccaletto, 2007). The opposite trend from marine to continental is only described on Naxos, where marine sandstones and pelites evolve into fluvial deposits (Bargnesi et al., 2013; Kuhlemann et al., 2004), implying that accommodation is decreasing with time, unless regional parameters actually prevail on local processes. Even in the case of detachments initiated in syn-orogenic conditions, such as in Crete, where possibly lacustrine breccias are described at the base of Neogene basins (Jolivet et al., 1996), no proper intra-montane collapse basins developed during activation of detachments in the Mediterranean realm. This

implies that initial topography was close to or rapidly lowered to sea-level while MCC developed.

2.2.2 Evolution of strain localization

Complex detachment systems evolution can be traced in the three end-member classification proposed here. As for the Cyclades, the north-verging North Cycladic Detachment System (table 10, Jolivet et al., 2010a) and south-verging Western Cycladic Detachment System (table 1R, Grasemann et al., 2011) are composed of several mylonite zones that successively localized strain during regional extension. In Northern Cyclades, the Tinos Detachment first localized along a former thrust at the top of the Cycladic Blueschist Unit and beneath the Upper Cycladic Nappe, constituted of a thick and resistant greenschist facies ophiolitic complex . Strain sequentially localized upward on the Livada detachment and then Mykonos detachment, while granites progressively intruded the lower plate and thermal reequilibration of the nappe pile occurred. On Mykonos, syn-tectonic granite directly roots in the highly strained migmatites of the lower plate (Denèle et al., 2011), evidencing a sharp contrast in thermal history between the upper and lower plates. In Ikaria, the Fanari ductile to brittle detachment located at the top of the syn-kinematic I type Raches granite and the ductile Agios Kyrkos shear zone show strain and age patterns similar to the North Cycladic Detachment System, and can be considered as the eastward extension of the same structure (Beaudoin et al., 2015; Laurent et al., 2015). The North Cycladic Detachment System therefore evolves from lower bound toward the HT end-member in the ternary classification proposed, with a perennial detachment zone relocalizing stepwise at the base of the upper plate while the lower plate exhumes. The Western Cycladic Detachment System described on

315 Kea, Kythnos and Serifos islands possibly followed the same evolution with initial
316 detachment located at the base of the Pelagonian serpentinites and schists and
317 progressively moving upward in the nappe pile as the thermal effect of syn-tectonic
318 intrusions (on Serifos and Lavrion) modified the crustal rheological properties
319 (Rabillard et al., 2015).

320 In Menderes, the initial structure that accommodated extension during Hellenic slab
321 retreat is the Simav Detachment (SiD, table 1T, Isik and Tekeli, 2001). This large scale
322 high strain zone is localized at the base of the Izmir - Ankara Ophiolite, within a HP
323 nappes complex preserving eclogites and blueschists (Afyon, Tavsanlı and Oren nappes
324 (Plünder et al., 2013). Rocks in its footwall mainly consist of amphibolite facies gneiss
325 with the metasedimentary Selimiye unit in uppermost position. Early northern
326 Menderes granites are considered as syntectonic to SiD (Isik and Tekeli, 2001), due to
327 development of high temperature foliation in these in the vicinity of SiD and overlap
328 between their U-Pb zircon crystallization ages and cooling ages of biotites within
329 derived mylonites (Dilek et al., 2009) in the early miocene . The younger Alasehir and
330 Kucuk-Menderes Detachment systems, Central Menderes, Turkey (AlKuD, table 1T),
331 subsequently cut across Menderes and divide it in Northern, Central and Southern
332 domains. Rocks beneath and above the Alasehir and Buyuk Menderes detachments
333 therefore derive from the same amphibolite facies metamorphic sequence (Bozkurt and
334 Oberhänsli, 2001). The north-verging Alasehir detachment located at the top of the
335 Salihli syn-tectonic granodiorite (Isik et al., 2003) in the late Miocene (Dilek et al., 2009)
336 and the antithetic detachment at the base of the Buyuk Menderes graben also relates to
337 the late Miocene intrusion of Menderes granites (Dilek et al., 2009). Even if recent
338 migmatites are actually present within Menderes (Bozkurt, pers. comm.) the sole
339 localization mechanism proposed so far for these detachments is the weakening due to

granite intrusion. Strain indeed localized at the top of these plutons that evolve from isotropic in their core toward mylonites in the vicinity of detachment (Dilek et al., 2009). Mylonitic foliation is marked by biotites within granites and greenschist facies assemblages within mylonites. Strain therefore progressively localized during granite cooling and exhumation. The central detachments accommodated the late stages of intra-Menderes extension in middle miocene times, while the top Menderes northern Simav detachment progressively ceased localizing strain (van Hinsbergen, 2010). These complex long-life detachment systems systematically show an initial localization along intra-crustal rheological contrasts inherited from nappe stacking stages, the base of ophiolites and HP-bearing cover units being prone to such localization. The later stages are associated to HT metamorphism and/or syntectonic granite intrusions. Strain is then localized higher up in the tectonic pile (as in the Cyclades) on a structure synthetic to deeper precursors or symmetrically neoformed in the core of the already exhumed lower plate (as in Menderes).

2.2.3 Relationships with brittle deformation

When evidenced, relationships between brittle strain in the upper and lower plates and the ductile to brittle strain along the detachment system were systematically noticed (Table 1, items 2.4, and 4.3). In most cases (Catena Costiera, Alpi Apuane, Rechnitz, Northern Cyclades, Northern Menderes) brittle strain is expressed as high- to low-angle normal faults, developed in the upper plate and rooting in the detachment zone. More rarely brittle deformation cuts across and offsets the main detachment system. When expressed as late stage high-angle normal faults (as in Balagne, Corsica, in the Motajica

window, in southern and eastern Rhodope or Crete), this brittle strain can be viewed as late features accommodating the late stages of extension or unrelated later tectonics. The case of Elba island, with the development of the Zuccale fault, is a rare described case of a low angle brittle normal fault localizing independently of ductile inheritance and cutting across the extended nappe pile (Collettini and Holdsworth, 2004). This feature is also recognized today along the Apennine detachment system, with the low angle normal active Alto Tiberina fault cross-cutting all the thrust structures (Chiaraluce et al., 2007), which system the Zuccale fault zone also pertains to (Collettini et al., 2006).

Interpretation of relationships between ductile and brittle strain is not unique: depending on whether faults bounding basins are rooted on or cut across the detachment system, Menderes basin system can be viewed as syn-tectonic collapse structures (according to VanHinsbergen, 2010 for instance) or a wide rift zone with periodic rift basins obliterating the core complex structure (Ring et al., 2003).

2.3 Rheological implications of model end-members

The three end-members proposed in this review are not equally straightforward in terms of rheological implications. The HT end-member actually represents the most widely accepted definition of MCC, with a hot and hence low-viscosity lower crust. It has been quantitatively considered in many thermo-mechanical models (e.g. Rey et al., 2009; Tirel et al., 2006). A theoretical HS end-member with resistant upper crust, constituted of massive mafic or ultramafic lithologies thrust on top of a less resistant crust, can also be considered. Intra-crustal rheological contrast is here produced by the tectonic

inversion of the rheological profile (Huet et al., 2010) and has been tested via thermo-mechanical modelling. The third HP end-member, representing detachments developing at the top or within HP cover units is less straightforward in terms of strength profile. Why would strain preferentially localize in or along such units in the post-thickening nappe pile ? (Huet et al., 2010) considered the Cycladic Blueschist Unit as a weak layer intercalated between the more resistant Upper Cycladic Unit ophiolite and the Cycladic Basement Unit constituted of granitoids mainly. Before testing the effect of such a layering, its relevance must be discussed. HP cover nappes can be considered weak due to the nature of their protolith. Initially constituted of shales and carbonates, they are fine grained, their water content is high and they are strongly foliated, thus being prone to localize deformation. Nevertheless, metamorphic transformations may significantly change their mechanical behaviour. In order to quantitatively discuss this point we performed thermochemical modelling of the impact of exhumation on average crustal lithologies (Figure 7). Modal evolution of a metabasalt, a metagranite and a metapelite with 10 wt% added water (dry compositions basalt and granite : Le Maitre (1976), shale : Boggs 1995) have been modeled with Theriak-Domino suite (de Capitani and Petrakakis, 2010) along different pressure-temperature paths from blueschist (1.2 GPa, 500°C) to lower greenschist facies (0.3 GPa, 200°C) using Holland & Powell database (Holland and Powell (2004) with Diener et al. (2007) activity model for amphiboles). Two cases can be considered : a path with thermal reequilibration coeval with decompression (path 1, Figure 7), and a path with decompression in the greenschist facies prior to substantial cooling (Path 2, Figure 7), more representative of the PT paths documented in HP units exhumed in Mediterranean MCC (Figure 3). In the case of a basalt, exhumation is marked by decrease in amphibole amount (due to glaucophane destabilization) whatever the path. Epidote is retrogressed into lawsonite along path 1

only. No substantial change in phyllosilicate content is observed whatever the considered path. In the case of a granite, the only substantial modal change modeled is a decrease then increase in phyllosilicate content along path 2. Eventually, in the case of a metapelite, epidote-lawsonite transition is also observed, and phyllosilicate content rises close to 50 vol% along path 2. Considering the amount of phyllosilicates as one of the first order mineralogical control on rocks strength in the blueschist/greenschist temperature range (Gueydan, 2004), the comparison of its evolution along paths for the different lithologies can yield the evolution of strength contrasts within a composite nappe pile (Figure 7D&H). As expected metapelites exhibit the highest phyllosilicates contents. No substantial contrast appear along path 1, while the more realistic two-stage PT path 2 exhibits a strong contrast between metapelites, reaching 40 vol. % phyllosilicates and other lithologies between 500 and 400°C i.e. within the ductile strain domain. It seems therefore that initially weak metasedimentary sequences are furthermore weakened by retrogression in the greenschist facies. Fluid transfers implied by these transformations also can lead to strength changes during exhumation. HP cover units can therefore be considered at first order as weaker zones interlayered in the nappe pile, and therefore prone to localize deformation during syn-to post-orogenic extension. It seems from the presented review that the development of a perennial detachment during extension of a beforehand-thickened crust is therefore possible when a critical intra-crustal strength contrast appears, due to thermal reequilibration (HT MCC case) or due to inheritance (strong upper crust and weak middle crust). This conceptual model, can be quantitatively explored with thermo-mechanical models in order to estimate the order of magnitude of the required intra-crustal strength contrast and to assess how the different end-members can yield a detachment structure.

436

437 **3. Numerical modelling**

438 **3.1 Numerical code used**

439 The numerical code used in the present study is based on the FLAC algorithm (Cundall,
440 1989) and its subsequent evolutions (Burov and Poliakov, 2001; Poliakov et al., 1993;
441 Yamato et al., 2007). Conservation of momentum equation and heat equation are
442 iteratively solved. No radiogenic internal heating was implemented in the present
443 simulations, mainly due to uncertainties on its value in the case of a crust resulting from
444 complex nappe stacking. Thermomechanical coupling is enforced using Boussinesq
445 approximation for the computation of density, including thermal stresses. Temperature
446 is advected with the mesh within the Lagrangian formulation of the code. The rock
447 behaviour is approximated by explicit visco-elasto-plastic rheology. Ductile deformation
448 is modeled with a power-law and brittle deformation with a Mohr-Coulomb plastic flow
449 law (table 3). The effective rheological behaviour is determined by current strain-rate,
450 state of stress and temperature (Le Pourhiet et al., 2004). A marker-based remeshing
451 procedure allows for handling of very large strains and displacements (Yamato et al.,
452 2007). To focus on the influence of the rheological stratification, as in (Huet et al., 2010),
453 the model setups have been deliberately simplified. Erosion, shear-heating, partial
454 melting and mineral phase transitions are not considered in the computation.

455

456 **3.1 Initial set-up and initial boundary conditions**

457 Experimental design (Figure 8) directly derives from Huet et al. (2010). Thermal
458 boundary conditions are 0°C at the surface and fixed initial temperatures at the bottom

(1000, 1200 and 1400 °C at 90 km, implying initial Moho temperatures between 398 and 1015°C, as listed in table 4). Lateral heat flow is set to zero. These thermal conditions allow to cover the complete crustal thermal gradient range (dashed curves on Figure 3) from low (13°C/km), as inferred from PT paths in the Cyclades for instance (Figure 3 and Huet et al., 2010) to moderate (18°C/km) as explored in previous thermo-mechanical modelling studies (Tirel et al., 2008 for instance).

Both sides of the models are assigned free slip condition. As in Huet et al., 2010, asymmetric lateral velocity is applied with 1 cm/yr on the lefthand side and 0 cm/yr on the righthand side.. Initial stretching rate for a 210 km long crustal segment is therefore $2 \cdot 10^{-15} \text{ s}^{-1}$. This velocity boundary conditions corresponds to a mean value in the sensitivity tests to strain rates in numerical studies (Tirel et al., 2008), or in the velocity range used for analytical modeling (Buck, 1990). Upper surface is free, and bottom behaves as a Winkler foundation, ensuring local isostatic compensation (Burov and Poliakov, 2001). The initial velocity field is linearly interpolated from the boundary conditions and the stress is initially lithostatic and isotropic. No resistance or density anomaly is prescribed to localize strain, a random noise of 5 MPa has been instead added to the mean cohesion value (20 MPa, table 3) in the upper crust, except on a 20 km wide zone on the left hand side of the model where cohesion is set to 30 MPa in order to prevent artificial strain localization on the fast moving boundary (Huet et al., 2010).

The model is initially composed of a 210 km wide and 30, 45 or 60 km thick two-layers continental crust resting on top of lithospheric mantle. Initial mesh grid size is 0.75 x 1.25 km. All crustal layers have the same nominal density and thermal expansion coefficient (table 3), so that only temperature contrasts can yield density contrasts.

Mantle properties are the same in all simulations, its viscous behaviour is modelled as the one of dry dunite (Chopra and Paterson, 1984). Viscosity parameters for crustal layers derive from experimental data from quartz-diorite (Ranalli and Murphy (1987) as used in Tirel et al., 2008), dry quartzite (Ranalli and Murphy (1987) as used in Huet et al., 2011) and wet quartzite (Kirby and Kronenberg, 1987). Crustal structures can be rheologically homogeneous (27 simulations: d/d, q/q and w/w series in table 4) or heterogeneous. Among heterogeneous crust simulations, 18 were performed with upper and lower crust the same thickness (evenly layered crust, table 4). 6 simulations were performed with total crustal thickness 45 km composed with quartz-diorite upper crust and dry quartzite lower crust, and 15 or 30 km thick upper crust.

3.2 Results

The 51 simulations run in this study are compared on the basis of the strain geometries produced by extension as well as P-T paths followed by the possibly exhumed lower crust. The different resulting geometries, exemplified in Figures 9, 10 and 11 are briefly described before a more statistical approach is proposed for all simulations. Only 3 cases out of 51 (X in table 4) yielded numerically unstable computation, with mantle rupturing from the very first stretching steps. These simulations, performed with cold conditions (bottom temperature 1000°C) are considered as geologically meaningless. Narrow rifts simulations all exhibit localized strain zones in the upper crust and upper mantle that connect across the ductile lower crust to stretch the complete crust to complete disruption after less than 10 Myr extension. They are characterized by high amplitude Moho deflection and limited lower crustal flow. Margin geometries produced exhibit complex geometries due to simple or complex rift systems (Figure 9C) isolating

507 upper crustal blocks. Wide rift geometries (Figure 9B, D and F) show no transcrustal
508 shear zones developing, upper crust disruption is compensated by lower crustal flow
509 and Moho deflection is limited. Nevertheless, as analytically predicted by Buck (1991),
510 lower crust resistance to flow is higher than plastic yield strength of the upper crust and
511 strain relocalizes laterally, before an extended lower crust dome is formed. Strain
512 localization shifts laterally in time with a wavelength determined by initial set-up.

513 Metamorphic core complex geometries can be sub-divided into two sub-types :
514 asymmetrical domes with a perennial detachment system assisting exhumation of the
515 lower crust over the whole stretching event (Figure 10), and more symmetrical
516 geometries with lower crust exhumation and strain localizing between two sub-domes
517 (double-dome core complexes, Figure 11, DDCC in table 4). In both cases, the upper
518 crust disruption is compensated by lateral and upward flow of the lower crust into a
519 single perennial core. In the perennial detachment metamorphic core complex case
520 (Figure 11, PDCC in table 4), strain relocalizes at the upper lower crust boundary
521 yielding an asymmetric flow pattern in the lower crust and the persistence of a
522 detachment system at the top of the dome structure with younging upward strain ages .
523 The youngest rocks exhumed are then found beneath this detachment system, and
524 exhibit "cool" PT path with no heating associated to decompression (Figure 10 B, F & G).

525 In the double-dome core complex case (Figure 11) the high strain zone is located at the
526 core of the exhumed lower crust, implying a symmetric flow pattern and exhumation
527 ages younging from the edges to the core (Figure 11B & F). The later rocks exhume, the
528 higher the amplitude of their decompressional heating (Figure 11G). Rocks exhumed
529 within or close to detachments systematically show decompression paths along lower
530 thermal gradients (Figures 10G & 11G). This remarkable feature is due to progressive
531 cooling by the overriding upper unit, and might be less pronounced with viscous heating

532 implemented in the code used. Thermal impact of viscous heating along extensional
533 detachment systems has been estimated about 50°C temperature excess during
534 decompression of hanging-wall units (Souche et al., 2013), and would therefore affect
535 PT path shapes only on the second order.

536 Resulting geometries occur logically according to initial set-up (Figure 12). Narrow rifts
537 are produced by thin crust simulations whatever their thermal state, except when lower
538 crust is drastically weak. Wide rift geometry is indeed produced for wet quartzite lower
539 crust simulation with initial crustal thickness 30 km and initial Moho temperature above
540 400°C (Figure 12C). On the other hand double-dome core complexes geometries are
541 produced in thick crusts, whatever their thermal state, except when especially strong.
542 Simulation with 60 km thick quartz-diorite homogeneous crust and $T_{\text{Moho}} = 752^{\circ}\text{C}$
543 indeed yield a wide rift pattern. The most sensible crustal thickness is 45 km. With this
544 thickness, weakening of the whole crust promotes a switch from wide rift to double
545 dome MCC pattern (Figure 12 A, B & C). Weakening of the lower crust only has the same
546 effect but intermediate strength lower crust simulations (Figure 12D) yield wide-rift
547 geometry when cold and perennial detachment MCCs pattern when warmer. The
548 position of the intra-crustal strength contrast within the crust also has an impact.
549 Thinner lower crust geometries (line 2/3 on Figure 12F) yield a narrow rift pattern
550 when cold, which can be explained by an easier connection between localized strain in
551 the upper crust and in the mantle and an overall less ductile rheology (higher
552 ductile/brittle ratio). Thicker lower crust simulations (lower ductile/brittle ratio, line
553 1/3 on Figure 12F) yield more distributed extension mode (wide rift or double dome
554 MCCs according to thermal state). Perennial detachment MCCs are produced for
555 intermediate and maximal initial Moho temperatures and low to intermediate lower
556 crust thicknesses. From this analysis, several general comments arise: i- thermal state is

557 not the dominant controlling factor on extension modes, at least in the range explored
558 here, ii- crustal thickness has a critical impact in the 30-60 km range, iii- perennial
559 detachment MCCs are developed only in specific conditions with intermediate initial
560 crustal thicknesses and intermediate intra-crustal strength contrasts.

561 In order to explain them with unified criteria, occurrences of the four final geometries
562 described here (narrow and wide rifts and metamorphic cores with perennial
563 detachment or double dome geometry) have systematically been reported against initial
564 thermal and rheological parameters for the tested lithospheres (Table 4 and Figure 13).
565 Intra-crustal strength contrast, i.e. ratio between the strength at the brittle-ductile
566 transition depth and viscous strength of the crust at Moho depth as well as the
567 integrated lithospheric strength expressed in MPa.km, have been calculated from the
568 initial conditions of all models. These parameters have been chosen for their relevance,
569 as well as for their easy calculation for already published comparable studies (cf
570 discussion). Tested integrated strengths range over more than 1 order of magnitude,
571 cold and thin crusts leading as expected to stronger lithospheres than warm and thick
572 ones. Intra-crustal strength contrasts range over 4 orders of magnitudes, with the thin
573 and cold crusts having contrast ratios lower than 10 and warm and thick crust
574 exhibiting ratios higher than 10^4 . Rift geometries are produced for high lithospheric
575 strength and low strength ratios. Only such lithosphere can indeed reach their plastic
576 yield strength and connect localized upper crust strain to localized strain in the mantle.
577 Wide rift geometries are produced over a wide range of integrated strength but a
578 narrow range of strength ratio. This reflects the close balance between upper crust
579 strength and lower crust ability to flow implied by the development of wide rifts. Double
580 dome MCCs are develop over a wide range of integrated strengths for strength ratios
581 higher than 10^3 . The specific conditions prone to the development of perennial

detachment appear as a narrow intra-crustal strength contrast range close to 10^3 . In this range, all the possible extension modes are actually produced; only rifts are produced for contrasts lower than $10^{2.5}$ (i.e. ~ 300) and only double dome MCCs are produced for contrasts higher the $10^{3.5}$ (i.e. ~ 3000).

The effect of each tested parameter can be precised with series having one varied parameter only (dashed lines on Figure 13). As expected, the initial crustal thickness and initial thermal state impact both parameters. Thick and/or warm crusts yield high intra-crustal strength contrast for low integrated strength. Layering has an impact on contrast more than on integrated strength. The intra-crustal strength contrast parameter is actually irrelevant to investigate the effect of the depth of inherited strength discontinuity within the crust (crosses on Figure 13). More sophisticated parameters would help, but would necessarily be more design-dependent.

The parametric study presented here yields some qualitative and quantitative conclusions about the boundary conditions and lithosphere properties prone to the development of MCC patterns. First, MCCs can form in moderately thickened crust along geotherms as low as $13^\circ\text{C}/\text{km}$. Depending mainly on the intrinsic strength ratio between the upper and lower crust, two types of MCCs can develop: double domes for crusts with a high strength ratio, MCCs with a perennial detachment at their top for crusts with a strength ration close to 10^3 . Exhumed lower crust material shows limited heating during decompression in MCCs with a perennial detachment, and only material exhumed late in double-domes exhibits substantial heating. MCC pattern development is therefore thermally compatible with the preservation of HP-LT paragenesis in their core, especially on their limb.

4. Discussion

4.1 Parameter study and published MCC models : critical intra-crustal strength contrast for MCC formation

In order to test our parameter study against published analogue studies, the same parameters (intra-crustal strength contrast and integrated lithospheric strength) were computed for published simulations with comparable initial set-up (Figure 14). 32 simulations from Tirel et al. (2008) and 9 cases study in Huet et al (2011) were performed with the same numerical code as present and layered or homeogeneous crustal geometries. Among the 4 simulations presented in Rey et al. (2009) and performed with the Ellipsis finite element code (Moresi et al., 2003), the 2 melt-free cases with varied stretching rate ($2 \cdot 10^{-16}$ and $2 \cdot 10^{-15} \text{ s}^{-1}$) can yield representative initial crustal strength constrast and integrated strength values. In the study performed by Schenker et al. (2012), with the I2VIS code (Gerya and Yuen, 2003) 12 initial geometries were tested with a rheologically homogeneous crust and varied initial thickness and Moho temperatures. In these two studies, partial melting is implemented as a viscosity drop down to a threshold value (10^{18} Pa.s in Rey et al., 2009, 10^{17} Pa.s in Schenker et al., 2012). Eventually, the two-layer wedge shape crust simulations without prescribed weak zone (A and B type models) from Wu et al. (2015), performed with a modified version of PARAVOZ (Tan et al., 2012) can be plotted on the same diagram as segments between their thin and thick ends. Compilation of these simulations with ours give a 108 simulations dataset, produced by different numerical codes, different crust and mantle representative materials, and a variety of boundary conditions. They all plot along the same trend in the intra-crustal strength contrast vs integrated lithosphere strength. Tirel et al. (2008) explored a linear domain across the whole possible field, while Huet et

630 al. (2011) and Rey et al. (2009) focused on weak lithospheres and intermediate strength
631 profiles. The equivalence between the present-study classification and other studies is
632 not straightforward. The "double-domes" produced by Rey et al. (2009), the spreading
633 domes in Huet et al., (2011) and the double-dome MCCs in the present study can be
634 considered as equivalents. The "true rifts", "wide rifts" and "MCCs" produced in Huet et
635 al. (2011) are respectively synonyms in terms of dynamics with rifts, wide rifts and
636 perennial detachment MCCs produced here. Even if Schenker et al. (2012) mainly
637 focused on the impact of the depth of the heat source (within the crust or within the
638 mantle) on the lithospheric response, their classification also has a morphological basis.
639 The "dysharmonic dome" and "lower crustal dome" types distinguished actually
640 respectively correspond to the perennial detachment mode and the double-dome core
641 complex types used here. "Gneiss domes" and "narrow rifts" structures produced in Wu
642 et al. (2015) with 2 layer crusts show some periodicity that could relate them to wide
643 rifts produced in the present study. In first approximation, "narrow rifts" and
644 "continental margins" types can be considered as localized extension modes with crustal
645 necking, while the "gneiss dome" mode compares with the double-dome MCCs produced
646 here. In Tirel et al. (2008) the only available categorization is "crustal necking" vs "MCC".
647 Even if presented examples of MCCs rather exhibit perennial detachment core
648 complexes geometries, we keep the classification as simple as published. This
649 compilation eventually shows that rifts or MCCs can be produced over a wide range of
650 integrated lithospheric strength. Intra-crustal strength contrast seems a sensitive
651 parameter for extension mode prediction: here again, only rifts are produced for
652 contrasts lower than $10^{2.5}$ (i.e. ~ 300) and only double dome MCCs are produced for
653 contrasts higher the $10^{3.5}$ (i.e. ~ 3000).

654 Tirel et al. (2008) extensively explored the effect of strain rate on extension mode, and
655 Rey et al. (2009) also varied this parameter. Their results on this diagram show that an
656 increase of 1 (for Rey et al., 2009) or close to 1 (for Tirel et al., 2008) order of magnitude
657 only slightly modifies the crustal properties, and only changes extension mode for
658 lithospheres in the critical domain next to 10^3 . Wu et al. (2015) who studied the effect of
659 layered wedge shape come to the same conclusion as here: intrinsic strength contrast
660 due to inherited layering is a first order parameter for changing extension mode. The
661 realistic values used in that study actually cover a 2 orders of magnitude range in
662 strength contrast between the strongest (plagioclase above dry quartz) and the weakest
663 (plagioclase above wet quartz with 0.12 % added water).

664 Scaling of analogue models in Figure 14 diagram is impossible, but viscosity and strain
665 rates values used in (Brun et al., 1994) allow the calculation of a $10^{1.2}$ intra-crustal
666 strength contrast in the tilted-block mode extension and $10^{2.39}$ contrast in MCC mode.
667 Here again, an increase about 1 order or magnitude is sufficient to promote MCC pattern
668 development.

669 Since the first order parameter often considered as critical for MCC pattern development
670 is initial thermal state (Buck, 1991, Tirel et al., 2008) to such an extent that MCCs are
671 sometimes considered as restricted to warm lithospheres, the effect of intra-crustal
672 strength contrast can be evaluated in regard to initial Moho temperature (Figure 15). All
673 simulations considered here cover a wide domain of initial thermal conditions between
674 400°C and 1340°C Moho temperatures, i.e. much wider than the expected possible
675 values, even in the most severe post-orogenic thermal relaxation. The present study
676 produces MCCs with initial Moho temperature as cold as 680°C . Huet et al. (2009) and
677 Wu et al. (2015) yielded the same result, highlighting that cold MCCs constitute a

thermo-mechanically sound concept. The rift patterns with the initially warmest lithosphere were actually produced by Tirel et al. (2008). The domain in which the different models can give any of the distinguished extension modes actually span from 470°C to 950°C initial Moho temperature, meaning that thermal state, in its admitted range, is not a critical parameter for predicting lithospheric extension mode.

4.2 Numerical insights versus Mediterranean examples

4.2.1 Natural vs numerical categorization

The parametric study presented here, together with a compilation of similar numerical modelling of continental extension modes leads on one hand to a 4 end-member model, with double domes MCCs, perennial detachment MCCs, wide rift and narrow rift as possible resulting geometries. On the other hand, the review of Mediterranean extensional MCCs points to a 3 end-member categorization of MCCs : HT MCCs with a high temperature lower unit, HP MMCs with an interlayered high-pressure nappe preserved in the vicinity of the detachment, and High Strength (HS) upper crust MCCs localizing strain at its base. The juxtaposition of both reviews enables discussing the mechanical properties behind the three natural types defined here. Actually each of the numerical studies also explored a ternary diagram with 3 end-members : high temperature, weak middle crust and strong upper crust (Figure 15 inset) that resembles the ternary diagram used for classification of natural Mediterranean cases. Weak middle crust relate to the HP natural end-member, and strong upper crust to HS end-member. Tirel et al. (2008), Schenker et al. (2012) as well as the high temperature experiments conducted here confirm the acknowledged behaviour of continental lithosphere with

high thermal gradient (HT end-member). Within this category the distinction of double-dome vs perennial detachment MCCS will be discussed further. Huet et al. (2009) explored the effect of an inter-layered weak zone, representing a HP metasedimentary nappe in natural cases. In this respect, they explored the left joint of the ternary diagram. The present study, with 2-layer crust lithospheres, as well as Wu et al. (2015) *pro parte* explored the right-hand joint of the diagram. Numerical modelling therefore evidences the mechanical feasibility of all natural end-members.

4.2.2 Natural double domes and perennial detachment MCCs in nature

Not all natural cases can be plotted in the diagram derived from numerical simulations (Figure 16), since their initiation thermal conditions and intra-crustal strength contrasts remain uneasy to assess. Nevertheless, some of them can be qualitatively localized and tend to confirm the relevance of this categorization. The description of double-domes by Rey et al. (2009) is based on a two dimensional projection of Naxos core complex main features. Schenker et al. (2012) also consider Naxos as a natural case for their collisional heat induced migmatitic core complex, while Rhodope represents the asthenospheric heat induced migmatitic core complex. These cases (table 1) all illustrate the high temperature-high strength (Figure 16) end of our simulations, with low-viscosity migmatites exhumed in the core of the lower plate beneath a detachment located at the base of an unmetamorphosed ophiolite-bearing unit. Among cold MCCs, the Catena Costiera detachment (Rossetti et al., 2004) exhumed blueschist continental metaclastics and marbles beneath a greenschist ophiolitic melange. Mylonites and cataclasites derived from weak marbles and metapelites, implying a strong inheritance effect. Strain indicators show a pervasive asymmetric sense of shear toward NNW, related to the

activation of a single detachment zone. Other large MCCs such as Northern Cyclades, Motajica, Menderes or Betic cordilleras actually show evolution through time that can relate to the evolution of the mechanical properties of the crust they develop in.

Compilation of PT paths in metamorphic units from the Aegean domain {Jolivet:2008eb} shows variations in decompression paths similar to synthetic paths computed from numerical modelling (Figures 10G & 11G), with cooler paths for the latest rocks exhumed in Crete or in Peloponnese. PT paths shapes are therefore relevantly reproduced without viscous heating in our models testifying that this process is a second order phenomenon in detachment system, especially when weak lithologies are implied over time scales about 10 Myr (Souche et al., 2013) or that shear heating is counterbalanced by local cooling processes such as fluid advection within detachment (Famin et al., 2004, Mezri et al., 2015).

4.2.3 Temporal evolution of multiple detachment systems

Detachment systems (§ 2.2.2) as well as smaller structures with several detachments (Betics and Motajica for instance) can be described as a sequence of changes in strain localization levels. Menderes and Cycladic systems evolve toward the high temperature end-member in the ternary diagram proposed for natural cases (Figure 5). Nevertheless, their mechanical evolution differ and they do not follow the same trend in the intra-crustal strength contrast vs thermal state diagram (Figure 16). In Menderes (Figure 17A), strain is first localized at the base of or within the high-pressure metamorphic rocks and the Izmir Ankara Ophiolite, along the Simav Detachment, which actually reactivates an older thrust. Whether the equivalent thrust at the base of the Lycian

748 nappes effectively accommodated extension remains unknown. Even though the
749 intrusion of the syn-tectonic Egrigoz granite (crystallization at 20-21 Ma and cooling at
750 20 Ma, Dilek, 2009) is expected to lower the effective strength of the detachment zone,
751 mylonites along Simav Detachment exhibit similar ages (22.8 Ma, Tekeli, 2001) implying
752 a short activation duration for the Simav Detachment. Strain is accommodated in a
753 second time within the lower plate, between 16 and 7 Ma according to syntectonic
754 granites ages (Figure 17A), basin infills in hanging-walls of detachments (16.7 to 14.9
755 Ma) and dating of white micas in mylonites (7 ± 1 Ma Lips, 2001). The Alasehir and
756 Buyuk detachments (Table 1) constitute a synchronous and antithetic detachment pair,
757 that relocates strain at the core of the Menderes massif while stretching goes on. This
758 implies that this domain is eventually weaker than the previously activated Simav
759 Detachment zone, probably due to crustal thinning, rise of thermal state and intrusion of
760 granites. Intra-crustal strength contrast is probably rising with thermal state during
761 stretching and extension mode evolves from perennial asymmetric detachment to a
762 double-dome symmetric MCC.

763 In Northern Cyclades (Figure 17B) the successive Tinos, Livada and Mykonos
764 detachment zones (Jolivet et al., 2010) progressively relocate strain higher within the
765 upper plate while stretching develops from 30 to 9 Ma and thermal state increases with
766 consecutive intrusions of syn-tectonic granites at 14-15 and 11-13 Ma. Late reactivation
767 of the Vari detachment on Syros at 10 Ma is limited and the North Cycladic Detachment
768 System can be considered as a perennial asymmetric detachment system responsible for
769 most of the exhumation in northern Cyclades. It seems that intra-crustal strength
770 contrast remained governed by inheritance and that intrusion of granites only promoted
771 upward migration of the shear zone within the upper plate. For this reason, the North

772 Cycladic Detachment System evolution can be viewed as a rise in thermal gradient
 773 without drastic change in intra-crustal contrast through time.

774 The Betic cordilleras (Figure 17C) clearly show a double detachment system, with the
 775 older Malaguides-Alpujarrides Detachment (activated from 23 to 20 Ma) on top and the
 776 younger Mecina-Filabres Shear Zone (activated from 16 to 7 Ma, Augier et al, 2005).
 777 While the first one is highly asymmetric with only top-to-the-North shear sense
 778 indicators developed in the northern mylonitic edge of the Sierra de las Estancias massif
 779 (Lonergan and Platt, 1995), the Mecina Filabres shear zone wraps a multiple dome
 780 structure (Sierra de los Filabres and Sierra Alhamilla) with divergent shear sense
 781 indicators (Augier et al., 2005a). Even if highly tridimensional, this pattern implies that
 782 the older detachment system (MAD, Figure 17C) progressively hardens while lower
 783 detachment activates.

784 In the Carpathian (Figure 17D), the Motajica dome is also the result of a two stage
 785 extension between 25 and 13 Ma (Ustaszewski et al., 2010). Until 18 Ma, the main
 786 structure accommodating strain is the former Tisza-Dacia unit basal thrust reactivated
 787 as a detachment, strain is then accommodated along a symmetric listric faults system
 788 (Figure 17D) that cuts across the Motajica Detachment. The Motajica granite, that
 789 yielded a 27 Ma U-Pb zircon age (Ustaszewski et al., 2010) predates the regional
 790 extension stage so that relocalization of strain within the metamorphic Sava zone cannot
 791 be ascribed to intrusion. Nevertheless, this example is a supplemental illustration of the
 792 recurrent evolution from perennial detachment MCCs toward double-dome MCCs with
 793 increasing stretching of the crust. This conclusion is apparently opposite to the temporal
 794 evolution of extension modes speculated from short term modelling (Buck, 1991) and
 795 exemplified by the Basin and Range: from MCC to wide-rift and then narrow-rift. Ring et

al. (2003) also view Menderes as a wide-rift structure overprinting a former MCC pattern. The main difference between these two evolutions is the heat budget and time scale. The perennial detachment toward double-dome MCC evolution implies heat input at the base of the crust higher than heat dissipation due to stretching, due to lithospheric stretching for instance, while the MCC toward rift evolution implies a decreasing heat budget due to efficient heat dissipation during MCC extension phase. While the first evolution seems supported by natural evidences over short time periods (15 to 20 Myrs in examples developed here, Figure 17), the later could be relevant for longer time-scale evolutions (over more than 40 Myrs for the Great Basin MCC cluster (Dickinson, 2002)

4.3 Thermal state, inheritance, strain rate and metamorphism as controlling parameters for intra-crustal strength contrast

The present study after other numerical modelling of extension modes of the continental lithosphere (Huet et al., 2010; 2011; Rey et al., 2009; 2011; Tirel et al., 2008; 2006; Wu et al., 2015) enables to discuss the balance between parameters controlling the intra-crustal strength contrast within the crust. The present compilation of models shows that an exaggeration of one order of magnitude of the intra-crustal strength contrast is sufficient to switch extension mode from rifting to MCC. In a crude approximation, the impact of the different parameters invoked is to lower the effective strength of the lower crust. Maximum strength within the crust being controlled by frictional behaviour, it is unlikely to vary over one order of magnitude for material governed by Byerlee's law. Using the different lithologies considered as relevant for crustal materials (diorite, quartzite and wet quartzite), a simple calculation of effective strength for various

temperatures and strain rates (Figure 18) shows that for a quartz-diorite at 650°C, strained at $10^{-13.5} \text{ s}^{-1}$, strain rate must decrease of more than 2 orders of magnitude (A in Figure 18), or temperature must rise of 200°C (B in Figure 18) to lower its effective strength of 1 order of magnitude. Alternatively, a wet quartzite in the same conditions is also 10 times less resistant (C in Figure 18). Strain rate and temperature therefore must change drastically in order to promote high intra-crustal strength contrast. Layering due to inheritance can therefore be a first order parameter responsible for such contrasts. These three effects actually can add up: a warm and layered crust stretched at a low strain rate will definitely develop a MCC pattern.

Together with layering, inner complex structures can be inherited from preliminary thickening stages. Dipping lithological contrast have a strong impact on extensional behavior of the crust (Huet et al., 2011; Le Pourhiet et al., 2004; Wu et al., 2015). Lateral strength contrast gradients produced with realistic geometries (Wu et al., 2015) are also about 1 order of magnitude (Figure 14). Dipping interfaces therefore also constitute a first order inheritance effect. Transcurrent structures, prone to develop during thickening

Metamorphic reactions responsible for destabilization or crystallization of weak phases or fluid influx were not taken into account in all the studies considered here, that all focused on materials with continuous rheology. According to experimental studies (Jaoul et al., 1984), a quartz aggregate with 0.28 wt% H_2O has an effective viscosity 100 times lower than a dry aggregate. This severe effect implies that the destabilization or crystallisation of a limited percentage of hydroxyl-bearing minerals during the exhumation of the lower plate (Figure 7) can induce high intra-crustal strength contrast. This effect has also been implemented in numerical modelling with complex feedback

between strain, permeability and metamorphic reactions (Mezri et al., 2015). It appears that models implementing water-controlled phase change develop an asymmetric detachment MCC evolving into a double-dome MCC, while reference model (TMoho 810°C, stretching rate 0.8 cm/yr) yields a narrow rift extension pattern. Second invariant of strain rate tensor in the high fluid flux domains is more than one order of magnitude higher than in the low fluid flux domains, implying a weakening effect of more than one order of magnitude. Partial melting, observed in most of high temperature MCCs cores, has also been considered in models as a sharp viscosity drop within the lower crust (Rey et al., 2009, Schenker et al., 2012). When implemented as such, solidus envelope at depth constitutes a sharp intra-crustal strength contrast and promotes the development of double-dome MCCs. Intra-crustal strength contrasts in these cases are close to 3 orders of magnitude higher than the critical value defined here (Figure 14, 15).

Conclusion

The understanding of MCC development and the localization of strain on a perennial detachment zone was initially based on models with homogeneous crust, and highlighted the major role of crustal thickness and temperature in the mode of extension of continental lithosphere. The examination of natural cases in the Mediterranean realm emphasizes a second type of MCCs exhuming "cold" metamorphic rocks beneath a detachment system localized along sharp intra-crustal strength contrast inherited from thickening stages of orogenic crusts. Interlayered HP parametamorphic nappes and basal contacts of resistant upper units most of the time localize extensional strain. The comparison of new and published numerical models of extending continental

lithosphere confirms the mechanical feasibility of core-complex in continental lithosphere with low thermal gradients and marked layering. It also shows that intra-crustal strength contrast is a key parameter for predicting the extension mode of continental lithosphere. Beyond a critical value of 1000, MCCs develop, whatever the initial thermal state, thickness or strain rate. For high values of this contrast, double-dome core complex develop, with strain progressively re-localizing at the center of the footwall during stretching, while asymmetric dome geometries with one perennial detachment are produced for intra-crustal strength contrast close to 1000. Natural examples of Mediterranean detachment systems also show these two MCC types, and evolution of multiple detachments system show that in most cases inheritance controls the first extension steps and asymmetric strain pattern, and that further stretching is accommodated by double dome dynamics, driven by thermal effects mainly.

The present study is based on mediterranean that share a common tectonic setting and possibly some specificities of the post-variscan crust and alpine cover. Nevertheless the proposed critical parameter for MCC development proposed here, i. e. the intra-crustal strength contrast and the gradation of extension modes and core-complex types could be used in comparable settings. The Cordilleran Core Complexes developed during post-orogenic extension or the numerous comparable studies recently described in China (Whitney et al., 2013) could also be considered in a broader review.

Acknowledgement

This study was initiated during the ANR EGEO project (PI L. Jolivet). This work has received funding from the European Research Council (ERC) under the seventh

891 Framework Programme of the European Union (ERC Advanced Grant, grant agreement
892 No 290864, RHEOLITH). Authors thank C. Doglioni, editor, M. Barchi, reviewer and an
893 anonymous reviewer for their constructive reviews and careful handling of the present
894 manuscript.

895

896

897 **Table and Figure captions**

898 **Table 1 :** Characteristic features of Mediterranean detachments and associated domes
899 and basins. P and T estimates for metamorphic peak conditions (4.2) are given in
900 supplemental table A1. A: amphibolite, BS: blueschist, E: eclogite; GS: greenschist,
901 HANF: high angle normal fault, HPA: high pressure amphibolite, HPG: high pressure
902 granulite, LANF: low angle normal fault, LPA: low pressure amphibolite,

903

904 **Table 2 :** Main features of Mediterranean detachments tectonostratigraphies. C for calc-
905 mylonites, m/c for marine vs continental basins, S and P for syn- and post-orogenic
906 initiations. c continental, m marine, off off-shore. The relative weighing of ticks between
907 lines 1,2 and 3, line 4 and line 5 was used to place natural cases on figure 5.

908 AA Alpi Apuane, ABMDS Alasehir-Buyuk Menderes Detachment System, Asp
909 Aspromonte, ASZ Alakeci Shear Zone, Bal Balagne basal contact, BB Beni Bousera, CaD
910 Capanne Detachment, CCD Calcare Cavernoso Detachment, CMe Central Menderes, CR
911 Corinth Rift, CrD Cretan Detachment, Ed Edough, GK Grande Kabylie, Ik Ikaria, Kaz
912 Kazdağ, KD Kabylia Detachment, KnD Kerdyllion Detachment, LL Lubenik Line, MAD
913 Malguides-Alpujarrides Detachment, MeSZ Messaria Shear Zone, MFSZ Mecina-Filabres
914 Shear Zone, Mo Motajica, NCDS Northern Cycladic Detachment System, NCy Northern
915 Cyclades, Ni Niğde, NP Naxos-Paros, NPD Naxos Paros Detachment, SD Simav
916 Detachment, SD Simav Detachment, Sy Syros, Sif Sifnos, TeSZ Tenda Shear Zone, TKD
917 Tokachka-Kesebir Detachment, VD Vari Detachment, Ve Veporic, ZFZ Zaouia Fault Zone.

918

Table 3 : Rheological parameters used and varied for the different models. Dislocation creep law for the mantle is dry dunite (Chopra and Paterson, 1984), and for the crust quartz-diorite and dry quartzite (Ranalli and Murphy, 1987) and wet quartzite (Kirby and Kronenberg, 1987).

Table 4 : Simulation results. Each simulation is named after its layering (d: quartz-diorite, q: quartzite, w: wet quartzite), the initial crust thickness, in km, and initial bottom temperature in Celsius. Evenly layered crusts have two half-crust layers. All parameters are computed with initial geometry and a 1 cm.yr^{-1} stretching rate for a 210 km long crustal segment, i.e. strain rate = $2.10^{-15} \text{ s}^{-1}$. Intra-crustal strength contrast is the ratio of maximum strength at the brittle-ductile transition over viscous strength at Moho depth, Integrated lithospheric strength is calculated on the 90 km height of models. Result types are listed as : R, narrow rifts, WR, wide rifts, PDC perennial detachment core complexes, DDC double dome core complexes. X refers to geologically meaningless simulations. Bold lines refer to simulations used as examples in figures 8, 9 and 10.

Figure 1 : Principle sketch of a Metamorphic Core Complex, showing the association of structures and fault-rock types considered as symptomatic for ductile extensional mode. Rheological envelope associated to these structures is also shown. BDT : Brittle-ductile transition.

Figure 2 : Map of Mediterranean core complexes on top of Moho isobaths, homogenized and smoothed after (Dèzes and Ziegler, 2001; Di Luccio and Pasyanos, 2007; Koulakov and Sobolev, 2006; Marone et al., 2003; Tiberi et al., 2001) The large Mendere and Betics domes have been subdivided into units. Shoreline in white.

Figure 3 : Pressure temperature conditions estimated for Mediterranean detachment footwalls compared to initial Moho temperatures ascribed in numerical models for MCC formation. Curves 1 : wet solidus of granite (Huang & Wyllie, 1981), 2 and 3 are maximum and minimum initial thermal gradients tested in the present parametric study.. AA Alpi Apuane (1 Mass, 2 Autochthon), Alpu Alpujarrides (1 Herradura u., 2 Sadobreña & Adra u. 3 Escalate u., 4 Lujar Gador u.), And Andros, Asp Aspromonte, BB Beni Bousera (1 mafic rocks, 2 kinzigites, 3 Filali schists), CC Catena Costiera, Ed Edough, ERho Eastern Rhodope (1 HP peak, 2 HT,), GK Greater Kabylie (1 Sidi Ali Bou Nab u., 2 migmatites), Ik Ikaria,, Kaz Kazdag, Me Menderes (1 Cine u. HP , 2 Cine u. HT - 3 Bozdag u., 4 Selimye u.), Mot Motajica, Nax Naxos (1 HP, 2 HT)NF Nevado-Filabrides (1 Bedar Macael u. , 3 Phase D3), Ni Nigde, , Re Rechnitz (1 & 2) SRho Southern Rhodope (Thasos), Sy Syros (1 high pressure, 2 late overprint) Eclogite peak 2 GS retrogression), Te Tenda massif, Corsica, Tin Tinos (1 Eclogite peak, 2 BS stage, 3 transition, 4 GS retrogression), Ve Veporic.

P,T values and references are given in supplemental material table A1

Figure 4 : Tectonostratigraphic positions of Mediterranean detachments. Colors refer to metamorphic facies: pale yellow, undistinct low grade, green: GS greenschist, orange: A amphibolite, brown: G granulite, blue: BS blueschist, purple: E, eclogite. Relevant lithologies are precised. Extensional detachments are abbreviated as in Table 1.

Adria. Adriatic plate, Alpu. Alpujarrides, Aspro. Aspromonte u., APU Amphibolites-Peridotites u., BMD Buyuk Menderes detachment, CBS Cycladic Blueschists, Eg Egrigoz granodiorite, FD Fanari Detachment, Ghom. Ghomarides, GT Gavrovo Tripolitza, Gum.

964 Gumusler, HGMU High grade metasedimentary u., HPU High pressure u., IAO Izmir-
 965 Ankara ophiolite, LANF Low Angle Normal Fault, LiD Livada detachment, Malag.
 966 Malaguides, MGU Migmatite-granite u., Mul Mulhacen, MyD Mykonos detachment, N.F.
 967 Nevado-Filabrides, NMM, CMM & SMM North Central and South Menderes massifs,
 968 Pelag. Pelagonian, Phyllites-Q. Phyllites-quartzites, Pi Pindos, Ro Rochovce, S. L. Schistes
 969 Lustrés, Seb. Sebtides, TB Tyros beds, Tell. Tellian u., TiD Tinos Detachment, Tu Turgutlu
 970 granite, UOU, LOU Upper and Lower ophiolite u., ZF Zaroukla fault

971 **Figure 5 :** Ternary diagram representing the balance of thermal and inheritance effects
 972 on the different Mediterranean extensional detachments, abbreviated as in Table 1.
 973 Positions in the triangle come from the relative weighting of characteristic features listed
 974 in table 1. Regular case : detachments inherited from syn-orogenic tectonics, bold case :
 975 post-orogenic detachments. HT high temperature footwall, HP high pressure unit
 976 preserved in the footwall, HS high strength hanging wall. White arrows represent time
 977 evolution of detachment systems.

978 **Figure 6 :** Synoptic comparative time evolution or relative position of syn-detachment
 979 collapse basins preserved in the Mediterranean realm according to their sedimentary
 980 infill. Arrow senses represent time evolution : most of basins evolve toward marine
 981 (downward arrows), only NaPaD basin is evolving from marine to continental with time
 982 (upward bold arrow). Color represent paleobathymetry for marine basins (light grey:
 983 shallow, darker grey: deep) Refers to line 2.1 in table 1 and references therein.
 984 Abbreviations as in table 1.

985 **Figure 7 :** Thermochemical modelling of mode evolution of three protoliths
 986 representative for pelites, granites and basalts as detailed in table I B95, Boggs, 1995,
 987 LM76 : Le Maitre, 1976. A, B, C and E,F,G : modal evolutions in volume fractions along

paths 1 and 2 as shown on J. Micas are phengite and paragonite together. More chaotic curves in C and G are due to poorly constrained complex local equilibria along main reactions in mafic rocks. D and H : Phyllosilicates volume fraction evolution along the same paths. Shaded areas represent brittle quartz domain.

Figure 8 : Set-up of numerical experiments initial conditions for a 45 km thick double layered crust (d/q_45_1200 cases in table 4) See Huet et al., 2011 for details. Bottom temperatures are varied between 1000, 1200 and 1400°C. Resulting initial Moho temperatures are listed in table 3. Hatched column on the left represent a 20 km wide area with no noise on cohesion and a higher cohesion (30 MPa instead of 20 MPa).

Figure 9 : Compared output of numerical experiments for simulation d/q_30_1400, representative of a rift-type extension mode (A & C) after 6 Myr stretching and simulation q/q_45_1200 representative for the wide rift extension mode after 12 Myr stretching. White curves show foliation trajectories from finite strain computed as in Huet et al., 2011. C & D represent instantaneous strain rates. Initial strength profiles computed for $2 \cdot 10^{-15} \text{ s}^{-1}$ and parameters detailed in table 2 are depicted in E and F. Values are precised in table 3.

Figure 10 : Simulation d/q_45_1200 output representative for perennial detachment core complex extension mode. A & B : geometry after 4 and 14 Myr stretching, with position of markers used for PT and exhumation calculations. White curves show foliation trajectories from finite strain computed as in Huet et al., 2011. C & D : instantaneous strain rates profiles, E Initial strength profile computed for $2 \cdot 10^{-15} \text{ s}^{-1}$ and parameters detailed in table 2F & G: exhumation history and PT paths for markers 1, 2 and 3.

Figure 11 : Simulation d/w_45_1200 output representative for double dome core complex extension mode. A & B: geometry after 4 and 14.1 Myr stretching, with position of markers used for PT and exhumation calculations. White curves show foliation trajectories from finite strain computed as in Huet et al., 2011. C & D : instantaneous strain rates profiles, E Initial strength profile computed for $2 \cdot 10^{-15} \text{ s}^{-1}$ and parameters detailed in table 2F & D: exhumation history and PT paths for markers 1, 2 and 3.

Figure 12 : Resulting geometries of the simulations run in this study. Series are referred to as in table 3. A, B & C represent homogeneous crust simulations, D and E evenly layered crust simulations and F unevenly layered simulations. Line 45 km on D represents the same simulations as line 1/2 on F.

Figure 13 : Intra-crustal strength contrast vs integrated lithosphere strength logarithm plot for all 48 meaningful simulations listed in table 3. Effect of different tested parameters is highlighted by series tied by dash lines. Triangles: series d/q_X_1000, squares: series d/d_60_X, diamonds: d/X_45_1200, crosses: d/q_45_1200. 1/3, 1/2, 2/3 refer to the relative size of the upper crust in unevenly layered crust simulations.

Figure 14 : Intra-crustal strength contrast vs integrated lithosphere strength log-log plot for 46 published numerical simulations compared to present study. TS This Study, H11 (Huet et al., 2011), all 9 simulations are reported, T08 (Tirel et al., 2008) all 32 simulations reported, R09 (Rey et al., 2009) 2 simulations reported, with no melt effect at slow ($2 \cdot 10^{-16} \text{ s}^{-1}$) and fast ($2 \cdot 10^{-15} \text{ s}^{-1}$) stretching rate. S12 (Schenker et al., 2012) all 12 models reported. W15 (Wu et al., 2015) 3 models reported, with wedge-shape layered crust and no prescribed weak interface. Calculations were made at each initial model end (thick and thin ends). Bold squares : stretching rate effect, as explored by Tirel et al.

(2008) on the series with initial crustal thickness 60 km and initial Moho temperature 830°C (mantle heat flux 30mW.m⁻²)

Figure 15 : Intra-crustal strength contrast vs initial Moho temperature logarithm plot for 46 published numerical simulations compared to present study. Legend as in figure 14. Triangle inset refers to ternary end-member MCC types analogue to ternary diagram used for Mediterranean examples (Figure 5). Symbols at triangle edges refer to where different studies plot in this diagram.

Figure 16 : Synthetic diagram showing preponderance of intra-crustal strength contrast over thermal state of the crust on extension mode in mediterranean natural cases on the basis of thermal evolutions recorded.

Figure 17 : Examples of time evolution of multiple detachment systems in the Mediterranean. Detachment named as in Table 1. Moho depth as in Figure 2. A Menderes, IAO : Izmir Ankara Ophiolite, modified after VanHinsbergen, 2010 and Ring, 2003, B- North Cycladic Detachment System, after Jolivet et al., (2010), C Betic Cordilleras after Augier et al. (2005) and Lonergan and Platt (1995), D Motajica after Ustazjewski et al. (2010). g. for granite

Figure 18 : Strength contours in MPa (presented with a logarithmic scale) for different crustal lithologies used in the present study. A, B and C are responsible for the same strength drop of 1 order of magnitude. A: strain rate drop, B: temperature increase, C: lithological change from quartz-diorite to wet quartzite.

References

- Augier, R., Agard, P., Monié, P., and Jolivet, L., 2005, Exhumation, doming and slab retreat in the Betic Cordillera (SE Spain): in situ $^{40}\text{Ar}/^{39}\text{Ar}$ ages and P-T-d-t paths for the Nevado-Filabride complex: *Journal of Metamorphic Geology*, doi: 10.1111/j.1525-1314.2005.00581.x.
- Augier, R., Jolivet, L., Robin, C., 2005b. Late Orogenic doming in the eastern Betic Cordilleras: Final exhumation of the Nevado-Filabride complex and its relation to basin genesis. *Tectonics* 24, n/a–n/a. doi:10.1029/2004TC001687
- Avigad, D., 1998, High-pressure metamorphism and cooling on SE Naxos (Cyclades, Greece): *European Journal of Mineralogy*, v. 10, no. 6, p. 1309–1319.
- Azañón, J.M., and Crespo-Blanc, A., 2000, Exhumation during a continental collision inferred from the tectonometamorphic evolution of the Alpujarride Complex in the central Betics (Alboran Domain, SE Spain): *Tectonics*, doi: 10.1029/2000TC900005.
- Bargnesi, E.A., Stockli, D.F., Mancktelow, N., and Soukis, K., 2013, Miocene core complex development and coeval supradetachment basin evolution of Paros, Greece, insights from (U-Th)/He thermochronometry: *Tectonophysics*, v. 595-596, p. 165–182, doi: 10.1016/j.tecto.2012.07.015.
- Beaudoin, A., Augier, R., Laurent, V., Jolivet, L., Lahfid, A., Bosse, V., Arbaret, L., Rabillard, A., Menant, A., 2015. The Ikaria high-temperature Metamorphic Core Complex (Cyclades, Greece): Geometry, kinematics and thermal structure. *Journal of Geodynamics* 92, 18–41. doi:10.1016/j.jog.2015.09.004
- Beccaletto, L., and Steiner, C., 2005, Evidence of two-stage extensional tectonics from the northern edge of the Edremit Graben, NW Turkey: *Geodinamica Acta*, v. 18, no. 3-4, p. 283–297, doi: 10.3166/ga.18.283-297.
- Beccaluva, L., Bianchini, G., and Siena, F., 2004, Tertiary-Quaternary volcanism and tectono-magmatic evolution in Italy: Special Volume of the Italian Geological Society for the IGC 32 - Florence.
- Block, L., Royden, L.H., 2010. Core complex geometries and regional scale flow in the lower crust. *Tectonics* 9, 557–567. doi:10.1029/TC009i004p00557

1085 Bonardi, G., Cavazza, W., Perrone, V., Rossi, S., 2001. Calabria-Peloritani terrane and
 1086 northern Ionian Sea, in: *Anatomy of an Orogen: the Apennines and Adjacent*
 1087 *Mediterranean Basins*. Springer Netherlands, Dordrecht, pp. 287–306. doi:10.1007/978-
 1088 94-015-9829-3_17

1089 Bonardi, G., Giunta, G., Perrone, V., Russo, M., Zuppetta, A., and Ciampo, G., 1980,
 1090 Osservazioni sull'evoluzione dell'arco Calabro-Peloritano nel Miocene inferiore; la
 1091 formazione di Stilo-Capo d'Orlando: *Bollettino della Societa Geologica Italiana*, v. 99, no.
 1092 4, p. 365–393.

1093 Bonev, N., Beccaletto, L., 2007. From syn- to post-orogenic Tertiary extension in the
 1094 north Aegean region: constraints on the kinematics in the eastern Rhodope Thrace,
 1095 Bulgaria Greece and the Biga Peninsula, NW Turkey. *Geological Society, London, Special*
 1096 *Publications* 291, 113–142. doi:10.1144/SP291.6

1097 Bonev, N., Burg, J.P., Ivanov, Z., 2006. Mesozoic–Tertiary structural evolution of an
 1098 extensional gneiss dome—the Kesebir–Kardamos dome, eastern Rhodope (Bulgaria–
 1099 Greece). *Int J Earth Sci*. doi:10.1007/s00531-005-0025-y

1100 Bonneau, M., 1984, Correlation of the Hellenic nappes in the south-east Aegean and their
 1101 tectonic reconstruction: *Geological Society, London, Special Publications*, v. 17, no. 1, p.
 1102 517–527, doi: 10.1144/gsl.sp.1984.017.01.38.

1103 Bosch, D., Hammor, D., Mechat, M., Fernandez, L., Bruguier, O., Caby, R., Verdoux, P.,
 1104 2014. Geochemical study (major, trace elements and Pb–Sr–Nd isotopes) of mantle
 1105 material obducted onto the North African margin (Edough Massif, North Eastern
 1106 Algeria): Tethys fragments or lost remnants of the Liguro–Provençal basin?
 1107 *Tectonophysics* 626, 53–68. doi:10.1016/j.tecto.2014.03.031

1108 Bozkurt, E., Oberhansli, R., 2001. Menderes Massif (Western Turkey): structural,
 1109 metamorphic and magmatic evolution - a synthesis. *Int J Earth Sci* 89, 679–708.
 1110 doi:10.1007/s005310000173

1111 Bruguier, O., Hammor, D., Bosch, D., Caby, R., 2009. Miocene incorporation of peridotite
 1112 into the Hercynian basement of the Maghrebides (Edough massif, NE Algeria):
 1113 Implications for the geodynamic evolution of the Western Mediterranean. *Chemical*
 1114 *Geology* 261, 172–184. doi:10.1016/j.chemgeo.2008.11.016

1115 Brun, J.-P., and Sokoutis, D., 2007, Kinematics of the Southern Rhodope Core Complex
 1116 (North Greece): *International Journal of Earth Sciences*, v. 96, no. 6, p. 1079–1099, doi:
 1117 10.1007/s00531-007-0174-2.

1118 Brun, J.P., Sokoutis, D., Van Den Driessche, J., 1994. Analogue modeling of detachment
 1119 fault systems and core complexes. *Geol* 22, 319–322. doi:10.1130/0091-
 1120 7613(1994)022<0319:AMODFS>2.3.CO;2

1121 Buck, W.R., 1991. Modes of continental lithospheric extension. *Journal of Geophysical*
 1122 *Research: Solid Earth* (1978–2012) 96, 20161–20178. doi:10.1029/91JB01485

1123 Burg, J.-P., Ricou, L.-E., Ivano, Z., Godfriaux, I., Dimov, D., and Klain, L., 1996, Syn-
 1124 metamorphic nappe complex in the Rhodope Massif. Structure and kinematics: *Terra*
 1125 *Nova*, v. 8, no. 1, p. 6–15, doi: 10.1111/j.1365-3121.1996.tb00720.x.

1126 Burov, E., Poliakov, A., 2001. Erosion and rheology controls on synrift and postrift
 1127 evolution: Verifying old and new ideas using a fully coupled numerical model. *J.*
 1128 *Geophys. Res.* 106, 16461–16481. doi:10.1029/2001JB000433

1129 Boggs S Jr 1995 *Principles Of Sedimentology And Stratigraphy*, Prentice Hall Inc., 774p

1130 Caby, R., Hammor, D., and Delor, C., 2001, Metamorphic evolution, partial melting and
 1131 Miocene exhumation of lower crust in the Edough metamorphic core complex, west
 1132 Mediterranean orogen: *Tectonophysics*, v. 342, no. 3-4, p. 239–273, doi: 10.1016/S0040-
 1133 1951(01)00166-4.

1134 Cao, S., Neubauer, F., Bernroider, M., Liu, J., Genser, J., 2013. Structures, microfabrics and
 1135 textures of the Cordilleran-type Rechnitz metamorphic core complex, Eastern Alps.
 1136 *Tectonophysics* 608, 1201–1225. doi:10.1016/j.tecto.2013.06.025

1137 Carmignani, L., Decandia, F.A., Disperati, L., Fantozzi, P.L., Kligfield, R., Lazzarotto, A.,
 1138 Liotta, D., and Meccheri, M., 2001, Inner Northern Apennines, in *Anatomy of an Orogen:*
 1139 *the Apennines and Adjacent Mediterranean Basins*, Springer Netherlands, Dordrecht, p.
 1140 197–213.

1141 Carmignani, L., Kligfield, R., 1990. Crustal extension in the northern Apennines: The
 1142 transition from compression to extension in the Alpi Apuane Core Complex. *Tectonics* 9,
 1143 1275–1303. doi:10.1029/TC009i006p01275

1144 Chiaraluce, L., Chiarabba, C., Collettini, C., Piccinini, D., and Cocco, M., 2007, Architecture
 1145 and mechanics of an active low-angle normal fault: Alto Tiberina Fault, northern
 1146 Apennines, Italy: *Journal of Geophysical Research*, v. 112, no. B10, p. B10310, doi:
 1147 10.1029/2007JB005015.

1148 Chopra, P.N., Paterson, M.S., 1984. The role of water in the deformation of dunite. *J.*
 1149 *Geophys. Res.* 89, 7861–7876. doi:10.1029/JB089iB09p07861

1150 Collettini, C., Holdsworth, R.E., 2004. Fault zone weakening and character of slip along
 1151 low-angle normal faults: insights from the Zuccale fault, Elba, Italy. *Journal of the*
 1152 *Geological Society* 161, 1039–1051. doi:10.1144/0016-764903-179

1153 Collettini, C., De Paola, N., Holdsworth, R.E., and Barchi, M.R., 2006, The development and
 1154 behaviour of low-angle normal faults during Cenozoic asymmetric extension in the
 1155 Northern Apennines, Italy: *Journal of Structural Geology*, v. 28, no. 2, p. 333–352, doi:
 1156 10.1016/j.jsg.2005.10.003.

1157 Cundall, P.A., 1989. Numerische Simulation der Lokalisierung in reibungsbehaftetem
 1158 Material. *Ingenieur-Archiv* 59, 148–159. doi:10.1007/BF00538368

1159 Daniel, J.-M., Jolivet, L., 1995. Detachment faults and pluton emplacement; Elba Island
 1160 (Tyrrhenian Sea). *Bulletin de la Societe Geologique de France* 166, 341–354.
 1161 doi:10.2113/gssgfbull.166.4.341

1162 Davis, G.A., Lister, G.S., Reynolds, S.J., 1986. Structural evolution of the Whipple and
 1163 South mountains shear zones, southwestern United States. *Geol* 14, 7.
 1164 doi:10.1130/0091-7613(1986)14<7:SEOTWA>2.0.CO;2

1165 de Capitani, C., Petrakakis, K., 2010. The computation of equilibrium assemblage
 1166 diagrams with Theriak/Domino software. *American Mineralogist* 95, 1006–1016.
 1167 doi:10.2138/am.2010.3354

1168 de Galdeano, C.S., and Vera, J.A., 1992, Stratigraphic record and palaeogeographical
 1169 context of the Neogene basins in the Betic Cordillera, Spain: *Basin Research*, v. 4, no. 1, p.
 1170 21–35.

1171 Denèle, Y., Lecomte, E., Jolivet, L., Lacombe, O., Labrousse, L., Huet, B., Le Pourhiet, L.,
 1172 2011. Granite intrusion in a metamorphic core complex: The example of the Mykonos

1173 laccolith (Cyclades, Greece). *Tectonophysics* 501, 52–70.
 1174 doi:10.1016/j.tecto.2011.01.013

1175 Dèzes, P., Ziegler, P., 2001. Map of the European Moho. EUCOR URGENT.

1176 Di Luccio, F., Pasyanos, M.E., 2007. Crustal and upper-mantle structure in the Eastern
 1177 Mediterranean from the analysis of surface wave dispersion curves. *Geophysical Journal*
 1178 *International* 169, 1139–1152. doi:10.1111/j.1365-246X.2007.03332.x

1179 Dickinson, W.R., 2002. The Basin and Range Province as a Composite Extensional
 1180 Domain. *International Geology Review* 44, 1–38. doi:10.2747/0020-6814.44.1.1

1181 Diener, J.F.A., Powell, R., White, R.W., Holland, T.J.B., 2007. A new thermodynamic model
 1182 for clino- and orthoamphiboles in the system Na₂O-CaO-FeO-MgO-Al₂O₃-SiO₂-H₂O.
 1183 *Journal of Metamorphic Geology* 25, 631–656. doi:10.1111/j.1525-1314.2007.00720.x

1184 Dilek, Y., Altunkaynak, S., Oner, Z., 2009. Syn-extensional granitoids in the Menderes core
 1185 complex and the late Cenozoic extensional tectonics of the Aegean province. *Geological*
 1186 *Society, London, Special Publications* 321, 197–223. doi:10.1144/SP321.10

1187 Ferrandini, M., Ferrandini, J., Lojze-Pilot, M.-D., Butterlin, J., Cravatte, J., and Janin, M.-C.,
 1188 1998, Le Miocène du bassin de Saint-Florent(Corse): Modalités de la transgression du
 1189 Burdigalien supérieur et mise en évidence du Serravallien: *Geobios*, v. 31, no. 1, p. 125–
 1190 137, doi: 10.1016/S0016-6995(98)80102-2.

1191 Famin, V., Philippot, P., Jolivet, L., and Agard, P., 2004, Evolution of hydrothermal regime
 1192 along a crustal shear zone, Tinos Island, Greece: *Tectonics*, v. 23, no. 5, p. n/a–n/a, doi:
 1193 10.1029/2003TC001509.

1194 Fournier, M., Jolivet, L., Goffé, B., and Dubois, R., 1991, Alpine Corsica Metamorphic Core
 1195 Complex: *Tectonics*, v. 10, no. 6, p. 1173–1186, doi: 10.1029/91TC00894.

1196 Gautier, P., Bozkurt, E., Bosse, V., Hallot, E., and Dirik, K., 2008, Coeval extensional
 1197 shearing and lateral underflow during Late Cretaceous core complex development in the
 1198 Niğde Massif, Central Anatolia, Turkey: *Tectonics*, v. 27, no. 1, p. n/a–n/a, doi:
 1199 10.1029/2006TC002089.

1200 Gautier, P., Bozkurt, E., Hallot, E., Dirik, K., 2002. Dating the exhumation of a
 1201 metamorphic dome: geological evidence for pre-Eocene unroofing of the Nigde Massif
 1202 (Central Anatolia, Turkey). *Geol. Mag.* 139. doi:10.1017/S0016756802006751

1203 Gautier, P., Brun, J.-P., Jolivet, L., 1993. Structure and kinematics of Upper Cenozoic
 1204 extensional detachment on Naxos and Paros (Cyclades Islands, Greece). *Tectonics* 12,
 1205 1180–1194. doi:10.1029/93TC01131

1206 Gerya, T.V., Yuen, D.A., 2003. Characteristics-based marker-in-cell method with
 1207 conservative finite-differences schemes for modeling geological flows with strongly
 1208 variable transport properties. *Physics of the Earth and Planetary Interiors* 140, 293–
 1209 318. doi:10.1016/j.pepi.2003.09.006

1210 Goncuoglu, M.C., Toprak, V., Kuscu, U., Erler, A., Olgun, E., 1991. Geology of the Western
 1211 part of the Central Anatolian Massif, Part 1: Southern Part. Turkish Petroleum
 1212 Corporation

1213 Grasemann, B., Schneider, D.A., Stockli, D.F., Iglseder, C., 2011. Miocene bivergent crustal
 1214 extension in the Aegean: Evidence from the western Cyclades (Greece). *Lithosphere* 4,
 1215 L164.1–39. doi:10.1130/L164.1

1216 Gueydan, F., 2003, Analysis of continental midcrustal strain localization induced by
 1217 microfracturing and reaction-softening: *Journal of Geophysical Research*, v. 108, no. B2,
 1218 p. 2064, doi: 10.1029/2001JB000611.

1219 Gueydan, F., 2004. Mechanics of low-angle extensional shear zones at the brittle-ductile
 1220 transition. *J. Geophys. Res.* 109, B12407. doi:10.1029/2003JB002806

1221 Heymes, T., Bouillin, J.P., Pêcher, A., Monié, P., and Compagnoni, R., 2008, Middle
 1222 Oligocene extension in the Mediterranean Calabro-Peloritan belt (southern Italy):
 1223 Insights from the Aspromonte nappes pile: *Tectonics*, v. 27, no. 2, p. n/a–n/a, doi:
 1224 10.1029/2007TC002157.

1225 Heymes, T., Monié, P., Arnaud, N., Pêcher, A., Bouillin, J.-P., Compagnoni, R., 2010. Alpine
 1226 tectonics in the Calabrian–Peloritan belt (southern Italy): New $^{40}\text{Ar}/^{39}\text{Ar}$ data in the
 1227 Aspromonte Massif area. *Lithos* 114, 451–472. doi:10.1016/j.lithos.2009.10.011

1228 Hoinkes, G., Koller, F., Rantitsch, G., Dachs, E., Hock, V., Neubauer, F., Schuster, R., 1999.
 1229 Alpine metamorphism of the Eastern Alps. Schweizerische Mineralogische Und
 1230 Petrographische Mitteilungen 79, 155–181.

1231 Holland, T.J.B., Powell, R., 2004. An internally consistent thermodynamic data set for
 1232 phases of petrological interest. Journal of Metamorphic Geology 16, 309–343.
 1233 doi:10.1111/j.1525-1314.1998.00140.x

1234 Huang, W.L., and Wyllie, P.J., 1981, Phase relationships of S-type granite with H₂O to 35
 1235 kbar: Muscovite granite from Harney Peak, South Dakota: Journal of Geophysical
 1236 Research: Solid Earth (1978–2012), v. 86, no. B11, p. 10515–10529, doi:
 1237 10.1029/JB086iB11p10515.

1238 Huet, B., Le Pourhiet, L., Labrousse, L., Burov, E., Jolivet, L., 2010. Post-orogenic
 1239 extension and metamorphic core complexes in a heterogeneous crust: the role of crustal
 1240 layering inherited from collision. Application to the Cyclades (Aegean domain).
 1241 Geophysical Journal International 184, 611–625. doi:10.1111/j.1365-
 1242 246X.2010.04849.x

1243 Huet, B., Le Pourhiet, L., Labrousse, L., Burov, E.B., Jolivet, L., 2011. Formation of
 1244 metamorphic core complex in inherited wedges: A thermomechanical modelling study.
 1245 Earth and Planetary Science Letters 309, 249–257. doi:10.1016/j.epsl.2011.07.004

1246 Isik, V., Tekeli, O., 2001. Late orogenic crustal extension in the northern Menderes massif
 1247 (western Turkey): evidence for metamorphic core complex formation. Int J Earth Sci 89,
 1248 757–765. doi:10.1007/s005310000105

1249 Isik, V., Tekeli, O., and Seyitoğlu, G., 2004, The ⁴⁰Ar/³⁹Ar age of extensional ductile
 1250 deformation and granitoid intrusion in the northern Menderes core complex:
 1251 implications for the initiation of extensional tectonics in western Turkey: Journal of
 1252 Asian Earth Sciences, v. 23, no. 4, p. 555–566, doi: 10.1016/j.jseaes.2003.09.001.

1253 Isik, V., Seyitoğlu, G., Çemen, İ., 2003. Ductile–brittle transition along the Alaşehir
 1254 detachment fault and its structural relationship with the Simav detachment fault,
 1255 Menderes massif, western Turkey. Tectonophysics 374, 1–18. doi:10.1016/S0040-
 1256 1951(03)00275-0

1257 Janák, M., Plašienka, D., Frey, M., Cosca, M., Schmidt, S.T., Lupták, B., Méres, Š., 2001.
1258 Cretaceous evolution of a metamorphic core complex, the Veporic unit, Western
1259 Carpathians (Slovakia): P-T conditions and in situ $^{40}\text{Ar}/^{39}\text{Ar}$ UV laser probe dating of
1260 metapelites. *Journal of Metamorphic Geology* 19, 197–216. doi:10.1046/j.0263-
1261 4929.2000.00304.x

1262 Jansen, J., 1977, The geology of Naxos.

1263 Jaoul, O., Tullis, J., Kronenberg, A., 1984. The effect of varying water contents on the
1264 creep behavior of Heavitree quartzite. *Journal of Geophysical Research: Solid Earth*
1265 (1978–2012) 89, 4298–4312. doi:10.1029/JB089iB06p04298

1266 Jolivet, L., Augier, R., Faccenna, C., Negro, F., Rimmelé, G., Agard, P., Robin, C., Rossetti, F.,
1267 and Crespo-Blanc, A., 2008, Subduction, convergence and the mode of backarc extension
1268 in the Mediterranean region: *Bulletin de la Societe Geologique de France*, v. 179, no. 6, p.
1269 525–550, doi: 10.2113/gssgfbull.179.6.525.

1270 Jolivet, L., Brun, J.-P., 2008. Cenozoic geodynamic evolution of the Aegean. *Int J Earth Sci*
1271 99, 109–138. doi:10.1007/s00531-008-0366-4

1272 Jolivet, L., Dubois, R., Fournier, M., Goffé, B., Michard, A., Jourdan, C., 1990. Ductile
1273 extension in alpine Corsica. *Geol* 18, 1007. doi:10.1130/0091-
1274 7613(1990)018<1007:DEIAC>2.3.CO;2

1275 Jolivet, L., Faccenna, C., Goffé, B., Burov, E., and Agard, P., 2003, Subduction tectonics and
1276 exhumation of high-pressure metamorphic rocks in the Mediterranean orogens:
1277 *American Journal of Science*, v. 303, no. 5, p. 353–409, doi: 10.2475/ajs.303.5.353.

1278 Jolivet, L., Faccenna, C., Piromallo, C., 2009a. From mantle to crust: Stretching the
1279 Mediterranean. *Earth and Planetary Science Letters* 285, 198–209.
1280 doi:10.1016/j.epsl.2009.06.017

1281 Jolivet, L., Goffé, B., Monié, P., and Luxey, C.T., 1996, Miocene detachment in Crete and
1282 exhumation P-T-t paths of high-pressure metamorphic rocks: *Tectonics*, v. 15, no. 6, p.
1283 1129–1153.

1284 Jolivet, L., Goffé, B., Monié, P., Luxey, C.T., 1996. Miocene detachment in Crete and
1285 exhumation P-T-t paths of high-pressure metamorphic rocks. *Tectonics* 15, 1129–1153.

1286 Jolivet, L., Labrousse, L., Agard, P., Lacombe, O., Bailly, V., Lecomte, E., Mouthereau, F.,
1287 Mehl, C., 2009b. Rifting and shallow-dipping detachments, clues from the Corinth Rift
1288 and the Aegean. *Tectonophysics* 1–18. doi:10.1016/j.tecto.2009.11.001

1289 Jolivet, L., Lecomte, E., Huet, B., Denèle, Y., Lacombe, O., Labrousse, L., Le Pourhiet, L.,
1290 Mehl, C., 2010a. The North Cycladic Detachment System. *Earth and Planetary Science*
1291 *Letters* 289, 87–104. doi:10.1016/j.epsl.2009.10.032

1292 Jolivet, L., Trotet, F., Monié, P., Vidal, O., Goffé, B., Labrousse, L., Agard, P., Ghorbal, B.,
1293 2010b. Along-strike variations of P–T conditions in accretionary wedges and syn-
1294 orogenic extension, the HP–LT Phyllite–Quartzite Nappe in Crete and the Peloponnese.
1295 *Tectonophysics* 480, 133–148. doi:10.1016/j.tecto.2009.10.002

1296 Karabulut, H., Paul, A., Ergün, T.A., Hatzfeld, D., Childs, D.M., and Aktar, M., 2013, Long-
1297 wavelength undulations of the seismic Moho beneath the strongly stretched Western
1298 Anatolia: *Geophysical Journal International*, v. 194, no. 1, p. 450–464, doi:
1299 10.1093/gji/ggt100.

1300 Kiliyas, A., Fassoulas, C., Mountrakis, D., 1994. Tertiary extension of continental crust and
1301 uplift of Psiloritis metamorphic core complex in the central part of the Hellenic Arc
1302 (Crete, Greece). *Geol Rundsch* 83, 417–430. doi:10.1007/BF00210555

1303 Kirby, S.H., Kronenberg, A.K., 1987. Rheology of the lithosphere: Selected topics. *Reviews*
1304 *of Geophysics* 25, 1219–1244. doi:10.1029/rg025i006p01219

1305 Koulakov, I., Sobolev, S.V., 2006. Moho depth and three-dimensional P and S structure of
1306 the crust and uppermost mantle in the Eastern Mediterranean and Middle East derived
1307 from tomographic inversion of local ISC data. *Geophysical Journal International* 164,
1308 218–235. doi:10.1111/j.1365-246X.2005.02791.x

1309 Krohe, A., and Mposkos, E., 2002, Multiple generations of extensional detachments in the
1310 Rhodope Mountains (northern Greece): evidence of episodic exhumation of high-
1311 pressure rocks, in Blundell, D.J., Quadf, von, A., and Gosen, von, W. eds., *Spec. Pub. Geol.*
1312 *Soc.* p. 151–178.

1313 Kuhlemann, J., Frisch, W., Dunkl, I., Kázmér, M., Schmiedl, G., 2004. Miocene siliciclastic
1314 deposits of Naxos Island: Geodynamic and environmental implications for the evolution

1315 of the southern Aegean Sea (Greece). Geological Society of America Special Papers 378,
1316 51–65. doi:10.1130/0-8137-2378-7.51

1317 Laurent, V., Beaudoin, A., Jolivet, L., Arbaret, L., Augier, R., Rabillard, A., Menant, A., 2015.
1318 Interrelations between extensional shear zones and synkinematic intrusions: The
1319 example of Ikaria Island (NE Cyclades, Greece). Tectonophysics 651-652, 152–171.
1320 doi:10.1016/j.tecto.2015.03.020

1321 Le Maitre, R.W., 1976. The Chemical Variability of some Common Igneous Rocks. Journal
1322 of Petrology 17, 589–598. doi:10.1093/petrology/17.4.589

1323 Le Pourhiet, L., Burov, E., Moretti, I., 2004. Rifting through a stack of inhomogeneous
1324 thrusts (the dipping pie concept). Tectonics 23, n/a–n/a. doi:10.1029/2003TC001584

1325 Liati, A., 2005, Identification of repeated Alpine (ultra) high-pressure metamorphic
1326 events by U–Pb SHRIMP geochronology and REE geochemistry of zircon: the Rhodope
1327 zone of Northern Greece: Contributions to Mineralogy and Petrology, v. 150, no. 6, p.
1328 608–630, doi: 10.1007/s00410-005-0038-3.

1329 Lips, A.L.W., Cassard, D., Sebilir, H., Yilmaz, H., and Wijbrans, J.R., 2001, Multistage
1330 exhumation of the Menderes Massif, western Anatolia (Turkey): International Journal of
1331 Earth Sciences, v. 89, no. 4, p. 781–792, doi: 10.1007/s005310000101.

1332 Lister, G.S., Davis, G.A., 1989. The origin of metamorphic core complexes and detachment
1333 faults formed during Tertiary continental extension in the northern Colorado River
1334 region, USA. Journal of Structural Geology 11, 65–94. doi:10.1016/0191-
1335 8141(89)90036-9

1336 Lonergan, L., Platt, J.P., 1995. The Malaguide-Alpujarride boundary: a major extensional
1337 contact in the Internal Zone of the eastern Betic Cordillera, SE Spain. Journal of
1338 Structural Geology. doi:10.1016/0191-8141(95)00070-T

1339 Marone, F., van der Meijde, M., van der Lee, S., Giardini, D., 2003. Joint inversion of local,
1340 regional and teleseismic data for crustal thickness in the Eurasia–Africa plate boundary
1341 region. Geophysical Journal International 154, 499–514. doi:10.1046/j.1365-
1342 246X.2003.01973.x

1343 Marroni, M., Pandolfi, L., 2003. Deformation history of the ophiolite sequence from the
 1344 Balagne Nappe, northern Corsica: insights in the tectonic evolution of Alpine Corsica.
 1345 *Geol. J.* 38, 67–83. doi:10.1002/gj.933

1346 Mattei, M., Cipollari, P., Cosentino, D., Argentieri, A., Rossetti, F., Speranza, F., and Di
 1347 Bella, L., 2002, The Miocene tectono-sedimentary evolution of the southern Tyrrhenian
 1348 Sea: stratigraphy, structural and palaeomagnetic data from the on-shore Amantea basin
 1349 (Calabrian Arc, Italy): *Basin Research*, v. 14, no. 2, p. 147–168, doi: 10.1046/j.1365-
 1350 2117.2002.00173.x.

1351 Mezri, L., Le Pourhiet, L., Wolf, S., Burov, E., 2015. New parametric implementation of
 1352 metamorphic reactions limited by water content, impact on exhumation along
 1353 detachment faults. *LITHOS* 236–237, 287–298. doi:10.1016/j.lithos.2015.08.021

1354 Michard, A., Negro, F., Saddiqi, O., 2006. Pressure–temperature–time constraints on the
 1355 Maghrebide mountain building: evidence from the Rif–Betic transect (Morocco, Spain),
 1356 Algerian correlations, and geodynamic implications. *Comptes Rendus Geoscience* 338,
 1357 92–114. doi:10.1016/j.crte.2005.11.011

1358 Miller, E.L., Gans, P.B., Garing, J., 1983. The Snake Range Décollement: An exhumed Mid-
 1359 Tertiary ductile-brittle transition. *Tectonics* 2, 239–263.
 1360 doi:10.1029/TC002i003p00239

1361 Moresi, L., Dufour, F., Mühlhaus, H.B., 2003. A Lagrangian integration point finite
 1362 element method for large deformation modeling of viscoelastic geomaterials. *Journal of*
 1363 *Computational Physics* 184, 476–497. doi:10.1016/S0021-9991(02)00031-1

1364 Mposkos, E., 1998, Cretaceous and tertiary tectonometamorphic events in Rhodope zone
 1365 (Greece): petrological and geochronological evidences: *Bull Geol Soc Greece*.

1366 Okay, A.I., Satir, M., 2000. Coeval plutonism and metamorphism in a latest Oligocene
 1367 metamorphic core complex in northwest Turkey. *Geol. Mag.* 137, 495–516.

1368 Pamić, J., Balen, D., Herak, M., 2012. Origin and geodynamic evolution of Late Paleogene
 1369 magmatic associations along the Periadriatic-Sava-Vardar magmatic belt.
 1370 <http://dx.doi.org/10.1080/09853111.2002.10510755> 15, 209–231.
 1371 doi:10.1080/09853111.2002.10510755

1372 Pavelić, D., 2001, Tectonostratigraphic model for the North Croatian and North Bosnian
 1373 sector of the Miocene Pannonian Basin System: *Basin Research*, v. 13, no. 3, p. 359–376,
 1374 doi: 10.1046/j.0950-091x.2001.00155.x.

1375 Piller, W.E., Harzhauser, M., and Mandic, O., 2007, Miocene Central Paratethys
 1376 stratigraphy—current status and future directions: *Stratigraphy*, v. 13, p. 359–376.

1377 Platt, J.P., Behrmann, J.H., Martínez, J.M.M., Vissers, R.L.M., 1984. A zone of mylonite and
 1378 related ductile deformation beneath the alpujarride nappe complex, betic cordilleras, S.
 1379 Spain. *Geol Rundsch* 73, 773–785. doi:10.1007/BF01824981

1380 Platzman, E., and Platt, J.P., 2004, Kinematics of a twisted core complex: Oblique axis
 1381 rotation in an extended terrane (Betic Cordillera, southern Spain): *Tectonics*, doi:
 1382 10.1029/2003TC001549.

1383 Plunder, A., Agard, P., Chopin, C., Okay, A.I., 2013. Geodynamics of the Tavşanlı zone,
 1384 western Turkey: Insights into subduction/obduction processes. *Tectonophysics* 608,
 1385 884–903. doi:10.1016/j.tecto.2013.07.028

1386 Poliakov, A.N.B., Podladchikov, Y., Talbot, C., 1993. Initiation of salt diapirs with
 1387 frictional overburdens: numerical experiments. *Tectonophysics* 228, 199–210.
 1388 doi:10.1016/0040-1951(93)90341-G

1389 Rabillard, A., Arbaret, L., Jolivet, L., Le Breton, N., Gumiaux, C., Augier, R., Grasemann, B.,
 1390 2015. Interactions between plutonism and detachments during metamorphic core
 1391 complex formation, Serifos Island (Cyclades, Greece). *Tectonics* 34, 1080–1106.
 1392 doi:10.1002/2014TC003650

1393 Ranalli, G., Murphy, D.C., 1987. Rheological stratification of the lithosphere.
 1394 *Tectonophysics* 132, 281–295. doi:10.1016/0040-1951(87)90348-9

1395 Ratschbacher, L., Frisch, W., Linzer, H.G., and Merle, O., 1991, Lateral extrusion in the
 1396 eastern Alps, Part 2: Structural analysis: *Tectonics*, v. 10, no. 2, p. 257–271, doi:
 1397 10.1029/90TC02623.

1398 Rey, P.F., Teyssier, C., Kruckenberg, S.C., Whitney, D.L., 2011. Viscous collision in channel
 1399 explains double domes in metamorphic core complexes. *Geol* 39, 387–390.
 1400 doi:10.1130/G31587.1

1401 Rey, P.F., Teyssier, C., Whitney, D.L., 2009. Extension rates, crustal melting, and core
1402 complex dynamics. *Geol* 37, 391–394. doi:10.1130/G25460A.1

1403 Ring, U., Johnson, C., Hetzel, R., and Gessner, K., 2003, Tectonic denudation of a Late
1404 CretaceousTertiary collisional belt: regionally symmetric cooling patterns and their
1405 relation to extensional faults in the Anatolide belt of western Turkey: *Geological*
1406 *Magazine*, v. 140, no. 4, p. 421–441, doi: 10.1017/S0016756803007878.

1407 Rohais, S., Eschard, R., Ford, M., Guillocheau, F., and Moretti, I., 2007, Stratigraphic
1408 architecture of the Plio-Pleistocene infill of the Corinth Rift: Implications for its
1409 structural evolution: *Tectonophysics*, v. 440, no. 1-4, p. 5–28, doi:
1410 10.1016/j.tecto.2006.11.006.

1411 Rossetti, F., Goffé, B., Monié, P., Faccenna, C., Vignaroli, G., 2004. Alpine orogenic P-T-t-
1412 deformation history of the Catena Costiera area and surrounding regions (Calabrian Arc,
1413 southern Italy): The nappe edifice of north Calabria revised with insights on the
1414 Tyrrhenian-Apennine system formation. *Tectonics* 23, n/a–n/a.
1415 doi:10.1029/2003tc001560

1416 Royden, L.H., 1993, Evolution of retreating subduction boundaries formed during
1417 continental collision: *Tectonics*, v. 12, no. 3, p. 629–638, doi: 10.1029/92TC02641.

1418 Saadallah, A., Caby, R., 1996. Alpine extensional detachment tectonics in the Grande
1419 Kabylie metamorphic core complex of the Maghrebides (northern Algeria).
1420 *Tectonophysics* 267, 257–273. doi:10.1016/S0040-1951(96)00101-1

1421 Sanchez-Gomez, M., Avigad, D., and Heimann, A., 2002, Geochronology of clasts in
1422 allochthonous Miocene sedimentary sequences on Mykonos and Paros Islands:
1423 implications for back-arc extension in the Aegean Sea: *Journal of the Geological Society*,
1424 v. 159, no. 1, p. 45–60, doi: 10.1144/0016-764901031.

1425 Schenker, F.L., Gerya, T., and Burg, J.P., 2012, Bimodal behavior of extended continental
1426 lithosphere: Modeling insight and application to thermal history of migmatitic core
1427 complexes: *Tectonophysics*, v. 579, no. C, p. 88–103, doi: 10.1016/j.tecto.2012.07.002.

1428 Sorel, D., 2000, A Pleistocene and still-active detachment fault and the origin of the
1429 Corinth–Patras rift, Greece: *Geology*, v. 28, p. 83–86.

- 1430 Souche, A., Medvedev, S., Andersen, T.B., and Dabrowski, M., 2013, Shear heating in
1431 extensional detachments: Implications for the thermal history of the Devonian basins of
1432 W Norway: *Tectonophysics*, v. 608, p. 1073–1085, doi: 10.1016/j.tecto.2013.07.005.
- 1433 Soukis, K., and Stockli, D.F., 2013, Structural and thermochronometric evidence for
1434 multi-stage exhumation of southern Syros, Cycladic islands, Greece: *Tectonophysics*, v.
1435 595-596, p. 148–164, doi: 10.1016/j.tecto.2012.05.017.
- 1436 Sümer, Ö., İnci, U., and Sözbilir, H., 2013, Tectonic evolution of the Söke Basin: Extension-
1437 dominated transtensional basin formation in western part of the Büyük Menderes
1438 Graben, Western Anatolia, Turkey: *Journal of Geodynamics*, doi: 10.3166/ga.19.363-
1439 390.
- 1440 Tan, E., Lavier, L.L., Van Avendonk, H.J.A., Heuret, A., 2012. The role of frictional strength
1441 on plate coupling at the subduction interface. *Geochem. Geophys. Geosyst.* 13, n/a–n/a.
1442 doi:10.1029/2012GC004214
- 1443 Tiberi, C., Diament, M., Lyon-Caen, H., King, T., 2001. Moho topography beneath the
1444 Corinth Rift area (Greece) from inversion of gravity data. *Geophysical Journal*
1445 *International* 145, 797–808. doi:10.1046/j.1365-246x.2001.01441.x
- 1446 Tirel, C., Brun, J.-P., Burov, E., 2008. Dynamics and structural development of
1447 metamorphic core complexes. *J. Geophys. Res.* 113, B04403.
1448 doi:10.1029/2005JB003694
- 1449 Tirel, C., Brun, J.P., Sokoutis, D., 2006. Extension of thickened and hot lithospheres:
1450 Inferences from laboratory modeling. *Tectonics* 25, n/a–n/a.
1451 doi:10.1029/2005TC001804
- 1452 Trotet, F., Jolivet, L., Vidal, O., 2001. Tectono-metamorphic evolution of Syros and Sifnos
1453 islands (Cyclades, Greece). *Tectonophysics* 338, 179–206. doi:10.1016/S0040-
1454 1951(01)00138-X
- 1455 Ustaszewski, K., Kounov, A., Schmid, S.M., Schaltegger, U., Krenn, E., Frank, W.,
1456 Fügenschuh, B., 2010. Evolution of the Adria-Europe plate boundary in the northern
1457 Dinarides: From continent-continent collision to back-arc extension. *Tectonics* 29.,
1458 doi:10.1029/2010TC002668

1459 van Hinsbergen, D.J.J., 2010. A key extensional metamorphic complex reviewed and
 1460 restored: The Menderes Massif of western Turkey. *Earth-Science Reviews* 102, 60–76.
 1461 doi:10.1016/j.earscirev.2010.05.005

1462 Vanderhaeghe, O., 2004. Structural development of the Naxos migmatite dome.
 1463 *Geological Society of America Special Papers* 380, 211–227. doi:10.1130/0-8137-2380-
 1464 9.211

1465 Whitney, D.L., Dilek, Y., 1997. Core complex development in central Anatolia, Turkey.
 1466 *Geol* 25, 1023. doi:10.1130/0091-7613(1997)025<1023:CCDICA>2.3.CO;2

1467 Whitney, D.L., Teyssier, C., Rey, P., and Buck, W.R., 2013, Continental and oceanic core
 1468 complexes: *Geological Society of America Bulletin*, v. 125, no. 3-4, p. 273–298, doi:
 1469 10.1130/B30754.1.

1470 Wu, G., Lavier, L.L., Choi, E., 2015. Modes of continental extension in a crustal wedge.
 1471 *Earth and Planetary Science Letters* 421, 89–97. doi:10.1016/j.epsl.2015.04.005

1472 Yamato, P., Agard, P., Burov, E., Le Pourhiet, L., Jolivet, L., Tiberi, C., 2007. Burial and
 1473 exhumation in a subduction wedge: Mutual constraints from thermomechanical
 1474 modeling and natural P-T-t data (Schistes Lustrés, western Alps). *J. Geophys. Res.* 112,
 1475 B07410. doi:10.1029/2006JB004441

1. Locality		A- Betics, Spain		B- Beni Bousera (B.B.), Morocco
2. Upper plate	2.1 Syntectonic basin		continental to marine (de Galdeano and Vera, 1992)	
	2.2 Basal lithologies	Malaguides : palaeozoic sandstones and slates	Alpujarrides : phyllites, quartzites and graphitic micaschists	Ghomarides : paleozoic metasediments
	2.3 Peak metamorphic conditions	lower anchizone (variscan)	HPA in Herradura to GS in Lujar Gador (Azañón and Crespo-Blanc, 2000; Jolivet et al., 2010b)	GS ?
	2.4 Brittle deformation		2 sets of LANF (Platzman and Platt, 2004; Augier et al., 2005)	
3. Detachment Zone	3.1 Abbreviation/name	MAD/Malaguides-Alpujarrides Detachment (Platzman and Platt, 2004)	MFSZ/Mecina-Filabres Shear Zone (Platt et al., 1984)	ZaFZ/Zaouia Fault Zone
	3.2 Associated lithologies	Calc-mylonites and fault gouge (Lonergan and Platt, 1995)	mylonites and gouges	« brittle-ductile fault » (Michard et al., 2006)
	3.3 Relationships to heritage	reactivated thrust	reactivated thrust	oblique to thrusts
4. Lower plate	4.1 Top lithologies	Alpujarrides : phyllites, quartzites and graphitic micaschists	Nevado-Filabrides : permo-triassic graphitic schists (Bedar Macael)	Sebtides : permo-triassic metasediments (Filali schists & Federico)
	4.2 Peak metamorphic conditions*	HPA in Herradura (Alpu1) to GS in Lujar Gador (Alpu4) (Azañón and Crespo-Blanc, 2000)	E in Bedar Macael (NF1) to GS in Ragua (NF3, Augier et al., 2005)	HPA in Federico (BB3) HPG in B.B. kinzigites (BB2)
	4.3 Brittle deformation			
5. Migmatites		at the base of Herradura	no HT overprint	
6. Intrusives	6.1 Type	mafic dikes in upper and lower plates		leucogranite dikes in B.B. lower u.
	6.2 Relation to ductile deformation	late ? magmatism		undeformed
7. Ophiolitic units	7.1 Type			peridotites
	7.2 Position			in the dome core
8. Timing in orogeny		syn-orogenic initiation (Jolivet et al., 2008)	mainly post-orogenic (Jolivet et al., 2008)	possibly syn-orogenic (Michard et al., 2006)
9. Geodynamic boundary conditions		change of slab retreat direction from southward to westward		Alboran slab retreat

Table 1a

1.	C- Grande Kabylie, Algeria	D- Edough, Algeria	E- Aspromonte, S. Italy	F- Catena Costiera, S. Italy
2.1	Kabylian oligo-miocene marine conglomerates (OMK)	none	continental to marine Stilo-Capo d'Orlando Formation (SCOF) (Bonardi et al., 1980)	Miocene to Q. marine b. (e. g. Amantea b. (Mattei et al., 2002)
2.	2.2 pre-permian phyllites	Tellian and Green Schist facies Unit GFU (Caby et al., 2001)	Stilo u. : para- and orthogneiss	Upper Ophiolitic U. (UOU) : ophiolitic melange & Verbicaro carbonates (Bonardi et al., 2001)
	2.3 variscan GS	GS	variscan GS to A, no Alpine overprint (Heymes et al., 2008)	GS
	2.4 normal faults in Central and Eastern Grande Kabylie			tilted blocks at the base of basin
3.1	KaD/Kabylian Detachment (Saadallah and Caby, 1996)	Ed/~KaD, major metamorphic gap	Asp/	CatCo/
3.	3.2 mylonites and younger cataclastes	GS mylonites, fibrolites bearing ultra-mylonites at base of APU	mylonitic gneiss	mylonites and cataclasites (Rossetti et al., 2004)
	3.3	reactivated Tellian thrust	reactivated thrust	
4.	4.1 Sidi Ali Boud Nab massif: Micaschists and orthogneiss	high grade metasedimentary U. (HGMU) Amphibolite-Peridotite U. (APU) Migmatite-Granite U. (MGU)	Aspromonte u. : para and orthogneiss	Continental metaclastics and marbles Lower Ophiolitic U. (LOU)
	4.2 A (GK1 & GK2)	HPG	Alpine BS over variscan LPA	BS (Rossetti et al., 2004)
	4.3			
5.	anatectic metapelites	syn-tectonic anatexis in MGU	none	none
6.	6.1 aplo-pegmatites and leucogranites	leucogranites	none	none
	6.2 syn-kinematic aplites / late granites	syn-kinematic in MGU		
7.	7.1	peridotites at the base of APU (Bosch et al., 2014)	none	ophiolitic melanges, pillow-lavas and picrites in Verbicaro
	7.2	top lower plate and lenses in core		in lower and upper plates
8.	post-orogenic	late post-orogenic	syn-to immediate post-o. (Heymes et al., 2010)	post-orogenic (Jolivet et al., 2008)
9.	Alboran slab retreat	Alboran slab retreat or later slab dynamics (Bruguier, 2009)	stretching oblique to inferred slab retreat direction	E-W stretching parallel to local slab retreat in miocene

Table 1b

1.	G- Alpi Apuane, N. Italy	H- Elba, Thyrrhenian sea	I- Corsica, W. Mediterranean
2.1	marine « Breccie delle Versilia » (Carmignani and Kligfield, 1990)	-	-
2.2	Tuscan nappes : carbonates with flysch on top	Upper Toscan ophiolite and flysch u.	Balagne-Nebbio unit : ophiolitic unit
2.3	no alpine metamorphism	weak thermometamorphism	non metamorphic
2.4	H & LANF rooting on detachment	Zuccale Fault : LANF with cataclasites & calcmylonites crosscuts CaDZ (Collettini and Holdsworth, 2004)	late normal faults (D5, Marroni and Pandolfi, 2003)
3.1	CalCa/Calcare Cavernoso (Carmignani and Kligfield, 1990)	CaDZ/Capanne Detachment Zone (Daniel and Jolivet, 1995)	TeSZ/Tenda Upper Shear Zone (Jolivet et al., 1990; Gueydan, 2003; Fournier et al., 1991)
3.2	mylonites & cataclasites from anhydrite and dolostone	mylonites within thermometamorphic aureole	mylonites
3.3	reactivated thrust	oblique to thrusts	reactivated basal thrust (D4, Marroni and Pandolfi, 2003)
4.1	Massa & « Autochthonous » u. : paleozoic phyllites and metasandstones	Lower Tuscan u. : Calamite paleozoic schists	Schistes Lustrés (SL) : calcschists, ophiolitic and continental basement slices
4.2	BS (AA1 and AA2)	GS and thermometamorphism	BS to E
4.3			BS (Te)
5.	none	none	none
6.1	volcanism 7 Ma younger	granodiorite (Monte Capanne)	Sisco lamproïtes in SL
6.2	(Carmignani et al., 2001)		18 Ma younger than extension (Beccaluva et al., 2004)
7.1	Liguride ophiolites		Nebbio and SL ophiolites
7.2	top of upper plate, limited extensional reactivation		uppermost (Balagne)
8.	post-orogenic		post-orogenic
9.	Thyrrhenian slab retreat		Thyrrhenian slab retreat

Table 1c

1.	J- Motajica window, Bosnia and Hercegovina	K- Rechnitz window area, Austria/Hungary	L- Central Western Carpathians, Slovakia	M- Southern Rhodope, Greece
2.1	Ottangian syn-rifts sediments : alluvial to marine } (Pavelić, 2001)	Fürstenfeld (W) and Danube (E) basins : transitional Karpatian deposits (Piller et al., 2007)		
2.	Sava Zone metasediments and ophiolites at the base of Tisza-Dacia u. (Ustaszewski et al., 2010)	Wechsel u. (base of Autro-Alpine) gneiss and Palaeozoic carbonates (Cao et al., 2013)	Gemic u. : palaeozoic basement rocks	Vertiskos Gneiss Complex : gneiss + Therma and Volvi gabbros
2.3	A to GS	GS	low grade	(U)HP (Liati, 2005)
2.4	late brittle normal faults	HANF		upper Miocene faulting
3.1	MoD/Motajica Detachment (Ustaszewski et al., 2010)	ReW/	LuL/Lubenik Line (Janák et al., 2001)	KerD/Kerdylion Detachment (Brun and Sokoutis, 2007)
3.	Upper GS foliation in paragneiss	Serpentinite and quartz-micas mylonites (Cao et al., 2013)	mylonites from cover u. and topmost intrusive granitoid	mylonites and ultra-mylonites
3.3	Partly reactivated thrust	possibly reactivated basal AA thrust	reactivated thrust	
4.1	Accretionary wedge complex and Motajica granite	Rechnitz ophiolite u. Penninic greenschist and serpentinites.	Veporic u. : pre-Alpine basement and phyllite cover	Southern Rhodope Core Complex : marbles and orthogneiss
4.	A 5mot) and local contact metamorphism	BS (Re1) and subsequent GS (Re2)	A (Ve) and contact metamorphism	A (Barrovian gradient SRho)
4.3		Chlorite breccias in borehole		upper Miocene faulting
5.	none	none	hercynian remnants	in the Kerdylion massif
6.1	S- to I-type Motajica granite (Pamić et al., 2012)	none	Rochovce granite	Vrondou and Symvolon granodiorites
6.	crystallization 5 Myrs prior to extension onset		syn-tectonic	syn-kinematic
7.1	Western Vardar ophiolite complex	Serpentinite, metagabbro, ophicarbonates	none	Volvi and Therma metabasites : gabbros and basalts
7.2	Upper level of upper plate	within lower plate		base of the upper plate
8.	post-orogenic Pannonian extension	post-orogenic Pannonian extension ?	Cretaceous syn-orogenic	post-orogenic
9.	Carpathian subduction roll-back (Royden, 1993)	Eastward lateral extrusion (Ratschbacher et al., 1991) and Carpathian roll-back	Growth of collision wedge after Meliata suture	

TableId

1.	N- Eastern Rhodope, Greece	O- Northern Cyclades, Greece (Andros-Tinos-Mikonos-Ikaria)	P- Syros, Greece	Q- Central Cyclades, Greece (Naxos Paros)
2.1	Krumovgrad grp : colluvial to marine (Bonev et al., 2006)	marine molasses then conglomerates on Mykonos (Sanchez-Gomez et al., 2002)		from marine sanstones and pelites toward fluvial deposits (Kuhlemann et al., 2004; Bargnesi et al., 2013)
2.	Kimi Complex : high grade rocks + meta-ophiolitic lenses.	Upper Cycladic Nappe: ophiolitic complex (UCN)	Vari unit and "Upper Unit" metabasites, gneiss and serpentinites (Soukis and Stockli, 2013)	ophiolite nappe and neritic limestones
2.3	(U?)HP 73 Ma (Krohe and Mposkos, 2002)	GS	HT (Jolivet and Brun, 2008)	unmetamorphosed
2.4	HANF offsetting detachment	HANF branching on detachment	late brittle LANF	synthetic normal faults
3.	ToKeD/Tokachka and Kechros detachments	NCDS/North Cycladic Detachment System (Jolivet et al., 2010a)	VaD/Vari Detachment	NaPaD/Naxos Paros Detachment (Gautier et al., 1993)
3.2	mylonites	mylonites, cataclasites and breccias	mylonites and late breccias	mylonites and ultra-mylonites
3.3		partly reactivating UCN thrust		reactivated thrust
4.	Kesebir-Kardamos and Kechros complexes: migmatitic gneiss and orthogneiss (Bonev et al., 2006)	Cycladic Blueschist Unit: metapelites and marbles on granitic basement	Cycladic Blueschist Unit : metapelites, marbles metaflysch (Bonneau, 1984)	Pre-Alpine Gneiss basement with marbles and schists.
4.2	lower E (ERho1,Mposkos, 1998)	E-BS (Tin1 to 3) then A (Mykonos and Ikaria; Ik)	E (Syr1)-BS (Syr2) (Trotet et al., 2001)	E then A (Nax 1&2, Avigad, 1998)
4.3	HANF offsetting detachment			
5.	in Kardamos complex	on Dilos, Mykonos and Ikaria	none	pervasive melting (Vanderhaeghe, 2004)
6.	Papikion granitoid	I-type granites on Tinos, Mykonos and Ikaria (JOLIVET)	none	granodiorite (Naxos) and leucogranites (Paros)
6.2	presumably syn-tectonic	syn- to late-tectonic		syn-extensional leucogranite
7.	Metaophiolite lenses	serpentinites, gabbros and basalts	metabasites of the Upper Unit	gabbros and ultramafics (Jansen, 1977)
7.2	at the base of upper plate (Burg et al., 1996)	at the base of the upper plate	below early then above late detachment (Soukis and Stockli, 2013)	at the base of the upper plate
8.	syn-orogenic	post-orogenic	mainly syn-orogenic	post-orogenic
9.	Vardar ocean closure	Aegean back-arc extension		Aegean back-arc extension

Table1e

1.	R- SouthWestern Cyclades, Greece (Kea-Kythnos-Serifos)	S- Kazdag, Turkey	T- Menderes, Turkey		
2.1	possibly on Milos	Küçükkuyu Fm : fluvio-lacustrine to turbidites (Bonev and Beccaletto, 2007)	Late deposits only	lacustrine to shallow marine supra-detachment basins (Sümer et al., 2013)	
2.	Pelagonian unit : serpentinite and schists	Cetmi melange : opiolitic melange (Beccaletto and Steiner, 2005)	Ismir-Ankara Ophiolite, HP-LT nappes at the base	Northern and Southern Menderes	
2.3	A-GS	E blocks in the melange	E-BS in Afyon, Tavsanli and Oren nappes (Plunder et al., 2013)	A	
2.4	HANF	HANF crosscutting detachment	HANF bounding Simav graben	Half-grabens on both sides	
3.	3.1	WCDS/ Western Cycladic Detachment System (Grasemann et al., 2011)	AlSeD/ Alakeçi SZ (N) and Selale Detachment (S)	SiD/ Simav Detachment (Isik and Tekeli, 2001)	AlBuD/ Alasehir & Buyuk Detachments
	3.2	(ultra)mylonites from marbles and schists	mylonites, metaserpentinites and breccias	mylonites	rmylonites only along Alasehir (Isık et al., 2003) abundant cataclasites
	3.3	reactivated thrusts ?	thrust ?	reactivated thrusts	cut through Menderes units
4.	4.1	Cycladic Blueschist Unit	High grade basement	Northern Menderes : Selimiye metasedimentary unit on top	Central Menderes
	4.2	BS then GS	A (Kaz, Okay & Satir, 2000)	A (Me4)	A
	4.3	HANF			
5.	none	at the top of the dome	inherited in the core (Schuiling, 1962)	?	
6.	6.1	I-type granodiorite (Serifos and Lavrion)	Evciler calc-alkaline granite	Early northern MEG	Later central MEG
	6.2	syn- to late tectonic	intrudes ASZ	syntectonic to SiD (Isik et al., 2004)	syn-extensional (Dilek et al., 2009)
7.	7.1	serpentinites	Cetmi, Karakaya, and Denizgören	Ismir-Ankara Ophiolite	none
	7.2	within the upper plate	at the base of the upper plate	Within the upper plate	
8.	post-orogenic	post-orogenic	post-orogenic	later post-orogenic	
9.	Aegean back-arc extension	Anatolian back-arc extension	Eastern termination of NCDS (JOLIVET)	Intra-Menderes stretching (van Hinsbergen, 2010)	

Table1f

1.	U- Corinth Rift, Greece	V- Crete, Greece	W- Nigde, Turkey
2.1	fluvio-lacustrine toward marine (Rohais et al., 2007)	possibly lacustrine breccias at the base of Neogene basins (Jolivet et al., 1996)	shallow marine deposits in the late Ulukisla basin (Gautier et al., 2008)
2.	Gavrovo-Tripolitza (GT) carbonates with Tyros beds (TB) shales at the base	Gavrovo-Tripolitza carbonates	Massive ophiolite, mostly gabbros (Goncuoglu et al., 1991)
2.3	GS (Jolivet et al., 2010b)	unmetamorphosed	GS at the base
2.4	Zaroukla cataclastic devollement between GT and TB (Sorel, 2000)	HANF cutting across DZ	
3.1	CrD1/Cretan Detachment (Jolivet et al., 2009)	CrD2/ (Kilias et al., 1994)	NiD
3.2	high strain phyllites	mylonites from the upper plate	mylonites to ultracataclasites
3.3	reactivated thrust	reactivated thrust	recativated obduction thrust
4.1	Phyllites-Quartzites	Phyllites-Quartzites nappe on top of Plattenkalk	Gumusler Formation : migmatites
4.2	BS	BS	A (Nig, Whitney and Dilek, 1997)
4.3	Rift faults cutting across detachment		
5.	none	none	most of the lower plate
6.1	none	none	Uçkapili leucogranite
6.2			mainly post-tectonic (Gautier et al., 2002)
7.1	none	ophiolite Asterousia nappe	massive ophiolite in the upper plate,
7.2		on top of upper plate	dilacerated slivers in the lower plate
8.	syn- to post-orogenic	syn-orogenic (Jolivet et al., 1996)	syn-orogenic ?
9.	early Aegean extension	Aegean back-arc extension	Central Anatolian edge extension

Table1g

Locality	A	B	C	D	E	F	G	H	I	J	K	L	MN	O	P	Q	R	S	T	U	V	W				
Detachment	MAD	MFSZ	ZaFZ	KaD	Ed	Asp	CatCo	CalCa	CaDZ	Bal	TeSZ	MoD	ReW	LL	KnD	ToKeD	NCDS	VD	NPD	WCDS	AlSeD	SiD	AlBuD	CrD1	CrD2	NiD
Mylonites	c	✓	✓	✓	✓	✓	c	c	✓	?	✓	✓	✓	✓	✓	✓	✓	?	✓	✓	✓	✓	?		✓	✓
1-Migmatites				✓	✓										✓	✓			✓		✓	✓				✓
2-HT lower plate			✓	✓	✓							✓		✓	✓	✓			✓		✓	✓	✓			✓
3-synD intrusives									✓			✓			✓	✓	✓		✓			✓	✓			✓
4-HP-LT in the l. plate	✓	✓				✓	✓	✓		✓	✓		✓				✓	✓	✓	✓		✓		✓	✓	
5-Resistive u. plate								✓		✓	✓	✓				✓					✓	✓				✓
syn or post-orogenic initiation	S	P	S	P	P	P	P	P	P	P	P	S		S	P	S	P	S	P	P	P	P	P	P	S	S

		Crust		Mantle	
density at 25°C (kg.m⁻³)		2700		3300	
thermal conductivity (W.m⁻¹.K⁻¹)		2.5		3.3	
Thermal expansion coefficient (K⁻¹)			3.0 x 10 ⁻⁵		
Specific heat (J.kg⁻¹.K⁻¹)			10 ³		
Shear modulus (GPa)			30		
Poisson's ratio			0.25		
Mohr-Coulomb criterion			20		
Cohesion (MPa)			20		
Friction angle (°)			30		
A (MPa⁻ⁿ.s⁻¹)	1.26 10 ⁻³	6.7 10 ⁻⁷	2.51 10 ⁻⁴	2.5 x 10 ⁴	
n	2.4	2.4	2.3	3.5	
E (kJ.mol⁻¹)	219	156	154	532	
Lithology	quartz-diorite	quartzite	wet quartzite	dunite	

table 3

Model	Ductile-brittle transition depth	Initial Moho temperature	Intra-crustal Strength Contrast	Integrated Lithospheric Strength	Extension Mode
units	[km]	[°C]		[MPa x km]	
Homogeneous crust :					
upper/lower crust _thickness [km]_Tbottom [° C] d: quartz-diorite, q: quartzite, w: wet quartzite					
d/d_30_1000	26.4	398	4	33739	R
d/d_30_1200	22.3	477	17	20860	R
d/d_30_1400	19.4	557	61	12846	R
d/d_45_1000	27.5	569	103	23239	X
d/d_45_1200	23.3	683	413	9688	WR
d/d_45_1400	20.3	797	1232	5437	WR
d/d_60_1000	28.7	725	822	11564	WR
d/d_60_1200	24.3	870	2832	7244	DDC
d/d_60_1400	21.1	1015	7304	5426	DDC
q/q_30_1000	18.4	398	31	30412	R
q/q_30_1200	15.7	477	93	18245	R
q/q_30_1400	13.7	557	223	10834	R
q/q_45_1000	19.2	569	350	19165	WR
q/q_45_1200	16.4	683	914	6761	WR
q/q_45_1400	14.3	797	1924	3227	WR
q/q_60_1000	20.0	725	1559	7119	DDC
q/q_60_1200	17.0	870	3618	4062	DDC
q/q_60_1400	14.9	1015	6918	3026	DDC
w/w_30_1000	15.3	398	135	29119	R
w/w_30_1200	13.0	477	414	17274	WR
w/w_30_1400	11.4	557	1029	10079	WR
w/w_45_1000	16.0	569	1616	17720	WR
w/w_45_1200	13.6	683	4348	5697	DDC
w/w_45_1400	11.9	797	9443	2405	DDC
w/w_60_1000	16.6	725	7478	5554	DDC
w/w_60_1200	14.1	870	17909	2912	DDC
w/w_60_1400	12.3	1015	15913	2138	DDC
Even layered crust					
d/q_30_1000	18.4	398	31	30412	R
d/q_30_1200	15.7	477	93	18245	R
d/q_30_1400	15.1	557	244	10950	R
d/q_45_1000	22.6	569	409	19757	WR
d/q_45_1200	22.5	683	1229	8313	PDC
d/q_45_1400	20.3	797	2664	5123	PDC
d/q_60_1000	28.7	725	2195	10504	DDC
d/q_60_1200	24.3	870	5060	7066	DDC
d/q_60_1400	21.1	1015	9564	5385	DDC
d/w_30_1000	15.3	398	135	29119	X
d/w_30_1200	15.1	477	475	17506	R


d/w_30_1400	15.1	557	1329	10688	R
d/w_45_1000	22.6	569	2232	19419	X
d/w_45_1200	22.5	683	6943	8217	DDC
d/w_45_1400	20.3	797	15448	5089	DDC
d/w_60_1000	28.7	725	12530	10419	DDC
d/w_60_1200	24.3	870	29764	7041	DDC
d/w_60_1400	21.1	1015	26204	5375	DDC


Uneven layered crust :


upper crust/lower crust_upper thickness/total thickness [km]_Tbottom [°C]


d/q_15/45_1000	19.2	569	350	19165	WR
d/q_15/45_1200	16.4	683	914	6761	WR
d/q_15/45_1400	15.1	797	2023	3268	DDC
d/q_30/45_1000	27.5	569	492	22579	R
d/q_30/45_1200	23.3	683	1270	9576	PDC
d/q_30/45_1400	20.3	797	2664	5411	PDC


table 4


















Pressure (GPa)









I

	SiO ₂	Al ₂ O ₃	Fe ₂ O ₃	FeO	MgO	CaO	Na ₂ O	K ₂ O
Pelite (B95)	63.31	17.22	0.82	5.45	3.00	3.52	1.48	3.64
Granite (LM76)	71.84	14.43	1.22	1.65	0.72	1.85	3.71	4.00
Basalt (LM76)	49.97	15.99	3.85	7.00	8.84	9.82	2.96	1.12






















$t=4$ Ma / finite stretching = 0.19





$t=14$ Ma / finite stretching = 0.66



















Log(Intra-crustal strength contrast)







Site	minimum pressure (kb)	maximum pressure (kb)
SABN	8	10
GK	3	5
Edough	12	14
Tenda		
Beni Mesala	13	18
Filali		
BB kinzigites	9	13
BB mafic	16	20
Catena Costiera	11	13
Apuane Auto	7	9
Apuane Massa	8	11
argentario	11	13
aspromonte	11,2	12,4
alpu LG		7
alpu Es	6,5	8,5
alpu He	10	12
alpu S & A	7	11
NF BM	18	20
NF CA	11	15
NF beginning D3	3	4
Menderes Selimye	4	8
Menderes Bozdag		
Menderes Cine HP		
Menderes Cine HT		
Erhodope HP	13	15
Erhodope HT	3	9
Thasos	6,5	9,5
kazdag	4	6
Nigde Gumusler	5	6
andros post-orogen	6	8
Ikaria	6	8
tinos (transition)	8	10
tinos (late)	0	2
tinos HP	15	18
tinos BS	8	13
syros HP	16	20
syros retrogression	3	6
ios overprint		
ios peak basement	15	18
naxos	4,5	9,5
rechnitz HP	6	8
rechnitz late	3	4
veporic	9	10
Motajica	5	7

minimum T (°C)	maximum temperature (°C)	average pressure (kb)
660	700	
650	700	
600	800	
		9
450	550	
		8
800	850	
760	820	
	350	
410	430	
450	500	
350	400	
540	570	
	400	
	420	
540	620	
390	450	
550	650	
520	620	
		8
		15
		7
450	600	
550	620	
550	650	
590	690	
700		
600	650	
500	550	
400	500	
450	500	
375	525	
450	550	
300	450	
		8
480	520	
620	720	
330	370	
350	450	
550	630	

average temperature (°C)	Code	Timing	ref
	GK1	syn-detachment	saadallah96
	GK2	syn-detachment	"
	Ed	syn-detachment	caby01
400	Tn	syn-detachment	gueydan03
	BM	syn-detachment	negro 07
780	BB3	syn-detachment	"
	BB2	syn-detachment	"
	BB1	inherited	"
	CC	syn-detachment	rossetti04
	AA2	syn-detachment	jolivet98
	AA1	inherited	"
	Arg	syn-detachment	brunet00
	Asp	syn-detachment	ortolano05
	Alpu4	syn-detachment	azanon00
	Alpu3	inherited	"
	Alpu1	inherited	"
	Alpu2	inherited	"
	NF1	inherited	augier05
	NF2	inherited	"
500	NF3	syn-detachment	"
550	Me4	syn-detachment	whitney02 / Regnier 03
530	Me3	syn-detachment	okay01
640	Me1	inherited	oberhansli&al in press
620	Me2	syn-detachment	"
	Rho1	inherited	bonev & beccaletto07
	Rho2	syn-detachment	"
	Rho3	syn-detachment	brun & sokoutis07
	Kaz	syn-detachment	cavazza09
	Nig	syn-detachment	Whitney & Dilek, 98
450	And	syn-detachment	huet phd
		syn-detachment	Laurent 2016
	Tin3	syn-detachment	parra 02
	Tin4	syn-detachment	"
	Tin1	inherited	"
	Tin2	inherited	"
	Syr1	inherited	trotet 01
	Syr2	syn-detachment	huet phd
400	Ios1	syn-detachment	huet phd
	Ios2	inherited	huet phd
	Nax	syn-detachment	buick & holland 89
	Rech1	inherited	koller 85
	Rech2	syn-detachment	"
620	Vep	syn-detachment	janak 01
	Mot	syn-detachment	Ustaszewski10