



HAL
open science

Coastal upwelling areas as safe havens during climate warming

Zi-Min Hu, Marie-Laure Guillemin

► **To cite this version:**

Zi-Min Hu, Marie-Laure Guillemin. Coastal upwelling areas as safe havens during climate warming. Journal of Biogeography, 2016, 10.1111/jbi.12887 . hal-01383210

HAL Id: hal-01383210

<https://hal.sorbonne-universite.fr/hal-01383210>

Submitted on 18 Oct 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **COMMENTARY**

2 **Coastal upwelling areas as safe havens during climate warming**

3

4 Title: 9 words

5 References: 10 (max = 10)

6 Total word count: 1588 (max = 1600)

7

8 Seaweeds have long been recognized as “ecosystem engineers” and carbon sinks in
9 coastal marine ecosystems. However, recent ocean warming has had drastic
10 eco-physiological impacts on seaweed assemblages and hence coastal community
11 structure (Wernberg *et al.*, 2011, 2013). Seaweeds, particularly species that prefer
12 lower water temperatures, are facing imminent threat and will experience strong
13 population-size reductions at low-latitude range limits (Wernberg *et al.*, 2011, 2013). If
14 they fail to acclimatize/adapt eco-physiologically to these new stressful conditions or
15 fail to track more suitable habitats moving poleward, they will eventually become
16 extinct (Wernberg *et al.*, 2011, 2013). Understanding how climate change affects
17 seaweed distribution shifts may allow for the identification of potential refugia where
18 these species and their associated biota can retreat to and persist; this, in turn, will
19 offer pivotal conservation insights for sustaining biodiversity.

20 So, is there any chance for coastal temperate species to persist in global warming
21 scenarios, even at their low-latitude range limits? Fortunately, the answer seems to be
22 “yes”. In a recent issue of the *Journal of Biogeography*, Lourenço *et al.* (2016) found

23 that the brown alga *Fucus guiryi* Zardi, Nicastro, Serrão & Pearson persists in several
24 upwelling areas along the coasts of south-western Iberia and north-western Africa
25 where sea surface temperatures (SST) have undergone significant warming during the
26 last decades. These southern refugia have retained distinctive genetic pools,
27 highlighting the fundamental potential of coastal upwelling systems to maintain
28 regional or global marine biodiversity threatened by climate change. Interestingly, a
29 study involving both phylogeography and ecological niche modeling have indicated
30 that the southern upwelling centres of the Canary Current were stable refugia for
31 another temperate brown alga (*Saccorhiza polyschides* (Lightfoot) Batters) during the
32 warmer Mid-Holocene (Assis *et al.*, 2016).

33 Coastal upwelling regimes are located along continental margins where equatorial
34 winds push surface waters offshore and replace them with deeper, cold, nutrient-rich
35 waters, enabling higher primary production and colder SST than surrounding areas
36 (Wang *et al.*, 2015). These current systems play a crucial role in structuring the
37 abundance and richness of pelagic plankton, and hence coastal marine ecosystems and
38 fisheries. It has been proposed that global climate change might intensify upwelling
39 regimes (Wang *et al.*, 2015), allowing for the persistence of localized extensions of cold
40 water embedded in warmer coastal waters for the years ahead and leading to more
41 drastic gradients of oceanic conditions in these regions. However, these trends are not
42 spatially homogeneous; if upwelling is expected to effectively strengthen along the
43 poleward portions of these currents, it could diminish along equatorial coasts (e.g.
44 along north-western Africa for the Canary Current; Wang *et al.*, 2015). So, while some

45 evidence suggests that these systems could be resilient to anthropogenic climate
46 change, their fate remains uncertain due to the large number of physical and
47 biogeochemical factors that influence ecosystem processes (Byrnes *et al.*, 2011; Wang
48 *et al.*, 2015).

49 In spite of the clear importance of coastal upwelling systems as potential refugia for
50 marine biota facing ocean warming, little attention has been paid to the diversity and
51 dynamics of seaweeds in these areas. However, Ormond & Banaimoon (1994)
52 reported that along the coast of southern Yemen, the maximum growth of intertidal
53 macroalgal assemblages followed the onset of elevated nutrient levels driven by
54 intense southern Arabian coastal upwelling. On the other hand, Leliaert *et al.* (2000)
55 described a drastic change in seaweed community composition around the Cape
56 Peninsula of South Africa in relation to oceanic conditions (i.e. the Atlantic side is
57 dominated by the Benguela upwelling system while no upwelling occurs within False
58 Bay). Such evidence indicates that coastal upwelling systems greatly influence the
59 diversity and biogeographic patterns of coastal marine macroalgae.

60 The newly published work by Lourenço *et al.* (2016) has extended our understanding
61 of how upwelling systems have acted as climatic refugia and how these systems have
62 influenced the evolution and biodiversity of seaweeds. Comparing historical and
63 recently collected data of algal presence and coverage along the North Atlantic, the
64 authors were able to confirm that the range of *F. guiryi* is shrinking and that, south of
65 37°N, the species has disappeared except near upwelling centres that act as cold-water
66 refugia. Moreover, information from nine microsatellite loci shows clear genetic

67 uniqueness of the remaining southern populations, suggesting that these
68 southernmost populations should potentially be targeted in future conservation and
69 management plans. One could also wonder about the possible importance of the
70 southernmost populations for the evolutionary potential of the species. Indeed,
71 populations of genetic variants specifically encountered in the warmer areas of *F.*
72 *guiryi*'s range could potentially be undergoing adaptive evolution to ocean warming
73 conditions. However, even if upwelling centres show certain resilience to ocean
74 warming (Wang *et al.*, 2015; Assis *et al.*, 2016; Lourenço *et al.*, 2016), one might ask
75 how long they will act as refugia. Are these southern refugia sufficiently stable to allow
76 remnant populations to adapt to new climatic conditions or even only to survive until
77 conditions become favourable again - if ever? Lourenço *et al.* (2016) reported that
78 relict populations of *F. guiryi* are small and sparse, and processes linked to the Allee
79 effect (i.e. negative effects of decreasing population density on fitness) or genetic drift
80 could greatly increase this species' vulnerability to extinction. Future studies of
81 recruitment patterns in relict populations of *F. guiryi* would be interesting considering
82 that *Fucus* is a perennial genus and some areas where it is found could be suitable for
83 adult survival but not for successful reproduction; these areas could be mistaken for
84 refugia. Moreover, some non-climatic factors have also been demonstrated to impose
85 severe eco-physiological effects on the growth, reproduction, and distribution of
86 canopy-forming *Fucus* species (e.g. Vadas *et al.*, 1992). Knowledge of the effect of
87 these non-climatic factors is highly relevant when developing effective management
88 and conservation plans since they may further exacerbate the effect of oceanic

89 warming on seaweed populations.

90 It is yet unknown whether the patterns of rapid population shrinkage and isolation
91 at range limits reported by Lourenço *et al.* (2016) for *F. guiryi* can be generalized to
92 other marine organisms facing the imminent threat of global warming. Sequential field
93 surveys and genetic studies in other major coastal upwelling systems worldwide (i.e.
94 the California Current System, the Benguela Current, and the Peru–Humboldt Current)
95 can undoubtedly provide more insights into the role of upwelling systems in shaping
96 seaweed populations and communities. Assessing the environmental requirements for
97 life cycle completion in populations along entire species' ranges is a prerequisite to
98 predicting possible shift in seaweed distributions. This is also critical for characterizing
99 the underlying linear and/or non-linear relationships between seaweed performance
100 and key oceanic parameters (e.g. temperature, salinity or pH) that are expected to
101 change steadily during the 21st century (Harley *et al.*, 2012). Lastly, it is essential to
102 gather historical and current distribution records for a large number of sympatric
103 species in order to empirically identify coastal community changes under global
104 warming scenarios.

105

106 ZI-MIN HU^{1,2*}

107 MARIE-LAURE GUILLEMIN^{3,4*}

108 ¹*Key Laboratory of Experimental Marine Biology, Institute of Oceanology, Chinese*

109 *Academy of Sciences, Qingdao 266071, China,* ²*Qingdao National Laboratory for*

110 *Marine Science and Technology, Qingdao 266071, China,* ³*Instituto de Ciencias*

111 *Ambientales y Evolutivas, Facultad de Ciencias, Universidad Austral de Chile, Casilla*
112 *567, Valdivia, Chile,⁴CNRS, Sorbonne Universités, UPMC University Paris VI, UMI 3614,*
113 *Evolutionary Biology and Ecology of Algae, Station Biologique de Roscoff, CS 90074,*
114 *Place G. Teissier, 29680 Roscoff, France*

115 *E-mail: huzimin9712@163.com; marielaure.guillemine@gmail.com

116

117 **ACKNOWLEDGEMENTS**

118 We thank two anonymous referees and the editor Alistair Crame for providing useful
119 and constructive comments to improve the manuscript. Z.-M. H. gratefully
120 acknowledges financial support from National Natural Science Foundation of China
121 (31370264) and The Scientific and Technological Innovation Project from Qingdao
122 National Laboratory for Marine Science and Technology (No.2015ASKJ02), M-L. G.
123 acknowledges support from CONICYT FONDECYT/REGULAR No. 1130797 and Centro
124 FONDAP IDEAL No. 15150003.

125

126 **REFERENCES**

127 Assis, J., Coelho, N.C., Lamy, T., Valero, M., Alberto, F. & Serrão, E.Á. (2016) Deep reefs
128 are climatic refugia for genetic diversity of marine forests. *Journal of Biogeography*,
129 **43**, 833-844.

130 Byrnes, J.E., Reed, D.C., Cardinale, B.J., Cavanaugh, K.C., Holbrook, S.J. & Schmitt, R.J.
131 (2011) Climate-driven increases in storm frequency simplify kelp forest food webs.
132 *Global Change Biology*, **17**, 2513–2524.

-
- 133 Harley, C.D.G., Anderson, K.M., Demes, K.W., Jorve, J.P., Kordas, R.L., Coyle, T.A. &
134 Graham, M.H. (2012) Effects of climate change on global seaweed communities.
135 *Journal of Phycology*, **48**, 1064–1078.
- 136 Leliaert, F., Anderson, R.J., Bolton, J.J. & Coppejans, E. (2000) Subtidal understory algal
137 community structure in kelp beds around the Cape Peninsula (Western Cape, South
138 Africa). *Botanica Marina*, **43**, 359–366.
- 139 Lourenço, C.R., Zardi, G.I., McQuaid, C.D., Serrão, E.A., Peargon, G.A., Jacinto, R. &
140 Nicastro, K.R. (2016) Upwelling areas as climate change refugia for the distribution
141 and genetic diversity of a marine macroalga. *Journal of Biogeography*, **43**, 1595–
142 1607.
- 143 Ormond, R.F.G. & Banaimoon, S.A. (1994) Ecology of intertidal macroalgal assemblages
144 on the Hadramout coast of southern Yemen, an area of seasonal upwelling. *Marine
145 Ecology Progress Series*, **105**, 105–120.
- 146 Vadas, R.L., Johnson, S. & Norton, T.A. (1992) Recruitment and mortality of early
147 post-settlement stages of benthic algae. *British Phycological Journal*, **27**, 331–351.
- 148 Wang, D.W., Gouhier, T.C., Menge, B.A. & Ganguly, A. R. (2015) Intensification and
149 spatial homogenization of coastal upwelling under climate change. *Nature*, **518**,
150 390–394.
- 151 Wernberg, T., Russell, B.D., Thomsen, M.S., Guigel, C.F.D., Bradshaw, C.J.A.,
152 Poloczanska, E.S. & Connell, S.D. (2011) Seaweed communities in retreat from ocean
153 warming. *Current Biology*, **21**, 1828–1832.
- 154 Wernberg, T., Smale, D.A., Tuya, F., Thomsen, M.S., Langlois, T.J., de Bettignies,

-
- 155 T., Bennett, S. & Rousseaux, C.S. (2013) An extreme climatic event alters marine
156 ecosystem structure in a global biodiversity hotspot. *Nature Climate Change*, **3**, 78–
157 82.
- 158 Editor: Alistair Crame