Sea surface temperature variability in the North Western Mediterranean Sea (Gulf of Lion) during the Common Era

Marie-Alexandrine Sicre, Bassem Jalali, Belen Martrat, Sabine Schmidt,
Maria-Angela Bassetti, Nejib Kallel

► To cite this version:

Marie-Alexandrine Sicre, Bassem Jalali, Belen Martrat, Sabine Schmidt, Maria-Angela Bassetti, et al.. Sea surface temperature variability in the North Western Mediterranean Sea (Gulf of Lion) during the Common Era. *Earth and Planetary Science Letters*, 2016, 456, pp.124-133. 10.1016/j.epsl.2016.09.032 . hal-01386646

HAL Id: hal-01386646

<https://hal.sorbonne-universite.fr/hal-01386646>

Submitted on 24 Oct 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Sea surface temperature variability in the North Western Mediterranean Sea (Gulf of Lion) during the Common Era

Marie-Alexandrine Sicre^{a,*}, Bassem Jalali^{a,b}, Belen Martrat^c, Sabine Schmidt^d, Maria-Angela Bassetti^e, Nejib Kallel^b

^a Sorbonne Universités (UPMC, Univ. Paris 06)-CNRS-IRD-MNH, LOCEAN Laboratory, 4 place Jussieu, F-75005 Paris, France.

* Corresponding author : Marie-Alexandrine.Sicre@locean-ipsl.upmc.fr

^b Université de Sfax, Faculté des Sciences de Sfax, Laboratoire GEOGLOB, BP.802, 3038, Sfax, Tunisia.

^c Department of Environmental Chemistry, Spanish Council for Scientific Research (IDÆA-CSIC), 08034 Barcelona, Spain

^d UMR5805 EPOC, Université de Bordeaux, Avenue Geoffroy Saint-Hilaire, 33615 Pessac, France

^e CEFREM, CNRS UMR5110, Université de Perpignan, Avenue J.-P. Alduy, 66860 Perpignan, France.

ABSTRACT

This study investigates the multidecadal-scale variability of sea surface temperatures (SSTs) in the convection region of the Gulf of Lion (NW Mediterranean Sea) over the full past 2000 yr (Common Era) using alkenone biomarkers. Our data show colder SSTs by 1.7°C over most of the first millennium (200 - 800 AD) and by 1.3°C during the Little Ice Age (LIA; 1400-1850 AD) than the 20th century mean (17.9°C). Although on average warmer, those of the Medieval Climate Anomaly (MCA) (1000-1200 AD) were lower by 1°C. We found a mean SST warming of 2°C/100 yr over the last century in close agreement with the 0.22 and 0.26°C/decade values calculated for the western Mediterranean Sea from *in situ* and satellite data, respectively. Our results also reveal strongly fluctuating SSTs characterized by cold extremes followed by abrupt warming during the LIA. We suggest that the coldest decades of the LIA were likely caused by prevailing negative EA states and associated anticyclone blocking over the North Atlantic resulting in cold continental northeasterly winds to blow over Western Europe and the Mediterranean region.

Keywords: Mediterranean Sea; sea surface temperature; alkenones; Common Era; East Atlantic mode; atmospheric blocking

1. Introduction

In the past decade, major efforts have been done to document the multi-decadal variability of the sea surface temperatures during the Common Era (last 2,000 yr) and to explore the role of external forcings (solar, volcanism, greenhouse gases) by combining paleo records and numerical simulations of the last millennium climate (McGregor et al., 2015; Sicre et al., 2011). Although the number of pre-instrumental reconstructions of sea surface temperature (SST) resolving the decadal scale has increased significantly they are still insufficient to precisely describe the space-time climate variability both at global and regional scales. This is particularly true for the Mediterranean region for which very few records exist despite alarming future climate projections (Lionello et al., 2006). Indeed, the Mediterranean region is one of the most sensitive areas to climate change owing to its geographical location between the temperate climate of Europe and the arid climate of North Africa. Because of this, even minor modifications in the extension and intensity of these climate zones can substantially alter the Mediterranean climate making this region particularly vulnerable to global warming (Lionello et al., 2006). In its history, the Mediterranean region has undergone important changes that can be investigated to better understand present-day interactions between global and regional climate and the underlying driving mechanisms.

The Mediterranean climate is strongly influenced by the large-scale mid-latitude atmospheric circulation of the North Atlantic (NA) and primarily the East Atlantic pattern (EA) and the North Atlantic Oscillation (NAO; Hurrell, 1995). The NAO, the dominant mode of atmospheric variability in the NA, reflects the atmospheric pressure difference between the Azores High and Icelandic low. The NAO state determines the latitudinal position of the NA storm tracks driving the Mediterranean winter precipitation, but its role on Mediterranean SSTs is secondary (Lionello et al., 2006). Instead, the EA has been recently recognized as the main controlling factor of the SST variability of the Mediterranean Sea, particularly in the western basin (Josey et al., 2011). The EA mode has a similar North South dipole structure as the NAO but its centers of action are displaced southeastward, which results in a stronger link

57 with the subtropical climate than NAO. Teleconnections with El Nino Southern Oscillation
58 (ENSO) have also been suggested mainly to explain winter rainfall in some areas of the
59 Mediterranean region (Alpert et al., 2006). Finally, the influence of the Atlantic Multidecadal
60 variability (AMV) (Knight et al., 2006) has been recently pointed out, yet the dynamical links
61 between AMV and Mediterranean SSTs is still an open question. Indeed, oceanic processes
62 have been suggested based on the detection of AMV-like 70-yr period oscillations in the
63 Mediterranean SSTs (Marullo et al., 2011) while according to the study of Mariotti and
64 Dell'Aquila (2012) atmospheric transmission of AMV is more likely.

65 The complex topography of the Mediterranean basin modifies the large-scale atmospheric
66 flow and subsequently influences the climate characteristics at a local scale. In the
67 northwestern Mediterranean Gulf of Lion (GoL), interactions between the mid-latitude
68 westerly winds and the Alps result in northerly winds blowing offshore in the South of
69 France, called Mistral (Jiang et al., 2003). This cold and dry wind blows at all seasons
70 (climatologically 49% of wind frequency is in the northwest quadrant; Burlando, 2009), but
71 more strongly in winter and spring (Jiang et al, 2003) causing intense surface water cooling.
72 Najac et al. (2009) have shown that Mistral is favored by anticyclonic blocking over the
73 northeastern Atlantic and a low-pressure system in the central Mediterranean Sea (Fig. 1), a
74 synoptic configuration described by negative EA that focuses the northerly air flow over
75 France. Despite previous suggestions of a dynamical connection between negative NAO and
76 the occurrence of NA blocking (Shabbar et al., 2001), the most severe Mistral episodes show
77 only a modest correlation with NAO compared to the strong correlation with negative EA
78 (Skirris et al., 2012; Papadopoulos et al., 2012). Ultimately, while weak westerlies during
79 negative NAO result in cold temperature in Europe, the most severe winters occur during
80 negative EA due to the deflection of the maritime westerly flow to the North around the
81 anticyclonic cell returning as cold and dry northerly continental winds over Europe and the
82 Northwestern Mediterranean Sea (Fig. 1) (Häkkinen et al., 2011). Yet, interactions exist
83 between the two modes (Moore and Renfrew, 2011). Indeed, bivariate reconstructions have

shown that EA modulates the strength and position of NAO centers of action and that winter severity is enhanced when both modes are in negative phase. This would for instance explain that despite similar negative NAO values, winter in Europe was much cooler in 2010 (negative EA) than in 2009 (positive EA) (Moore and Renfrew, 2011).

Mistral exerts a strong control on the SSTs in the Northwestern Mediterranean Sea and is responsible for among the coldest values found in the Gulf of Lions (GoL). Mistral also triggers intense blooms in February, March and April (FMA) when it is stronger (Bosc et al., 2004; Durrieu de Madron et al., 2013). A relationship between primary production maxima, Mistral and negative EA has also been evidenced by Olita et al. (2011). Because Mistral concurrently causes surface cooling and high primary production, alkenone-derived SSTs in the GoL are expected to well capture past changes of atmospheric conditions promoting Mistral. These unique properties motivated the choice of GoL shelf sediment for generating a high-resolution SST reconstruction over the Common Era using alkenone as a temperature proxy. Based on this time series we investigate the links between mid-latitude atmospheric variability, Mistral and SSTs in the GoL with a focus on the strong amplitude SST fluctuations observed during the Little Ice Age (LIA).

2. Material and methods

2.1 Analytical procedure

A gravity core KSGC-31 (GMO2-Carnac cruise in 2002, R/V “Le Suroît”) and multi-core Gol-Ho1B (GolHo cruise in 2013, R/V “Néréis”) were retrieved at virtually the same location (43°0’23N; 3°17’56E, water depth 60 m, Fig. 1) in the Rhone river mud belt deposited onto the GoL continental mid-shelf.

The two sediment cores were sampled continuously at a sampling step of 1 cm and freeze-dried overnight. Between 2 - 3 g of dried sediments were extracted with a mixture of methanol/methylene chloride (1:2 v/v) to recover the lipid biomarkers. Alkenones were

further isolated from the total lipid extract by silica gel chromatography using solvent mixtures of increasing polarity. Gas chromatography (GC) analyses were performed on a Varian CX 3400 gas chromatograph equipped with a fused CP-Sil-5 silica capillary column (50 m x 0.32 mm i.d.) and a flame ionization detector. Helium was used as carrier gas. The oven was temperature programmed from 100°C to 300°C at a rate of 20°C min⁻¹. 5 α -cholestane was added prior GC analyses for quantitation. Details on the analytical procedure can be found in Ternois et al. (1996).

Alkenones are primarily biosynthesized by the ubiquitous haptophyte algae *Emiliania huxleyi*. This compound series comprises a suite of methyl and ethyl C₃₇ to C₃₉ ketones with two or three double bonds. The C₃₇ alkenones are used to calculate the C₃₇ unsaturation index U₃₇^K (U₃₇^K = C_{37:2}/(C_{37:2}+C_{37:3})) now a well-established temperature proxy in paleoceanographic studies. SSTs were calculated from the downcore U₃₇^K values and the most recent global calibration published by Conte et al. (2006) ($T = -0.957 + 54.293(U_{37}^K) - 52.894(U_{37}^K)^2 + 28.321(U_{37}^K)^3$, standard error estimate: 1.2°C). Precision based on triplicate alkenone analyses is 0.01 U₃₇^K unit, which in the temperature range considered here, translates into 0.3°C.

2.2 Chronology

Age control for the multi-core Gol-Ho1B is based on ²¹⁰Pb dating and for gravity core KSGC31 on both ²¹⁰Pb and ¹⁴C dating. For the gravity core, 10 radiocarbon dates were acquired on bivalve shells with the ARTEMIS accelerator mass spectrometer (AMS) operated in the Laboratoire de Mesure du Carbone 14, Saclay (France). Radiocarbon ages were corrected for local reservoir effect of $\Delta R = 23 \pm 71$ (Table 1) and converted into 1 σ calendar years using the CALIB7.1 software and the marine calibration curve Marine13 (Stuiver and Reimer, 1993; Reimer et al., 2013). Linear interpolation was performed between ¹⁴C-dated horizons to translate each sampling depth to age expressed here in Anno Domini (AD) years (Fig. 2). Because the two upper most ¹⁴C dates indicate post-bomb ages (calibrated using

OxCal 4.2; Ramsey and Lee, 2013) additional ^{210}Pb and ^{137}Cs measurements were performed in the first 10 cm of the gravity core KSGC31 (Table 2). Note that the ^{14}C date at 18.5 cm (1570 ± 78 yr AD) was considered as an outlier and discarded from the age model as it appeared incompatible with the ^{14}C modern age of the above horizons (5.5 and 11.5 cm) and the presence of ^{210}Pb and ^{137}Cs in the upper 10 cm.

Activities of ^{210}Pb , ^{226}Ra , ^{137}Cs and ^{232}Th were determined in the gravity core KSGC-31 and multi-core Gol-Ho1B by γ spectrometry using a low-background high-efficiency well-shaped germanium detector equipped of a Cryo-cycle (CANBERRA; Schmidt et al., 2014) (Fig. 3). The calibration of the detector was achieved using certified IAEA reference materials (RGU-1; RGTh; IAEA-135). Activities are expressed in mBq g^{-1} and errors based on 1σ counting statistics. ^{210}Pb excess ($^{210}\text{Pb}_{\text{xs}}$) was determined by subtracting the activity supported by its parent isotope ^{226}Ra from the total ^{210}Pb activity in the sediment. Errors in $^{210}\text{Pb}_{\text{xs}}$ were calculated by propagation of errors in the corresponding pair (^{226}Pa and ^{210}Pb). To compensate for potential effect of compositional changes of sediments and to optimize splicing of the multicore and gravity core records, $^{210}\text{Pb}_{\text{xs}}$ activities were normalized using ^{232}Th concentrations measured simultaneously by γ counting ($^{210}\text{Pb}_{\text{xs}}^{\text{Th}}$).

Depth profiles of ^{232}Th , $^{210}\text{Pb}_{\text{xs}}^{\text{Th}}$ and ^{137}Cs activities for multicore Gol-Ho1B were determined shortly after collection at sea and prior biomarker analyses. $^{210}\text{Pb}_{\text{xs}}^{\text{Th}}$ values show a mixed layer in the upper 4 cm, followed by an exponential decrease with increasing depth in the core (Table 2, Fig. 3B) from which sediment age is calculated. Sediment accumulation rates were derived from the $^{210}\text{Pb}_{\text{xs}}^{\text{Th}}$ profile assuming constant flux and constant sedimentation accumulation rates (referred to as the CF:CS model, Schmidt et al. 2014). To account for compaction effect, $^{210}\text{Pb}_{\text{xs}}^{\text{Th}}$ values were plotted against cumulative mass. A mean Mass Accumulation Rate (MAR) of $0.31 \text{ g cm}^{-2} \text{ yr}^{-1}$ was estimated. The deposition time (in years) of each sediment layer was obtained by dividing the cumulated dry mass per unit area by the MAR (Fig. 3D). The deposition year was subsequently estimated based on the sampling year of the core, in 2013. The ^{210}Pb chronology indicates that the multicore Gol-Ho1B ranges from

1960 (± 5.6) to 2013 AD. ^{137}Cs was detected throughout the multi-core in agreement with the well-known pulse inputs related to the nuclear weapon test fall-out in the early sixties (maximum atmospheric fallout is in 1963 in the Northern Hemisphere) (Fig. 3C, Table 2). Comparison between ^{232}Th , $^{210}\text{Pb}_{\text{xs}}$ and ^{137}Cs activities in KSGC31 and Gol-Ho1B cores was used to determine the material loss during gravity coring and to splice records. A shift in depth of 15 cm gave the best correspondence between the two profiles (Fig. 3) leading to a core-top age of 1971 (± 1.4) AD for the gravity core KSGC31.

For the multicore Gol-Ho1B sediment, the sedimentation rate decreases from 0.47 cm yr⁻¹ at the water-sediment interface to 0.32 cm yr⁻¹ at the base of the core. In the gravity core KSGC31, sedimentation rate determined for the upper 10 cm is estimated to be around 0.16 cm yr⁻¹ (without porosity values for the gravity core MAR could not be calculated). Sedimentation rate for the underlying layers derived from ^{14}C dating is on the order of 0.1 cm yr⁻¹. Lower sedimentation rates in the gravity core compared to the multicore is not unusual and explained by increasing natural compaction of sediment with depth and possibly also by compaction due to the gravity coring (Sicre et al., 2011).

3. Results

3.1. Industrial Era period

Fig. 4A shows the SST reconstruction since 1850 AD obtained for the upper gravity core KSGC31 (dark green solid line) and entire multicore Gol-Ho1B (light green dashed line) records against age, based on the ^{210}Pb and ^{14}C chronology. Values range from 16.5 to 18°C in the gravity core, and from 16.5 to 19°C in the multicore. Both records indicate comparable SST values in the overlapping interval except for the warmest value in the mid-1960s (~19°C) seen in multi-core sediment but not in the gravity core sediment. We interpreted this result by a probable artifact in the uppermost cms of the gravity core due to coring as suggested by smoothing in the exponential decay in the ^{210}Pb curve (Fig. 3B). Indeed, gravity

coring generally causes material loss and the disturbance of the uppermost sediment layers while a multi-corer typically recovers intact surface sediment. The two-fold difference in sedimentation rates (0.32 versus 0.16 cm yr⁻¹) between the two records in this interval translates into a difference of temporal resolution that could also account for this discrepancy. In any case, all values were considered to construct the spliced record (Fig. 4B, Fig. 5A). The full 2k SST reconstruction was thus obtained by combining the gravity core KSGC31 SST values from 0 to 1971 AD and the complete multicore Gol-Ho1B SST data spanning from 1960 to 2013 AD. As can be seen in Figure 4B, the SST signal depicts an overall warming ranging from ~16.5°C to ~19°C during the 20th century followed by cooling over the most recent decades. Multi-decadal scale fluctuations are superimposed to this trend. These values are among the lowest of the Mediterranean surface waters in agreement with the temperature field of the Mediterranean Sea shown in Figure 1 and previous alkenone SST data obtained from surface sediments distributed across the NW Mediterranean Sea pointing to colder SSTs in the GoL (Ternois et al., 1996).

3.2. The Common Era (last 2k)

Figure 6A plots the full past 2k SST record of the GoL. A 25-yr binning was applied (black line) in order to easily evaluate departures from the average value (16.7°C ± 0.7°C). The first millennium AD reveals several cold excursions of ~ 1.5°C and a mean value cooler by 1.7°C than the 20th century mean (17.9°C). During the MCA (ca. 1000 - 1200 AD) SSTs were warmer (~ 16.9°C) but ~ 1°C lower than the 20th century. Around 1200 AD, they progressively decline and display marked multi-decadal fluctuations with well-expressed minima (ca. ~ 16°C) around 1300 AD, 1500 AD, mid-16s AD and mid-17s AD, each followed by warm spells, the most outstanding by duration and amplitude (~ 2°C in few decades) centered at ~ 1700 AD. As a result, average SSTs for the LIA (1400 - 1850 AD) were 1.3°C colder than the 20th century and not as cold as the first millennium during which post-cooling temperature rebounds above average are absent. In the following section, we

compare the proxy and instrumental data and analyze the full 2k SST reconstruction with a focus on the high amplitude SST oscillations of the LIA.

4. Discussion

4.1. The Industrial Era SST signal

Figure 5 shows the Gol-Ho1B_KSGC-31 spliced SST reconstruction, the instrumental annual mean and late winter-early spring (FMA) temperature time series average over a 5°x 5° grid from Kaplan SST V2 data (http://www.esrl.noaa.gov/psd/data/gridded/data.kaplan_sst.html) since the late 19th century.

Both proxy and instrumental data show a long-term warming trend over the 20th century. The alkenone SSTs rise since 1900 AD is ~1.7°C but reaches ~2.2°C when considering the 20th century only, i.e. after removing the last cold decade (2000-2012). These estimates are close to the value of 0.026 °C yr⁻¹ obtained from satellite SSTs for the W Mediterranean Sea over the 1985 - 2008 period, and 0.022°C yr⁻¹ of the NOCS *in situ* SSTs from 1973 to 2008 (Skirris et al., 2012). Alkenone SSTs indicate values closer to the annual mean than FMA (Fig. 5) suggesting that alkenone production does not only take place during the months of intense Mistral (Bosc et al., 2004; Durrieu de Madron et al., 2013) but also during the warm season (late spring to summer). Vertical mixing induced by Mistral would thus sustain primary production over a longer season by replenishing surface waters with nutrients.

The SSTs decline since the beginning of the 21st century can be related to several severe winters such as the coldest on record in Europe and United Kingdom, in 2010/2011 that Moore and Renfrew (2011) attributed to combined negative NAO and EA states. These authors showed that EA⁻ NAO⁻ accounted for 50% of the winter temperature that year against 20% by NAO alone, outlining the value of considering both modes to better describe climate conditions. Winter 2012 was another exceptionally cold winter associated with a strong anticyclonic blocking in the NA (Durrieu de Madron et al., 2013) that could account together with winter 2010/2011 for the cold uppermost SST value. During the 20th century, cold SSTs in the 70s to early 80s are also noteworthy both in the proxy and instrumental data (Fig. 5A,

5B, 5C). They coincide with a period of strongly negative EA (Fig. 5D) and episodes of major dense water formation in the GoL (Béthoux et al., 2002) in accordance with the role of EA on SSTs in the GoL. Regression between EA index and SSTs on 5 yr binned values since 1950 were calculated to account for age model uncertainties. The obtained Pearson correlation coefficient, $r = 0.45$ (at the 95% confidence interval) is close to values calculated using instrumental data (0.58 for NOCS in situ SSTs and 0.50 for satellite SSTs in the W Mediterranean, at the 95% confidence interval) (Skliris et al., 2012).

4.2. General SST trends over the past 2000 yr

In this section, we compare the SST signal of the GoL with other Mediterranean and North Atlantic records that may have features in common (Fig. 6). Records that document centennial scale SST variability in the Mediterranean during the Common Era are very scarce. They belong to different sectors of the W-Mediterranean basin, i.e. the Alboran Sea (Fig. 6B ~ 30 yr time resolution; Nieto-Moreno et al., 2013) and Balearic basin (Fig. 6C; ~ 30 yr time resolution; Moreno et al., 2012) and cover all or part of the CE. None of them have a chronostratigraphic control during the instrumental period to evaluate the relationship between SSTs and climate processes. The high-resolution SST reconstruction from North Iceland (MD99-2275; Sicre et al., 2008) was also considered for a regional analysis of the data because its cross-analysis with instrumental data has shown a strong control of NAO on SSTs (Sicre et al., 2011). Precision on the age model of this core based on tephrochronology is on the order of the decade while for the other cores it is on the order of several decades (Table 1).

All records shown in Fig. 6 are based on alkenone SSTs, but primary production patterns have different characteristics depending on the location of each core in the Mediterranean Sea. While the GoL belongs to a blooming regime tightly linked to Mistral, the Alboran and Balearic Sea relate to different primary production patterns as defined by D'Ortenzio and Ribera d'Alcala (2009). In the Balearic Sea, chlorophyll concentrations show algal blooms of moderate amplitude and variable timing that occur in spring sporadically. This area thus

combines periods of enhanced production with oligotrophic conditions leading to an erratic regime classified as an intermittently blooming region (D'Ortenzio and Ribera d'Alcala, 2009). In the Alboran Sea, the seasonal cycle of phytoplankton production is the most chaotic of the Mediterranean (D'Ortenzio and Ribera d'Alcala, 2009). Primary production exhibits a strong inter-annual variability with a pronounced production peak in February - March and in fall (October). It is also strongly influenced by Atlantic inflow waters and meso-scale gyre induced upwelling which all together leads to confounding response of phytoplankton to physical forcings (Bosc et al., 2004; D'Ortenzio and Ribera d'Alcala, 2009). These cores thus belong to biogeographical clusters with temporal and dynamical regimes that are distinct from the GoL, including in its unique link to atmospheric forcing (see Fig. 4 in D'Ortenzio and Ribera d'Alcala, 2009). Considering age model uncertainties, lower temporal resolution and the lack of instrumental control to evaluate the ability of local alkenone SSTs to capture climatic information, only broad features of the records were examined for this comparison.

As can be seen in Fig. 6, the long-term cooling ending at ~ 1800 AD is shared by all almost sites and consistent with the recent finding of McGregor et al. (2015) of a global ocean cooling from 0 to 1800 AD that, according to model simulations of the pre-industrial millennium climate (801 – 1800 AD) would be imputable to volcanism. This trend reversed in the GoL and Alboran Sea around 1800 AD, but persisted off N. Iceland because of sustained sea ice occurrence, even in lower abundance during 20th century than the LIA (Marcias-Fauria et al., 2010). Unexpectedly, the absence of warming is also notable in the Balearic record as modern SSTs show rising values both in the Eastern and Western Mediterranean (Skirris et al. 2012; Papadopoulos et al., 2012) and in European land-based T reconstructions (Luterbacher et al., 2004). The lack of age control of the past ca. 500 yr (Fig. 6C, dashed lines) and extrapolation of the age model in the upper core is the probable explanation for this discrepancy. Another salient feature of these SST records, in common with the GoL, is the distinctly cooler first millennium as compared to the MCA off N. Iceland while in the Balearic basin it is almost muted. In contrast, the last millennium cooling at the onset of the

LIA is observed in all records, though not synchronously. It is much abrupt off N. Iceland due to the imprint of sea ice on SSTs associated with the southward shift of the polar front under weakened NAO (Trouet et al., 2009).

As earlier outlined, the most outstanding feature of the GoL SST signal is the high amplitude oscillations and in particular the rebounds above the mean observed during the LIA that are not seen during the first millennium. Interestingly, while sea ice was common in the North Atlantic (Massé et al., 2008) and the northward oceanic heat transport reduced (Lund et al., 2006) during the LIA, the first millennium is a period of minimum sea ice cover and enhanced northward advection of heat. In the next section we investigate causes for the outstanding SST fluctuations of the LIA in the GoL and explore past changes in atmospheric modes of variability and notably EA.

4.3 SSTs and large-scale climate variability modes over the last millennium

Multidecadal to centennial scale variability of the GoL SSTs over the past 1000 yr (Fig. 7E) is explored in light of high-resolution proxy data from the adjacent North Atlantic, i.e. Iceland (Sicre et al., 2011; Larsen et al., 2013; Ólafsdóttir et al., 2013; Fig. 7B, 7C), the Iceland Basin (Moffa-Sanchez et al., 2014, Fig. 7D), the Swiss Alp glaciers (Holzhauser et al., 2005, Fig. 7A), and western Europe land temperatures (Luterbacher et al., 2004, Fig. 7F), as well as the NAO index (Ortega et al., 2015; Fig. 7G) and solar activity reconstructed by Steinhilber et al. (2012) (Fig. 7H) to investigate the role of large-scale atmospheric low frequency variability and oceanic processes.

Previous proxy reconstructions have shown that during the MCA surface waters were generally warmer along the path of the North Atlantic Current (Keigwin, 1996; Sicre et al., 2008; Richter et al. 2009) due to stronger Atlantic Meridional Overturning Circulation (AMOC) consistent with the hypothesis that positive NAO phase (Fig. 7H) prevailed during this period and that strong NAO enhances AMOC (Delworth and Greatbach, 2000). Relatively warm and/or wet summers resulted in increase melting and glacier retreat in Iceland (Larsen et al., 2013; Ólafsdóttir et al., 2013) (Fig. 7B) and in the Alps (Holzhauser et

al., 2005; Denton and Broecker, 2008) (Fig. 7A). Off North Iceland, the stepwise SST increase ~ 1000 AD indicates a northward migration of the polar front within a few decades (Fig. 7C) a feature that is not seen in the Mg/Ca of *G. inflata* South of Iceland (RAPiD17-15, 6yr time resolution; Fig. 7D) where surface water properties are mainly shaped by the strength of the subpolar gyre (Moffa-Sanchez et al., 2014). In the GoL, SSTs also generally warm (Fig. 7E). The transition to colder conditions between 1200 and 1300 AD, depending on the records, is thought to be related to a weakening of NAO leading to colder climate in the North Atlantic and the adjacent Euro-Mediterranean region (Trouet et al., 2009).

All independent proxy records shown in Fig. 7 point to more severe conditions during the LIA and the return of seasonal sea ice but with differences in the decadal structure of the signals. SSTs in the GoL and S. Iceland basin (Fig. 7D, 7E) indicates that cooling occurred in parallel during the early LIA (1400 to 1550 AD), but tend to vary in opposite phase during the Late LIA. Notably, between 1530 and 1680 AD, cold SSTs in the GoL contrast with warm surface waters South of Iceland confirming previous findings of Richter et al. (2009) at Feni drift of unexpected mild conditions in the subpolar North Atlantic. According to Larsen et al. (2013) this warmth would have been responsible for the retreat of the Langjökull ice cap from 1550 to 1680 AD (Fig. 7B). Advection of temperature anomalies from the tropical Atlantic combined with reduced heat loss to the atmosphere under weakened NAO have been proposed as possible explanations for these unexpected warm conditions during the LIA under reduced AMOC (Richter et al., 2009). Concomitantly, coldest winters in Europe supported by the multiproxy reconstruction of Luterbacher et al. (2004) ((Fig. 7F) and the expansion of Swiss glacier (Fig. 7A; Holzhauser et al., 2005; Denton and Broecker, 2008) evidence a W-E temperature asymmetry between the cold Euro-Mediterranean climate and warm subpolar waters during this time interval. This spatial pattern contrasting warm subpolar surface waters upstream the NA anticyclonic cell with cold temperatures in downstream regions (Europe and Mediterranean Sea) is expected from blocked regimes during negative EA (Häkkinen et al., 2011). Co-eval enhanced sea ice and lower SSTs in the

Nordic Seas (off N. Iceland) is also coherent with EA driven spatial SST pattern today (Cassou et al., 2011). Severe conditions during the LIA in the Euro-Mediterranean region can thus be seen as resulting from enhanced frequency of blocking regimes associated with negative EA on longer time scale (Cassou et al., 2011). This hypothesis is supported by modeling experiments performed for the second half of the 20th century and the Late Maunder Minimum (1645 – 1715 AD) showing that NA blocking is favored by lower solar activity that prevailed during the LIA (Nesmé-Ribes et al., 1993; Barriopedro et al., 2008). Rebounds above average would in turn reflect rapid shifts of the EA state towards more positive values implying stronger links with the subtropical climate and weaker Mistral. Overall, this set of proxy records exhibits a spatial temperature pattern that is coherent in sign with persistent anticyclone blocking and negative EA promoted by low solar activity of the LIA (Fig. 7H), in particular towards the Late LIA. Our interpretation of the GoL data together with high-resolution records distributed across the NA/Euro-Mediterranean region thus supports the hypothesis that EA played an important role in the (late) LIA climate early formulated based on one single record in the NA (RAPiD17-15) by Moffa-Sanchez et al. (2014).

5. Conclusions

A unique SST reconstruction resolving decadal variability was developed over the full Common Era from shelf sediments in the convection region of the GoL (NW Mediterranean Sea). The tight link between alkenone-derived SSTs, Mistral and EA, the dominant mode of variability in the Mediterranean Sea, was used to investigate the climate of the Common Era in the NW-Mediterranean Sea and notably the strong fluctuations of the LIA. Comparison between instrumental and proxy data over the 20th century revealed a similar warming trend of about 2°C. SSTs during the MCA (1000 – 1200 AD) were among the warmest of the pre-industrial last millennium though ~ 1°C lower than those of the 20th century. Most of the first millennium (200 -800 AD) and the LIA both indicate cold climate conditions but the LIA differ by the presence of remarkable cold extremes followed by above average temperature rebounds. Cold decades were found to reflect strong heat loss caused by negative EA and

377 associated NA blocking regimes creating the conditions for intensified and cold Mistral flow.
378 Regional synthesis of high-resolution land and marine time series from the NA and Euro-
379 Mediterranean region further highlighted a W-E temperature asymmetry during the late LIA
380 that is spatially coherent with persistent blocked regimes under negative EA and weak NAO.
381 Reinforced northerly flow (Mistral) over Western Europe would have been favored by low
382 solar activity.

383 **ACKNOWLEDGMENTS:** We thank CNRS and the MISTRALS/PALEOMEX program for
384 financial support and the Ocean2k working group of the Past Global Changes (PAGES)
385 project for discussions. We are very grateful to Nabil Sultan, Ifremer for providing facilities at
386 sea and the crew operating the GMO2 Carnac cruise. We are also thankful for the crew of the
387 N/V Néréis of the “Observatoire Océanographique de Banuyls” (OOB). We also thank the
388 Laboratoire de Mesure du Carbone 14, UMS 2572, ARTEMIS for 14C measurements by
389 mass accelerator spectrometry in the frame of the National Service to CNRS, CEA, IRD,
390 IRSN and Ministère de la Culture et de la Communication. B.M. wishes to express her thanks
391 to the CSIC-Ramón y Cajal post-doctoral program RYC-2013-14073 and the Shackleton
392 Fellowship (Clare Hall College). This dataset is available on the NOAA database
393 (<http://www.esrl.noaa.gov/>).

394

- Alpert, P., Baldi, M., Ilani, R., Krichak, S., Price, C., Rodo, X., Saaroni, H., Ziv, B., Pavel Kishcha, P., Barkan, J., Mariotti, A., and Xoplaki, E., 2006. Relations between climate variability in the Mediterranean region and the tropics: ENSO, South Asian and African monsoons, hurricanes and Saharan dust. *Developments in Earth and Environmental Sciences*, 4, 149-177.
- Barriopedro, D., García-Herrera, R., Huth, R., 2008. Solar modulation of Northern Hemisphere winter blocking. *J. Geophys. Res. Atmos*, 113, D14118.
- Béthoux, J.P., Durieu de Madron, X., Nyffeler, F., Tailliez, D., 2002, Deep water in the western Mediterranean: peculiar 1999 and 2000 characteristics, shelf formation hypothesis, variability since 1970 and geochemical inferences. *J. Mar. Sys.*, 33-34, 117-131.
- Bosc, E., Bricaud, A., Antoine, D., 2004. Seasonal and interannual variability in algal biomass and primary production in the Mediterranean Sea, as derived from 4 years of SeaWiFS observations. *Glob. Biogeochem. Cycles*, 18, 1-17.
- Burlando, M., 2009. The synoptic-scale surface wind climate regimes of the Mediterranean Sea according to the cluster analysis of ERA-40 wind fields, *Theor Appl. Climatol.*, 96, 69-83.
- Cassou, C., Minvielle, M., Terray, L., Perigaud, C., 2011. A statistical-dynamical scheme for reconstructing ocean forcing in the Atlantic. Part I: weather regimes as predictors for ocean surface variables. *Climate Dynamics*, 36, 19-39.
- Conte, M.H., Sicre, M.-A., Rühlemann, C., Weber, J.C., Schulte, S., Schulz-Bull, D., Blanz, T., 2006. Global temperature calibration of the alkenone unsaturation index (UK37) in surface waters and comparison with surface sediments. *Geochem. Geophys. Geosyst.*, 7, Q02005.
- Delworth T.L., Greatbach, R., 2000. Multidecadal thermohaline circulation variability driven by atmospheric surface flux forcing. *J. of Clim.*, 13, 1481-1495.
- Denton, G.H., Broecker, W.S., 2008. Wobbly ocean conveyor circulation during the Holocene? *Quat. Sci. Rev.* 27, 1939-1950
- Durrieu de Madron, X., Houpert, L., Puig, P., Sanchez-Vidal, A., Testor, P., Bosse, A., Estournel, C., Somot, S., Bourrin, F., Bouin, M. N., Beauverger, M., Beguery, L., Calafat, A., Canals, M., Cassou, C., Coppola, L., Dausse, D., D'Ortenzio, F., Font, J., Heussner, S., Kunesch, S., Lefevre, D., Le Goff, H., Martín, J., Mortier, L., Palanques, A., Raimbault, P., 2013. Interaction of dense shelf water

424 cascading and open-sea convection in the northwestern Mediterranean during winter 2012. *Geophys.*
 425 *Res. Lett.* 40, 1379-1385.

426 D'Ortenzio, F., Ribera d'Alcala M., 2009. On the trophic regimes of the Mediterranean Sea: a satellite
 427 analysis. *Biogeosciences*, 6, 1-10.

428 Häkkinen, S., Rhines, P.B., Worthen, D.L., 2011. Atmospheric blocking and Atlantic multidecadal
 429 ocean variability. *Science*, v. 334, no. 6056, p. 655-659.

430 Holzhauser, H., Magny, M., Zumbühl, H.J., 2005. Glacier and lake-level variation in west-central
 431 Europe over the last 3500 years. *The Holocene* 15, 789–801.

432 Hurrell, J.W., 1995. Decadal trends in the North Atlantic Oscillation: regional temperatures and
 433 precipitation. *Science* 269, 676-679.

434 Jiang, Q., Smith, R.B., Doyle, J. 2003. The nature of the mistral: Observations and modelling of two
 435 MAP events. *Q.J.R. Meteorol. Soc.* 129, 857-875. doi:10.1256/qj.02.21

436 Josey, S.A., Somot, S., Tsimplis, M., 2011, Impacts of atmospheric modes of variability on
 437 Mediterranean Sea surface heat exchange. *Journal of Geophysical Research: Oceans*, v. 116, p.
 438 C02032.

439 Kaplan, A., Cane, M., Kushnir, Y. Clement, A., Blumenthal, M., Rajagopalan, B., 1998. Analyses of
 440 global sea surface temperature 1856-1991. *Journal of Geophysical Research*, 103, 18,567-18,589

441 Keigwin, L.D., 1996. The little ice age and medieval warm period in the Sargasso Sea. *Science*, 274,
 442 1504-1508.

443 Knight, J.R., Folland, C.K., Scaife, A.A., 2006. Climate impacts of the Atlantic Multidecadal
 444 Oscillation. *Geophys. Res. Lett.*, 33, L17706, doi:10.1029/2006GL026242.

445 Larsen, D.J., Miller, G.H., Geirsdóttir, Á., 2013. Asynchronous Little Ice Age glacier fluctuations in
 446 Iceland and European Alps linked to shifts in subpolar North Atlantic circulation. *Earth and Planet.*
 447 *Sci. Lett.* 380, 52-59.

448 Lionello, P., Malanotte-Rizzoli, P., Boscolo, R., Alpert, P., Artale, V., Li, L., Luterbacher, J., May, W.,
 449 Trigo, R., Tsimplis, M., Ulbrich, U., Xoplaki, E., 2006. The Mediterranean climate: An overview of
 450 the main characteristics and issues, *in* Lionello P. and Boscolo, R., eds. *Developments in Earth and*
 451 *Environmental Sciences*, Volume 4, Elsevier, p. 1-26.

452 Lund, D.C., Lynch-Steiglitz, J., Curry, W.B., 2006. Gulf Stream density structure and transport during
 453 the past millennium. *Nature*, 444, doi:10.1038/nature05277.

454 Luterbacher, J., Dietrich, D., Xoplaki, E., Grosjean, M., Wanner, H., 2004. European seasonal and
 455 annual temperature, trends and extremes since 1500. *Science*, 303, 1499-1503.

456 Massé, G., Rowland, S.J., Sicre, M.-A., Jacob, J., Jansen, E., Belt, S.T., 2008. Abrupt climate changes
 457 for Iceland during the last millennium: Evidence from high-resolution sea ice reconstructions. *Earth*
 458 *and Planet. Sci. Lett.* 269, 565-569.

459 Marcias-Fauria, M., Grinsted, A. Helama, S. Moore, J. Timonen, M. Martma, T. Isaksson, E. Eronen,
 460 M., 2010. Unprecedented low twentieth century winter sea ice extent in the Mestern Nordic Seas
 461 since A.D. 1200. *Clim. Dyn.* 34, 781-795.

462 Mariotti, A., Dell'Aquila, A., 2012. Decadal variability in the Mediterranean region: role of large scale
 463 forcing and regional processes. *Clim. Dyn.* 38, 1129-1145.

464 Marullo, S., Artale, V., Santoleri, R., 2011. The SST multidecadal variability in the Atlantic-
 465 Mediterranean region and its relation to AMO. *J. of Clim.*, 4385- 4401.

466 Moffa-Sanchez, P., Born, A., Hall, I.R., Thornalley, D.J.R., Barker, S., 2014. Solar forcing of North
 467 Atlantic surface temperature and salinity over the past millennium. *Nature Geosci*, 7, 275-278.

468 McGregor, H.V., Evans, M.N., Goose, H., Leduc, G., Martrat, B., Addison, J.A., Mortyn, P.G., Oppo,
 469 D.W., Seidenkrantz, M.-S., Sicre, M.-A., Phipps, S.J., Selvaraj, K., Thirumalai, K. , Filipsson, H.,
 470 Ersek, V., 2015. Robust global ocean cooling trend for the pre-industrial Common Era. *Nat. Geosci.*
 471 doi:10.1038/NGEO2510.

472 Moore, G.W.K., Renfrew, I.A., 2011. Cold European winters: interplay between the NAO and the East
 473 Atlantic mode. *Atmos. Sci. Lett.* 13, 1-8.

474 Moreno, A., Pérez, A., Frigola, J., Nieto-Moreno, V., Rodrigo-Gámiz, M., Martrat, B., González-
 475 Sampériz, P., Morellón, M., Martín-Puertas, C., Corella, J.P., Belmonte, Á., Sancho, C., Cacho, I.,
 476 Herrera, G., Canals, M., Grimalt, J.O., Jiménez-Espejo, F., Martínez-Ruiz, F., Vegas-Vilarrúbia, T.,
 477 Valero-Garcés, B.L., 2012. The Medieval Climate Anomaly in the Iberian Peninsula reconstructed
 478 from marine and lake records. *Quat. Sci. Rev.*, 43, 16-32.

479 Najac, J., Boè, J., Terray L., 2009. A multi-model ensemble approach for assessment of climate change
 480 impact on surface winds in France. *Clim. Dyn.* 32, 615-634.

481 Nesmé-Ribes, E., Ferreira E.N., Sadourny R., Le Treut H. Li, Z.L., 1993. Solar dynamics and its
 482 impact on solar irradiance and the terrestrial climate. *J. Geophys. Res.* 98, 18923-18935.

483 Nieto-Moreno, V., Martínez-Ruiz, F., Willmott, V., García-Orellana, J., Masqué, P., Sinninghe
484 Damsté, J.S., 2013. Climate conditions in the westernmost Mediterranean over the last two
485 millennia: An integrated biomarker approach. *Org. Geochem.* 55, 1-10.

486 Ólafsdóttir, K.B., Geirsdóttir, Á., Miller, G.H., Larsen, D.J., 2013. Evolution of NAO and AMO
487 strength and cyclicity derived from a 3-ka varve-thickness record from Iceland. *Quat. Sci. Rev.* 69,
488 142-154.

489 Olita, A., Sorgente, R., Ribotti, A., Fazioli, L., Perilli A., 2011. Pelagic primary production in the
490 Alghero-provençal basin by means of multisensory satellite data: focus on interannual variability and
491 its drivers. *Ocean Dyn.*, 61, 1005-1016.

492 Ortega, P., Lehner, F., Swingedouw, D., Masson-Delmotte, V., Raible, C.C., Casado, M., Yiou, P.,
493 2015. A model-tested North Atlantic Oscillation reconstruction for the past millennium. *Nature* 523,
494 doi:10.1038/nature14518.

495 Papadopoulos, V.-P., Josey, S.-A., Bartzokas, A., Somot, S., Ruiz, S., Drakopoulou, P., 2012. Large-
496 Scale Atmospheric Circulation Favoring Deep- and Intermediate-Water Formation in the
497 Mediterranean Sea. *J. Clim.* 25, 6079-6091.

498 Ramsey, C. B., Lee, S., 2013. Recent and planned developments of the program Oxcal. *Radiocarbon*
499 55, 720-730.

500 Reimer, P. J., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Bronk Ramsey, C., Buck, C. E.,
501 Cheng, H., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Haflidason, H., Hajdas,
502 I., Hatté, C., Heaton, T. J., Hoffmann, D. L., Hogg, A. G., Hughen, K. A., Kaiser, K. F., Kromer, B.,
503 Manning, S. W., Niu, M., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Staff, R. A.,
504 Turney, C. S. M., van der Plicht, J., 2013. IntCal13 and Marine13 Radiocarbon Age Calibration
505 Curves 0–50,000 Years cal BP. *Radiocarbon*, , 55, 1869–1887, doi: 10.2458/azu_js_rc.55.16947.

506 Richter, T.O., Peeters, F.J.C., van Weering, T.C.E., 2009. Late Holocene (0-2.4 ka BP) surface water
507 temperature and salinity variability, Feni Drift, NE Atlantic Ocean. *Quat. Sci. Rev.* 28, 1941–1955.

508 Schabbar, A., Huang, J., Higuchi, K., 2001. The relationship between the wintertime north Atlantic
509 Oscillation and blocking episodes in the North Atlantic. *Int. J. of Climatology*, 21, 355-369.

510 Schmidt, S., Howa, H., Diallo, A., Martín, J., Cremer, M., Duros, P., Fontanier, C., Deflandre, B.,
511 Metzger, E., Mulder, Th., 2014. Recent sediment transport and deposition in the Cap-Ferret Canyon,
512 South-East margin of Bay of Biscay. *Deep-Sea Res. II* 104, 134-144.

513 Sicre, M.-A., Hall, I.R., Mignot, J., Khodri, M., Ezat, U., Truong, M.X., Eiríksson, J., Knudsen, K.-L.,
 514 2011. Sea surface temperature variability in the subpolar Atlantic over the last two millennia.
 515 *Paleoceanography*, 26, PA4218.

516 Sicre, M.-A., Yiou, P., Eiriksson, J., Ezat, U., Guimbaut, E., Dahhaoui, I., Knudsen, K.-L., Jansen, E.,
 517 Turon, J.-L., 2008. A 4500-year reconstruction of sea surface temperature variability at decadal time
 518 scales off North Iceland. *Quat. Sci. Rev.* 27, 2041-2047.

519 Skliris, N., Sofianos, S., Gkanasos, A., Mantziafou, A., Vervatis, V., Axaopoulos, P., Lascaratos, A.,
 520 2012. Decadal scale variability of sea surface temperature in the Mediterranean Sea in relation to
 521 atmospheric variability. *Ocean Dyn.* 62, 13-30.

522 Steinhilber, F., Abreu, J.A., Beer, J., Brunner, I., Christl, M., Fischer, H., Heikkilä, U., Kubik, P.W.,
 523 Mann, M., McCracken, K.G., Miller, H., Oerter, H., Wilhelms, F., 2012. 9,400 years of cosmic
 524 radiation and solar activity from ice cores and tree rings. *Proceedings of the National Academy of*
 525 *Sciences* 109, 5967-5971, doi:10.1073/pnas.1118965109.

526 Stuiver, M., Reimer, P.J., 1993. Extended 14C database and revised CALIB 3.0 14C age calibration
 527 program. *Radiocarbon*, 35, 215-230.

528 Ternois, Y., Sicre, M.-A., Boireau, A., Marty, J.-C., Miquel, J.-C., 1996. Production pattern of
 529 alkenones in the Mediterranean Sea. *Geophys. Res. Lett.* 23, 3171-3174.

530 Trouet, V., Esper, J., Graham, N.E., Baker, A., Scourse, J.D., Frank, D.C., 2009. Persistent Positive
 531 North Atlantic Oscillation Mode Dominated the Medieval Climate Anomaly. *Science* 324, 78-80.

532

Figure captions

Fig. 1. Map showing the location of Gol-Ho1B_KSGC-31 site, and spatial field of Mediterranean annual mean SSTs (1955-2012). The insert in the upper right corner shows the monthly SSTs at the core location; the dashed line indicates the annual mean (17°C) from World Ocean Atlas Database (NOAA/NODC WOA13 (1° grid)). The anticyclonic blocking cell over the northeastern Atlantic and low-pressure system in the central Mediterranean associated to EA is also represented (adapted from Papadopoulos et al., 2012). All sites discussed in the text are shown from West to East: RAPiD-17-5P, South of Iceland (Moffa-Sanchez et al., 2014); MD99-2275, North of Iceland (Sicre et al., 2011); TTR17-1_384B, 436B, Alboran basin (Nieto-Moreno et al., 2013); MINMC-1,2, Balearic basin (Moreno et al., 2012). Mistral wind is shown by the black arrow.

Fig. 2. KSGC31 gravity core chronology based on ^{210}Pb (grey dots) and ^{14}C (dark squares) dating and associated error bar at $\pm 1\sigma$ (A). The insert B is an enlarged view of the uppermost 0-30 cm section of the core corresponding to the grey area. The outlier date shown as an empty square was not used in the age model.

Fig. 3. ^{232}Th (A), $^{210}\text{Pb}_{\text{xs}}$ (B) and ^{137}Cs (C) profiles determined in the multicore Gol-Ho1B (dark dots) and gravity core KSGC31 (red squares) (see data in Table 2). KSGC31 profiles have been shifted to meet the best correspondence of the plots, yielding a material loss of 15 ± 0.5 cm in the top of the gravity core. Chronology of the multicore Gol-Ho1B and the top 10 cm of the gravity core KSGC31 after adjustment based on ^{210}Pb dating and associated age model uncertainty at $\pm 1\sigma$ (D).

Fig. 4. SST reconstruction since 1850 AD combining SSTs from the multi-core Gol-Ho1B (light green dashed line and solid dots) and gravity core KSGC-31 (dark green and solid dots). Empty dots indicate the overlapping gravity core SST values with the multicore (A). Spliced Gol-Ho1B_KSGC-31 record built from the two core data (B) (see text for explanations).

Fig. 5. Alkenone-derived SST reconstruction at the Gol-Ho1B_KSGC-31 site over the instrumental period (A). ^{14}C and ^{210}Pb error bars on dating are shown for each SST data point. Kaplan instrumental data from 1866 AD to 2012 AD showing annual mean (in purple)(B), FMA months (in blue)(C) (Kaplan et al., 1998). Annual mean values of the East Atlantic (EA) index since 1950 are also shown for comparison with proxy data (D).

Fig. 6. Alkenone-SST time series from the Western Mediterranean Sea and North Atlantic over the last two millennia. (A) core Gol-Ho1B_KSGC-31 (NW Mediterranean Sea) (this study); (B) Cores TTR17-1_384B, 436B from the Alboran basin (Nieto-Moreno et al., 2013); (C) Cores MINMC-1,2 from the Balearic isles (Moreno et al., 2012); (D) core MD99-2275 off North Iceland (Sicre et al., 2011). For (B) and (C) SSTs were recalculated using the calibration of Conte et al. (2006). Dashed lines in (C) highlight the portion of the record that has been extrapolated to present day from the last dated sediment horizon around 1500 AD (see text). Filled areas of the curves indicate negative SST anomaly relative to mean values of each core. A 25-yr binning was applied to all records (dark and red lines) to better visualize anomalies or departures from average conditions. Triangles indicate the AMS ^{14}C control points for all cores. The grey vertical bars broadly highlight the most outstanding cold spells of core Gol-Ho1B_KSGC-31 during the LIA. MCA: Medieval Climate Anomaly; LIA: Little Ice Age.

Fig. 7. A detailed view of the last millennium. (A) Fluctuations of the Great Aletsch and Gorner glaciers in the Swiss Alps (Holzhauser et al., 2005); (B) Retreat/advance of the Langjökull glacier (Iceland) reconstructed from the Hvítárvatn lake varves (Larsen et al., 2013; Ólafsdóttir et al., 2013); (C) Sea ice occurrence traced by the IP25 abundances (Massé et al., 2008) and alkenone-SSTs from core MD99-2275, North Iceland (Sicre et al., 2011); (D) Mg/Ca-SSTs obtained from foraminifera calcite of *G. inflata* in core RAPiD-17-5P, South Iceland (Moffa-Sanchez et al., 2014); (E) Alkenone-SSTs from site Gol-Ho1B_KSGC-31 in the Gulf of Lion (this study); (F) 30-yr Gaussian low-pass filter of winter (DJF) European temperature anomaly (relative to the 1901 - 1995 calibration average) (black line) with two

585 standard errors (blue lines) (Luterbacher et al., 2004); (G) The NAO index anomalies
586 calculated in 25-yr averages from the palæo-reconstruction by Ortega et al. (2015) (black
587 line); 7-yr running average of the NAO instrumental (red line)
588 (http://www.esrl.noaa.gov/psd/gcos_wgsp/Timeseries/NAO/nao.long.data). (H) Solar activity
589 anomalies in W/m^2 (Total Solar Irradiance, TSI) calculated in 25-yr averages from the cosmic
590 ray intensity reconstruction (Steinhilber et al., 2012). From (B) to (E) dark and red lines
591 indicate a 25-yr binning. From (C) to (H) filled areas indicate negative anomaly relative to
592 average values, except for (F) whose anomaly was calculated relative to the 1901-1995
593 calibration interval. Triangles indicate the AMS- ^{14}C control points at 1σ uncertainty. The grey
594 vertical bars broadly highlight the major cold spells of the LIA period in the Gol-
595 Ho1B_KSGC-31 record.

Table 1. Radiocarbon dates and their calibrated ages along the KSGC31 sediment core. Results are reported with a 1 σ uncertainty.

Depth (cm)	Calibration	Material	¹⁴ C age	cal year BP	Year AD	± 1 σ
5.5	Oxcal 4.2, NH zone 1 post bomb ages curve	<i>Bittium</i> sp.	420± 30	24 ^a	1926	60
11.5	Oxcal 4.2, NH zone 1 post bomb ages curve	<i>Tellina</i> sp.	430±30	34 ^a	1916	60
18.5	CALIB 7.1, Marine13 curve (Reimer et al., 2013)	<i>Pecten</i> sp.	720± 40	380 ^b	1570	78
25.5	CALIB 7.1, Marine13 curve (Reimer et al., 2013)	<i>Venus</i> sp.	640 ± 30	235	1716	99.5
41.5	CALIB 7.1, Marine13 curve (Reimer et al., 2013)	<i>Pecten</i> sp.	700±30	339	1611	79
52	CALIB 7.1, Marine13 curve (Reimer et al., 2013)	Indet. bivalve	960±30	551	1399	59
71	CALIB 7.1, Marine13 curve (Reimer et al., 2013)	<i>Arca tetragona</i>	1340±30	852	1099	80.5
110.5	CALIB 7.1, Marine13 curve (Reimer et al., 2013)	<i>Venus</i> sp.	1465±30	992	958	85
185.5	CALIB 7.1, Marine13 curve (Reimer et al., 2013)	<i>Nucula</i> sp.	2235±40	1806	145	99.5
252	CALIB 7.1, Marine13 curve (Reimer et al., 2013)	juvenile bivalve shells (ind.)	2940±30	2676	-726	100.5

^a KSGC-31 core top ages are derived from 210 Pb xs profile (see methods, Table 2)

^b Reversal, not used for the interpolation

Depth (in cm)	²¹⁰ Pb _{xs} mBq g ⁻¹			²³² Th mBq g ⁻¹			¹³⁷ Cs mBq g ⁻¹			Age in years AD Mean		
Multi-core GOL-HO-1-1-B												
0.5	72	±	9	22	±	1	3	±	1	2012	±	0
1.5										2010	±	0
2.5	74	±	9	22	±	1	4	±	1	2008	±	1
3.5										2006	±	1
4.5	73	±	8	21	±	1	4	±	1	2003	±	1
5.5	64	±	8	22	±	1	4	±	1	2001	±	1
6.5	58	±	5	21	±	1	4	±	0	1998	±	2
7.5										1996	±	2
8.5	52	±	5	23	±	1	4	±	0	1993	±	2
9.5										1990	±	2
10.5	45	±	7	25	±	1	4	±	1	1987	±	3
11.5										1985	±	3
12.5	48	±	5	23	±	1	4	±	0	1982	±	3
13.5										1979	±	4
14.5	27	±	4	26	±	1	4	±	0	1976	±	4
15.5	30	±	5	25	±	1	3	±	0	1973	±	4
16.5	32	±	4	25	±	1	3	±	0	1970	±	5
17.5										1967	±	5
18.5	21	±	4	28	±	1	2	±	0	1963	±	5
Gravity core KS-GC-31												
0.5	28	±	5	26	±	1	3	±	0	1971	±	1
1.5										1965	±	2
2.5	17	±	6	28	±	1	2	±	1	1959	±	3
3.5										1953	±	5
4.5	21	±	6	30	±	1	2	±	1	1946	±	6
5.5	20	±	6	29	±	1	1	±	1	1940	±	8
6.5	14	±	6	31	±	1	1	±	1	1934	±	10
7.5	8	±	5	29	±	1	0	±	1	1928	±	11
8.5	5	±	5	29	±	1	0	±	0	1922	±	13

Figure 1

Age (years AD)

-1000

0

1000

2000

0

A/ Age:

---■--- ^{14}C

● ^{210}Pb

50

100

150

200

250

Depth (cm)

B/ Zoom 0-30 cm

0

Depth (cm)

10

20

30

1500

2000

Depth (cm)

Figure 4

Figure 5

cal year AD

0 200 400 600 800 1000 1200 1400 1600 1800 2000

MCA

LIA

Figure 6

Figure 7