Limited impact of ocean acidification on phytoplankton community structure and carbon export in an oligotrophic environment: Results from two short-term mesocosm studies in the Mediterranean Sea

Frédéric Gazeau, A. Sallon, P. Pitta, A. Tsiola, L. Maugendre, M. Giani, M. Celussi, M.L. Pedrotti, S. Marro, Cecile Guieu

► To cite this version:

Frédéric Gazeau, A. Sallon, P. Pitta, A. Tsiola, L. Maugendre, et al.. Limited impact of ocean acidification on phytoplankton community structure and carbon export in an oligotrophic environment: Results from two short-term mesocosm studies in the Mediterranean Sea. *Estuarine, Coastal and Shelf Science*, 2017, 186 (A), pp.72-88. 10.1016/j.ecss.2016.11.016 . hal-01401698

HAL Id: hal-01401698

<https://hal.sorbonne-universite.fr/hal-01401698>

Submitted on 23 Nov 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Limited impact of ocean acidification on phytoplankton community structure and carbon export in an oligotrophic environment: results from two short-term mesocosm studies in the Mediterranean Sea

Gazeau, F.^{1,2*}, Sallon, A.^{1,2}, Pitta, P.³, Tsiola, A.^{3,4}, Maugendre, L.^{1,2}, Giani M.⁵, Celussi M.⁵, Pedrotti, M. L.^{1,2}, Marro, S.^{1,2} and Guieu, C.^{1,2}

[1] Sorbonne Universités, UPMC Univ Paris 06, UMR 7093, LOV, Observatoire océanologique, F-06230, Villefranche/mer, France

[2] CNRS, UMR 7093, LOV, Observatoire océanologique, F-06230, Villefranche/mer, France

[3] Hellenic Centre for Marine Research, Institute of Oceanography, PO Box 2214, 71003 Heraklion, Crete, Greece

[4] University of Crete, Department of Biology, University Campus, 70013, Heraklion, Crete, Greece

[5] Oceanography Division, OGS (Istituto Nazionale di Oceanografia e di Geofisica Sperimentale), v. A. Piccard 54, I-34151 Trieste, Italy

* Correspondence:

Dr. Frédéric Gazeau

Laboratoire d'Océanographie de Villefranche

CNRS-UPMC, UMR 7093

06230 Villefranche-sur-mer

FRANCE

f.gazeau@obs-vlfr.fr

Highlights:

- Two large mesocosm experiments carried out in the Northwestern Mediterranean Sea
- Experiments conducted in the summer oligotrophic vs. winter mesotrophic periods
- Production limited by nutrient availability and community dominated by small species
- Organic matter export was not impacted by CO₂-enrichment
- In areas where nutrient availability exerts a strong pressure on phytoplankton growth, CO₂ addition will likely have very limited effects on phytoplankton diversity

Keywords:

Ocean acidification; Pelagic mesocosms; Mediterranean Sea; Oligotrophic area; Phytoplankton community

Abstract

Modifications in the strength of the biological pump as a consequence of ocean acidification, whether positive or negative, have the potential to impact atmospheric CO₂ and therefore climate. So far, most plankton community perturbation studies have been performed in nutrient-rich areas although there are some indications that CO₂-dependent growth could differ in nutrient-replete vs. -limited regions and with different community compositions. Two *in situ* mesocosm experiments were performed in the NW Mediterranean Sea during two seasons with contrasted environmental conditions: summer oligotrophic stratified waters in the Bay of Calvi vs. winter mesotrophic well-mixed waters in the Bay of Villefranche. Nine mesocosms were deployed for 20 and 12 d, respectively, and subjected to seven CO₂ levels (3 controls, 6 elevated levels). Both phytoplankton assemblages were dominated by pico- and nano-phytoplankton cells. Although haptophyceae and dinoflagellates benefited from short-term CO₂ enrichment in summer, their response remained small with no consequences on organic matter export due to strong environmental constraints (nutrient availability). In winter, most of the plankton growth and associated nutrient consumption occurred during the 4-day acidification period (before the experimental phase). During the remaining experimental period, characterized by low nutrient availability, plankton growth was minimal and no clear CO₂-dependency was found for any of the tested parameters. While there is a strong confidence on the absence of significant effect of short-term CO₂ addition under oligotrophic conditions, more investigations are needed to assess the response of plankton communities in winter when vertical mixing and weather conditions are major factors controlling plankton dynamics.

1. Introduction

During the last 150 years, human activities, through the combustion of fossil fuels (oil, gas and coal), have led to a dramatic release of carbon dioxide (CO₂) to the Earth's atmosphere. The accumulation of CO₂ impacts the radiative forcing, thereby warming the atmosphere and the ocean. The ocean acts as a climate integrator that absorbed 93% of Earth's additional heat since the 1970s, offsetting much atmospheric warming but increasing ocean temperature and sea level and captured 28% of anthropogenic CO₂ emissions since 1750 (Gattuso et al., 2015). Although providing a valuable human service by moderating the rate and severity of global warming, the consequence of this oceanic CO₂ pump is the ongoing increase in ocean acidity (i.e. decrease in pH). Surface ocean pH has already decreased by 0.1 units since the beginning of the industrial era (i.e. increased acidity of 30%; Ciais et al., 2013). According to recent projections and depending on the emission scenario considered, an additional decrease ranging between 0.06 and 0.32 units is expected by 2100 (Ciais et al., 2013).

The decrease in seawater pH leads to a decrease in the concentration of carbonate ions (CO₃²⁻), one of the building blocks of calcium carbonate (CaCO₃), and alters the ability of many calcifying organisms to precipitate CaCO₃ (e.g. Kroeker et al., 2013). In addition, a decrease in seawater pH leads to an increase in dissolved CO₂ and bicarbonate (HCO₃⁻) concentrations. Carbon fixation by marine photosynthetic organisms represents about 50% of global Earth primary production (Field et al., 1998), and the export of part of the produced organic matter from the sunlit surface layer to the deep-ocean (i.e. the biological or soft-tissue pump) is responsible for ~70% of surface to deep-ocean dissolved inorganic carbon (C_T) gradients (Sarmiento and Gruber, 2006). Therefore, modifications in the strength of this

biotically mediated carbon pump, whether positive or negative, have the potential to impact atmospheric CO₂ and therefore climate (Riebesell et al., 2007).

CO₂ rather than the much more abundant HCO₃⁻ is the substrate used in the carbon fixation step of photosynthesis and RubisCO, the enzyme catalyzing this reaction, has a low affinity for CO₂ (Badger et al., 1998; Giordano et al., 2005). As such, this enzyme is theoretically not saturated under current ambient CO₂ levels (Badger et al., 1998). However, nearly all marine species have developed various mechanisms (carbon concentration mechanisms or CCMs) to compensate for this low CO₂ availability through the energy-demanding use of carbonic anhydrase enzymes or active CO₂ and/or bicarbonate transports through membranes (Raven et al., 2014). There is evidence that both the RubisCO affinity for CO₂ as well as the efficiency of these CCMs differ widely among taxa, species or even strains (Tortell, 2000; Young et al., 2016), complicating the prediction of whether a cell's carbon fixation rate will respond directly to ambient changes in CO₂ availability through increased CO₂ diffusion and/or less energy expenditure needed to operate CCMs (Mangan et al., 2016; Raven and Beardall, 2014). Finally, although downregulation of CCMs at elevated CO₂ has been observed, the significance of this downregulation to overall cell physiology and growth is not currently well constrained due to the presence of other limiting factors in the oceans such as macro- or micro-nutrients and light (Hennon et al., 2015; Young and Morel, 2015). All of this can partly explain the very diverse findings that have been documented on the effect of increased ambient CO₂ availability on photosynthesis and growth of marine phytoplankton (Dutkiewicz et al., 2015).

Apart from the above-mentioned variability in RubisCO affinity for CO₂ and CCMs efficiency, a significant part of the observed discrepancies among available perturbation studies could be explained by differences in experimental setups and environmental conditions such as temperature, light conditions and nutrient availability. Phytoplankton

growth obviously does not only depend on carbon availability but on a combination of physico-chemico-biological drivers such as macro- and micro-nutrient availability, temperature, light, competition and grazing. It is therefore very likely that the response of phytoplankton will differ depending on these environmental conditions (Verspagen et al., 2014). Furthermore, as this is the amount of organic matter that can escapes the sunlit layer that determines the capacity of the surface ocean to pump atmospheric CO₂, there is a great need to evaluate the impact of CO₂, not only on phytoplankton growth but on the export of this organic matter to deeper layers. The build-up of organic matter and its potential export strongly depends on phytoplankton community composition (Eggers et al., 2014). Indeed, large cells (e.g. diatoms) account for a large proportion of export production and ultimate burial in sediments (Finkel et al., 2005). In contrast, small cells (nano- and pico-plankton) are particularly important in regions with limited nutrient availability with a close coupling between production and grazing through the microbial loop and a with low export capacity (Riebesell and Tortell, 2011). As already mentioned, differing responses to increased CO₂ availability between different functional groups, size classes and species (Dutkiewicz et al., 2015) have the potential to significantly alter community structure and functioning. In that sense, studies focused on plankton assemblages rather than on isolated single species and under very contrasted environmental conditions are very informative (Tarling et al., 2016).

During the last decade, there has been a noticeable increase in the number of experimental assessments of the sensitivity of plankton community compositions to the on-going increase in CO₂. These experiments were conducted in various areas of the world ocean using different approaches, from small bottle incubations to large mesocosm deployments, and over different time scales (few days to few weeks). Several of these experiments highlighted significant modifications of community compositions under elevated CO₂ levels. For instance, CO₂ enrichment has been shown to stimulate growth of large species such as

diatoms (e.g. Domingues et al., 2014; Feng et al., 2009; Reul et al., 2014; Tortell et al., 2002; Tortell et al., 2008; Wu et al., 2014). Several experiments suggested stimulating effects on small species (pico-phytoplankton; e.g. Newbold et al., 2012; Paulino et al., 2008; Schulz et al., 2013). In contrast, Richier et al. (2014) reported significant decrease in the growth of small phytoplankton species ($< 10 \mu\text{m}$) suggesting that small species are less adapted to changes in their local pH while larger cells must face larger pH variations at short time scales (Flynn et al., 2012). Other studies showed differential responses between species from the same taxa (e.g. Endo et al., 2016; Feng et al., 2010; Kim et al., 2006; Meakin and Wyman, 2011) and finally among different phylotypes and phenotypes of the same species (e.g. Brading et al., 2011; Rickaby et al., 2016).

Whether or not these modifications of community structure (e.g. increase or decrease in cell size) can modify the amount of organic matter sinking to deeper layers can be evaluated through the use of mesocosms. They are defined as experimental enclosures from 1 thousand to several thousands of litres that allow the maintenance of natural communities under close-to-natural conditions and the collection of sinking organic matter (Riebesell et al., 2008; Riebesell et al., 2013a). In recent years, plankton community studies performed using such experimental systems have led to very contrasted outcomes in terms of community composition and carbon export responses to CO_2 enrichment (see Table 1). Most of these experiments have been performed in nutrient-rich areas (or following artificial nutrient enrichment) dominated by large species and experiments conducted in areas limited by nitrate, phosphate and/or iron are currently lacking (Paul et al., 2015a). These areas represent a very large surface area of the ocean and are projected to expand in the coming decades because of enhanced thermal stratification and nutrient depletion (Irwin and Oliver, 2009; Polovina et al., 2008). As already mentioned, they are usually dominated by small cells adapted to low-nutrient conditions and have low export capacities. Recently, and in contrast

to theoretical considerations (Verspagen et al., 2014), two mesocosm experiments suggested that communities exposed to low nutrient concentrations may be more responsive to CO₂ enrichment than previously thought (Bach et al., 2016; Paul et al., 2015a). This was confirmed recently by Sala et al. (2016) based on indoor experiments in a coastal site of the Western Mediterranean Sea. During these experiments, effects of ocean acidification, i.e. positive effect on pico- and nano-phytoplankton, were more important when nutrient concentrations were low. However, it must be stressed that nutrient and chlorophyll levels observed during these experiments were representative of an urbanized coastal area and much higher than levels usually observed in the vast majority of the Mediterranean Sea.

The Mediterranean Sea is generally considered as oligotrophic but actually exhibits a gradient from mesotrophic-oligotrophic in the western basin to ultra-oligotrophic in the eastern basin (The Mermex group, 2011). These features are induced by the different localizations of the physical (the winter mixed layer) and nutrient (the nutricline) vertical interfaces, which are both determined by the large-scale circulation pattern (The Mermex group, 2011). Based on satellite-derived estimates, chlorophyll *a* concentrations exhibit low values (less than 0.2 µg L⁻¹) over most of the Mediterranean Sea, except for the Liguro-Provençal region where large blooms can be observed in late winter-early spring (D'Ortenzio and d'Alcala, 2009). Overall, phytoplankton communities are dominated by pico-phytoplankton (Siokou-Frangou et al., 2010). However, because of its very diversified (spatially and temporally) physical structure, localized higher nutrient availability can drive more intense biological activities and transient dominance of larger species such as diatoms and dinoflagellates (Bustillos-Guzmán et al., 1995). Diatoms are more abundant during the transition between mixed and stratified conditions (Claustre et al., 1994). These features make the Mediterranean Sea a perfect natural laboratory to study the effects of nutrient availability and community composition on the response of plankton community to CO₂ enrichment.

In the frame of the European project ‘Mediterranean Sea Acidification under changing climate’ (MedSeA; <http://medsea-project.eu>), for the first time, two short-term *in situ* mesocosm experiments were performed in the Northwestern Mediterranean Sea during two seasons with contrasted environmental conditions (i.e. summer oligotrophic stratified waters vs. winter mesotrophic well-mixed waters) and different phytoplankton community compositions (i.e. higher proportion of diatoms and lower proportion of pico-phytoplankton and cyanophyceae in winter compared to summer). In this paper, we report on the response of the phytoplankton community composition as well as of particulate organic matter dynamics and export to CO₂-enrichment.

2. Material and Methods

2.1. Study sites and experimental set-up

Two mesocosm experiments were conducted in the Northwestern Mediterranean Sea: the first one, in the Bay of Calvi (Corsica, France) in summer (June-July 2012), and the second one in the Bay of Villefranche (France) in winter (February-March 2013). The experimental set-up and mesocosm characteristics are fully described in Gazeau et al. (in press, this issue). Briefly, for each experiment, nine mesocosms of ca. 50 m³ (2.3 m in diameter and 12 m deep) were deployed for 20 and 12 days in the Bay of Calvi and the Bay of Villefranche, respectively. Once the bottom of the mesocosms was closed, CO₂ saturated seawater was added to obtain a *p*CO₂ gradient across mesocosms ranging from ambient levels to 1,250 µatm, with three control mesocosms (C1, C2 and C3) and six mesocosms with increasing *p*CO₂ (P1 to P6). In the Bay of Calvi, the six targeted elevated *p*CO₂ levels were P1: 550, P2: 650, P3: 750, P4: 850, P5: 1000 and P6: 1250 µatm. In the Bay of Villefranche, the levels were P1: 450, P2: 550, P3: 750, P4: 850, P5: 1000 and P6: 1250 µatm. Mesocosms were grouped in clusters of 3 with each cluster containing a control, a medium and a high

$p\text{CO}_2$ level (cluster 1: C1, P1, P4; cluster 2: C2, P2, P5 and cluster 3: C3, P3, P6). Acidification of the mesocosms was performed over four days by homogenous addition of various volumes of CO_2 -saturated seawater. Once targeted $p\text{CO}_2$ levels were reached, the experiment started (day 0 = 24 June 2012 and 22 February 2013 for the Bay of Calvi and the Bay of Villefranche, respectively). No further CO_2 additions were performed during the experiments and $p\text{CO}_2$ levels evolved in mesocosms as a consequence of air-sea fluxes, temperature changes and plankton net community production. Weather permitting, conductivity-temperature-depth (CTD) casts were performed on a daily basis in each mesocosm and in the external environment. Surface irradiance (photosynthetically active radiation; PAR) was measured continuously during the two experiments using a LI-COR LI-192SA 2-Pi sensor connected to a LI-1400 data logger (Gazeau et al., in press, this issue). Vertical attenuation coefficients were estimated daily in each mesocosms, based on PAR profiles (0-12 m) performed using a QSP-2200 4-Pi sensor (Biospherical Instruments Inc.) mounted on the CTD. Mean daily photon doses were calculated using surface PAR and estimated attenuation coefficients. In the Bay of Calvi, wind speed and direction were recorded with a THIES© anemometer deployed, by the University of Liège (Belgium), on top of one of buildings of the at 11.8 m height at a distance of about 400 m from the mesocosms. For the experiment in the Bay of Villefranche, wind speed data (daily averages) were obtained from the Météo France station at the Nice-Côte d'Azur International Airport (43°39'55" N, 7°12'48" E).

2.2. Sampling and analytical methods

Depth-integrated (0-10 m) samplings from the mesocosms and the external environment (referred thereafter to as OUT) were performed daily at 8:30 (local time) during both experiments. All three clusters were simultaneously sampled from a plastic platform by three teams of two scientists, each using an integrating water sampler (IWS; HYDRO-

BIOS©). The IWS units were hanged on a Kevlar cordage and downcasts were performed manually at a regular speed of 10 cm s^{-1} after rinsing it outside the mesocosms.

Samples for pigment determination were taken every day at 8:30 (local time) during both experiments. Two litres of sampled seawater were filtered onto GF/F. Filters were directly frozen in liquid nitrogen and stored at -80°C pending analysis at the Laboratoire d'Océanographie de Villefranche (France). Filters were extracted at -20°C in 3 mL methanol (100%), disrupted by sonication and clarified one hour later by vacuum filtration through GF/F filters. The extracts were rapidly analyzed (within 24 h) by high performance liquid chromatography (HPLC) with a complete Agilent Technologies system. The pigments were separated and quantified as described in Ras et al. (2008).

Synechococcus, *Prochlorococcus*, autotrophic pico-eukaryotes and nano-eukaryotes abundances were determined by flow cytometry analysis from samples taken every 2 days at 4:00 and 5:00 in the Bay of Calvi and the Bay of Villefranche, respectively (local times). Seawater samples (2 mL) from each mesocosm were immediately fixed with $0.2 \mu\text{m}$ pre-filtered 25% glutaraldehyde (0.5% final concentration), kept at 4°C for approximately 30 min, then flash frozen in liquid nitrogen and finally stored at -80°C until further processing (Troussellier et al., 1995; Vaulot et al., 1989). Single cell analysis was processed with a maximum flow rate of $65 \mu\text{L min}^{-1}$ through a Becton Dickinson, FACSCalibur flow cytometer, equipped with an air-cooled Argon laser emitting at 488 nm and analyzed with the Cell Quest Pro software (Becton Dickinson). The sample volume analyzed per time unit was accurately defined by systematically adding to the samples fluorescent latex beads suspensions of $1 \mu\text{m}$ (Polysciences Inc., Europe) at a concentration of $2.5 \times 10^5 \text{ beads mL}^{-1}$. The abundance of autotrophic prokaryotes and pico- and nano eukaryotes was assessed from unstained samples following the method described by Marie et al. (1999). In the Bay of Villefranche, four groups were determined based on the optical parameters characterizing

each cell. *Synechococcus* ($< 1.5 \mu\text{m}$) cells were detected by their signature in a plot of orange fluorescence (FL2, 565–592 nm wavelength ranges) vs. red fluorescence (FL3, $> 620 \text{ nm}$). *Prochlorococcus*, autotrophic pico-eukaryotes ($< 2 \mu\text{m}$) and nano-eukaryotes ($2 - 10 \mu\text{m}$) were detected in a plot of SSC vs. red fluorescence (FL3, $> 620 \text{ nm}$). In the Bay of Calvi, only abundances of *Synechococcus* spp. and autotrophic pico-eukaryotes were assessed.

For particulate element concentrations, sampled seawater (0.75 - 2 L) was filtered through pre-combusted glass-fiber filters. Particulate organic carbon (POC) and nitrogen (PON) were determined at the Istituto Nazionale di Oceanografia e di Geofisica Sperimentale (Italy) using a CHNO-S elemental analyzer (Costech ECS4010) after acidification with 1 N HCl and high-temperature combustion.

Collection of sediment traps was performed by a diver on a daily basis in the Bay of Calvi and less regularly as a consequence of bad weather conditions in the Bay of Villefranche (see Gazeau et al., in press, this issue). On each occasion and for each mesocosm, divers followed the same procedure: (1) hitting the cone of the mesocosms in case some sinking material was retained on the walls, (2) waiting for 15 minutes, (3) closing the collector, (4) collecting the 250 mL flask screwed to the trap system, (5) immediately replacing the sampled flask by a new empty one and (6) opening the collector again. All mesocosms were sampled within 30 min. Back in the laboratory, samples were immediately preserved in a pH buffered formalin solution (5%). Swimmers (i.e. opportunistic copepods and other zooplankton species that swim into the traps; Lee et al., 1988) larger than 1 mm were removed (and discarded) and the remaining material was rinsed, centrifuged, freeze-dried and grinded. In the Bay of Calvi, as a consequence of low amounts of material especially at the end of the experiment, daily sediment traps samples were pooled as follows: days 5-7, 8-10, 11-14 and 15-19. Total particulate matter was weighed for flux determination and subsamples were used for POC and PON measurements performed on elemental

analyzers after acidification with 1N HCl. Samples from the experiment in the Bay of Calvi were analyzed at NIOZ-Yerseke (The Netherlands) on a Thermo Electron Flash 1112. Samples from the experiment in the Bay of Villefranche were analyzed at the Laboratoire d'Océanographie de Villefranche (LOV, France) on a Elementar Vario Pyrocube.

2.3. Data analysis and statistics

The contribution of each phytoplankton group to total phytoplankton biomass (chlorophyll *a*) was estimated by using the CHEMTAX program with input ratios from Rodriguez et al. (2006) and Not et al. (2007). These pigment ratios established for open ocean plankton communities were modified from the original values by comparing microscopic and flow cytometry counts to HPLC analyses from samples collected in the NW Iberian coast, an area dominated by pico- and nano-eukaryotes, as observed in our study.

All data collected during the two experiments are freely available on Pangaea, Bay of Calvi: <http://doi.pangaea.de/10.1594/PANGAEA.810331> and Bay of Villefranche: <http://doi.pangaea.de/10.1594/PANGAEA.835117>.

For these two experiments, we chose to follow a CO₂ gradient approach rather than to replicate certain levels (ANOVA approach). As already done in several similar perturbation experiments (Paul et al., 2015a; Riebesell et al., 2013b), stepwise multiple linear regression analyses were performed to establish relationships between environmental/experimental conditions including *p*CO₂ and (1) POC and PON fluxes to the sediment traps as well as their ratios, (2) water column POC and PON concentrations and their ratios as well as (3) chlorophyll *a*-equivalent biomass or abundances of the different identified groups. Besides *p*CO₂, other environmental conditions that have been considered were temperature, salinity, daily photon doses, daily averaged wind speeds and nutrient concentrations (NO_x: NO₃⁻ + NO₂⁻, ammonium: NH₄⁺, phosphate: PO₄³⁻, and, only for diatoms, silicate: Si). Integrated levels of temperature and salinity were acquired through the daily CTD casts performed in

each mesocosm. NO_x and phosphate were measured using nanomolar techniques as described in Louis et al. (in press, this issue). Ammonium and silicate concentrations were determined as described by Gazeau et al. (in press, this issue). As fully described in Gazeau et al. (in press, this issue), daily $p\text{CO}_2$ levels in each mesocosm were determined from dissolved inorganic carbon, total alkalinity, temperature and salinity using the R package seacarb (Lavigne et al., 2014).

All analyses were performed using the R software (R Core Team, 2015) and were considered significant at a probability $p < 0.01$.

3. Results

3.1. Environmental and experimental conditions during both experiments

Conditions in each mesocosm at the start and at the end of both experiments (days 0 and 20 in the Bay of Calvi and days 0 and 12 in the Bay of Villefranche) are shown in Tables 2 and 3. For both experiments, $p\text{CO}_2$ values in CO_2 -enriched mesocosms (P1 to P6) were close to targeted levels. Ambient $p\text{CO}_2$ levels were higher in the Bay of Calvi in summer as compared to the Bay of Villefranche in winter (~ 450 vs. $350 \mu\text{atm}$ respectively). While $p\text{CO}_2$ levels slightly decreased (pH levels slightly increased) in the Bay of Calvi during the course of the experiment, especially for high CO_2 mesocosms (P5 and P6), drops in $p\text{CO}_2$ levels (increases in pH levels) were much stronger in the Bay of Villefranche with mesocosms P1 to P4 showing very similar levels by the end of the experiment (Fig. 1). Hydrological data (temperature and salinity) are fully described in Gazeau et al. (in press, this issue). Briefly, while temperature levels in the Bay of Calvi gradually increased from $\sim 22.1^\circ\text{C}$ on day 0 to $\sim 24.2^\circ\text{C}$ on day 20, they were constant in the Bay of Villefranche at around $\sim 13.2^\circ\text{C}$. Salinity increased roughly by 0.1-0.2 units during both experiments because of evaporation. In winter in the Bay of Villefranche, surface irradiance was generally constant during the entire experiment with minimal and maximal daily (sunrise to sunset) average values of 531 and $735 \mu\text{mol photons m}^{-2} \text{s}^{-1}$. Maximum irradiance levels (~ 1300 - $1400 \mu\text{mol photons m}^{-2} \text{s}^{-1}$) were reached at around 12:00 pm and the Light:Darkness (L:D) cycle was 16.5:7.5 and 16:8, respectively at the start and at the end of the experiment. In the Bay of Villefranche, minimal and maximal daily (sunrise to sunset) average values of 103 and $513 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ were recorded with a L:D regime of 11.5:12.5, and maximal irradiance levels (~ 300 - $1100 \mu\text{mol photons m}^{-2} \text{s}^{-1}$) reached at 1:00 pm. Light attenuation coefficients were generally constants during both experiments with higher values estimated in winter in the Bay of Villefranche than in summer in the bay of Calvi ($0.19 \pm 0.07 \text{ SD}$ and $0.14 \pm 0.05 \text{ SD m}^{-1}$,

respectively). Higher daily averaged wind speeds were recorded during the winter experiment in the Bay of Villefranche with very windy conditions experienced on day 8 that prevented sampling during that day, and even winder on day 12 and the following night that irreversibly damaged the bags.

In summer in the Bay of Calvi, NO_x concentrations initially decreased then increased again after day 14 to reach similar levels than at the start of the experiment (47 ± 20 on day 0 vs. $60 \pm 15 \text{ nmol L}^{-1}$ on day 20; average $\pm$ SD between the nine mesocosms). Dissolved inorganic phosphate (PO_4^{3-}) quickly decreased from day 0 to day 1 and remained constant during the rest of the experiment (23 ± 12 on day 0 vs. $7 \pm 2 \text{ nmol L}^{-1}$ on day 20). In winter in the Bay of Villefranche, PO_4^{3-} concentrations were generally similar to the ones encountered in summer in the Bay of Calvi and no strong variations could be observed in all mesocosms along the course of the experiment (global average: $9 \pm 1 \text{ nM}$). NO_x levels were much higher in the Bay of Villefranche than in the Bay of Calvi when bags were closed ($> 1 \text{ }\mu\text{M}$). However, during the acidification phase (day-4 to -1), due to favorable weather conditions (low wind, high irradiance levels, data not shown), chlorophyll *a* concentrations increased, consuming a large proportion of available nitrate and nitrite before the start of the experimental phase (day 0). As a consequence, while $[\text{NO}_x]$ in external waters remained high ($\sim 1 \text{ }\mu\text{M}$), all mesocosms were depleted in NO_x with an average concentration of $129 \pm 30 \text{ nM}$. NO_x to phosphate ratios were higher in the Bay of Villefranche than in the Bay of Calvi (2 ± 1 and 9 ± 4 on day 0 and on day 20 in the Bay of Calvi vs. 13 ± 4 and 33 ± 18 on day 0 and on day 12 in the Bay of Villefranche). More details on nutrient dynamics can be found in Louis et al. (in press, this issue).

The experiment in the Bay of Calvi was representative of summer conditions in the Ligurian Sea with low nutrient concentrations, low chlorophyll *a* concentrations (see below), warm waters and high irradiance levels. In the Bay of Villefranche in winter, while

hydrological and weather conditions were typical of winter conditions in the Northwestern Mediterranean Sea (low temperature and irradiance levels), nutrients were rapidly depleted inside the mesocosms before the start of the experiment, and reached levels not usually encountered during this period of the year.

3.2. Phytoplankton assemblages during the summer experiment in the Bay of Calvi

Total chlorophyll *a* concentrations in the Bay of calvi (Fig. 2) averaged $0.07 \pm 0.01 \mu\text{g L}^{-1}$ in the nine mesocosms along the experiment, a value much lower than that in the surrounding waters ($0.12 \pm 0.02 \mu\text{g L}^{-1}$). In mesocosms, chlorophyll *a* concentrations linearly increased during the experiment (GLM, $r^2 = 0.6$, $p < 0.001$) with a maximal concentration of $0.09 \pm 0.003 \mu\text{g L}^{-1}$ on day 14.

When pigment data for all mesocosms were pooled together (Fig. 3), the plankton community in the Bay of Calvi was found to be dominated at the start of the experiment by haptophyceae representing $36 \pm 5\%$ of the chlorophyll content, followed by cyanophyceae ($20 \pm 3\%$), chlorophyceae ($14 \pm 3\%$) and pelagophyceae ($11 \pm 2\%$). Important differences were identified along the experiment between concentrations of the different species inside the mesocosms and in the surrounding waters (Fig. 4). All species, except for diatoms, showed lower chlorophyll *a* biomass inside mesocosms. Diatoms were virtually absent in the surrounding waters, except at the end of the experiment. On day 20, while the contribution of cyanophyceae, dinophyceae, diatoms, pelagophyceae and cryptophyceae did not strongly change as compared to day 0, the contribution of chlorophyceae increased to $31 \pm 4\%$.

Based on flow cytometry measurements, *Synechococcus* abundances increased during the first days of the experiment, reached maximal values on day 10 (averaged between mesocosms of $29600 \pm 3000 \text{ cells mL}^{-1}$) and then decreased until the end of the experiment (Fig. 5). Similar dynamics, although with more variability among mesocosms, were observed

for autotrophic pico-eukaryotes, with abundances one order of magnitude lower than *Synechococcus*.

Table 4 shows total chlorophyll *a* concentrations were not correlated with $p\text{CO}_2$ but showed positive trends with salinity and to a lesser extent with NH_4^+ . Chlorophyll *a*-equivalent biomass of two groups of phytoplankton were significantly correlated with $p\text{CO}_2$, dinophyceae and haptophyceae. For these two groups, $p\text{CO}_2$ appeared as the most important contributor to the variance. Note that a maximum of 66% of the variance (i.e. for chlorophyceae) observed in total chlorophyll *a* or group-specific biomasses could be explained by these stepwise linear regression analyses using the tested environmental and/or experimental variables.

3.3. Phytoplankton assemblage during the winter experiment in the Bay of Villefranche

In the Bay of Villefranche (Fig. 2), total chlorophyll *a* concentrations averaged $0.98 \pm 0.15 \mu\text{g L}^{-1}$ in the nine mesocosms along the 12-day experiment. Chlorophyll *a* remained slightly above levels in the surrounding waters for the entire experimental period, except for the last day (day 12) when concentrations increased abruptly outside the mesocosms. HPLC data are available for the acidification phase of this experiment (day -4 to day -1), data show that chlorophyll *a* concentrations increased during that period, consuming a large proportion of available nutrients, notably nitrate and nitrite, before the start of the experimental phase (see 3.1). In all mesocosms, after this initial peak, chlorophyll *a* concentrations linearly decreased until the end (GLM, $r^2 = 0.8$, $p < 0.001$).

When pigment data for all mesocosms were pooled together (Fig. 3), the plankton community in the Bay of Villefranche was dominated at the start of the experiment by cryptophyceae representing $26 \pm 1\%$ of the chlorophyll *a* content and by haptophyceae at the end ($32 \pm 5\%$). Following total chlorophyll *a* dynamics, almost all groups declined in terms of chlorophyll *a* equivalent biomass during the 12-day experiment except for cyanophyceae

whose biomass almost doubled between days 0 and 12 (Fig. 6). Groups that increased during the acidification phase and consumed available nitrate and nitrite belonged to cryptophyceae, haptophyceae, pelagophyceae and cyanophyceae. While pelagophyceae biomass remained constant throughout the experiment, cryptophyceae biomass linearly declined and haptophyceae showed maximal biomass on days 2 and 4 and then slightly declined. Several groups did not follow the initial chlorophyll *a* increase during the acidification phase (diatoms, dinophyceae, prasinophyceae and chlorophyceae).

Consistently with pigment data, flow cytometry data showed that *Synechococcus* abundances significantly increased during the acidification phase and reached values much above environmental (external) levels (Fig. 7). After few days of stagnation (days 0 to 6), abundances further increased to maximal values on day 12 (averaged between mesocosms of 42600 ± 3000 cells mL⁻¹). In contrast, it appears that *Prochlorococcus* took less advantage of this initial acidification phase with abundances on day 0 similar to external levels. After a small initial decline, abundances increased during the entire experiment with increasing variability between mesocosms. While autotrophic nano-eukaryotes abundance increased before the start of the experiment to levels much higher on day 0 than in the surrounding waters, no difference could be observed for autotrophic pico-eukaryotes on day 0 between mesocosms and the surrounding waters. Autotrophic pico-eukaryote abundance decreased until day 5, with very low variability between mesocosms, and increased until the end of the experiment with much larger discrepancy between mesocosms. In contrast, abundances of autotrophic nano-eukaryotes decreased almost linearly between day 0 and 12 with a large inter-mesocosm variability throughout the experiment.

Table 4 shows that the chlorophyll *a*-equivalent biomass of haptophyceae and diatoms were significantly correlated with *p*CO₂, although for none of these species *p*CO₂ appeared as the most important contributor to the variance. While haptophyceae appeared negatively

correlated to $p\text{CO}_2$, diatoms were positively correlated to this variable. Note that for these two groups, less than half of the variance could be explained by these multiple regressions. For most of the tested variables, salinity appeared as the most important co-variable, being either positively or negatively correlated to them. As already mentioned, salinity increased gradually during the experiment and these correlations most likely reflect a time effect on these variables.

3.4. Particulate organic matter and export

In the Bay of Calvi, particulate C and N concentrations were very low and close to the analyzer detection limit (respectively, $2.9 - 7.4 \mu\text{mol C L}^{-1}$ and $0.4 - 1.3 \mu\text{mol N L}^{-1}$). C:N ratio of the particulate organic matter remained constant in the mesocosms throughout the experiment (7.0 ± 1.0) and very close to ambient conditions (7.1 ± 0.7 ; Fig. 2). In the Bay of Villefranche, higher POC and PON concentrations were measured (respectively, $7.9 - 20.2 \mu\text{mol C L}^{-1}$ and $1.0 - 2.3 \mu\text{mol N L}^{-1}$). As in the Bay of Calvi, C:N ratio of the particulate organic matter remained constant in the mesocosms throughout the experiment (7.5 ± 0.9) and lower than in the surrounding environment (12.9 ± 4.3 ; Fig. 2). During both experiments, none of the measured variables (POC, PON or POC:PON) displayed any observable dependence on seawater acidification (Table 4).

Particulate organic carbon and nitrogen export fluxes are presented in Fig. 8. During both experiments, although more visible in the Bay of Calvi, organic matter export rates were maximal at the start of the experiments and gradually decreased until the end of the experiments. Much more variability was observed in the Bay of Villefranche with higher exported quantities of organic matter. In the Bay of Calvi, exported C:N ratios were generally homogeneous between mesocosms at the start of the experiment and much more variability was observed towards the end. Stepwise linear regressions showed no $p\text{CO}_2$ effects on these export fluxes (Table 4).

4. Discussion

The overall objective of our study was to evaluate the response of the phytoplankton community, particulate organic matter dynamics and export to $p\text{CO}_2$ changes in the NW Mediterranean Sea under contrasted physico-chemical (e.g. hydrology, nutrients and irradiance) and biological conditions (assemblage composition and abundance). Unfortunately, both summer and winter experiments were conducted under nutrient limiting conditions on plankton communities dominated by small species.

The summer experiment in the Bay of Calvi was conducted under typical stratified summer conditions characterized by very low nutrient and chlorophyll concentrations and surface irradiance levels of $\sim 1,400 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$, corresponding to maximal yearly values in that area (data not shown). With respect to nutrient availability, as fully discussed by (Louis et al., in press, this issue), observed NO_x and phosphate depleted conditions during the experiment are in the range of usually observed values in the oligotrophic Mediterranean Sea in summer. At the start of the experiment, inorganic N:P ratio was 1.7 and increased up to ~ 4 in the mesocosms on day 20. Both a low N:P ratio and low nutrient concentrations suggest that this system experienced N and P co-limitation during this period (Louis et al., in press, this issue). During this experiment, the plankton community was clearly dominated by small phytoplankton cells such as haptophyceae, cyanobacteria and chlorophyceae. Similar conditions were reported in this area at this period of the year. Using the same mesocosm setup, Giovagnetti et al. (2013) showed that the summer plankton community was dominated by pico-phytoplankton, representing $\sim 70\%$ of total biomass and composed mostly of haptophyceae and cyanobacteria. The same experiment showed that nano- and micro-phytoplankton ($\sim 30\%$ of total biomass) were composed of haptophyceae, chlorophyceae and dinoflagellates. During our experiment, phytoplankton biomass decreased during the

acidification phase in all mesocosms, independently of $p\text{CO}_2$ conditions, as shown by fluorometric data acquired using daily CTD profiles (Gazeau et al., in press, this issue). This corresponded to important organic matter sedimentation at the start of the experiment (first few days) that further stabilized at low rates until the end of the experiment. No important changes in the proportions of the different groups investigated were observed, at the exception of chlorophyceae (see above)

At the end of our experiment and considering the averaged composition in all nine mesocosms, dominance shifted towards chlorophyceae, in contrast to the external water community which remained unchanged during the course of the experiment. This relative overgrowth of chlorophyceae in all mesocosms, independently of $p\text{CO}_2$ conditions, was potentially due to wall growth. Indeed, a strong wind and wave event prevented sampling on day 19 (Gazeau et al., in press, this issue). On day 20 (our final sampling day), concentrations in chlorophyceae (but also diatoms) increased significantly (+ 30%). This observed increase was likely due to mesocosm shaking from wave actions, that released periphyton (i.e. species growing on the wall of the mesocosms) in the water column. Large species such as diatoms represented less than ~10% of phytoplankton biomass by the end of the experiment, although biomasses were usually above those in external waters (~5 vs. < 1 ng L⁻¹). Obviously, this does not appear as a surprise, as it is well known that, during the summer stratified period, diatoms are outcompeted by small species, better adapted to low nutrient and high irradiance levels, and usually do not represent more than 10% of the phytoplankton biomass in surface waters of the Ligurian Sea (Navarro et al., 2014).

The winter experiment conducted in the Bay of Villefranche was carried out in order to test for CO_2 enrichment effects on a Mediterranean plankton community not limited by nutrient availability. However, as a consequence of very favorable weather conditions during a short time window, much of the temporal dynamics observed during the experiment was

concentrated during the first few days before the end of the acidification process and nutrients were rapidly consumed in the mesocosms. At the start of the experiment, when targeted $p\text{CO}_2$ levels were reached, most of the available NO_x was already consumed and irradiance conditions dropped significantly (Gazeau et al., in press, this issue) precluding the formation of a real bloom in the bags. Addition of nutrients would have then been necessary to activate plankton dynamics in the mesocosms but this strategy was not possible as we have been forced to end the experiment after 12 days as a consequence of very bad weather conditions. In the Bay, as a consequence of intense vertical mixing, chlorophyll concentrations have been maintained at a lower level while nutrients have been continuously replenished (Louis et al., in press, this issue). In addition to probably not reflecting properly light conditions (Gazeau et al., in press, this issue), the isolation of a water mass and the reduction of mixing certainly does appear as a pitfall of this mesocosm approach or of any incubation system in these ecosystems which dynamics is strongly linked to physico-chemical conditions (e.g. mixing, irradiance). As such, results obtained during this experiment must be taken with extreme caution because of conditions inside the mesocosms not fully reflecting winter conditions in this area. Nevertheless, in winter in the Bay of Villefranche, phytoplankton biomass was much higher than in summer in the Bay of Calvi with values around $1 \mu\text{g L}^{-1}$ and a clear dominance of small species such as haptophyceae, cryptophyceae and pelagophyceae (> 65% of chlorophyll *a*-equivalent biomass). Previous observations at the entrance of the Bay of Villefranche have shown that the spring phytoplankton bloom usually takes place in February-March and is dominated by pico-nano-phytoplankton (Thyssen et al., 2014). Although not always observed, this first bloom is followed by a second one in May that is dominated by diatoms and large dinoflagellates (Bustillos-Guzmán et al., 1995; Gomez and Gorsky, 2003). During this year 2013, the highest annual chlorophyll *a* concentrations were reached later than observed in previous years (Gazeau et al., in press, this issue). Our

experiment therefore coincided with pre-bloom conditions although, again, nutrients were rapidly consumed in the mesocosms. Community composition did not drastically change during the course of the experiment inside the mesocosms but small phytoplankton species took advantage of the first few days of the experiment and are likely responsible for the strong consumption of NO_x during the acidification phase (Louis et al., in press, this issue). In contrast, diatoms and dinoflagellates did not take advantage of the closing of the bags and of favorable weather conditions during these first few days and continuously decreased in abundance until the end of the experiment. This is not a surprise since these species are known to be outcompeted by smaller phytoplankton cells when nutrient limitation is temporally relieved after the winter vertical mixing (Bustillos-Guzmán et al., 1995). Instead of these species, autotrophic prokaryotes, especially *Synechococcus*, appeared to benefit from the closing of the bags, as their abundance was 3-fold higher than the ambient levels and kept increasing throughout the experiment. While autotrophic nano-eukaryotes decreased in abundance after the initial chlorophyll *a* increase during the acidification phase, autotrophic pico-eukaryotes benefited from the recycled nutrient pool, as a consequence of increasing bacterial abundance (Celussi et al., in press, this issue), and increased in number during the second part of the experiment.

During these two experiments, while total chlorophyll *a* concentrations appeared correlated to environmental conditions (e.g. nutrients, irradiance, salinity) and/or with time, no significant correlations were found with $p\text{CO}_2$. Similarly, we could not evidence any relationship between $p\text{CO}_2$ and POC or PON concentrations as well as organic carbon and nitrogen export to the sediment traps. When phytoplankton groups were analysed separately, positive effects were found for haptophyceae and autotrophic dinoflagellates in the Bay of Calvi during the oligotrophic summer period, similarly to what was found during a large *in situ* mesocosm experiment in the Arctic (Schulz et al., 2013). In winter in the Bay of

Villefranche, while haptophyceae were negatively correlated with $p\text{CO}_2$, diatoms appeared positively impacted, although for these two groups it must be stressed that CO_2 was not the first parameter driving their variance. Such positive CO_2 -effects as observed in summer on haptophyceae and autotrophic dinoflagellates are not surprising, as these species do not possess very efficient CCMs (Reinfelder, 2011). Although cyanobacteria (including *Synechococcus*) appeared to benefit from our experimental conditions and from the very limited amount of nutrients, they were not impacted by CO_2 -enrichment. These results are consistent to what was observed by Lomas et al. (2012) in the subtropical North Atlantic but stand in contrast to the negative impact of ocean acidification on *Synechococcus* abundance observed by Paulino et al. (2008) in a North Sea Fjord under very different trophic and experimental conditions compared to our experiments (i.e. higher chlorophyll levels as well as and enrichment with N and P). As suggested by Lomas et al. (2012), the response of cyanobacteria might be indirect and controlled by other variables such as nutrients.

All in all, the short-term addition of CO_2 in our nutrient-limited systems did not induce any clear effect on community composition based on pigment analysis. It must be stressed that these analyses do not allow detecting potential modifications/replacements at the specific or at the intra-specific level as suggested by several studies in the recent years (Brading et al., 2011; Rickaby et al., 2016). Nevertheless, scanning electron microscopy analyses reported by (Oviedo et al., in press, this issue) did not highlight any changes in coccolithophores and siliceous phytoplankton community compositions, and especially any changes in species size that could have an impact on sedimentation rates (Feng et al., 2010; Tortell et al., 2008; Wu et al., 2014). During our experiments, no phylogenetic studies have been conducted at the exception of diazotrophs during the summer in the Bay of Calvi (Rees et al., in press, this issue). For this group, no significant changes could be evidenced. In the present study, the small positive or negative effects that have been highlighted on selected

groups based on pigment analyses appear to be minimal and did not lead to significant changes in terms of community metabolism (Maugendre et al., in press, this issue-b), bacterial production (Celussi et al., in press, this issue), carbon transfer (Maugendre et al., in press, this issue-a) as well as carbon and nitrogen export (this study).

These results clearly stand in contrast to recent experiments conducted in a coastal site in the Western Mediterranean Sea, using indoor tanks (Sala et al., 2016). Similar to our planned experimental protocol, two experiments were conducted under contrasting conditions: winter, at the peak of the annual phytoplankton bloom, and summer, under low nutrient conditions. Their results suggested microbial communities will be considerably more affected by ocean acidification under oligotrophic conditions than in more productive waters. It must be stressed that even during their summer low-nutrient experiment, reported nitrate concentrations were almost ten times higher than concentrations observed in summer in the Bay of Calvi and four times higher than observed in the Bay of Villefranche in the mesocosms at the end of the acidification period. Similarly, chlorophyll concentrations during our summer experiment were three times lower than observed by Sala et al. (2016) in summer in the Bay of Blanes. Recently, two other experiments conducted using large *in situ* mesocosms also suggested that communities in nutrient-limited areas may be more responsive to changing carbonate chemistry than those having access to high inorganic nutrient concentrations (Bach et al., 2016; Paul et al., 2015a). These two experiments, sharing a similar experimental protocol than in the present study, were conducted over significantly longer time scales (> 43 days). During both experiments, impacts of elevated CO₂ were visible during the last phase when plankton communities were relying on remineralized nutrients. As both our experiments did not exceed ~20 days, the build-up of remineralized nutrients did not reach concentrations high enough to significantly relieve the nutrient limitation. Nutrient limitation can be episodically relieved in summer through atmospheric

inputs (The Mermex group, 2011 and references therein) and it is now well known that pulsed atmospheric nutrient inputs enhance phototrophic, heterotrophic and diazotrophic activities (Guieu et al., 2014). It appears therefore of the utmost importance to target future experimental efforts on the response of summer plankton communities to ocean acidification in the case of a transient relieve in nutrient limitation through of a dust deposition event.

5. Conclusion

To conclude, for the first time, short-term *in situ* pelagic mesocosm experiments have been conducted in LNLC areas of the Northwestern Mediterranean Sea to assess the response of phytoplankton communities to ocean acidification. In contrast to most previous mesocosm experiments, no nutrient addition took place during the experiments conducted in summer and winter. The summer plankton community was dominated by pico-phytoplankton and cyanobacteria and was strongly limited by NO_x and phosphate availability. Although, haptophyceae and autotrophic dinoflagellates appeared to be favored by increased CO_2 availability during this short-term experiment, this benefit remained very minimal with no impact on carbon export, as a consequence of strong environmental constraints. The winter community was also dominated by small species (especially haptophyceae and cryptophyceae) that reacted soon after closing the bags and during the acidification period, possibly due to favorable weather conditions and irradiance levels. During this experiment, no signs of short term CO_2 addition dependency were detected on plankton community structure based on pigment analyses and on organic matter export. As a consequence of the very dynamic nature of environmental conditions and therefore of plankton biomass and composition in the Mediterranean Sea, more investigations are needed to carefully assess the response of plankton communities in winter when vertical mixing and weather conditions are major factors controlling plankton dynamics in this area. Future experimental protocols might

643 consider maintaining nutrient and chlorophyll levels as close as possible to ambient
644 conditions over longer time scales. Although this might be experimentally challenging, we
645 believe this is the only way to investigate these very dynamic communities. Finally, as
646 atmospheric depositions in summer have the capacity to relieve nutrient limitations and to
647 enhance plankton productions, there is a great need to perform future experiments considering
648 these pulsed nutrient additions.

Acknowledgements

This work was funded by the EC FP7 project ‘Mediterranean Sea Acidification in a changing climate’ (MedSeA; grant agreement 265103), the project ‘European Free Ocean Carbon Enrichment’ (eFOCE; BNP-Paribas Foundation), the MISTRALS-MERMEX program (Institut des Sciences de l’Univers, INSU), the Corsican local authorities and the Rhone-Mediterranean and Corsica Water Agency (<http://www.eaurmc.fr>). It is a contribution to the Surface Ocean-Lower Atmosphere Study (SOLAS) and Integrated Marine Biogeochemistry and Ecosystem Research (IMBER) projects. The STARESO marine station in Corsica is gratefully acknowledged for its assistance and boat support carried out within the framework of the STARECAPMED project funded by the Rhone-Mediterranean and Corsica Water Agency. B. Cataletto, F. Cerino, C. Comici, W. Dellisanti and C. Fabbro are acknowledged for sample preparation and analyses of particulate organic carbon and nitrogen. The staff of the Observatoire Océanologique de Villefranche is gratefully acknowledged for their assistance and boat support, colleagues of the Laboratoire d’Océanographie de Villefranche for providing laboratory space. S. Alliouane, B. Hesse, D. Luquet, D. Robin, P. Mahacek and E. Cox are acknowledged for assistance with diving operations. Thanks are due to all the MedSeA mesocosm team for help during the experiments. Finally, C. Brunet is acknowledged for his help on CHEMTAX analyses.

References

- Bach, L.T., Taucher, J., Boxhammer, T., Ludwig, A., Achterberg, E.P., Algueró-Muñiz, M., Anderson, L.G., Bellworthy, J., Büdenbender, J., Czerny, J., Ericson, Y., Esposito, M., Fischer, M., Haunost, M., Hellemann, D., Horn, H.G., Hornick, T., Meyer, J., Sswat, M., Zark, M., Riebesell, U., The Kristineberg, K.C., 2016. Influence of Ocean Acidification on a Natural Winter-to-Summer Plankton Succession: First Insights from a Long-Term Mesocosm Study Draw Attention to Periods of Low Nutrient Concentrations. *PLoS ONE* 11, e0159068.
- Badger, M.R., Andrews, T.J., Whitney, S.M., Ludwig, M., Yellowlees, D.C., Leggat, W., Price, G.D., 1998. The diversity and coevolution of Rubisco, plastids, pyrenoids, and chloroplast-based CO₂-concentrating mechanisms in algae. *Canadian Journal of Botany-Revue Canadienne De Botanique* 76, 1052-1071.
- Brading, P., Warner, M.E., Davey, P., Smith, D.J., Achterberg, E.P., Suggett, D.J., 2011. Differential effects of ocean acidification on growth and photosynthesis among phylotypes of *Symbiodinium* (Dinophyceae). *Limnol. Oceanogr.* 56, 927-938.
- Brussaard, C.P.D., Noordeloos, A.A.M., Witte, H., Collenteur, M.C.J., Schulz, K.G., Ludwig, A., Riebesell, U., 2013. Arctic microbial community dynamics influenced by elevated CO₂ levels. *Biogeosciences* 10, 719-731.
- Bustillos-Guzmán, J., Claustre, H., Marty, J.C., 1995. Specific phytoplankton signatures and their relationship to hydrographic conditions in the coastal Northwestern Mediterranean Sea. *Marine Ecology Progress Series* 124, 247-258.
- Calbet, A., Sazhin, A.F., Nejstgaard, J.C., Berger, S.A., Tait, Z.S., Olmos, L., Sousoni, D., Isari, S., Martinez, R.A., Bouquet, J.M., Thompson, E.M., Bamstedt, U., Jakobsen, H.H., 2014. Future climate scenarios for a coastal productive planktonic food web

resulting in microplankton phenology changes and decreased trophic transfer efficiency.

Plos One 9.

- Celussi, M., Malfatti, F., Franzo, A., Gazeau, F., Giannakourou, A., Pitta, P., Tsiola, A., Del Negro, P., in press, this issue. Ocean acidification effect on prokaryotic metabolism in two different locations in the Mediterranean Sea. *Estuarine, Coastal and Shelf Science*.
- Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R.B., Piao, S., Thornton, P., 2013. Carbon and Other Biogeochemical Cycles, in: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (Eds.), *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Claustre, H., Kerherve, P., Marty, J.C., Prieur, L., Videau, C., Hecq, J.H., 1994. Phytoplankton dynamics associated with a geostrophic front - ecological and biogeochemical implications. *Journal of Marine Research* 52, 711-742.
- Czerny, J., Schulz, K.G., Boxhammer, T., Bellerby, R.G.J., Büdenbender, J., Engel, A., Krug, S.A., Ludwig, A., Nachtigall, K., Nondal, G., Niehoff, B., Silyakova, A., Riebesell, U., 2013. Implications of elevated CO₂ on pelagic carbon fluxes in an Arctic mesocosm study – an elemental mass balance approach. *Biogeosciences* 10, 3109-3125.
- D'Ortenzio, F., d'Alcala, M.R., 2009. On the trophic regimes of the Mediterranean Sea: a satellite analysis. *Biogeosciences* 6, 139-148.
- Domingues, R.B., Guerra, C.C., Barbosa, A.B., Brotas, V., Galvao, H.M., 2014. Effects of ultraviolet radiation and CO₂ increase on winter phytoplankton assemblages in a temperate coastal lagoon. *Journal of Plankton Research* 36, 672-684.

- Dutkiewicz, S., Morris, J.J., Follows, M.J., Scott, J., Levitan, O., Dyhrman, S.T., Berman-Frank, I., 2015. Impact of ocean acidification on the structure of future phytoplankton communities. *Nature Climate change* 5, 1002-1006.
- Eggers, S.L., Lewandowska, A.M., Ramos, J.B.E., Blanco-Ameijeiras, S., Gallo, F., Matthiessen, B., 2014. Community composition has greater impact on the functioning of marine phytoplankton communities than ocean acidification. *Global Change Biology* 20, 713-723.
- Endo, H., Sugie, K., Yoshimura, T., Suzuki, K., 2016. Response of spring diatoms to CO₂ availability in the Western North Pacific as determined by next-generation sequencing. *PLoS ONE* 11, e0154291-e0154291.
- Engel, A., Schulz, K.G., Riebesell, U., Bellerby, R., Delille, B., Schartau, M., 2008. Effects of CO₂ on particle size distribution and phytoplankton abundance during a mesocosm bloom experiment (PeECE II). *Biogeosciences* 5, 509-521.
- Engel, A., Zondervan, I., Aerts, K., Beaufort, L., Benthien, A., Chou, L., Delille, B., Gattuso, J.P., Harlay, J., Heemann, C., Hoffmann, L., Jacquet, S., Nejstgaard, J., Pizay, M.D., Rochelle-Newall, E., Schneider, U., Terbrueggen, A., Riebesell, U., 2005. Testing the direct effect of CO₂ concentration on a bloom of the coccolithophorid *Emiliania huxleyi* in mesocosm experiments. *Limnol. Oceanogr.* 50, 493-507.
- Feng, Y., Hare, C.E., Rose, J.M., Handy, S.M., DiTullio, G.R., Lee, P.A., Smith, W.O., Peloquin, J., Tozzi, S., Sun, J., Zhang, Y., Dunbar, R.B., Long, M.C., Sohst, B., Lohan, M., Hutchins, D.A., 2010. Interactive effects of iron, irradiance and CO₂ on Ross Sea phytoplankton. *Deep-Sea Research Part I-Oceanographic Research Papers* 57, 368-383.
- Feng, Y.Y., Hare, C.E., Leblanc, K., Rose, J.M., Zhang, Y.H., DiTullio, G.R., Lee, P.A., Wilhelm, S.W., Rowe, J.M., Sun, J., Nemcek, N., Gueguen, C., Passow, U., Benner, I., Brown, C., Hutchins, D.A., 2009. Effects of increased pCO₂ and temperature on the

- North Atlantic spring bloom. I. The phytoplankton community and biogeochemical response. *Marine Ecology Progress Series* 388, 13-25.
- Field, C.B., Behrenfeld, M.J., Randerson, J.T., Falkowski, P., 1998. Primary production of the biosphere: integrating terrestrial and oceanic components. *Science* 281, 237-240.
- Finkel, Z.V., Katz, M.E., Wright, J.D., Schofield, O.M.E., Falkowski, P.G., 2005. Climatically driven macroevolutionary patterns in the size of marine diatoms over the Cenozoic. *Proceedings of the National Academy of Sciences of the United States of America* 102, 8927-8932.
- Flynn, K.J., Blackford, J.C., Baird, M.E., Raven, J.A., Clark, D.R., Beardall, J., Brownlee, C., Fabian, H., Wheeler, G.L., 2012. Changes in pH at the exterior surface of plankton with ocean acidification. *Nature Climate Change*, 510-513.
- Gattuso, J.P., Magnan, A., Bille, R., Cheung, W.W.L., Howes, E.L., Joos, F., Allemand, D., Bopp, L., Cooley, S.R., Eakin, C.M., Hoegh-Guldberg, O., Kelly, R.P., Poertner, H.O., Rogers, A.D., Baxter, J.M., Laffoley, D., Osborn, D., Rankovic, A., Rochette, J., Sumaila, U.R., Treyer, S., Turley, C., 2015. Contrasting futures for ocean and society from different anthropogenic CO₂ emissions scenarios. *Science* 349, 45-+.
- Gazeau, F., Sallon, A., Maugendre, L., Louis, J., Dellisanti, W., Gaubert, M., Lejeune, P., Gobert, S., Borges, A.V., Harlay, J., Champenois, W., Alliouane, S., Taillandier, V., Louis, F., Obolensky, G., Grisoni, J.M., Guieu, C., in press, this issue. First mesocosm experiments to study the impacts of ocean acidification on plankton communities in the NW Mediterranean Sea (MedSeA project). *Estuarine, Coastal and Shelf Science*.
- Giordano, M., Beardall, J., Raven, J.A., 2005. CO₂ concentrating mechanisms in algae: Mechanisms, environmental modulation, and evolution, *Annual Review of Plant Biology*, pp. 99-131.

- Giovagnetti, V., Brunet, C., Conversano, F., Tramontano, F., Obernosterer, I., Ridame, C., Guieu, C., 2013. Assessing the role of dust deposition on phytoplankton ecophysiology and succession in a low-nutrient low-chlorophyll ecosystem: a mesocosm experiment in the Mediterranean Sea. *Biogeosciences* 10, 2973-2991.
- Gomez, F., Gorsky, G., 2003. Annual microplankton cycles in Villefranche Bay, Ligurian Sea, NW Mediterranean. *Journal of Plankton Research* 25, 323-339.
- Guieu, C., Aumont, O., Paytan, A., Bopp, L., Law, C.S., Mahowald, N., Achterberg, E.P., Maranon, E., Salihoglu, B., Crise, A., Wagener, T., Herut, B., Desboeufs, K., Kanakidou, M., Olgun, N., Peters, F., Pulido-Villena, E., Tovar-Sanchez, A., Voelker, C., 2014. The significance of the episodic nature of atmospheric deposition to Low Nutrient Low Chlorophyll regions. *Global Biogeochemical Cycles* 28, 1179-1198.
- Hennon, G.M.M., Ashworth, J., Groussman, R.D., Berthiaume, C., Morales, R.L., Baliga, N.S., Orellana, M.V., Armbrust, E.V., 2015. Diatom acclimation to elevated CO₂ via cAMP signalling and coordinated gene expression. *Nature Clim. Change* 5, 761-765.
- Hopkins, F.E., Turner, S.M., Nightingale, P.D., Steinke, M., Bakker, D., Liss, P.S., 2010. Ocean acidification and marine trace gas emissions. *Proceedings of the National Academy of Sciences of the United States of America* 107, 760-765.
- Irwin, A.J., Oliver, M.J., 2009. Are ocean deserts getting larger? *Geophys. Res. Lett.* 36.
- Kim, J.H., Kim, K.Y., Kang, E.J., Lee, K., Kim, J.M., Park, K.T., Shin, K., Hyun, B., Jeong, H.J., 2013. Enhancement of photosynthetic carbon assimilation efficiency by phytoplankton in the future coastal ocean. *Biogeosciences* 10, 7525-7535.
- Kim, J.M., Lee, K., Shin, K., Kang, J.H., Lee, H.W., Kim, M., Jang, P.G., Jang, M.C., 2006. The effect of seawater CO₂ concentration on growth of a natural phytoplankton assemblage in a controlled mesocosm experiment. *Limnol. Oceanogr.* 51, 1629-1636.

- 789 Kim, J.M., Lee, K., Shin, K., Yang, E.J., Engel, A., Karl, D.M., Kim, H.C., 2011. Shifts in
 790 biogenic carbon flow from particulate to dissolved forms under high carbon dioxide and
 791 warm ocean conditions. *Geophys. Res. Lett.* 38.
- 792 Kim, J.M., Lee, K., Yang, E.J., Shin, K., Noh, J.H., Park, K.T., Hyun, B., Jeong, H.J., Kim,
 793 J.H., Kim, K.Y., Kim, M., Kim, H.C., Jang, P.G., Jang, M.C., 2010. Enhanced
 794 production of oceanic dimethylsulfide resulting from CO₂-induced grazing activity in a
 795 high CO₂ world. *Environmental Science & Technology* 44, 8140-8143.
- 796 Kroeker, K.J., Kordas, R.L., Crim, R., Hendriks, I.E., Ramajo, L., Singh, G.S., Duarte, C.M.,
 797 Gattuso, J.P., 2013. Impacts of ocean acidification on marine organisms: quantifying
 798 sensitivities and interaction with warming. *Global Change Biology* 19, 1884-1896.
- 799 Lavigne, H., Epitalon, J.M., Gattuso, J.-P., 2014. seacarb: seawater carbonate chemistry with
 800 R. [ran.r-project.org/package=seacarb](http://cran.r-project.org/package=seacarb).
- 801 Lee, C., Wakeham, S.G., Hedges, J.I., 1988. The measurement of oceanic particle flux - Are
 802 "swimmers" a problem? *Oceanography* 1, 34-36.
- 803 Lomas, M.W., Hopkinson, B.M., Losh, J.L., Ryan, D.E., Shi, D.L., Xu, Y., Morel, F.M.M.,
 804 2012. Effect of ocean acidification on cyanobacteria in the subtropical North Atlantic.
 805 *Aquatic Microbial Ecology* 66, 211-222.
- 806 Louis, J., Guieu, C., Gazeau, F., in press, this issue. Nutrient dynamics under different ocean
 807 acidification scenarios in a low nutrient low chlorophyll system: the Northwestern
 808 Mediterranean Sea. *Estuarine, Coastal and Shelf Science*.
- 809 Mangan, N.M., Flamholz, A., Hood, R.D., Milo, R., Savage, D.F., 2016. pH determines the
 810 energetic efficiency of the cyanobacterial CO₂ concentrating mechanism. *Proceedings*
 811 *of the National Academy of Sciences* 113, E5354-E5362.

- 812 Marie, D., Brussaard, C.P.D., Thyrhaug, R., Bratbak, G., Vaulot, D., 1999. Enumeration of
 813 marine viruses in culture and natural samples by flow cytometry. *Applied and*
 814 *Environmental Microbiology* 65, 45-52.
- 815 Maugendre, L., Gattuso, J.-P., de Kluijver, A., Soetaert, K., van Oevelen, D., Middelburg,
 816 J.J., Gazeau, F., in press, this issue-a. Carbon-13 labelling studies show no effect of
 817 ocean acidification on Mediterranean plankton communities. *Estuarine, Coastal and*
 818 *Shelf Science*.
- 819 Maugendre, L., Gattuso, J.P., Poulton, A.J., Dellisanti, W., Gaubert, M., Guieu, C., Gazeau,
 820 F., in press, this issue-b. No detectable effect of ocean acidification on plankton
 821 metabolism in the NW oligotrophic Mediterranean Sea: Results from two mesocosm
 822 studies. *Estuarine, Coastal and Shelf Science*.
- 823 Meakin, N.G., Wyman, M., 2011. Rapid shifts in picoeukaryote community structure in
 824 response to ocean acidification. *Isme Journal* 5, 1397-1405.
- 825 Navarro, G., Alvain, S., Vantrepotte, V., Huertas, I.E., 2014. Identification of dominant
 826 phytoplankton functional types in the Mediterranean Sea based on a regionalized
 827 remote sensing approach. *Remote Sensing of Environment* 152, 557-575.
- 828 Newbold, L.K., Oliver, A.E., Booth, T., Tiwari, B., DeSantis, T., Maguire, M., Andersen, G.,
 829 van der Gast, C.J., Whiteley, A.S., 2012. The response of marine picoplankton to ocean
 830 acidification. *Environmental Microbiology* 14, 2293-2307.
- 831 Not, F., Zapata, M., Pazos, Y., Campana, E., Doval, M., Rodriguez, F., 2007. Size-
 832 fractionated phytoplankton diversity in the NW Iberian coast: a combination of
 833 microscopic, pigment and molecular analyses. *Aquatic Microbial Ecology* 49, 255-265.
- 834 Oviedo, A.M., Ziveri, P., Gazeau, F., in press, this issue. Coccolithophore dynamics in
 835 response to increasing $p\text{CO}_2$ in Mediterranean oligotrophic waters. *Estuarine, Coastal*
 836 *and Shelf Science*.

- 837 Paul, A.J., Bach, L.T., Schulz, K.G., Boxhammer, T., Czerny, J., Achterberg, E.P.,
 838 Hellemann, D., Trense, Y., Nausch, M., Sswat, M., Riebesell, U., 2015a. Effect of
 839 elevated CO₂ on organic matter pools and fluxes in a summer Baltic Sea plankton
 840 community. *Biogeosciences* 12, 6181-6203.
- 841 Paul, C., Matthiessen, B., Sommer, U., 2015b. Warming, but not enhanced CO₂
 842 concentration, quantitatively and qualitatively affects phytoplankton biomass. *Marine*
 843 *Ecology Progress Series* 528, 39-51.
- 844 Paulino, A.I., Egge, J.K., Larsen, A., 2008. Effects of increased atmospheric CO₂ on small
 845 and intermediate sized osmotrophs during a nutrient induced phytoplankton bloom.
 846 *Biogeosciences* 5, 739-748.
- 847 Polovina, J.J., Howell, E.A., Abecassis, M., 2008. Ocean's least productive waters are
 848 expanding. *Geophys. Res. Lett.* 35.
- 849 Ras, J., Claustre, H., Uitz, J., 2008. Spatial variability of phytoplankton pigment distributions
 850 in the Subtropical South Pacific Ocean: comparison between *in situ* and predicted data.
 851 *Biogeosciences* 5, 353-369.
- 852 Raven, J.A., Beardall, J., 2014. CO₂ concentrating mechanisms and environmental change.
 853 *Aquatic Botany* 118, 24-37.
- 854 Raven, J.A., Beardall, J., Giordano, M., 2014. Energy costs of carbon dioxide concentrating
 855 mechanisms in aquatic organisms. *Photosynthesis Research* 121, 111-124.
- 856 Rees, A.P., Turk-Kubo, K., Al-Moosawi, L., Alliouane, S., Gazeau, F., Hogan, M.E., Zehr,
 857 J.P., in press, this issue. Ocean acidification impacts on Mediterranean nitrogen fixation
 858 at pCO₂ levels indicated for the end of the 21st Century. *Estuarine, Coastal and Shelf*
 859 *Science*.

- 860 Reinfelder, J.R., 2011. Carbon concentrating mechanisms in Eukaryotic marine
861 phytoplankton, in: Carlson, C.A., Giovannoni, S.J. (Eds.), Annual Review of Marine
862 Science, Vol 3, pp. 291-315.
- 863 Reul, A., Munoz, M., Bautista, B., Neale, P.J., Sobrino, C., Mercado, J.M., Segovia, M.,
864 Salles, S., Kulk, G., Leon, P., van de Poll, W.H., Perez, E., Buma, A., Blanco, J.M.,
865 2014. Effect of CO₂, nutrients and light on coastal plankton. III. Trophic cascade, size
866 structure and composition. Aquatic Biology 22, 59-76.
- 867 Richier, S., Achterberg, E.P., Dumousseaud, C., Poulton, A.J., Suggett, D.J., Tyrrell, T.,
868 Zubkov, M.V., Moore, C.M., 2014. Phytoplankton responses and associated carbon
869 cycling during shipboard carbonate chemistry manipulation experiments conducted
870 around Northwest European shelf seas. Biogeosciences 11, 4733-4752.
- 871 Rickaby, R.E.M., Hermoso, M., Lee, R.B.Y., Rae, B.D., Heures, A.M.C., Balestreri, C.,
872 Chakravarti, L., Schroeder, D.C., Brownlee, C., 2016. Environmental carbonate
873 chemistry selects for phenotype of recently isolated strains of *Emiliana huxleyi*. Deep
874 Sea Research Part II: Topical Studies in Oceanography 127, 28-40.
- 875 Riebesell, U., Bellerby, R.G.J., Grossart, H.P., Thingstad, F., 2008. Mesocosm CO₂
876 perturbation studies: from organism to community level. Biogeosciences 5, 1157-1164.
- 877 Riebesell, U., Czerny, J., von Brockel, K., Boxhammer, T., Budenbender, J., Deckelnick, M.,
878 Fischer, M., Hoffmann, D., Krug, S.A., Lentz, U., Ludwig, A., Muehe, R., Schulz,
879 K.G., 2013a. Technical Note: A mobile sea-going mesocosm system - new
880 opportunities for ocean change research. Biogeosciences 10, 1835-1847.
- 881 Riebesell, U., Gattuso, J.P., Thingstad, T.F., Middelburg, J.J., 2013b. Preface "Arctic ocean
882 acidification: pelagic ecosystem and biogeochemical responses during a mesocosm
883 study". Biogeosciences 10, 5619-5626.

- 884 Riebesell, U., Schulz, K.G., Bellerby, R.G.J., Botros, M., Fritsche, P., Meyerhofer, M., Neill,
 885 C., Nondal, G., Oschlies, A., Wohlers, J., Zollner, E., 2007. Enhanced biological carbon
 886 consumption in a high CO₂ ocean. *Nature* 450, 545-U510.
- 887 Riebesell, U., Tortell, P.D., 2011. Effects of ocean acidification on pelagic organisms and
 888 ecosystems, in: Gattuso, J.-P., Hansson, L. (Eds.), *Ocean acidification*. Oxford
 889 University Press, Oxford, pp. 99-121.
- 890 Rodriguez, F., Garrido, J.L., Crespo, B.G., Arbones, B., Figueiras, F.G., 2006. Size-
 891 fractionated phytoplankton pigment groups in the NW Iberian upwelling system: impact
 892 of the Iberian Poleward Current. *Marine Ecology Progress Series* 323, 59-73.
- 893 Sala, M.M., Aparicio, F.L., Balagué, V., Boras, J.A., Borrull, E., Cardelús, C., Cros, L.,
 894 Gomes, A., López-Sanz, A., Malits, A., Martínez, R.A., Mestre, M., Movilla, J.,
 895 Sarmiento, H., Vázquez-Domínguez, E., Vaqué, D., Pinhassi, J., Calbet, A., Calvo, E.,
 896 Gasol, J.M., Pelejero, C., Marrasé, C., 2016. Contrasting effects of ocean acidification
 897 on the microbial food web under different trophic conditions. *ICES Journal of Marine*
 898 *Science* 73, 670-679.
- 899 Sarmiento, J.L., Gruber, N., 2006. *Ocean biogeochemical dynamics*. Princeton University
 900 Press, Princeton, Woodstock.
- 901 Schulz, K.G., Bellerby, R.G.J., Brussaard, C.P.D., Budenbender, J., Czerny, J., Engel, A.,
 902 Fischer, M., Koch-Klavsen, S., Krug, S.A., Lischka, S., Ludwig, A., Meyerhofer, M.,
 903 Nondal, G., Silyakova, A., Stühr, A., Riebesell, U., 2013. Temporal biomass dynamics
 904 of an Arctic plankton bloom in response to increasing levels of atmospheric carbon
 905 dioxide. *Biogeosciences* 10, 161-180.
- 906 Schulz, K.G., Riebesell, U., Bellerby, R.G.J., Biswas, H., Meyerhofer, M., Muller, M.N.,
 907 Egge, J.K., Nejstgaard, J.C., Neill, C., Wohlers, J., Zollner, E., 2008. Build-up and
 908 decline of organic matter during PeECE III. *Biogeosciences* 5, 707-718.

- Siokou-Frangou, I., Christaki, U., Mazzocchi, M.G., Montresor, M., Ribera d'Alcalá, M.,
 Vaqué, D., Zingone, A., 2010. Plankton in the open Mediterranean Sea: a review.
 Biogeosciences 7, 1543-1586.
- Sommer, U., Paul, C., Moustaka-Gouni, M., 2015. Warming and ocean acidification effects
 on phytoplankton - From species shifts to size shifts within species in a mesocosm
 experiment. PLoS ONE 10, e0125239.
- Tarling, G.A., Peck, V., Ward, P., Ensor, N.S., Achterberg, E., Tynan, E., Poulton, A.J.,
 Mitchell, E., Zubkov, M.V., 2016. Effects of acute ocean acidification on spatially-
 diverse polar pelagic foodwebs: insights from on-deck microcosms. Deep Sea Research
 Part II: Topical Studies in Oceanography 127, 75-92.
- The Mermex group, 2011. Marine ecosystems' responses to climatic and anthropogenic
 forcings in the Mediterranean. Progress in Oceanography 91, 97-166.
- Thyssen, M., Gregori, G.J., Grisoni, J.-M., Pedrotti, M., Mousseau, L., Artigas, L.F., Marro,
 S., Garcia, N., Passafiume, O., Denis, M.J., 2014. Onset of the spring bloom in the
 Northwestern Mediterranean Sea: influence of environmental pulse events on the *in situ*
 hourly-scale dynamics of the phytoplankton community structure. Frontiers in
 Microbiology 5.
- Tortell, P.D., 2000. Evolutionary and ecological perspectives on carbon acquisition in
 phytoplankton. Limnol. Oceanogr. 45, 744-750.
- Tortell, P.D., DiTullio, G.R., Sigman, D.M., Morel, F.M.M., 2002. CO₂ effects on taxonomic
 composition and nutrient utilization in an Equatorial Pacific phytoplankton assemblage.
 Marine Ecology Progress Series 236, 37-43.
- Tortell, P.D., Payne, C.D., Li, Y.Y., Trimborn, S., Rost, B., Smith, W.O., Riesselman, C.,
 Dunbar, R.B., Sedwick, P., DiTullio, G.R., 2008. CO₂ sensitivity of Southern Ocean
 phytoplankton. Geophys. Res. Lett. 35, 5.

- Troussellier, M., Courties, C., Zettelmaier, S., 1995. Flow cytometric analysis of coastal lagoon bacterioplankton and picophytoplankton - fixation and storage effects. *Estuarine Coastal and Shelf Science* 40, 621-633.
- Vaulot, D., Courties, C., Partensky, F., 1989. A simple method to preserve oceanic phytoplankton for flow cytometric analyses. *Cytometry* 10, 629-635.
- Verspagen, J.M.H., Van de Waal, D.B., Finke, J.F., Visser, P.M., Huisman, J., 2014. Contrasting effects of rising CO₂ on primary production and ecological stoichiometry at different nutrient levels. *Ecology Letters* 17, 951-960.
- Wu, Y., Campbell, D.A., Irwin, A.J., Suggett, D.J., Finkel, Z.V., 2014. Ocean acidification enhances the growth rate of larger diatoms. *Limnol. Oceanogr.* 59, 1027-1034.
- Young, J.N., Heures, A.M.C., Sharwood, R.E., Rickaby, R.E.M., Morel, F.M.M., Whitney, S.M., 2016. Large variation in the Rubisco kinetics of diatoms reveals diversity among their carbon-concentrating mechanisms. *Journal of Experimental Botany*.
- Young, J.N., Morel, F.M.M., 2015. Biological oceanography: The CO₂ switch in diatoms. *Nature Climate change* 5, 722-723.

Figure legends

Figure 1. Carbonate chemistry conditions in the nine mesocosms and in the external environment (OUT) during the experiment in the Bay of Calvi in summer 2012 (left panels) and in the Bay of Villefranche in winter 2013 (right panels). Partial pressure of CO_2 ($p\text{CO}_2$, upper panels) and pH on the total scale (pH_T , lower panels) were calculated using seacarb, based on dissolved inorganic carbon concentrations (C_T , not shown) and total alkalinity (A_T , not shown), measured daily from depth-integrated (0-10 m) samples. Vertical dotted lines show the start of the respective experiments (day 0).

Figure 2. Depth-integrated (0-10 m) chlorophyll *a* concentrations as measured by high performance liquid chromatography (HPLC; upper panels) and particulate organic carbon (POC) to particulate organic nitrogen (PON) ratio (POC:PON; lower panels) in the nine mesocosms and in the external environment during the experiment in the Bay of Calvi in summer 2012 (left panels) and in the Bay of Villefranche in winter 2013 (right panels). Vertical dotted lines on the right panels show the start of the experiment (day 0). No chlorophyll *a* and POC data are available before day 0 in the Bay of Calvi.

Figure 3. Averaged contribution (%) between all nine mesocosms of the main phytoplanktonic taxonomic groups to total chlorophyll *a* concentrations at the start (day 0) and at the end (day 20 or 12) of the experiments in the Bay of Calvi in summer 2012 (left panel) and in the Bay of Villefranche in winter 2013 (right panel).

Figure 4. Temporal evolution of chlorophyll *a* (chl *a*) -equivalent concentrations of eight taxonomic groups of phytoplankton during the experiment in the Bay of Calvi in summer 2012, in the nine mesocosms and in the external environment (OUT). Prasino: prasinophyceae, Dino: dinophyceae, Crypto: cryptophyceae, Hapto: haptophyceae, Pelago: pelagophyceae, Chloro: chlorophyceae, Cyano: cyanophyceae.

Figure 5. Temporal evolution of *Synechococcus* and pico-eukaryotes abundances as measured by flow cytometry, during the experiment in the Bay of Calvi in summer 2012, in the nine mesocosms.

Figure 6. Temporal evolution of chlorophyll *a* (chl *a*) -equivalent concentrations of 8 taxonomic groups of phytoplankton determined from high performance liquid chromatography (HPLC) measurements using modified CHEMTAX, during the experiment in the Bay of Villefranche in winter 2013, in the 9 mesocosms and in the external environment (OUT). Prasino: prasinophyceae, Dino: dinophyceae, Crypto: cryptophyceae, Hapto: haptophyceae, Pelago: pelagophyceae, Chloro: chlorophyceae, Cyano: cyanophyceae.

Figure 7. Temporal evolution of *Synechococcus*, *Prochlorococcus*, pico-eukaryotes and nano-eukaryotes abundances as measured by flow cytometry, during the experiment in the Bay of Villefranche in winter 2013, in the nine mesocosms and in the external environment (OUT).

Figure 8. Upper panel: temporal evolution of particulate organic carbon (POC) fluxes to the sediment traps during the experiment in the Bay of Calvi in summer 2012 (left panel) and in the Bay of Villefranche in winter 2013 (right panel). Lower panel: particulate organic carbon (POC) to particulate organic nitrogen (PON) ratio in the sediment traps during the experiment in the Bay of Calvi in summer 2012. No PON data available during the experiment in the Bay of Villefranche in winter 2013. Vertical dotted lines show the start of the respective experiment (day 0).

Table 1. Summary of past mesocosm (volume between 1 and 1000 m³) ocean acidification experiments results on phytoplankton communities. ⇔, ↑ and ↓ refer to neutral, positive and negative effects on chlorophyll *a* concentrations (Chl *a*) as well as concentrations of diatoms (diat), dinophyceae (Dino), nano-eukaryotes (Nano), pico-eukaryotes (Pico) and cyanophyceae (Cyano). Impacts on carbon export are also reported when available. “√” indicates that mesocosms were enriched with nutrients (Nut: nitrate, phosphate and sometimes silicate). “-” indicates that no information is available. Cryp and Chlo refer to cryptophyceae and chlorophyceae respectively.

Reference	Study location	Season	Nut	Major group	Chl	Diat	Dino	Nano	Pico	Cyano	Export	Notes
Indoor												
Sommer et al. (2015) Paul et al. (2015b)	Kiel Bight	Fall		Diat/Dino	⇔	↓	⇔	⇔	⇔	⇔	-	
Outdoor - Floating raft												
Engel et al. (2005)	Norwegian Fjord	Spring	√	Pico/Cyano	⇔	-	-	⇔	⇔	⇔	-	Decrease of coccolithophore calcification
Kim et al. (2006)	Korean coast	Fall		Micro/Diat	-	⇔	-	-	-	-	-	
Engel et al. (2008)	Norwegian Fjord	Spring	√	Pico	⇔	⇔	-	↓	↑	-	-	
Paulino et al. (2008) Schulz et al. (2008)	Norwegian Fjord	Spring	√	-	⇔	⇔	⇔	⇔	↑	↓	-	
Hopkins et al. (2010) Meakin and Wyman (2011) Newbold et al. (2012)	Norwegian Fjord	Spring	√	Micro	↓	-	-	↑	↑	↑	-	Increase in large pico-eukaryotes
Kim et al. (2010) Kim et al. (2011) Kim et al. (2013)	Korean coast	Fall	√	Diat/Dino	⇔	↑	⇔	↑	⇔	-	-	Shift from weakly to heavily silified diatoms
Calbet et al. (2014)	Norwegian Fjord	Spring	√	Nano	⇔	↑	↓	↑	↑	-	-	
Outdoor - Free floating												

Schulz et al. (2013)	Arctic Fjord	Spring	√	Nano	↔	↔	↔	↔	↔	↔	↔	Phase 1 before nutrient enrichment
Brussaard et al. (2013)				-	↑	↔	↑	↑	↑	↔	↔	Phase 2
Czerny et al. (2013)				-	↓	↔	↑	↔	↑	↔	↓	Phase 3
Paul et al. (2015a)	Baltic Sea	Spring		Cryp	↔	↔	-	↔	↑	↔	↔	Phase 1
				Chlo	↑	↔	-	↔	↑	↔	↔	Phase 2
				Chlo	↑	↓	-	↓	↑	↔	↔	Phase 3
Bach et al. (2016)	Swedish Fjord	Winter		-	-	↔	↔	↔	↔	↔	-	
		Spring		Diatoms	-	↔	↔	↔	↔	↔	-	First chlorophyll build-up
		Spring		Diatoms	-	↔	↔	↔	↑	↔	-	Second chlorophyll build-up on remineralized nutrients
		Spring		-	-	↔	↔	↔	↔	↔	-	

Table 2. Environmental and experimental conditions in the nine mesocosms and in the external environment (OUT) during the experiment in the Bay of Calvi in summer 2012. Levels of temperature (T in °C), salinity (S), partial pressure of CO₂ (*p*CO₂ in µatm), nitrate + nitrite (NO_x in nmol L⁻¹) and phosphate (PO₄³⁻ in nmol L⁻¹), ammonium (NH₄⁺ in nmol L⁻¹) and silicate (Si in µmol L⁻¹) at the end of the acidification period (day 0) and at the end of the experiment (day 20) are reported. NO₃⁻ and PO₄³⁻ data from Louis et al. (in press, this issue). NH₄⁺ and Si data are from Gazeau et al. (in press, this issue). NA: not available.

	Day 0							Day 20						
	T	S	<i>p</i> CO ₂	NO _x	PO ₄ ³⁻	NH ₄ ⁺	Si	T	S	<i>p</i> CO ₂	NO _x	PO ₄ ³⁻	NH ₄ ⁺	Si
OUT	22.2	38.0	442	50	35	150	1.9	24.3	38.2	489	NA	NA	660	1.8
C1	22.1	38.0	455	59	NA	450	NA	NA	NA	456	77	4	190	1.1
C2	22.1	38.0	447	53	25	550	NA	24.2	38.2	472	61	6	230	1.4
C3	22.1	38.0	444	69	21	210	NA	24.2	38.1	473	59	7	210	1.3
P1	22.2	38.0	583	NA	NA	330	NA	24.3	38.2	544	45	6	130	1.3
P2	22.1	38.0	698	37	23	400	NA	24.3	38.2	609	41	4	290	1.4
P3	22.1	38.0	753	36	20	225	1.7	24.2	38.2	655	42	10	100	1.3
P4	22.1	38.0	875	30	19	770	1.7	24.3	38.2	764	75	8	230	1.2
P5	22.1	38.0	1134	37	31	260	1.7	24.3	38.1	754	76	9	350	1.3
P6	22.1	38.0	1279	57	NA	130	1.7	24.2	38.2	738	61	8	180	1.4

Table 3. Environmental and experimental conditions in the nine mesocosms and in the external environment (OUT) during the experiment in the Bay of Villefranche in winter 2013. Levels of temperature (T in °C), salinity (S), partial pressure of CO₂ (*p*CO₂ in µatm), nitrate (NO₃⁻ in nmol L⁻¹), phosphate (PO₄³⁻ in nmol L⁻¹), ammonium (NH₄⁺ in nmol L⁻¹) and silicate (Si in µmol L⁻¹) at the end of the acidification period (day 0) and at the end of the experiment (day 12) are reported. *No data are available for day 12 therefore levels on day 11 are reported. NO₃⁻ and PO₄³⁻ data from Louis et al. (in press, this issue). NH₄⁺ and Si data are from Gazeau et al. (in press, this issue). NA: not available.

	Day 0							Day 12						
	T	S	<i>p</i> CO ₂	NO ₃ ⁻	PO ₄ ³⁻	NH ₄ ⁺	Si	T	S	<i>p</i> CO ₂	NO ₃ ⁻	PO ₄ ³⁻	NH ₄ ⁺	Si
OUT	13.2	38.1	354	1166	10	62	1.3	13.2	38.2	391	1307	12	40	1.2
C1	13.2	38.1	378	167	10	79	NA	13.2	38.2	388	394	9	49	1.0
C2	13.2	38.1	347	118	12	57	1.1	13.2	38.2	354	194	11	31	1.1
C3	13.2	38.1	350	110	9	81	1.2	NA	NA	376	127	10	26	1.2
P1	13.2	38.1	494	135	10	73	NA	13.2	38.2	429	491	10	68	1.1
P2	13.2	38.1	622	133	9	64	1.2	13.2	38.2	413*	NA	10	NA	NA
P3	13.2	38.1	691	NA	9	64	1.2	NA	NA	451	236	NA	26	1.2
P4	13.2	38.1	744	72	12	80	NA	13.2	38.2	436	491	9	36	0.9
P5	13.2	38.1	932	134	15	60	1.2	13.2	38.2	497	226	10	30	1.1
P6	13.2	38.1	1250	156	8	60	1.1	NA	NA	579*	NA	NA	NA	NA

Table 4. Stepwise multiple regression analysis between environmental/experimental variables (T: temperature, S: salinity, $p\text{CO}_2$: partial pressure of CO_2 , NO_3^- : nitrate concentrations, PO_4^{3-} : phosphate concentrations, NH_4^+ : ammonium concentrations, Si: silicate concentrations (only for diatoms), I: daily integrated photon doses and w: daily averaged wind speeds) and total chlorophyll *a* (chl *a*) concentrations or chlorophyll *a*-equivalent concentrations of eight taxonomic groups of phytoplankton determined from high performance liquid chromatography (HPLC) measurements using modified CHEMTAX or *Synechococcus*, *Prochlorococcus*, pico-eukaryotes and nano-eukaryotes abundances as measured by flow cytometry, during the experiment in the Bay of Calvi in summer 2012 and in the Bay of Villefranche in winter 2013. Note that *Prochlorococcus* and nano-eukaryotes abundances are not available for the Bay of Calvi and that PON fluxes are not available for the Bay of Villefranche (denoted as NA). Bold text denotes significant correlations ($p < 0.01$) between the considered variable and $p\text{CO}_2$ and the sign (+ or -) refers to the sign of the relationship between the considered variable and the environmental/experimental parameters considered. NS: not significant.

Bay of Calvi	F	Adj. r^2	df	Overall p	Variable	Sign	p	Bay of Villefranche	F	Adj. r^2	df	Overall p	Variable	Sign	p
Particulate matter								Particulate matter							
POC	NS							POC	34.5	0.57	97	< 0.001	NO_3^-	-	< 0.001
													S	-	< 0.001
PON	7.0	0.06	177		I	-	< 0.001	PON	17.3	0.32	101	< 0.001	NO_3^-	-	< 0.001
					S	-	0.003								
POC:PON	5.2	0.09	174	< 0.001	T	+	< 0.001	POC:PON	9.5	0.14	99	< 0.001	NH_4^+	+	0.006
Pigments								Pigments							

Total chl <i>a</i>	27.2	0.45	156	< 0.001	S NH ₄ ⁺	+ +	< 0.001 0.006	Total chl <i>a</i>	22.5	0.51	100	< 0.001	I S NO ₃ ⁻	- - -	< 0.001 < 0.001 < 0.001
Prasinophyceae	13.8	0.24	157	< 0.001	NO ₃ ⁻ S I	+ + +	< 0.001 < 0.001 < 0.001	Prasinophyceae	18.7	0.50	99	< 0.001	T NO ₃ ⁻ NH ₄ ⁺	+ + +	< 0.001 < 0.001 < 0.001
Dinophyceae	19.5	0.36	156	< 0.001	<i>p</i> CO ₂ I S NO ₃ ⁻	+ + + -	< 0.001 < 0.001 < 0.001 0.001	Dinophyceae	14.5	0.39	100	< 0.001	S w	- -	< 0.001 < 0.001
Cryptophyceae	32.3	0.49	156	< 0.001	NO ₃ ⁻ T I	+ - +	< 0.001 0.001 0.01	Cryptophyceae	58.5	0.77	99	< 0.001	S NO ₃ ⁻ T	- - -	< 0.001 < 0.001 < 0.001
Haptophyceae	11.4	0.28	155	< 0.001	<i>p</i> CO ₂ I NO ₃ ⁻	+ + +	< 0.001 < 0.001 0.002	Haptophyceae	20.3	0.48	100	< 0.001	T NO ₃ ⁻ <i>p</i> CO ₂	- - -	< 0.001 < 0.001 < 0.001
Pelagophyceae	19.4	0.48	153	< 0.001	S I T PO ₄ ³⁻	+ + + +	< 0.001 < 0.001 0.001 0.005	Pelagophyceae	10.6	0.27	101	< 0.001	NO ₃ ⁻	-	< 0.001
Chlorophyceae	104	0.66	158	< 0.001	S T	+ +	< 0.001 0.002	Chlorophyceae	12.2	0.39	99	< 0.001	NO ₃ ⁻ S T I	+ - + -	0.001 0.004 0.004 0.005

Cyanophyceae	15.6	0.27	157	< 0.001	NO ₃ ⁻	-	< 0.001	Cyanophyceae	132	0.83	101	< 0.001	S	+	< 0.001
					PO ₄ ³⁻	-	< 0.001						T	+	< 0.001
													NO ₃ ⁻	+	< 0.001
Diatoms	12	0.29	155	< 0.001	I	-	< 0.001	Diatoms	40	0.45	100	< 0.001	S	-	< 0.001
					T	-	< 0.001						pCO ₂	+	< 0.001
					PO ₄ ³⁻	-	< 0.001						NH ₄ ⁺	+	< 0.001
													I	-	0.006
Flow cytometry								Flow cytometry							
<i>Prochlorococcus</i>	NA							<i>Prochlorococcus</i>	17.8	0.49	66	< 0.001	S	+	< 0.001
													T	+	< 0.001
<i>Synechococcus</i>	18.7	0.44	85	< 0.001	NO ₃ ⁻	-	< 0.001	<i>Synechococcus</i>	98.2	0.89	64	< 0.001	S	+	< 0.001
					I	-	< 0.001						T	+	< 0.001
													NO ₃ ⁻	+	< 0.001
													NH ₄ ⁺	-	0.003
Pico-eukaryotes	12.8	0.40	84	< 0.001	NO ₃ ⁻	-	< 0.001	Pico-eukaryotes	22.8	0.61	65	< 0.001	T	+	< 0.001
													w	+	< 0.001
													S	-	< 0.001
													NO ₃ ⁻	+	< 0.001
													NH ₄ ⁺	+	0.005
Nano-eukaryotes	NA							Nano-eukaryotes	27.4	0.65	65	< 0.001	S	-	< 0.001
													NO ₃ ⁻	-	< 0.001
													w	+	< 0.001
													NH ₄ ⁺	+	0.006
Particle flux								Particle flux							

POC	33.7	0.63	74	< 0.001	T	-	< 0.001	POC	NS
PON	34.6	0.63	74	< 0.001	T NO ₃ ⁻	- +	< 0.001 0.009	PON	NA
POC:PON	22.9	0.46	75	< 0.001	S	+	< 0.001	POC:PON	NA

1019

1020 Fig. 1

1021

1022 Fig. 2

Bay of Calvi

Bay of Villefranche

1023

1024 Fig. 3

Bay of Calvi

1025

1026 Fig. 4

1027

1028 Fig. 5

1029

Bay of Villefranche

1030

1031 Fig. 6

Bay of Villefranche

1032
1033 Fig. 7

1034

1035 Fig. 8