



Silicimonas algicola gen. nov., sp. nov., a member of the Roseobacter clade isolated from the cell surface of the marine diatom Thalassiosira delicatula

Klevi Crenn, Delphine Serpin, Cendrella Lepleux, Jörg Overmann, Christian
Jeanthon

► To cite this version:

Klevi Crenn, Delphine Serpin, Cendrella Lepleux, Jörg Overmann, Christian Jeanthon. *Silicimonas algicola* gen. nov., sp. nov., a member of the Roseobacter clade isolated from the cell surface of the marine diatom *Thalassiosira delicatula*. *International Journal of Systematic and Evolutionary Microbiology*, 2016, 66 (11), pp.4580 - 4588. 10.1099/ijsem.0.001394 . hal-01444937

HAL Id: hal-01444937

<https://hal.sorbonne-universite.fr/hal-01444937>

Submitted on 24 Jan 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

International Journal of Systematic and Evolutionary Microbiology

NEW TAXA - Proteobacteria

Silicimonas algicola gen. nov., sp. nov., a novel member of the *Roseobacter* clade isolated from the cell surface of the marine diatom *Thalassiosira delicatula*

Klervi Crenn^{1,2}, Delphine Serpin^{1,2}, Cendrella Lepleux³, Jörg Overmann³ and Christian Jeanthon^{1,2}

¹CNRS, Station Biologique de Roscoff, Adaptation and Diversité en Milieu Marin, Marine Phototrophic Prokaryotes Team, Roscoff, France.

²Sorbonne Universités, UPMC Univ Paris 06, Station Biologique de Roscoff, Adaptation et Diversité en Milieu Marin, Oceanic Plankton Group, Roscoff, France.

³Leibniz-Institut DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen, Braunschweig, Germany

Running title: *Silicimonas algicola* gen. nov., sp. nov.

Correspondence: C. Jeanthon, UMR 7144, Station Biologique, Place Georges Teissier, CS90074, 29688 Roscoff cedex, France; Phone : + 33 298 292 563 ; Fax : + 33 298 292 324 ; E-mail: jeanthon@sb-roscoff.fr

Key words: *Silicimonas algicola*, *Roseobacter* clade, algal-bacterial interactions, *Thalassiosira*.

The GenBank/EMBL/DDBJ accession numbers for the 16S rRNA gene sequence of strain KC90^T is KU926270.

Abstract

A Gram-negative, aerobic, non-motile bacterium, designated strain KC90B^T, was isolated from the surface of a cell of the marine diatom *Thalassiosira delicatula*. The bacterial cells were pleomorphic and formed very small beige colonies on marine agar. Optimal growth was obtained at 25°C, at pH 6.5-7.5 and in the presence of 1.5-2.0% (w/v) NaCl. Phylogenetic analyses based on its 16S rRNA gene sequence revealed that strain KC90B^T belonged to the *Roseobacter* clade and formed a monophyletic cluster with the sequences of *Boseongicola aestuarii*, *Profundibacterium mesophilum*, *Hwanghaeicola aestuarii*, *Maribius pelagius* and *M. salinus*, showing 91.4-95.7% sequence similarities. Ubiquinone Q-10 was the predominant lipoquinone but a significant amount of ubiquinone Q-9 was also detected. The major cellular fatty acids were C_{18:1} ω7c, 11-methyl C_{18:1} ω7c and C_{18:0}. Strain KC90B^T also contained specific fatty acids (C_{17:0}, anteiso C_{15:0} and anteiso C_{17:0}) that were not detected in its closest described relatives. The major polar lipids of strain KC90B^T comprised phosphatidylglycerol, phosphatidylcholine, diphosphatidylglycerol and an unidentified aminolipid. The DNA G+C content of strain KC90B^T was 65.2 mol%. The phylogenetic analysis of strain KC90B^T, together with the differential phenotypic and chemotaxonomic properties demonstrate that strain KC90B^T is distinct from type strains of *B. aestuarii*, *P. mesophilum*, *H. aestuarii*, *M. pelagius* and *M. salinus*. Based on the data presented in this study, strain KC90B^T represents a novel genus and species within the family *Rhodobacteraceae*, for which the name *Silicimonas algicola* gen. nov., sp. nov is proposed. The type strain is KC90B^T (=DSM 103371^T=RCC 4681^T).

Alphaproteobacteria are the most abundant heterotrophic bacteria found in marine pelagic environments (Zinger *et al.*, 2011) with a high contribution of the *Roseobacter* clade (family *Rhodobacteraceae*) (Buchan *et al.*, 2005; Luo & Moran, 2014). Members of the *Roseobacter* clade are often dominant in natural assemblages with marine algae and have been shown to increase in abundance during phytoplankton blooms (Amin *et al.*, 2012; Buchan *et al.*, 2014; Gonzalez *et al.*, 2000; Mayali *et al.*, 2008; Zubkov *et al.*, 2001). They also are often found in laboratory cultures of marine phytoplankton (Alavi *et al.*, 2001; Amin *et al.*, 2012; Grossart *et al.*, 2005; Jasti *et al.*, 2005) and both mutualistic (Geng & Belas, 2010; Wagner-Döbler *et al.*, 2010) and pathogenic (Boettcher *et al.*, 2005; Seyedsayamdost *et al.*, 2011) lifestyles have been suggested. To date, numerous *Roseobacter* clade genomes have been sequenced, revealing versatile metabolic capabilities that partly explain the success of the clade in marine environments. They gain energy from the oxidation of a multitude of organic compounds, and some members are also capable of phototrophy. Light utilization involving bacteriochlorophyll *a* (BChl *a*) by aerobic anoxygenic phototrophs (Moran *et al.*, 2004; Swingley *et al.*, 2007; Wagner-Döbler *et al.*, 2010) and based on rhodopsins (Newton *et al.*, 2010; Voget *et al.*, 2015) is found in phylogenetically diverse strains. Recently, Pujalte *et al.* (2014) divided the *Roseobacter* clade into 68 genera that correspond to 164 species but new genera and species have been described afterwards, including the genera *Boseongicola* (Park *et al.*, 2014), *Pseudoseohaecicola* (Park *et al.*, 2015), and *Xuhuaishuia* (Wang *et al.*, 2016). However, many other *Roseobacter* lineages do not have cultivated members.

In a study investigating the specificity of bacteria attached to marine diatom cells in laboratory cultures, we isolated bacteria attached to the cell walls of *Thalassiosira delicatula* RCC 2560 (Roscoff Culture Collection, France). This microalgal culture isolated from surface water at the coastal long-term monitoring station SOMLIT-Astan site (48°45' N, 3°57' W, north off Roscoff, Western English Channel) is maintained in the RCC since its isolation in January 2011. To isolate attached bacteria, single diatom cells were isolated under sterile conditions in a laminar flow hood. Algal cells were first gently separated by gravity using a 47 mm diameter, 11 µm pore-size nylon filter (Millipore) and washed three times with 50 mL of autoclaved seawater in order to lower the number of free-living bacteria in the algal culture. Single diatom cells were then picked with a sterile glass capillary micropipette and washed 3-4 times with filter-sterilized seawater. Controls were performed for each diatom cell isolated by checking

the absence of bacteria in the last drop of seawater used in the washing series. For cultivation of diatom epibionts, single isolated algal cells and controls were directly transferred in 48-well plates containing low-nutrient heterotrophic medium (LNHM) (Rappé *et al.*, 2002) prepared by dissolving 35 g.l⁻¹ of commercial sea salts (Red Sea Europe) instead of using natural seawater. Bacterial cultures were incubated at 19°C for 3 to 4 weeks and growth was analysed by flow cytometry using a BD Accuri C6 cytometer (BD Biosciences). Cultures that contained bacteria were streaked on LNHM agar for purification at least two times. Strain KC90B^T was one of the resulting isolates. Strain KC90B^T was further cultivated routinely in modified Marine Agar (1:10; 0.5 g peptone, 0.1 g yeast extract, 35 g sea salts dissolved in 1 l of Milli-Q water and 15 g agar) and in modified Marine Broth (MB) (1:2; 2.5 g peptone, 0.5 g yeast extract, 35 g sea salts dissolved in 1 l of Milli-Q water). The bacterial culture was then stored at -80 °C in the presence of 7.5% (v/v) DMSO.

Phenotypic characteristics of strain KC90B^T including growth, physiological and biochemical properties were tested as follows. Cell morphology and motility were examined using phase-contrast light microscopy (BX51; Olympus) and transmission electron microscopy (TEM) (JEM-1400, JEOL). TEM was performed after negative staining of cells with 2% uranyl acetate on Formvar-carbon-coated 400 mesh copper grids. Gram staining was performed according to (Smibert & Krieg, 1994). Growth at various temperatures (4-45 °C) and pH (4.5-10.5) were determined in MB (1:2). Media used to determine pH range for growth were adjusted using the following buffers: CH₃COONa 2M/acetic acid 2M for pH 4.5 to 5.5, Na₂HPO₄ 2M/NaH₂PO₄ 2M for pH 6 to 8.5 and Na₂CO₃ 1M/NaHCO₃ 1M for pH 9 to 10.5. The media were sterilized by filtration using 0.1µm pore size PES membrane filter units (Nalgene™ Rapid-Flow™). The requirement and tolerance to NaCl was tested in MB (1:2) using increasing concentrations of NaCl from 0 to 3 % (w/v) in increments of 0.5% and from 3 to 8 % in increments of 1%. Bacterial growth was assessed by flow cytometry. For flow cytometry, 100 µl cultures were fixed with glutaraldehyde (0.25%, final concentration) and stained with Sybr Green (Life Technologies) (Marie *et al.* 1997). Susceptibility to antibiotics was evaluated by spreading a bacterial suspension (200 µl) with a turbidity of 1-2 McFarland on MA (1:2) plates using susceptibility disks (bioMérieux) containing ampicillin (10 µg), chloramphenicol (30 µg), penicillin G (10 IU), gentamicin (10 µg), kanamycin (30 µg), streptomycin (10 µg), tetracycline (30 µg), nalidixic acid (30 µg), rifampicin (5 µg), erythromycin (15 µg) and neomycin (30 µg). Enzyme activities were determined after incubation at optimal growth temperature for 4 days, by using the API

ZYM system (bioMérieux). Assimilation tests were performed using the API 20 NE and API 50 CH systems incubated at optimal growth temperature for 15 days. All API test kits were used following the manufacturer's instructions except that the inoculating medium consisted of sterile Red Sea salts (35 ppt salinity) supplemented with mix of trace metals and vitamins solutions used in Carini *et al.* (2013). Catalase and oxidase activities were determined as described by Smibert & Krieg (1994).

Genomic DNA was extracted using lysis and neutralization buffers prepared as described in Humily *et al.* (2014). Briefly, after addition of 0.5 µl of lysis buffer, the mixture was incubated at 4°C for 10 min in a thermocycler. The lysate was further incubated at 95°C for 1 min, cooled at 4°C before adding 0.5 µL of neutralization buffer, and kept 3 min on ice until amplification by PCR. The 16S rRNA gene of KC90B^T was amplified using the primers 8F and 1492R (Turner *et al.*, 1999). The reaction mixture (12.5µL) contained 1µL of cell lysate, 0.1 mM of each deoxynucleoside triphosphate, 1X Green GoTaq Flexi Buffer, 2.0 mM MgCl₂, 0.2 µM of each primer, and 0.75 U of GoTaq G2 Flexi DNA polymerase (Promega). Conditions for PCR were as follows: 95°C for 10 min followed by 35 cycles (95°C for 30 s, 55°C for 1 min and 72°C for 1 min), and a final extension step for 10 min at 72°C. Sequencing was carried out using an Applied Biosystem 3100 automated DNA sequencer (Biogenouest platform, Station Biologique de Roscoff). The resulting 16S rRNA gene sequence (1395 nt) was compared by BLASTn with sequences available in GenBank. Phylogenetic analysis was performed using the neighbor joining, maximum parsimony and maximum likelihood inference approaches implemented in MEGA6 software (Tamura *et al.*, 2013). To amplify partial sequences of the *pufM* gene, coding for of the M subunit of the photosynthetic reaction centre, the PufMF forward (5'-TACGGSAACTGTWCTAC-3', Béjà *et al.*, 2002) and Puf-WAW reverse primers (5'-AYNGCRAACCACCANGCCCA-3', Yutin *et al.*, 2005) were used according to Lehours *et al.* (2010). For proteorhodopsin detection, the set of degenerated primers PR-1aF (5'-GATCGAGCGNTAYRTHGAYTGG-3') and PR-1aR (5'-GATCGAGCRTADATNGCCCANCC-3') was employed using conditions described by Campbell *et al.* (2008).

For genome analyses, genomic DNA was isolated from 500 mg harvested cells grown in MB (1:2) at 20°C after 15 days. The genome size and DNA G+C content were directly calculated from the complete genome sequence of the strain KC90B^T. Complete genome sequencing was carried out using the PacBio RSII System (Pacific Biosciences, Menlo Park, CA) at the Leibniz-Institut DSMZ. This calculation method differs from

conventional indirect methods used for the five reference strains [HPLC according to Tamaoka & Komagata (1984) or Mesbah *et al.* (1989)], but calculation of G+C content directly from genome is more accurate (Meier-Kolthoff *et al.*, 2014) and differences between two methods are between 1.2 and 2% (Mesbah *et al.*, 2011).

Cells (0.2 to 0.5 μm wide and 0.2 to 17 μm long) are Gram-negative, aerobic, non-flagellated and pleomorphic (few coccoids, some ovoids and mainly rod-shaped cells of various lengths) (Supplementary Fig. 1). Colonies on MA are circular, slightly convex, glistening, beige and 0.3–1 mm in diameter after incubation for 14 days at 25°C. The distinctive morphological, cultural, physiological and biochemical characteristics of strain KC90B^T are given in the genus and species descriptions (see below) and in Table 1.

Phylogenetic analysis based on the 16S rRNA gene sequence showed that strain KC90B^T formed a distinct lineage within the *Roseobacter* clade in the family *Rhodobacteraceae* of the *Alphaproteobacteria* (Figure 1), the nearest described relatives being *Boseongicola aestuarii* (95.7%), *Maribius pelagius* (94.0%), *M. salinus* (94.0%), *Profundibacterium mesophilum* (93.5%), and *Hwanghaeicola aestuarii* (91.4%). The lineage did not associate significantly with any of the currently described genera in the family. Interestingly, strain KC90B^T shared a higher sequence similarity (96.6%) with undescribed strain DG981 isolated from a culture of the toxic dinoflagellate *Gymnodinium catenatum* GCTRA14, originating from Spring Bay in Tasmania (Green *et al.*, 2004). The branching orders and phylogenetic relationships between strain KC90B^T and DG981-*Boseongicola*-*Profundibacterium*-*Maribius*-*Hwanghaeicola* were well conserved in the phylogenetic trees reconstructed using neighbor-joining, maximum-parsimony and maximum-likelihood algorithms.

The robustness of the phylogenetic relationships and the low sequence similarities between the strains and the other genera demonstrate that the novel isolate represents a new genus in the family *Rhodobacteraceae*.

The estimated genome size, based on genome sequencing data, was approximately 4.4 Mbp. The DNA G+C content of strain KC90B^T was 65.2 mol% as computed from genome sequences. Conclusively, no genes for *pufM* and proteorhodopsin could be detected for KC90B^T using PCR (data not shown).

For fatty acid analysis, cells were grown in liquid Marine broth for 10 days at 25°C. Data taken from the literature were obtained under growth conditions comparable to those

used for strain KC90B^T (Park *et al.*, 2014). After harvesting the biomass, cells were extracted according to the standard protocol (Sasser, 1990) of the Microbial Identification System (MIDI Inc.; version 6.1). The fatty acids were identified by comparison to the TSBA40 peak-naming table database. Strain KC90B^T has straight-chain, methyl- or hydroxy-branched saturated and monounsaturated fatty acids. The major fatty acid (>10% of the total fatty acids) detected in strain KC90B^T was C_{18:1}ω7*c* (60.0%). The fatty acid profile of the reference strain BS-W15^T showed the same prevalence of the fatty acid C_{18:1}ω7*c* (Park *et al.*, 2014). However, the fatty acid profile of KC90B^T is distinguishable from BS-W15 due to differences in fatty acid composition (Table 2). KC90B^T contains 2 anteiso fatty acids (*anteiso*-C_{15:0} and *anteiso*-C_{17:0}) while BS-W15^T does not have any. In addition, the fatty acid *cyclo*C_{19:0}ω8*c* (0.9%) and the unknown fatty acid 11.799 (2.8%) were detected in KC90B^T but not in BS-W15^T. Isoprenoid quinones were extracted from dried biomass with chloroform/methanol (2:1, v/v; Collins & Jones, 1981) and analysed via HPLC (Tindall, 1990). A large amount of ubiquinone Q-10 was detected (81.5%) which is typical of the *Alphaproteobacteria* class. In addition, a significant amount of ubiquinone Q-9 (18.4%) was detected. This profile differs from the one of BS-W15^T where ubiquinone Q-10 (predominant), Q-8 (16.0 %) and Q-9 (2.0 %) were detected (Park *et al.*, 2014). The polar lipid composition of strain KC90B^T was analysed by two-dimensional TLC (modified after Bligh & Dyer, 1959, Tindall *et al.*, 2007). The major polar lipids detected were phosphatidylglycerol, phosphatidylcholine, diphosphatidylglycerol and an unidentified aminolipid (Figure 2). In addition, minor amounts of three unidentified glycolipids, three unidentified phospholipids, one unidentified aminolipid and one unidentified lipid were detected. Compared to closely related genera, the polar lipid profile of strain KC90B^T is quite distinguishable. The strain *Boseongicola aestuarii* BSW15^T, *Profundibacterium mesophilum* JCM 17812^T, *Hwanghaeicola aestuarii* KACC 13705^T, *Maribius pelagius* KCCM 42336^T and *Maribius salinus* KCCM 42113^T do not present any glycolipids except for *H. aestuarii* KACC 13705^T. Except for *M. pelagius* KCCM 42336^T and *M. salinus* KCCM 42113^T, they all present low amounts of aminolipids (Park *et al.*, 2014). The polar lipid profile of strain KC90B^T is also distinguishable from other phylogenetically related genera of the *Roseobacter* clade such as *Marivita*, *Roseovarius* and *Litoreibacter* because of the absence of phosphatidylethanolamine as a major component (Hwang *et al.*, 2009; Kim *et al.*, 2012; Park & Yoon, 2013).

Strain KC90B^T was differentiated from the type strains *B. aestuarii*, *P. mesophilum*, *H. aestuarii*, *M. pelagius* and *M. salinus* by differences in its phenotypic characteristics, including cell morphology, motility, optimal temperature, salinity and pH for growth, assimilation of some substrates, susceptibility to antibiotics and some enzymatic activities. The phylogenetic and chemotaxonomic analyses and the different tested properties conclusively demonstrated that strain KC90B^T represents a novel genus and species in the *Roseobacter* clade (family *Rhodobacteraceae*, order *Rhodobacterales*), for which the name *Silicimonas algicola* gen. nov., sp. nov. is proposed.

Description of *Silicimonas* gen. nov.

Silicimonas [Si.li.ci.mo'nas L. n. *silex*, silica; L. fem. n *monas*, a monad, a unit; N.L. fem. n. *Silicimonas*, a monad isolated from silica]

Cells are Gram-negative, aerobic, non-flagellated and pleomorphic (few coccoids, some ovoids and mainly rod-shaped cells of various lengths). Catalase and oxidase positive. The major fatty acid is C_{18:1} ω7c. The predominant ubiquinone is Q-10. The major polar lipids are phosphatidylglycerol, phosphatidylcholine, diphosphatidylglycerol and an unidentified aminolipid. The genus is a member of the class *Alphaproteobacteria*, order *Rhodobacterales*, family *Rhodobacteraceae*. The type, and only species is *Silicimonas algicola*.

Description of *Silicimonas algicola* sp. nov.

Silicimonas algicola (al.gi'co.la. L. fem. n. *alga* alga or seaweed; L. suff. *-cola* from L. n. *incola* an inhabitant or dweller; N. L. fem. n. *algicola* alga dweller)

Cells are 0.2 to 0.5 μm wide and 0.2 to 17 μm long. Colonies on MA are circular, slightly convex, glistening, beige and 0.3–1 mm in diameter after incubation for 14 days at 25°C. Growth occurs at 10–40 °C (optimum 25°C), pH 6 to 9 (optimum 6.5–7.5), and 0.5–4% (w/v) NaCl (optimum 1.5–2%). No growth was obtained at 4°C or 45°C, at pH 5.5 and 9.5, and at NaCl concentrations of 0 and 4.5% (w/v). Nitrate reduction is negative. D-mannose, arbutin, esculine ferric citrate and potassium 2-ketoglutanate are utilized, but not glycerol, erythritol, D-arabinose, L-arabinose, D-ribose, D-xylose, L-xylose, D-adonitol, methyl-βD-xylopyranoside, D-galactose, D-glucose, D-fructose, D-mannose, L-sorbose, L-rhamnose, dulcitol, inositol, D-mannitol, D-sorbitol, methyl-αD-mannopyranoside, methyl-αD-glucopyranoside, N-acetylglucosamine, amygdalin,

salicin, D-cellobiose, D-maltose, D-lactose (bovine origin), D-melibiose, D-saccharose, D-trehalose, inulin, D-melezitose, D-raffinose, starch, glycogen, xylitol, gentiobiose, D-turanose, D-lyxose, D-tagatose, D-fucose, L-fucose, D-arabitol, L-arabitol, potassium gluconate and potassium 5-ketogluconate. Alkaline phosphatase, esterase (C4), esterase lipase (C8), leucine arylamidase, valine arylamidase, acid phosphatase, naphthol-AS-BI-phosphohydrolase, β -galactosidase, α -glucosidase and β -glucosidase activities are present, but lipase (C14) is weakly present and cystine arylamidase, trypsin, α -chymotrypsin, α -galactosidase, β -glucuronidase, N-acetyl- β -glucosaminidase, α -mannosidase and α -fucosidase activities are absent.

The major fatty acids are C_{18:1} ω 7c, 11-methyl C_{18:1} ω 7c and C_{18:0}.

The predominant ubiquinone is Q-10.

The major polar lipids are phosphatidylglycerol, phosphatidylcholine, diphosphatidylglycerol and an unidentified aminolipid.

The DNA G+C content of the type strain is 65.2 mol% by whole genome sequencing.

The type strain, KC90B^T (=DSM 103371^T=RCC 4681^T), was isolated from the silica cell wall of *Thalassiosira delicatula* RCC 2565, a marine diatom originating from Roscoff offshore seawater in the western English Channel.

Acknowledgements

This work was supported by the french national programme EC2CO-Microbien (project MICROMAR) and the MaCuMBA project funded by the European Union's Seventh Framework Programme (grant agreement no 311975). We are grateful to G. Tanguy (Biogenouest sequencing platform-FR2424-Station Biologique de Roscoff) and Sophie Le Panse (Merimage platform-FR2424-Station Biologique de Roscoff) for help with sequencing and transmission electron microscopy, respectively. We also warmly thank the following DSMZ members, Boyke Bunk and Cathrin Spröer for genome sequencing, Gabriele Pötter for analysis of cellular fatty acids, Susanne Verbag and Brian Tindall for the analysis of polar lipids and respiratory quinones. Klervi Crenn received a doctoral grant funded by Région Bretagne and CNRS.

References

Alavi, M., Miller, T., Erlandson, K., Schneider, R. & Belas, R. (2001). Bacterial

- community associated with *Pfesteria*-like dinoflagellate cultures. *Environ Microbiol* **3**, 380–396.
- Amin, S. A., Parker, M. S. & Armbrust, E. V. (2012).** Interactions between diatoms and bacteria. *Microbiol Mol Biol Rev* **76**, 667–684.
- Béjà, O., Suzuki, M. T., Heidelberg, J. F., Nelson, W. C., Preston, C. M., Hamada, T., Eisen, J. A., Fraser, C. M. & DeLong, E. F. (2002).** Unsuspected diversity among marine aerobic anoxygenic phototrophs. *Nature* **415**, 630–633.
- Bligh, E. G. & Dyer, W. J. (1959).** A rapid method of total lipid extraction and purification. *Can J Biochem Physiol* **37**, 911–917.
- Boettcher, K. J., Geaghan, K. K., Maloy, A. P. & Barber, B. J. (2005).** *Roseovarius crassostreae* sp. nov., a member of the *Roseobacter* clade and the apparent cause of juvenile oyster disease (JOD) in cultured Eastern oysters. *Int J Syst Evol Microbiol* **55**, 1531–1537.
- Buchan, A., González, J. M. & Moran, M. A. (2005).** Overview of the marine *Roseobacter* lineage. *Appl Environ Microbiol* **71**, 5665–5677.
- Buchan, A., LeCleir, G. R., Gulvik, C. A. & González, J. M. (2014).** Master recyclers: features and functions of bacteria associated with phytoplankton blooms. *Nat Rev Microbiol* **12**, 686–698. Nature Publishing Group.
- Campbell, B. J., Waidner, L. A., Cottrell, M. T. & Kirchman, D. L. (2008).** Abundant proteorhodopsin genes in the North Atlantic Ocean. *Environ Microbiol* **10**, 99–109.
- Carini, P., Steindler, L., Beszteri, S. & Giovannoni S. J. (2013).** Nutrient requirements for growth of the extreme oligotroph '*Candidatus* Pelagibacter ubique' HTCC1062 on a defined medium. *ISME J* **7**, 592–602.
- Choi, D. H., Cho, J. C., Lanoil, B. D., Giovannoni, S. J. & Cho, B. C. (2007).** *Maribius salinus* gen. nov., sp. nov., isolated from a solar saltern and *Maribius pelagius* sp. nov., cultured from the Sargasso Sea, belonging to the *Roseobacter* clade. *Int J Syst Evol Microbiol* **57**, 270–275.
- Collins, M. D. & Jones, D. (1981).** Distribution of isoprenoid quinone structural types in bacteria and their taxonomic implication. *Microbiol Rev* **45**, 316–354.
- Geng, H. & Belas, R. (2010).** Molecular mechanisms underlying *Roseobacter*-phytoplankton symbioses. *Curr Opin Biotechnol* **21**, 332–338. Elsevier Ltd.
- Gonzalez, J. M., Simo, R., Casamayor, E. O., Pedro, C. & Moran, M. A. (2000).** Bacterial community structure associated with a dimethylsulfoniopropionate-producing North Atlantic algal bloom. *Appl Environ Microbiol* **66**, 4237–4246.
- Green, D. H., Llewellyn, L. E., Negri, A. P., Blackburn, S. I. & Bolch, C. J. S. (2004).**

- Phylogenetic and functional diversity of the cultivable bacterial community associated with the paralytic shellfish poisoning dinoflagellate *Gymnodinium catenatum*. *FEMS Microbiol Ecol* **47**, 345–357.
- Grossart, H. P., Levold, F., Allgaier, M., Simon, M. & Brinkhoff, T. (2005).** Marine diatom species harbour distinct bacterial communities. *Environ Microbiol* **7**, 860–873.
- Humily, F., Farrant, G. K., Marie, D., Partensky, F., Mazard, S., Perennou, M., Labadie, K., Aury, J. M., Wincker, P. & other authors. (2014).** Development of a targeted metagenomic approach to study a genomic region involved in light harvesting in marine *Synechococcus*. *FEMS Microbiol Ecol* **88**, 231–249.
- Hwang, C. Y., Bae, G. D., Yih, W. & Cho, B. C. (2009).** *Marivita cryptomonadis* gen. nov., sp. nov. and *Marivita litorea* sp. nov., of the family *Rhodobacteraceae*, isolated from marine habitats. *Int J Syst Evol Microbiol* **59**, 1568–1575.
- Jasti, S., Sieracki, M. E., Poulton, N. J., Giewat, M. W. & Rooney-Varga, J. N. (2005).** Phylogenetic diversity and specificity of bacteria closely associated with *Alexandrium* spp. and other phytoplankton. *Appl Environ Microbiol* **71**, 3483–3494.
- Kim, J. M., Jung, J. Y., Chae, H. B., Park, W. & Jeon, C. O. (2010).** *Hwanghaeicola aestuarii* gen. nov., sp. nov., a moderately halophilic bacterium isolated from a tidal flat of the Yellow Sea. *Int J Syst Evol Microbiol* **60**, 2877–2881.
- Kim, Y. O., Park, S., Nam, B. H., Kang, S. J., Hur, Y. B., Kim, D. G., Oh, T. K. & Yoon, J. H. (2012).** Description of *Litoreibacter meonggei* sp. nov., isolated from the sea squirt *Halocynthia roretzi*, reclassification of *Thalassobacter arenae* as *Litoreibacter arenae* comb. nov. and emended description of the genus *Litoreibacter romanenko* et al. 2011. *Int J Syst Evol Microbiol* **62**, 1825–1831.
- Lai, P. Y., Miao, L., Lee, O. O., Liu, L. L., Zhou, X. J., Xu, Y., Al-Suwailem, A. & Qian, P. Y. (2013).** *Profundibacterium mesophilum* gen. nov., sp. nov., a novel member in the family *Rhodobacteraceae* isolated from deep-sea sediment in the Red Sea, Saudi Arabia. *Int J Syst Evol Microbiol* **63**, 1007–1012.
- Luo, H. & Moran, M. A. (2014).** Evolutionary ecology of the marine *Roseobacter* clade. *Microbiol Mol Biol Rev* **78**, 573–587.
- Marie, D., Partensky, F., Jacquet, S. & Vaultot, D. (1997).** Enumeration and cell cycle analysis of natural populations of marine picoplankton by flow cytometry using the nucleic acid stain SYBR Green 1. *Appl Environ Microbiol* **63**, 186–193.
- Mayali, X., Franks, P. J. S. & Azam, F. (2008).** Cultivation and ecosystem role of a marine *Roseobacter* clade-affiliated cluster bacterium. *Appl Environ Microbiol* **74**,

383 2595–2603.

384 **Meier-Kolthoff, J. P., Klenk, H. P. & Göker, M. (2014).** Taxonomic use of DNA G+C
385 content and DNA-DNA hybridization in the genomic age. *Int J Syst Evol Microbiol*
386 **64**, 352–356.

387 **Mesbah, M., Premachandran, U. & Whitman, W. B. (1989).** Precise measurement of
388 the G + C content of deoxyribonucleic acid by high-performance liquid
389 chromatography. *Int J Syst Bacteriol* **39**, 159–167.

390 **Mesbah, N. M., Whitman, W. B. & Mesbah, M. (2011).** Determination of the G+C
391 content of prokaryotes. In *Taxon Prokaryotes*, pp. 299–324. Edited by F. Rainey & A.
392 Oren. Academic Press.

393 **Moran, M. A., Buchan, A., González, J. M., Heidelberg, J. F., Whitman, W. B., Kiene, R.**
394 **P., Henriksen, J. R., King, G. M., Belas, R. & other authors. (2004).** Genome
395 sequence of *Silicibacter pomeroyi* reveals adaptations to the marine environment.
396 *Nature* **432**, 910–913.

397 **Newton, R. J., Griffin, L. E., Bowles, K. M., Meile, C., Gifford, S., Givens, C. E., Howard,**
398 **E. C., King, E., Oakley, C. A. & other authors. (2010).** Genome characteristics of a
399 generalist marine bacterial lineage. *ISME J* **4**, 784–798. Nature Publishing Group.

400 **Park, S., Park, J. M., Lee, K. C., Bae, K. S. & Yoon, J. H. (2014).** *Boseongicola aestuarii*
401 gen. nov., sp. nov., isolated from a tidal flat sediment. *Int J Syst Evol Microbiol* **64**,
402 2618–2624.

403 **Park, S., Park, J. M., Kang, C. H., Kim, S. G. & Yoon, J. H. (2015).** *Pseudoseohaecicola*
404 *caenipelagi* gen. nov., sp. nov., isolated from a tidal flat. *Int J Syst Evol Microbiol* **65**,
405 1819–1824.

406 **Park, S. & Yoon, J. H. (2013).** *Roseovarius sediminilitoris* sp. nov., isolated from
407 seashore sediment. *Int J Syst Evol Microbiol* **63**, 1741–1745.

408 **Pujalte, M. J., Lucena, T., Ruvira, M. A., Arahal, D. R. & Macian, M. C. (2014).** The
409 family *Rhodobacteraceae*. In *Prokaryotes - Alphaproteobacteria Betaproteobacteria*,
410 pp. 439–512. Edited by E. Rosenberg, E. F. DeLong, S. Lory, E. Stackebrandt & F.
411 Thompson. Springer-Verlag Berlin Heidelberg.

412 **Rappé, M. S., Connon, S. a, Vergin, K. L. & Giovannoni, S. J. (2002).** Cultivation of the
413 ubiquitous SAR11 marine bacterioplankton clade. *Nature* **418**, 630–633.

414 **Sasser, M. (1990).** Identification of bacteria by gas chromatography of cellular fatty
415 acids. In *MIDI Tech Note no 101 Microb ID, Inc, Newark*.

416 **Seyedsayamdost, M. R., Carr, G., Kolter, R. & Clardy, J. (2011).** Roseobacticides:
417 Small molecule modulators of an algal-bacterial symbiosis. *J Am Chem Soc* **133**,

- Smibert, R. M. & Krieg, N. R. (1994).** Phenotypic characterization. In *Methods Gen Mol Bacteriol*, American S., pp. 607–654. Edited by P. Gerhardt, R. G. EMurray, W. A. Wood & N. R. Krieg. Washington, DC.
- Swingley, W. D., Sadekar, S., Mastrian, S. D., Matthies, H. J., Hao, J., Ramos, H., Acharya, C. R., Conrad, A. L., Taylor, H. L. & other authors. (2007).** The complete genome sequence of *Roseobacter denitrificans* reveals a mixotrophic rather than photosynthetic metabolism. *J Bacteriol* **189**, 683–690.
- Tamaoka, J. & Komagata, K. (1984).** Determination of DNA base composition by reversed-phase high-performance liquid chromatography. *FEMS Microbiol Lett* **25**, 125–128.
- Tamura, K., Stecher, G., Peterson, D., Filipski, A. & Kumar, S. (2013).** MEGA6: Molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* **30**, 2725–2729.
- Tindall, B. J. (1990).** Lipid composition of *Halobacterium lacusprofundi*. *FEMS Microbiol Lett* **66**, 199–202.
- Tindall, B. J., Sikorski, J., Smibert, R. M. & Krieg, N. R. (2007).** Phenotypic characterization and the principles of comparative systematics. In *Methods Gen Mol Microbiol 3rd edn*, pp. 330–393. Edited by C. A. Reddy, T. J. Beveridge, J. A. Breznak, G. Marzluf, T. M. Schmidt & L. R. Snyder. Washington, DC: American Society for Microbiology.
- Turner, S., Pryer, K. M., Miao, V. P. & Palmer, J. D. (1999).** Investigating deep phylogenetic relationships among cyanobacteria and plastids by small subunit rRNA sequence analysis. *J Eukaryot Microbiol* **46**, 327–338.
- Voget, S., Wemheuer, B., Brinkhoff, T., Vollmers, J., Dietrich, S., Giebel, H.-A., Beardsley, C., Sardemann, C., Bakenhus, I. & other authors. (2015).** Adaptation of an abundant *Roseobacter* RCA organism to pelagic systems revealed by genomic and transcriptomic analyses. *ISME J* **9**, 371–384. Nature Publishing Group.
- Wagner-Döbler, I., Ballhausen, B., Berger, M., Brinkhoff, T., Buchholz, I., Bunk, B., Cypionka, H., Daniel, R., Drepper, T. & other authors. (2010).** The complete genome sequence of the algal symbiont *Dinoroseobacter shibae*: a hitchhiker's guide to life in the sea. *ISME J* **4**, 61–77.
- Wang, L., Liu, Y., Shi, X., Wang, Y., Zheng, Y., Dai, X. & Zhang, X.-H. (2016).** *Xuhuaishuia manganoxidans* gen. nov. sp. nov., a manganese-oxidizing bacterium isolated from deep-sea sediment of Pacific polymetallic nodule province. *Int J Syst Evol Microbiol* **In press**.

Yutin, N., Suzuki, M. T. & Be, O. (2005). Novel primers reveal wider diversity among marine aerobic anoxygenic phototrophs. *Appl Environ Microbiol* **71**, 8958–8962.

Zinger, L., Amaral-Zettler, L. A., Fuhrman, J. A., Horner-Devine, M. C., Huse, S. M., Welch, D. B. M., Martiny, J. B. H., Sogin, M., Boetius, A. & Ramette, A. (2011). Global patterns of bacterial beta-diversity in seafloor and seawater ecosystems. *PLoS One* **6**, e24570.

Zubkov, M. V., Fuchs, B. M., Archer, S. D., Kiene, R. P., Amann, R. & Burkill, P. H. (2001). Linking the composition of bacterioplankton to rapid turnover of dissolved dimethylsulphoniopropionate in an algal bloom in the North Sea. *Environ Microbiol* **3**, 304–311.

Figure captions

Figure 1. Neighbour-joining phylogenetic tree based on 16S rRNA gene sequences showing the position of strain KC90B^T and representatives of some related taxa. Only bootstrap values (expressed as percentages of 1000 replications) of > 40% are shown. Filled circles indicate that the corresponding nodes were also recovered using the maximum-likelihood and maximum-parsimony algorithms, while open circles indicate that the corresponding nodes were also recovered using the maximum-likelihood method. *Stappia stellulata* IAM 12621^T was used as an outgroup. Bar, 0.01 substitutions per nucleotide position.

Figure 2. Thin layer chromatograms of polar lipids of strain KC90B^T. GL1-GL3, unidentified glycolipids ; PL1-PL3, unidentified phospholipids; PC, phosphatidylcholine ; PG, phosphatidylglycerol ; AL, unidentified aminolipid ; DPG, diphosphatidylglycerol ; L, unidentified lipid.

Supplementary Figure S1. Transmission electron micrograph showing the pleomorphic forms [coccoid (a), ovoid (b), and rod-shaped (c)] of negatively stained cells of strain KC90B^T after growth for 10 days at 25°C in MB (1:2). Bar, 5 µm.

487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523

Table 1. Differential phenotypic characteristics of strain KC90B^T and the type strains of phylogenetically related species. Strains: 1. KC90B^T ;2. *Boseongicola aestuarii* BS-W15^T;3. *Profundibacterium mesophilum* JCM 17872^T;4. *Hwanghaeicola aestuarii* KACC 13705^T;5. *Maribius pelagius* KCCM 42336^T; 6. *Maribius salinus* KCCM 42113^T. Data obtained from this study and from Choi *et al.* (2007), Kim *et al.* (2010), Lai *et al.* (2013) and Park *et al.* (2014). +, positive reaction; -, negative reaction; w, weakly positive reaction; ND, not determined. All strains are positive for the following enzymatic activities: activity of esterase lipase (C8), leucine arylamidase, oxidase and catalase. All strains are negative for the following activities: acid production from D-melibiose, activity of trypsin, α -galactosidase, α -mannosidase, α -fucosidase, nitrate reduction.

Characteristics	1	2	3	4	5	6
	Culture of <i>Thalassiosira delicatula</i>	Tidal flat sediment at Boseong (South Korea)	Deep-sea sediment (Red Sea)	Tidal flat (Yellow Sea)	Surface water (Sargasso Sea)	Hypersaline water of a solar saltern (Korea)
Cell morphology	Pleomorphic	Pleomorphic	Coccoid	Coccoid	Rod-shaped	Rod-shaped
Motility	-	-	-	+	-	-
Optimal growth temperature (°C)	25	25	20–25	25–30	30–35	30–35
Growth temperature range (°C)	10–40	10–30	15–25	15–35	10–40	10–35
Optimal growth pH	6.5–7.5	7–8	7–8	6.5–7.5	ND	ND
Growth pH range	6–9	6.5–9.5	6–8.5	6–8	6–9	7–8
Optimal growth NaCl (%)	1.5–2	2	2–6	2–3	ND	ND
Growth NaCl range (%)	0.5–4	0.5–5	0.4–24	1.5–6	2–15	1–10
Colony size (mm)	0.3–1	0.4–0.8	0.1–0.3	ND	ND	ND
Colony color	beige	Yellowish-white	transparent	pale pink	beige	beige
Growth time on MA (days)	15	10	10	3–5	15–30	15–30
Assimilation of:						
Glycerol	-	ND	+	ND	+	-
L-arabinose	-	-	-	-	+	w
D-ribose	-	-	-	+	+	-
D-xylose	-	-	+	+	+	+
D-galactose	-	-	+	-	-	-
D-glucose	-	+	+	-	+	-
D-fructose	-	+	-	w	+	+
D-mannose	+	+	-	-	-	-
L-rhamnose	-	+	-	+	-	-
Inositol	-	-	-	-	-	+
D-mannitol	-	-	-	-	-	+
D-sorbitol	-	-	-	-	-	+
N-acetylglucosamine	-	ND	ND	ND	+	+
D-cellobiose	-	+	-	w	-	+
D-maltose	-	+	-	-	-	-
D-lactose (bovine origin)	-	+	-	-	-	-
D-trehalose	-	+	-	-	-	-
D-raffinose	-	+	-	-	-	+
L-tryptophane	-	ND	-	+	-	-
L-arginine	-	ND	+	ND	+	+
Urea	-	-	-	+	+	+
Gelatin	-	-	-	+	-	-
D-glucose	-	+	-	-	-	-
L-arabinose	-	-	-	-	+	w
D-mannose	-	+	-	-	-	-
D-mannitol	-	-	-	+	-	-
N-acetylglucosamine	-	ND	ND	-	+	+
D-maltose	-	+	-	-	-	-
Trisodium citrate	-	ND	+	-	+	+
Susceptibility to:						
Ampicillin	+	-	+	-	+	+
Chloramphenicol	+	-	+	+	+	+
Penicillin G	+	-	+	+	+	+
Gentamicin	w	-	+	-	+	+
Kanamycin	+	-	-	-	+	+
Streptomycin	+	-	+	-	+	-
Tetracycline	+	+	-	-	+	+
Nalidixic acid	-	ND	-	ND	-	-
Erythromycin	+	ND	ND	ND	+	+
Neomycin	+	+	+	+	+	+
Enzyme activity (API ZYM):						
Alkaline phosphatase	+	+	+	+	-	-
Esterase (C4)	+	+	-	+	+	+
Lipase (C14)	w	-	-	w	-	-
Valine arylamidase	+	-	+	w	-	-
Cystine arylamidase	-	-	-	w	-	-
α-chymotrypsin	-	-	-	w	-	-
Acid phosphatase	+	+	+	w	-	-
Naphthol-AS-BI-phosphohydrolase	+	-	+	w	-	-
β-galactosidase	+	-	+	-	+	+
β-glucuronidase	-	-	-	w	-	-
α-glucosidase	+	-	-	-	-	-
β-glucosidase	+	-	-	-	-	-
N-acetyl-β-glucosaminidase	-	-	+	-	-	-
DNA G+C content (mol%)	65.2	58.7	64.0	61.0	66.7	70.0

Table 2. Cellular fatty acid composition (%) of strain KC90B^T and its closest validly named relative BS-W15^T (data from Park *et al.*, 2014).

Fatty acid	KC90B ^T	BS-W15 ^T
Straight- chain		
C _{16:0}	3.6	1.8
C _{17:0}	0.5	-
C _{18:0}	5.9	5.5
Unsaturated		
C _{18:1} ω 7 <i>c</i>	60.0	73.1
C _{18:1} ω 9 <i>c</i>	1.8	1.7
C _{20:1} ω 7 <i>c</i>	-	0.9
Hydroxy		
C _{10:0} 3-OH	2.5	2.2
C _{12:0} 3-OH	0.7	<0.5
Methyl-branched		
<i>anteiso</i> -C _{15:0}	1.3	-
<i>anteiso</i> -C _{17:0}	0.7	-
11-methyl C _{18:1} ω 7 <i>c</i>	8.4	12.9
<i>cyclo</i> C _{19:0} ω 8 <i>c</i>	0.9	-
Unknown 11.799	2.8	-
Summed features		
3 (C _{16:1} ω 7 <i>c</i> / C _{16:1} ω 6 <i>c</i>)	0.6	0.8
7(C _{19:1} ω 6 <i>c</i> / unknown		
18.846 / <i>cyclo</i> -C _{19:1}	0.6	0.7
ω 10 <i>c</i>)		




Figure 1





Figure 2



Supplementary Figure 1. Transmission electron micrograph showing the pleomorphic forms [coccoid (a), ovoid (b), and rod-shaped (c)] of negatively stained cells of strain KC90B^T after growth for 10 days at 25°C in MB (1:2). Bar, 5 μ m.