Anthropogenic charcoal-rich soils of the XIX century reveal that biochar leads to enhanced fertility and fodder quality of alpine grasslands

I. Criscuoli, S. Baronti, G. Alberti, C. Rumpel, M. Giordan, F. Camin, L. Ziller, C. Martinez, E. Pusceddu, F. Miglietta

► To cite this version:

I. Criscuoli, S. Baronti, G. Alberti, C. Rumpel, M. Giordan, et al.. Anthropogenic charcoal-rich soils of the XIX century reveal that biochar leads to enhanced fertility and fodder quality of alpine grasslands. *Plant and Soil*, 2017, 411 (1), pp.499-516. 10.1007/s11104-016-3046-3 . hal-01469413

HAL Id: hal-01469413

<https://hal.sorbonne-universite.fr/hal-01469413>

Submitted on 16 Feb 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Anthropogenic charcoal-rich soils of the XIX century reveal that biochar leads to enhanced fertility and fodder quality of alpine grasslands

I. Criscuoli^{1,6*}, S. Baronti², G. Alberti^{3,4}, C. Rumpel⁵, M. Giordan⁷, F. Camin⁷, L. Ziller⁷, C. Martinez¹, E. Pusceddu², F. Miglietta^{1,2,8}

¹ Foxlab Joint CNR-FEM Initiative, Via E. Mach 1, 38010 San Michele all'Adige, Trento, Italy

² Institute of Biometeorology, National Research Council (IBIMET-CNR), Via Caproni 8, 50145 Firenze, Italy

³ University of Udine, Dipartimento di Scienze Agro-Alimentari, Animali ed Ambientali, via delle Scienze, 206, 33100, Udine, Italy, Currently at Chair of Silviculture, Faculty of Environment and Natural Resources, University of Freiburg, Tennenbacher str. 4, 79085 Freiburg, Germany

⁴ The EFI Project Centre on Mountain Forests (MOUNTFOR), Edmund Mach Foundation, via Edmund Mach 1, 38010 San Michele all'Adige, Trento, Italy

⁵ CNRS, IEES, Campus AgroParisTech, Thiverval-Grignon, France

⁶ Institute of Ecology and Environmental Sciences of Paris (UMR 7618), Université Pierre et Marie Curie Paris06 - Sorbonne (UPEC, UPMC, CNRS, IRD, INRA, Paris Diderot), 7 quai Saint Bernard, 75005, Paris, France

⁷ Research and Innovation Centre, Fondazione Edmund Mach (FEM), Via E. Mach 1, 38010 San Michele all'Adige, Trento, Italy

⁸ ECHO-EPFL, Laboratory of Ecohydrology, Ecole Polytechnique Fédérale de Lausanne, Lausanne, Switzerland

*Email: irene.criscuoli@gmail.com, tel. +39 0461 615389, fax. +39 0461 650183

Abstract

Background and Aims

Soil incorporation of charcoal (biochar) has been suggested as practice to sequester carbon, improve soil properties and crop yields but most studies have been done in the short term. Old anthropogenic charcoal-rich soils in the Alps enable to explore the long-term impact of charcoal addition to alpine grassland on seed germination, fertility and fodder nutritive value.

Methods

A germination test and a growth experiment in pots with *Festuca nigrescens* Lam. and *Trifolium pratense* L. were performed using three different substrates: control soil (i.e. sandy-loam brown acid soils with some podsolization), charcoal hearth soil (i.e. charcoal-enriched anthropogenic soils derived from the carbonization of larch wood on flat terraces) and control soil mixed with a fraction of fresh larch wood charcoal to reach the soil-charcoal ratio of 0.6.

Results

Both aged and fresh charcoal improved germination and markedly increased plant growth of the two plant species. The addition of fresh charcoal had an initial detrimental effect that disappeared in the second and third growth cycles. Plant Nitrogen:Phosphorus ratio revealed that growth was N-limited in the anthropogenic soils and P-limited in the control

41 and freshly amended soils demonstrating that biochar aging is critical to obtain a significant growth stimulation. Plant
42 nutrient contents revealed an improved fodder quality in both the charcoal amended soils.

43

44 *Conclusions*

45 Despite the occurrence of limited toxic effects on seedlings, larch wood charcoal appears to have positive effects on
46 fertility and fodder quality of alpine grasslands in the long term.

47

48 **Keywords:**

49 biochar, charcoal, alpine grasslands, fertility, N:P ratio, fodder nutritional value

50

51 **Acknowledgments**

52 The authors would like to thank the Stelvio National Park for allowing sampling campaigns; Ce.Spe.Vi. s.r.l.,
53 Pistoia for the experiment in the greenhouse; Dr. Luisa Andrenelli and Dr. Adriano Baglio (DISPAA Dipartimento di
54 Scienze delle Produzioni Agroalimentari e dell'Ambiente - University of Florence) for laboratory analysis; Roberta
55 Franchi (Fondazione Edmund Mach) for her important advise on alpine pastures and meadows management and the
56 Italian Biochar Association (ICHAR).

57 **Introduction**

58 Alpine grasslands are considered critical biodiversity hotspots (Väre et al. 2003) providing a series of
59 ecosystem services (Fontana et al. 2013) and sustaining the production of typical food resources with added value
60 (Bovolenta et al. 2011). The persistence of these semi-natural ecosystems is based on ancient and specific management
61 techniques adapted to the local needs ensuring their productivity for thousands of years (Poschlod and Wallisdevries
62 2002) and the accumulation of one of the highest soil carbon stock (Gamper et al. 2007; Ciais et al. 2010). Traditional
63 management techniques include intense land uses based on different stocking rates (different breeds and grazing period
64 lengths), mowing frequency, organic and inorganic fertilization (Maurer 2005), periodic re-sowing (Tozer et al. 2013)
65 and more rarely irrigation (Riedener et al. 2013). Major threats to these ecosystems arise from climate change, land
66 abandonment and inadequate intensification. In particular, abandonment has substantially reduced the surface area
67 occupied by mountain grasslands throughout Europe (Tasser et al. 2007), mainly leading to the expansion of secondary
68 forests (Tasser and Tappeiner 2002). Inadequate management techniques are instead increasing the risks of soil erosion,
69 landslides and avalanches (Tasser et al. 2003), often causing the diffusion of unpalatable plant species (Krahulec et al.
70 2001), loss of biodiversity (Dullinger et al. 2003), decreasing touristic attractiveness (Hunziker 1995) and, in some
71 instances, the loss of important soil organic carbon stocks (Poepflau and Don 2013). In the context of these changes,
72 appropriate management of mountain pastures and meadows is becoming a societal priority that should be based on
73 novel techniques ensuring the socio-economic viability of mountain communities, the conservation of their beauty as
74 well as their ecosystem services.

75 The use of biochar, a carbon rich co-product of pyrolysis, as soil amendment is currently receiving a lot of
76 attention as it can improve the physico-chemical properties of soils, reduce nutrient leaching (Glaser et al. 2001),
77 increase water infiltration and water holding capacity (Lim et al. 2016; Novak et al. 2016), enhance cation exchange
78 capacity (Liang et al. 2006), soil aeration (Case et al. 2012), thus stimulating biomass yields (Jeffery et al. 2011;
79 Biederman and Harpole 2012). Moreover, because of its substantial recalcitrance to microbial degradation, biochar
80 application to soil is also considered as an effective way to sequester atmospheric carbon (Lehmann et al. 2006; Sohi et
81 al. 2009; Criscuoli et al. 2014; Wang et al. 2015). On the other hand, the use of biochar as a soil amendment entails also
82 some risks which are not yet fully understood (Kuppusamy et al. 2016). In fact, biochar can be a source of toxicants
83 such as polycyclic aromatic hydrocarbon (PAHs) (Kloss et al. 2012), can retain heavy metals, can suppress the efficacy
84 of applied pesticides and may induce changes in soil microbial community composition and structure (Jenkins et al.
85 2016) with possible influence on the microbial-mediated transformation of nutrients. Moreover, its role in offsetting C
86 and the other greenhouse gases' emissions has been questioned because of the possible priming effect on the native soil
87 organic matter (Ventura et al. 2015; Fang et al. 2015), the induced changes in the surface albedo (Genesio et al. 2012)
88 and the possible associated C aerosol emissions (Genesio et al. 2016).

89 The large majority of studies involving biochar applications, including those made on lowland grasslands (Van
90 de Voorde et al. 2014; Schimmelpfennig et al. 2014; Schimmelpfennig et al. 2015), are based on short-term
91 experiments, mostly made immediately after biochar application. Only the work of Hernandez-Soriano et al. (2015)
92 showed that 150 years old charcoal kiln areas located in agricultural fields were still able to increase the productivity of
93 maize (+10%). As biochar properties were shown to change with time of exposure (i.e. ageing), in particular through
94 surface oxidation (Liang et al. 2006), short-term studies may be insufficient to assess its impact on soil fertility in the
95 long-term. The productivity of mountain grassland ecosystems is the result of a complex set of interactions between
96 grazing pressures and nutrients export, plant species composition and the establishment of soil microorganisms and soil

micro-fauna communities, which are related to specific functional traits, whose dynamics necessarily require long-term studies (Grigulis et al. 2013).

To properly address this critical issue, this paper assesses the effects of 158 years old anthropogenic charcoal rich soils (Criscuoli et al. 2014), derived from carbonization of larch wood (*Larix decidua* L.) on flat terraces (Backmeroff 2013), on germination, growth and nutritive value of a typical grass and mountain leguminous species (*Festuca nigrescens* Lam. and *Trifolium pratense* L. subsp. *nivale* (Koch)) in comparison to native (control) soil and control soil amended with fresh biochar derived from larch wood. The presence of multiple charcoal rich soils gave us the opportunity to work on a fully replicated scheme, to be considered as an analogue of a deliberate centennial time-scale application of biochar to a mountain grassland soil. The hypothesis is that plant productivity and nutritional value of both plant species, and especially of *T. pratense*, are higher in charcoal hearth soils compared to control soils and to soils amended with fresh larch charcoal because of char ageing (Pusceddu et al. 2013). The effects on seeds germination are expected to be less evident.

Materials and methods

Study site, soil sampling and charcoal production

Several charcoal-enriched anthropogenic soils (charcoal hearths) derived from the carbonization of larch (*Larix decidua* L.) wood on flat terraces, dating back to between 1500 and 1858 (Backmeroff 2013), were identified in Val di Pejo (Trentino, Northern Italy) at an altitude of 2150 m a.s.l., in a larch alpine grassland grazed in summer. A complete description of the site, soil characteristics and historical charcoal production can be found in Criscuoli et al. (2014). In brief, the charcoal hearths' soils are made of a surface organic horizon (~2 cm) and a thicker black anthropogenic layer (~20 cm) rich in charcoal residues left after the carbonization 158 years ago and today well mixed with the pre-existing soil layer. The control soils are sandy-loam brown acid soils with some podsolization (Lithic Dystrudept and Entic Haplorthod) (Smith and Atkinson 1975). Both control and hearths' soils have a pH of 5.1. The hearths' soils are very rich in carbon ($26.2 \pm 5.3 \text{ kg C m}^{-2}$), 90% of which is contained in the charcoal, and are also richer in nutrients than the surrounding control soils (Tab. 1).

Three paired sites (i.e. charcoal hearth and native soil as control) were selected on the basis of common aspect (SE) and conservation state (no significant geo-morphological dynamics or recent anthropogenic disturbances). In September 2014, 20 L of soil were sampled with shovels from the center of the three hearths and in the corresponding adjacent control areas, after removing the top organic layer.

Fresh charcoal was produced from larch wood at 450°C, the average temperature occurring in traditional carbonization wood piles (FAO 1987). Fragments of larch wood were carefully wrapped in Aluminum foil and placed in a preheated muffle furnace at a temperature of 450°C for 10 minutes. The carbon content of the freshly produced charcoal was $76 \pm 7 \text{ gC kg}^{-1}$, the nutrient composition is reported in the Tab. 1 and the specific surface area (total BET) was $239.5 \text{ m}^2 \text{ g}^{-1}$.

Germination test

Seeds of *Festuca nigrescens* Lam. and *Trifolium pratense* L. subsp. *nivale* (Koch) were obtained from a nursery. Total germination and germination rates for each species were assessed through a test in petri dishes. Three different substrates were considered: control soil (C), charcoal hearth soil (H) and control soil mixed with a fraction of

fresh larch wood charcoal (CC) to reach the soil-charcoal ratio of 0.6 equivalent to a charcoal dose of 39 kg m⁻², which was estimated as the initial char input at the three charcoal hearths 158 years ago (Criscuoli et al. 2014). 54 dishes were prepared (2 species x 3 soil types x 3 sampling areas x 3 replicates). In each petri dish, 10 seed of one species were sown and dishes were maintained in the dark. Germination was determined when the coleoptile was ≥ 2 mm for *F. nigrescens* and when the two cotyledons were visible and free from the seed coat for *T. pratense*.

Growth experiment

The three soil types were also used as substrates for a plant growth experiment in small pots of 135 cm³. Prior to sowing, the pots were left for two months in a greenhouse with regular irrigation in order to allow the seeds contained in the seed bank to germinate spontaneously. Naturally germinated plants were then periodically removed manually. To eliminate possible confounding effects due to differences in the soil temperature, caused by different light reflection/absorbance properties (albedo), the soil surface of each pot was covered with an uniform thin layer of black inert quartz granules of 1.5 mm diameter (Granulati Zandobbio, Bergamo). The plants were grown in monoculture for a total of 144 pots (2 species x 3 soil types x 3 sampling areas x 8 replicates). Seeds were sown at an initial density of 20 seeds pot⁻¹ on 7th January 2015 and the pots were stored on trays in a greenhouse, following a fully randomized experimental design. Air temperature and light in the greenhouse were left to fluctuate following external conditions, however temperature never dropped below 15°C by means of an automated heating system, while a minimum day length of 12 hours was ensured using artificial illumination during the winter months. Soil temperature was regularly checked at 3 cm depth using a digital thermal probe (109SS-L, Campbell scientific, Inc.) and no significant difference was observed among the soil treatments.

Nebulized irrigation provided water automatically at regular intervals, four times a day. Seven weeks after sowing, plant density was reduced to six plants per pot. Subsequently, three growth cycles were considered with complete aboveground biomass harvest made 17 (1st cycle), 22 (2nd cycle) and 30 weeks (3rd cycle) after sowing. At each harvest, the plants were oven dried at 80°C for 48 hours and weighed. At the end of the 3rd cycle, the roots were manually separated from the soil, washed and dried for 48 hours at 80°C and weighed.

Plant, charcoal and soil physico-chemical analysis

Plant, soil and charcoal samples were dried and finely ground prior to analysis.

Isotopic ratios were determined with an Isotope Ratio Mass Spectrometer (Thermo Fischer Scientific, Delta V Plus). Natural abundances of stable carbon (δ¹³C) and nitrogen (δ¹⁵N) isotopes were measured for both plant species and the fine soil fraction (<2 mm) of hearth and control soils as well as for both the ancient and the freshly produced charcoal fragments. The carbon isotope discrimination (Δ) was then calculated according to Farquhar and Richards (1984):

$$\Delta = \frac{\delta_a - \delta_p}{1 + \delta_p}$$

where δ_a is the δ¹³CO₂ in the air and δ_p is that of plant carbon.

Total Ca²⁺, K⁺, Mg²⁺, Na⁺, P, Cu, Fe were determined on oven-dried plant subsamples (105°C for 24 h) from the second harvest, according to the EPA method 3052 (USEPA 1996) using an ICP-OES spectrophotometer (Varian

Inc., Vista MPX). Total carbon and nitrogen concentrations were determined via total combustion in an elemental analyzer (EA Flash 1112 ThermoFinnigan).

Soil and charcoal pH were measured in a soil-charcoal/distilled water solution (1:4 ratio as reported in Di Lonardo et al. 2013; Vaccari et al. 2015).

The surface area of fresh charcoal was calculated by using the BET (Brunauer-Emmet-Teller) method and Langmuir method applied to nitrogen adsorption data in the relative pressure (P/P^0) range of 0.1-0.44.

Statistical analyses

Data were analyzed using R (version 2.15.3). The effects of the 3 treatments (C, H and CC) and of the 3 blocks (3 charcoal hearths and 3 control soils) were analyzed by a two-way ANOVA for each plant species. Individual comparisons were based on the Tukey's HSD post-hoc test. The data were checked for normality and homogeneity of variances by inspection of the residuals vs. fitted values and the Normal Q-Q plot. When data did not fulfill these requirements, they were log-transformed or square root-transformed. In the case of negative data ($\delta^{13}\text{C}$), the data were standardized. Germination rates were compared using a binomial test (test for equality of proportions). The p-values of individual comparisons were corrected with the Holm method. All data in the text, figures and tables are reported as mean $\pm$ standard deviation, if not differently stated.

Results

Germination test

Plants started to germinate after 96 hours of incubation. The percentage of germinated seeds after 96 and 120 hours from sowing was higher in CC compared to C and H both for *F. nigrescens* and *T. pratense* (Tab. 2), while early germination rates were similar or the same in C and H. Looking at the total number of germinated plants, almost all seeds germinated in all soil types (between 78 and 94%). The plant germination rates at the end of the experiment for both plants species were higher in both CC and H than in C (*F. nigrescens*: CC vs. C +14%, $p=0.08$ and H vs. C +13%, $p=0.09$; *T. pratense*: CC vs. C +13%, $p=0.05$ and H vs. C +12%, $p=0.07$).

Plant growth

At the end of the first growth cycle (week 1-17), both plant species grew more in H than in C and CC (Fig. 1) and the lower growth observed in the CC was highly statistically significant if compared to both H and C. The relative changes in plant biomass in comparison to C were +21, +9% for H and -35, -71% for CC, for *T. pratense* and *F. nigrescens*, respectively. The mean plant growth rate (MGR) was much lower in CC (*T. pratense*: $1 \text{ mg day}^{-1} \text{ pot}^{-1}$; *F. nigrescens*: $1.5 \text{ mg day}^{-1} \text{ pot}^{-1}$) than in C (*T. pratense*: $2.3 \text{ mg day}^{-1} \text{ pot}^{-1}$; *F. nigrescens*: $3.2 \text{ mg day}^{-1} \text{ pot}^{-1}$) and H (*T. pratense*: $2.8 \text{ mg day}^{-1} \text{ pot}^{-1}$; *F. nigrescens*: $3.5 \text{ mg day}^{-1} \text{ pot}^{-1}$).

At the end of the second growth cycle, biomass was again significantly higher in H than in C and CC for both species (Fig. 1). *T. pratense* and *F. nigrescens* plants grown on H produced almost 3 and 1.5 times more than on C, respectively. Plants grown on CC performed better than on the control soil (both *T. pratense*, *F. nigrescens* = +37%), even though the difference was significant only in the case of *T. pratense* ($p=0.01$). MGR was accelerated compared to the first cycle and rose to 10.7 and 41.1 $\text{mg day}^{-1} \text{ pot}^{-1}$ for *T. pratense* and to 16.6 and 40.7 $\text{mg day}^{-1} \text{ pot}^{-1}$ for *F. nigrescens* grown in C and H, respectively.

Similarly to the second cycle, at the end of the third cycle plant biomass was significantly higher in H than in the other two treatments (Fig. 1). *F. nigrescens* plants grown on H produced almost 1.5 times more than plants grown on C and *T. pratense* 0.6 times more. Plants grown on CC performed better than on C (*F. nigrescens* = +51%, *T. pratense* = +22%) even though the difference was significant only in the case of *F. nigrescens* ($p=0.01$). The MGR decreased compared to the second cycle to 24.1, 9.9 and 15 mg day⁻¹ pot⁻¹ for *F. nigrescens* and to 19.1, 12.1 and 14.9 mg day⁻¹ pot⁻¹ for *T. pratense* grown in H, C and CC, respectively.

Overall, the total amount of biomass produced over the three cycles (above and belowground) was 88% and 108% higher in H compared to CC ($p<0.001$) and 114% and 148% higher in H compared to C ($p<0.001$) for *F. nigrescens* and *T. pratense*, respectively (Fig. 2). Total biomass was slightly higher in CC compared to C for both plant species, but the differences were not significant (*F. nigrescens*: $p=0.29$, *T. pratense*: $p=0.25$).

The root:shoot ratio, was significantly higher for *T. pratense* plants grown on H than C ($p=0.02$), while for *F. nigrescens* no differences in the ratio were detected among treatments (Fig. 3).

Soil and plant nutrients

The content of Ca²⁺, Mg²⁺, Na⁺ and P (mg g⁻¹) in the plants tissues (Tab. 3) was the highest in *F. nigrescens* and *T. pratense* grown on H followed, in most cases, by plants grown on CC. The differences in the nutrient content of *T. pratense* between H and C were significant at $p=0.10$ for all the nutrients, while no significant differences between H and CC in *T. pratense* and among all treatments for *F. nigrescens* were detected (Tab. 3). The K⁺ content of *T. pratense* was higher in plants grown on H (H vs. CC: $p=0.18$; H vs. C: $p=0.06$), but for *F. nigrescens* plants it was higher when plants were grown on CC, even though not significantly. An opposite trend was observed for N, which was higher for plants grown on C for both species, followed by CC and H. Fe content showed a very high variability in plants grown on the same soil type, while soil Cu content was very similar for all the three treatments. The nutrient content of *T. pratense* was usually higher than that of *F. nigrescens* with the exception of P that was higher for *F. nigrescens* plants when grown on H and C.

The N:P ratio of *F. nigrescens* and *T. pratense* plants grown on H was 10.4 and 10.7, respectively, while plants grown on C showed a ratio of 18.3 and 21.5 and plants grown on CC soils 23.3 and 16.3 (Fig. 4).

Soil nutrient content data were taken from Criscuoli et al. (2014) and are here reported in the Tab. 1. The soil N:P ratio in soil was 13.3, 5.7 and 11 in C, H and CC, respectively (Fig. 4).

Plant P concentration increased with increasing soil concentration. The highest values were observed for the H soil and in the plants grown on it, while soil and plants grown on CC and C gave very similar results (Fig. 5). Plant K⁺ concentration correlated positively with the soil nutrient contents. Concentrations of both soils and plants decreased in the order H>CC>C (Fig. 5). We observed a similar behavior for Ca²⁺ in the case of *T. pratense*, while the content of *F. nigrescens* was insensitive to the soil content variations (Fig. 5). Na⁺ content was the lowest for H soils. The concentration of the plants tissues decreased for *T. pratense* with increasing soil concentrations, while values observed for *F. nigrescens* did not reflect the Na⁺ soil content (Fig. 5). Mg²⁺ concentration of the soil was not reflected in the content of plants for both species. The Mg²⁺ content of *T. pratense* was double compared to *F. nigrescens* (Fig 5, Tab. 1 and Tab. 3).

Isotopic signature

The ^{13}C discrimination (Δ) at the end of the first growth cycle was significantly higher for *F. nigrescens* and *T. pratense* plants grown on CC (24.5 and 23.6‰, respectively) than on C (21.8‰ with $p=0.0006$ and 22.1‰ with $p=0.051$) and H (21.5‰ with $p=0.0004$ and 22.1‰ with $p=0.051$).

At the end of the second growth cycle, the Δ of both plant species grown on H and C slightly increased in comparison with the first one (*T. pratense*: H +0.74‰, C +0.42‰; *F. nigrescens*: H +1.02‰, C +0.56‰), while, when they were grown on CC, the Δ decreased (*T. pratense*: -0.56‰; *F. nigrescens*: -1.55‰) to similar values for both species (22.99‰) and to the other two soil treatments. At the third harvest, plants showed smaller Δ values and no differences among treatments (Fig. 6).

At the end of the first growth cycle, the $\delta^{15}\text{N}$ was very similar for *T. pratense* and *F. nigrescens* plants grown on C soils (6.9 and 7‰, respectively). The isotopic signature of plants grown on H was lower and very similar for the two plant species as well (*T. pratense*: 4.9‰; *F. nigrescens*: 5.1‰). Similar data were observed at the end of the second growth cycle (*T. pratense* C: 6.5‰, H: 4.8‰; *F. nigrescens* C: 6.4‰, H: 5.8‰), while, at the third harvest, the isotopic signature rose in both species and became slightly higher for plants grown on H as compared to those grown on C (*T. pratense* C: 7.2‰, H: 7.7‰; *F. nigrescens* C: 7.1, H: 7.3‰; Fig. 7). A very different trend was observed for plants grown on CC where, at the end of the first growth cycle, *T. pratense* plants showed a $\delta^{15}\text{N}$ of -0.07‰, much lower than the other treatments, especially compared to C (CC vs. H $p=0.19$; CC vs. C $p=0.08$). *F. nigrescens* had a $\delta^{15}\text{N}$ equal to 3.3‰ (CC vs. H $p=0.24$; CC vs. C $p=0.04$). After the second growth cycle, the isotopic signature of both *T. pratense* and *F. nigrescens* increased (1.4‰ and 4.8‰, respectively), but did not equal the values observed for the other two soil treatments (*T. pratense*: CC vs. H $p=0.06$, CC vs. C $p=0.02$; *F. nigrescens*: CC vs. H $p=0.25$, CC vs. C $p=0.07$). The $\delta^{15}\text{N}$ of both species increased also during the third growth cycle (*T. pratense*: 5.3‰ and *F. nigrescens*: 5.8‰), getting closer to the signature measured in the other treatments (ranging between 7.1 and 7.7‰; Fig. 7).

The control soil had a $\delta^{15}\text{N}$ equal to 3.9 ± 1.3 ‰, markedly higher than H (2.1 ± 1.1 ‰), while the ancient and the freshly produced charcoal had very similar $\delta^{15}\text{N}$ signatures (1.1 ± 1.6 ‰ and 0.9 ± 0.2 ‰, respectively).

Discussion and conclusions

The three substrates had differential effects on germination. For both plant species, early germination rates were higher in the control soils amended with fresh charcoal (CC) than in hearths soils (H) and control soils (C). Moreover, the total number of germinated seeds 10 days after sowing was stimulated for both species by the presence of old (H) and fresh charcoal (CC) amendment (Tab.2). Previous research on *T. pratense* by Van de Voorde et al. (2014) found no effect of biochar addition to the soil substrate on the germination and Solaiman et al. (2011) found a dose-dependent negative germination effect on another *Trifolium* species. In contrast, we did not observe any inhibitory effect on seeds germination, but a tendency to an increase in germination in the presence of both fresh and old charcoal. These positive effects could be related to the presence of specific compounds such as karrikins, a family of butenolides, which are known to be contained in charcoal (Flematti et al. 2008; Nelson et al. 2012). Karrikins act as germination stimulants and have been recently isolated, identified (Flematti et al. 2004) and successfully synthesized (Flematti et al. 2011) even though the mechanisms by which they might trigger seed germination are far from being completely described. Another possible mechanism explaining the higher germination rates in CC and H could be related to the ability of charcoal to adsorb biogenic phyto-toxins eventually present into the soil (Garnett et al. 2004; Hille and Den Ouden 2005).

In the pot experiment, at the end of the first growth cycle, both species grew better on H than on C and CC (Fig. 1). Reduced growth on CC was remarkable and attributable to a stress effect caused by the addition of fresh char. Such a stress caused a decrease in the photosynthetic capacity of plant leaves, which was reflected in an increase in their $\Delta^{13}\text{C}$ (Fig. 6) (Brugnoli et al. 1989). A possible source of stress affecting plant growth on CC could have been the difference in the pH between the fresh larch charcoal (pH=7.1) and the control soil to which it was added to (pH=5.1). Soil alkalization is known to have detrimental effects on plants as high pH generally causes metal ions to precipitate, thus affecting the absorption of inorganic ions and disrupting the ionic balance of tissues (Chen et al. 2009). In our case, the bulk soil pH of CC rose to 5.8 and we cannot exclude that plant roots were in direct contact with charcoal fragments with a higher pH. However, 7.1 is a value close to neutrality and, moreover, Gell et al. (2011) did not find a clear relationship between pH and short-term phyto-toxicity on lettuce, radish and wheat plants. Thus, the toxic effect observed in the first growth cycle is very likely not to be linked to a pH stress. On the contrary, there could be a direct phytotoxic effect of charcoal on plant growth, both on germination and root and shoot growth as observed in previous studies that related the phyto-toxicity to the presence of chemicals in the biochar. For example, Rombolà et al. (2015) made the hypothesis that the phyto-toxicity was due to NH_3 , Volatile Fatty Acids and Benzoic acids contained in the charcoal, while Gell et al. (2011) associated the toxic effect of different kinds of biochar with their high water-soluble salts content, such as Chloride and Sodium, and possibly aliphatic and aromatic hydrocarbons such as phenols.

Observations made during the first growth cycle have important implications regarding the possibility of using biochar as a soil amendment in the framework of mountain pastures and meadows management and restoration. The data showed that while long-term exposure of pyrogenic carbon in soil, as occurred in H, buffers any possible toxic effect, the sudden addition of biochar could have detrimental effects on plant growth. Even if the exact temporal dynamics of the transition between toxic and non-toxic conditions following biochar incorporation remains unknown, these data already suggest that the negative effect of charcoal incorporation into the soil is transitory as better environmental conditions certainly develop during biochar ageing. Moreover, transitory toxic effects can be avoided if is a) washed with water or an organic solvent, b) degraded via a biological activity through composting or mixing it with activated sludge (Bargmann et al. 2013; Rombolà et al. 2015) or c) dried at temperatures between 100 and 300°C for 24 hours (Koltowski and Oleszczuk 2015) before being amended to soil. It is also likely that lower doses than used in the present experiment, could reduce detrimental effects on the plant growth, even in the very early stages.

The hypothesis of a transitory toxic effect of biochar on plant growth is also confirmed when looking at biomass production during the second and the third growth cycles (Fig.1). In fact, at the end of the second cycle, biomass was again significantly higher for plants grown on H compared to those grown on C. *T. pratense* plants grown on H produced almost threefold if compared to C and *F. nigrescens* 1.5-times more. On the other hand, conversely to what was observed in the first cycle, plants grown on CC performed better than those grown on C (both *T. pratense* and *F. nigrescens*= +37%). The recovery of both species grown in CC was also confirmed by the $\Delta^{13}\text{C}$, which was not significantly different from the other two treatments (Fig. 6). The mean plant growth rate measured for plants grown on H was 4 and 3 times higher than those of plants grown on C for *T. pratense* and *F. nigrescens* and 2 and 3 times higher compared to those grown on CC, respectively, indicating a clearly higher fertility in charcoal hearths soils than in their natural counterparts and soils with fresh biochar. Similar results were obtained by Naisse et al. (2014), who observed much higher (micro-)biological activity in charcoal hearths as compared to control soils without ancient charcoal. The increase in the charcoal hearth soils' fertility of our study site can be explained by the accumulation over time of soil

nutrients, their higher plant availability as well as improved physical characteristics of charcoal hearth soils compared to control (Criscuoli et al., 2014, Tab. 1).

Similarly to the second cycle, also at the third harvest, plant biomass was significantly higher for H than for the other two treatments (Fig. 1). However, the biomass measured for *T. pratense* grown on H at the end of the third cycle was lower in comparison to the second one. This could be due to the higher temperatures registered during the third growth cycle (summer) compared to the second one (spring), which caused the spread of necrotic tissues in the leguminous species (data not shown).

Overall, the total plant biomass produced over the three cycles (above and belowground) was significantly higher when plants were grown on H compared to CC and C both for *F. nigrescens* and *T. pratense* (Fig. 2). No significant differences were detected between plants grown on CC and C, making the positive impact of charcoal on the long term productivity of alpine grasslands even more evident than the results of one single growth cycle.

The root:shoot ratio, a widely used indicator of plants health, gave differential results according to the species. It was significantly higher for *T. pratense* plants grown on H than for those grown on C, while for *F. nigrescens* no difference in the ratio was detected among treatments (Fig. 3). The different behavior between the two species is related to the high interspecific variation in root:shoot ratios as reported by Koerner & Renhardt (1987). The results observed for *T. pratense* seem to be in contrast with previous literature. In fact, for another species of clover (Solaiman et al. 2012), inconsistent effects of biochar on root:shoot ratio were observed. At high application rates charcoal had a negative effect. Moreover, it is known that the development of the root compartment may be smaller compared to aboveground biomass in nutrient rich soils (Agren and Franklin 2003), but according to our results *T. pratense* had a much higher root to shoot ratio in H soil, which shows the highest nutrient contents, both total and available (Criscuoli et al., 2014, Tab. 1). A possible explanation of high root to shoot ratios for *T. pratense* grown on H may be that roots development was more driven by hormonal factors rather than nutrients concentration. In fact, an increase in ethylene production, a plant hormone with important implication for plant growth and development, has been previously observed from biochar and biochar-amended soil (Spokas et al. 2010).

Soil and plants nutrient contents provided useful information to further examine and interpret our results. The idea that the ratios of N, P and K in plant tissues provides an indication of the relative availability of these nutrients in the soil has often been discussed since the Von Liebig's Law of the Minimum (Von Liebig 1840). Koerselman & Meuleman (1996), among others, assumed that the N:P ratio of plant tissues is a reliable indicator of soil fertility and proposed, on the basis of a meta-analysis of experimental data, that N:P thresholds of plant tissues can be used as indicators of N-limitations ($N:P < 14$) or P limitations ($N:P > 16$) or their co-limitation ($14 < N:P < 16$) to plant growth. Accordingly, plants of both *F. nigrescens* and *T. pratense* species grown on H soils can be considered mostly N-limited (leaves N:P ratios of 10.4 and 10.7, respectively), while they were P-limited when grown on C and CC soils ($N:P$ ratios > 16 ; Tab. 3 and Fig. 4). Thus, the lower plant biomass production in C and CC could be related to a P-limitation. In fact, total and available P-concentrations of these soils were similar and about one third compared to that of H soils (Criscuoli et al., 2014; Tab. 1). The soil P concentration for C and CC was reflected in the plants concentration which was similar (Tab. 3 and Fig. 5), showing that P added via charcoal was available for plants. This has been previously demonstrated specifically for larch wood (*Larix gmelinii* Rupr.) charcoal produced at 400°C (available P: 42.7 mg kg⁻¹ biochar) and for the larch seedlings grown on a mixture of sand larch biochar which had higher foliar P concentrations in the presence of higher rates of char (Makoto et al., 2011). Thus, the higher P content observed in the H soils and plants growing on it have to be explained via other processes taking place in the long term other than the initial P input

via charcoal addition. It has been shown that P availability in charcoal amended soils is enhanced also via an increased mycorrhizal colonization and a change in soil P fractionation (Graber et al., 2015). Moreover, charcoal hearth soils might have accumulated the P atmospheric depositions due to desert dust transport and atmospheric pollution (Bergametti et al. 1992) or dung released from cattle over the 158 years of exposure in soil.

The higher plant N:P ratio observed in CC soils for *F. nigrescens* compared to *T. pratense* (Fig. 4) is due to the higher N content in *T. pratense* (19.2 vs. 14.2 mg g⁻¹ in *F. nigrescens*), while P content is almost the same in the two species (0.9 vs. 0.8 mg g⁻¹, respectively; Tab. 3). The higher N content of the leguminous species might be explained by its nitrogen fixing capacity and this is also confirmed by the lower $\delta^{15}\text{N}$ that we observed for plants grown on CC in the first and second growth cycle, much closer to the atmospheric value (0 ‰) as compared to *F. nigrescens* (Fig. 7). These results are in line with several previous studies reporting an increase in nitrogen fixation in biochar amended soils (Rondon et al. 2006; Ogawa and Okimori 2010; Güereña et al. 2015; Van Zwieten et al. 2015) which has been mainly related to a greater B, Mo and P availability. The $\delta^{15}\text{N}$ signature of *F. nigrescens* plants grown on CC was higher than the signature of *T. pratense* grown on the same soil, but was lower than the signature of plants of the same species grown on the other soil types along the three growth cycles. This might be explained by the uptake of nitrogen contained in the fresh charcoal fragments (De la Rosa and Knicker, 2011), as the nitrogen isotopic signature of charcoal fragments is close to 1‰, while control soils have a signature of 3.9 ± 1.3 ‰.

Similar to P, the concentration of K⁺ in tissues of both plant species and of Ca²⁺ in *T. pratense* plants correlated positively with soil nutrient content (Fig. 5), with higher concentrations observed in the case of higher biomass production (H>CC>C). Similar observations were made by Schimmelpfennig et al. (2015). Even if the Ca²⁺ content is much lower in C compared to H and CC, it cannot be considered as limiting factor as its concentration is well above the critical deficiency levels (0.5-1.5 cmol_c kg⁻¹) reported by Kopittke and Menzies (2007). The concentration of Ca²⁺ was doubled in the leguminous species compared to grass, in agreement with literature results (Juknevičius and Sabiené 2007).

Na⁺ content of the plants did not reflect the concentration of the nutrient in the soil (Fig. 5). Sodium is toxic at high concentrations for most of the plant species (Greenway and Munns 1980), so the adsorption through the plant roots' cell is limited as much as possible by mechanisms of selective uptake and ion exclusions (Schachtman and Liu 1999). The behavior of Na⁺ is opposite to what was observed for K⁺. The two ions are very similar for their ionic radius and ion hydration energies, factors determining their movement through the cell's membrane (Hille 1992), but the competition between the two ions in soils which are not saline is usually in favor of K⁺ because it is not toxic. On the contrary, it is fundamental for plant growth of all species and it is the cation with the highest concentration in plant tissues (Mäser et al. 2002). As for Ca²⁺, the concentration of *T. pratense* plants is double, or more, compared to *F. nigrescens* even if *T. pratense* is less tolerant to salinity than *F. nigrescens* (FAO 2002). However Na⁺ concentration in the leaves of *T. pratense* did not reach toxic levels, in fact no detrimental effects were observed on the plant growth.

Similarly to Na⁺, higher soil concentrations of Mg²⁺ did not correspond to any increase in the magnesium content in plant tissues. The concentrations of Na⁺ and Mg²⁺ were lower (H<CC<C) in the soils with elevated K⁺ and Ca²⁺ concentrations (H>CC>C). This may be related to the competition between cations given a certain cation exchange capacity of the soil. The fact that lower Mg²⁺ soil content was not reflected by a lower concentration in plants tissues might be seen as the evidence of a proper cations balance in the soil. As for Na⁺ and Ca²⁺, Mg²⁺ content of *T. pratense* plants is twice than that of *F. nigrescens* and this is in line with previous works on these two plant species (Juknevičius and Sabiené, 2007).

The micro- and macro-nutrient contents of pasture and meadow plants are fundamental for the quality of the fodder. The German Society of Nutritional Physiology, among others, provides specific recommendations at this regard (Flachowsky et al. 2001, Online Resource Tab. 1). In our experiment the P content of fodder improves when plants were grown on H compared to those grown on C and CC but did not fulfill the cattle requirements.

K⁺ content of plants increased for both charcoal treatments compared to C where both plant species had lower concentrations than recommended. K⁺ content in *F. nigrescens* was in line with recommendations when grown on H and CC, while K⁺ concentrations in *T. pratense* plants exceeded the recommendations when grown on H. However, the value did not reach the limit of $\geq 35 \text{ g kg}^{-1}$ dry biomass, which is considered to be the cause of the so called “grass tetany”, a Mg²⁺ deficiency in ruminants which implies cattle to be fed with supplementary magnesium (Kessler, 2001).

Ca²⁺ and Mg²⁺ concentrations were very high for both plant species grown on all soil types compared to the recommendations, especially for those grown on soils amended with charcoal and for *T. pratense* plants. But an adequate nutritive balance between K⁺, Ca²⁺ and Mg²⁺ was guaranteed, as the ratio K:(Ca+Mg) was lower than 2.2 (Reid and Horvath 1980).

Fe measurements showed a large variability in our experiment so they have to be considered with caution, but it is clear that Fe content of plant tissues exceeded the recommendation in all treatments and species, with highest values for those grown on C. It is very common for pastures and meadows plants to exceed the iron supply recommendations for cattle and values up to 500 mg kg^{-1} are usually tolerated (Flachowsky et al. 2001). This threshold is only exceeded in the case of *F. nigrescens* plants grown on C, showing that charcoal application has been beneficial.

Na⁺ content was higher than recommended for *T. pratense*, especially when grown on soils amended with charcoal, but these concentrations are not considered harmful for cattle. The concentrations we found were lower than 2.5 g kg^{-1} Dry Matter (DM) (Tab. 3), while the maximum level of sodium tolerated in forage for dairy cattle is $15.73 \text{ g Na kg}^{-1}$ DM (Johansson 2008)

Cu was higher than recommended only for *T. pratense* plants but with no difference among the treatments. Chronic copper poisoning is possible in cattle with a dietary concentration of 40 mg kg^{-1} (National Research Council 2001), but this threshold was far from being reached in our samples with values around 17 mg kg^{-1} (Tab. 3).

Charcoal application to the alpine grasslands in this study, both in the short and in the long term (CC and H), overall improved fodder quality in terms of P, K⁺ and Fe content compared to the control soils. Ca²⁺, Mg²⁺, Na⁺ and Cu contents were found to be higher than recommendations in all treatments, including control, but they did not reach harmful levels, thus guaranteeing cattle health.

The positive results observed for charcoal amended soils in terms of germination, plant growth, root:shoot ratio and fodder quality, both directly after application and after ageing, support the idea that larch wood charcoal is a soil amendment suitable for alpine pastures and meadows even at an application rate of 390 t ha^{-1} . From a management point of view, biochar incorporation into soil implies ploughing, a technique which is very rarely used in alpine management because of multiple factors: the soil layer is thin to bedrock, with very irregular profiles (Stanchi et al. 2012); ploughing will increase soil erosion which is already a main problem in mountain soils because of slope, soil depth, climate and soil low resilience which makes them almost no renewable (Tasser et al. 2003; Stanchi et al. 2012); most of the grasslands are located in areas very difficult to reach with machines that is needed for biochar spreading and incorporation into soil. Thus, biochar cannot be used for standard management, but can be helpful in the framework of grassland restoration and rehabilitation, as defined by Aronson et al. (1993). The restoration and rehabilitation of abandoned pastures and meadows in the Alps is usually based on the manual removal of trees and invasive species,

450 sometimes applying herbicides locally, and very rarely destroying the soil layer because of the reasons listed above
451 (Belleri 2014). For these reasons, biochar application is worth considering in cases of severely damaged grasslands (ski
452 tracks openings) or where the modifications to the environment have been so profound that the reconstitution of the
453 former ecosystem is no longer possible (roads or dumps construction) (Muller et al. 1998). However, possible impacts
454 on plant biodiversity remain to be explored and a deeper examination of nutrient cycles, microbial biodiversity and the
455 role played by hormones should be a research priority for next experiments in this unique environment to better
456 understand and quantify the overall impact of biochar application on nature conservation and the important ecosystem
457 services that mountain grasslands provide.

458 TABLES
459

Element	Control soil	Charcoal hearth soil	Fresh char	Control soil + fresh char
Ca	993±233* (278±61)*	3301±321* (1006±274)*	5909±86	2870±150
K	1604±201* (147±52)	2464±120* (279±208)	2906±54	2100±130
Mg	2739±128 (80±21)*	2379±665 (245±7)*	1522±56	2274±82
Na	298±88* (34±2)	93±15* (33±45)	210±4	264±54
P	321±13* (7±3)*	921±357* (12±7)*	308±1	316±8
N	4307±1402	5189±1756	2139±340	3480±870

460
461 **Tab. 1** Total nutrient content (mg kg⁻¹) in control soil, charcoal hearth soil, fresh char and control soil + fresh char at the
462 beginning of the experiment. In parenthesis, data on available nutrient concentration (mg kg⁻¹) for control and charcoal
463 hearths soils are reported. Data for hearths, control soils and fresh charcoal are taken from Criscuoli et al. (2014), while
464 data for the control soil + fresh charcoal are calculated. Asterisks indicate significant differences between control and
465 charcoal hearth soils at p≤0.05.
466

plant species	time	C	CC	H
<i>T. pratense</i>	96 hours	2a	7a	2a
	120 hours	18ab	29a	16b
	10 days	83a	94b	93b
<i>F. nigrescens</i>	96 hours	6a	13a	6a
	120 hours	11a	19a	12a
	10 days	78a	90b	89b

Tab. 2 Germinated seeds (%) of *T. pratense* and *F. nigrescens* after 96 and 120 hours and 10 days from sowing in control (C), control + fresh charcoal (CC) and charcoal hearth (H) soils. Different letters indicate significant differences among soil treatments within each plant species and time frame at $p \leq 0.10$.

<i>Trifolium pratense</i>			
Element (mg g ⁻¹)	C	CC	H
Cu	0.017 ± 0.002a	0.016 ± 0.001ab	0.017 ± 0.001b
Fe	0.4 ± 0.3a	0.3 ± 0.3a	0.2 ± 0.3a
Ca ²⁺	23.9 ± 3.3ab	26.7 ± 1.8a	28.6 ± 2.1b
K ⁺	8.4 ± 0.5ab	10.1 ± 1.9a	13.6 ± 3.5a
Mg ²⁺	5.8 ± 0.7a	4.8 ± 0.3b	6.6 ± 0.7b
Na ⁺	1.6 ± 0.2ab	0.2 ± 0.3a	2.5 ± 0.7b
P	0.9 ± 0.1ab	0.9 ± 0.3a	1.4 ± 0.2b
N	19.6 ± 0.3a	19.2 ± 3.7a	15.1 ± 2a

<i>Festuca nigrescens</i>			
Element (mg g ⁻¹)	C	CC	H
Cu	0.008 ± 0.0002a	0.007 ± 0.0a	0.008 ± 0.002a
Fe	0.5 ± 0.6a	0.2 ± 0.0a	0.2 ± 0.1a
Ca ²⁺	12.8 ± 0.3a	13.2 ± 0.1a	15.2 ± 0.1a
K ⁺	7.6 ± 2.5a	10.6 ± 1.6a	10.1 ± 2.5a
Mg ²⁺	2.6 ± 0.2a	2.7 ± 0.2a	2.9 ± 0.4a
Na ⁺	1.2 ± 0.2a	1.2 ± 0.05a	1.3 ± 0.2a
P	1.1 ± 0.3a	0.8 ± 0.2a	1.5 ± 0.5a
N	18.5 ± 0.9a	14.2 ± 2.5b	14.1 ± 2.1b

Tab. 3 Nutrient content (mg g⁻¹) in *T. pratense* (top) and *F. nigrescens* plants (bottom) at the end of the second growth cycle in control (C), control + fresh charcoal (CC) and charcoal hearth (H) soils. Different letters indicate significant differences among soil treatments at p≤0.10.

FIGURES

Fig. 1 Aboveground biomass (g pot^{-1}) at the end of the three growth cycles for *T. pratense* (plot a) and *F. nigrescens* (plot b) in control (C; light grey), control + fresh charcoal (CC; grey), and charcoal hearths (H; black) soils. Different letters indicate significant differences among soil treatments within each plant species.

Fig. 2 Total belowground (black) and aboveground biomass (grey) per pot accumulated over the three growth cycles for *T. pratense* and *F. nigrescens* plants grown on control (C), control + fresh charcoal (CC) and charcoal hearth (H) soils. Different letters indicate significant differences among soil treatments within each plant species.

Fig. 3 Root:shoot ratio of total biomass accumulated over the three growth cycles for *T. pratense* and *F. nigrescens* grown on C (light grey), CC (grey) and H (black) soils. Different letters indicate significant differences among soil treatments within each plant species.

Fig. 4 N:P ratio of soils (plot a) and of *T. pratense* and *F. nigrescens* (plot b) grown in C (light grey), CC (grey) and H (black) soils. The two lines correspond to N:P ratios of 14 (dashed) and 16 (continuous). Different letters indicate significant differences among soil treatments.

Fig. 5 Relationship between plant and soil P, K^+ , Ca^{2+} , Na^+ , Mg^{2+} content (mg g^{-1}) (plot a, b, c, d and e, respectively) in *T. pratense* (circles) and *F. nigrescens* (triangles) grown in C (white), CC (grey) and H (black) soils.

Fig. 6 ^{13}C discrimination (Δ) of *T. pratense* and *F. nigrescens* plants after the first (a), second (b) and third growth cycle (c) in C (light grey), CC (grey) and H (black) soils. Different letters indicate significant differences among soil treatments within each plant species.

Fig. 7 Isotopic signature ($\delta^{15}\text{N}$) of *F. nigrescens* and *T. pratense* plants after the first (a), second (b) and third growth cycle (c) in C (light grey), CC (grey) and H (black) soils. P-values of the individual comparisons are reported in the two tables. Different letters indicate significant differences among soil treatments within each plant species.

505 **Online Resource**

506

507 **Tab. 1** Recommended nutrients supply of a dairy cow with a mean performance of 30 kg milk day⁻¹ and a daily intake
508 of 20 kg biomass dry matter (Flachowsky et al. 2001).

509 **Funding:** This study was financed by: EuroChar project (FP7-ENV-478 2010ID-265179) funded by the European
510 Commission and AgroPyroGas funded by Regione Toscana, Italy. G.A. was partially supported by a German Academic
511 Exchange Service (DAAD, Germany) scholarship for a three-month research period at University of Freiburg,
512 Germany. I.C. was supported by a Ph.D. grant from Fondazione Edmund Mach, Trento. The funders had no role in
513 study design, data collection and analysis, decision to publish, or preparation of the manuscript.

514
515 **Conflict of Interest:** The authors declare that they have no conflict of interest.

516

517 **References**

518

- 519 Agren GI, Franklin O (2003) Root : Shoot Ratios , Optimization and Nitrogen Productivity. *Ann Bot* 92:795–800. doi:
520 10.1093/aob/mcg203
- 521 Aronson J, Floret C, LeFloc’h E, et al. (1993) Restoration and Rehabilitation of Degraded Ecosystems in Arid and
522 Semi-Arid Lands . I . A View from the South. *Restor Ecol* 8–17.
- 523 Backmeroff C (2013) Precisely dating iron-ore mining and its effects on an alpine valley: Summary of a
524 dendrochronological investigation of charcoal hearths and relict woodland stands. In: Silvertant J ed. (ed) *Min.*
525 *Cult. Landsc. – 8th Int. Symp. Archaeol. Min. Hist.* pp 218–251
- 526 Bargmann I, Rillig MC, Buss W, et al. (2013) Hydrochar and Biochar Effects on Germination of Spring Barley. *J*
527 *Agron Crop Sci* 199:360–373. doi: 10.1111/jac.12024
- 528 Belleri C (2014) L’Oasi di Protezione faunistica del Baremone: analisi e sviluppi.
- 529 Bergametti G, Remoudaki E, Losno R, et al. (1992) Source, transport and deposition of atmospheric phosphorus over
530 the northwestern Mediterranean. *J Atmos Chem* 14:501–513.
- 531 Biederman L a., Harpole WS (2013) Biochar and its effects on plant productivity and nutrient cycling: a meta-analysis.
532 *GCB Bioenergy* 5:202–214. doi: 10.1111/gcbb.12037
- 533 Bovolenta S, Dovier S, Parente G (2011) Dairy production systems in the Italian alpine area. *Proceeding 16 th Meet.*
534 *FAO CIHEAM Mt. Pastures Netw. Kraków, POLAND*, pp 137–140
- 535 Brugnoli E, Hubick KT, von Caemmerer S, et al. (1989) Correlation between the Carbon Isotope Discrimination in Leaf
536 Starch and Sugars of C3 Plants and the Ratio of Intercellular and Atmospheric Partial Pressures of Carbon
537 Dioxide. *Plant Physiol* 88:1418–1424. doi: 10.1104/pp.88.4.1418
- 538 Case SDC, McNamara NP, Reay DS, Whitaker J (2012) The effect of biochar addition on N2O and CO2 emissions
539 from a sandy loam soil – The role of soil aeration. *Soil Biol Biochem* 51:125–134. doi:
540 10.1016/j.soilbio.2012.03.017
- 541 Chen W, Cui P, Sun H, et al. (2009) Comparative effects of salt and alkali stresses on organic acid accumulation and
542 ionic balance of seabuckthorn (*Hippophae rhamnoides* L .). *Ind Crops Prod* 30:351–358. doi:
543 10.1016/j.indcrop.2009.06.007
- 544 Ciais P, Soussana JF, Vuichard N, et al. (2010) The greenhouse gas balance of European grasslands. *Biogeosciences*
545 *Discuss* 7:5997–6050. doi: 10.5194/bgd-7-5997-2010
- 546 Criscuoli I, Alberti G, Baronti S, et al. (2014) Carbon sequestration and fertility after centennial time scale
547 incorporation of charcoal into soil. *PLoS One* 9:e91114. doi: 10.1371/journal.pone.0091114
- 548 Dullinger S, Dirnbock T, Greimler J, Grabherr G (2003) A resampling approach for evaluating effects of pasture
549 abandonment on subalpine plant species diversity. *J Veg Sci* 14:243–252.
- 550 Fang Y, Singh B, Singh BP (2015) Effect of temperature on biochar priming effects and its stability in soils. *Soil Biol*
551 *Biochem* 80:136–145. doi: 10.1016/j.soilbio.2014.10.006
- 552 FAO (1985) Simple technologies for charcoal making. *Mechanical Wood Products Branch Forest Industries Division*
553 *FAO Forestry Department, Rome*
- 554 FAO (2002) *FAO IRRIGATION AND DRAINAGE PAPER 61 Agricultural drainage water management in arid and*
555 *semi-arid areas. Rome, Italy*
- 556 Farquhar G, Richards R (1984) Isotopic Composition of Plant Carbon Correlates With Water-Use Efficiency of Wheat
557 Genotypes. *Aust J Plant Physiol* 11:539. doi: 10.1071/PP9840539
- 558 Flachowsky G, Jeroch H, Kirchgessner M, et al. (2001) Empfehlungen zur Energie- und Nährstoffversorgung der
559 Milchkühe und Aufzuchttrinder. (Recommendations for the energy- and nutrient supply of dairy cows and
560 breeding cattle). *DLG-Verlag GmbH, Frankfurt am Main, Germany*
- 561 Flematti GR, Ghisalberti EL, Dixon KW, Trengove RD (2008) Germination stimulant in smoke: isolation and
562 identification. *Bioact. Nat. Prod. Detect. Isol. Struct. Elucidation*
- 563 Flematti GR, Ghisalberti EL, Dixon KW, Trengove RD (2004) A compound from smoke that promotes seed
564 germination. *Science* 305:977. doi: 10.1126/science.1099944
- 565 Flematti GR, Scaffidi A, Dixon KW, et al. (2011) Production of the seed germination stimulant karrikinolide from

566 combustion of simple carbohydrates. *J Agric Food Chem* 59:1195–8. doi: 10.1021/jf1041728

567 Fontana V, Radtke A, Bossi Fedrigotti V, et al. (2013) Comparing land-use alternatives: Using the ecosystem services
568 concept to define a multi-criteria decision analysis. *Ecol Econ* 93:128–136. doi: 10.1016/j.ecolecon.2013.05.007

569 Gamper SM, Tasser E, Tappeiner U (2007) Short-time effects of land-use changes on O-horizon in subalpine
570 grasslands. *Plant Soil* 299:101–115. doi: 10.1007/s11104-007-9366-6

571 Garnett E, Jonsson LM, Dighton J, Murner K (2004) Control of pitch pine seed germination and initial growth exerted
572 by leaf litters and polyphenolic compounds. *Biol Fertil Soils* 40:421–426. doi: 10.1007/s00374-004-0801-z

573 Gell K, van Groenigen J, Cayuela ML (2011) Residues of bioenergy production chains as soil amendments: immediate
574 and temporal phytotoxicity. *J Hazard Mater* 186:2017–2025. doi: 10.1016/j.jhazmat.2010.12.105

575 Genesio L, Miglietta F, Lugato E, et al. (2012) Surface albedo following biochar application in durum wheat. *Environ*
576 *Res Lett* 7:014025. doi: 10.1088/1748-9326/7/1/014025

577 Genesio L, Vaccari FP, Miglietta F (2016) Black Carbon aerosol from biochar threatens its negative emission potential.
578 *Glob Chang Biol*. doi: 10.1111/gcb.13254

579 Glaser B, Haumaier L, Guggenberger G, Zech W (2001) The “Terra Preta” phenomenon: a model for sustainable
580 agriculture in the humid tropics. *Naturwissenschaften* 88:37–41. doi: 10.1007/s001140000193

581 Graber ER, Tsechansky L, Mayzlish-Gati E, et al. (2015) A humic substances product extracted from biochar reduces
582 *Arabidopsis* root hair density and length under P-sufficient and P-starvation conditions. *Plant Soil* 395:21–30. doi:
583 10.1007/s11104-015-2524-3

584 Greenway H, Munns R (1980) MECHANISMS OF SALT TOLERANCE IN NONHALOPHYTES. *Annu Rev Plant*
585 *Physiol* 31:149–190.

586 Grigulis K, Lavorel S, Krainer U, et al. (2013) Relative contributions of plant traits and soil microbial properties to
587 mountain grassland ecosystem services. 47–57. doi: 10.1111/1365-2745.12014

588 Güereña DT, Lehmann J, Thies JE, et al. (2015) Partitioning the contributions of biochar properties to enhanced
589 biological nitrogen fixation in common bean (*Phaseolus vulgaris*). *Biol Fertil Soils* 51:479–491. doi:
590 10.1007/s00374-014-0990-z

591 Hernandez-Soriano MC, Kerré B, Goos P, et al. (2015) Long-term effect of biochar on the stabilization of recent
592 carbon : soils with historical inputs of charcoal. *Glob Chang Biol Bioenergy* 1–11. doi: 10.1111/gcbb.12250

593 Hille B (1992) *Ionic Channels of Excitable Membranes*, 2nd editio. Sinauer

594 Hille M, Den Ouden J (2005) Charcoal and activated carbon as adsorbate of phytotoxic compounds a comparative
595 study. *OIKOS* 108:202–207.

596 Hunziker M (1995) The spontaneous reforestation in abandoned agricultural lands : perception and aesthetic
597 assessment by locals and tourists. *Landsc Urban Plan* 1:399–410.

598 Jeffery S, Verheijen FG a., van der Velde M, Bastos a. C (2011) A quantitative review of the effects of biochar
599 application to soils on crop productivity using meta-analysis. *Agric Ecosyst Environ* 144:175–187. doi:
600 10.1016/j.agee.2011.08.015

601 Jenkins JR, Viger M, Arnold EC, et al. (2016) Biochar alters the soil microbiome and soil function: results of next
602 generation amplicon sequencing across Europe. *GCB Bioenergy*. doi: 10.1111/gcbb.12371

603 Johansson K (2008) *Salt to ruminants and horses*. Uppsala

604 Juknevičius S, Sabiené N (2007) The content of mineral elements in some grasses and legumes. *EKOLOGIJA* 53:44–
605 52.

606 Kessler J (2001) Mineralstoffversorgung der Milchkuh auf einen Blick rap aktuell (Mineral nutrient supply of dairy
607 cows at a glance). Villars-sur-Glane

608 Kloss S, Zehetner F, Dellantonio A, et al. (2012) Characterization of slow pyrolysis biochars: effects of feedstocks and
609 pyrolysis temperature on biochar properties. *J Environ Qual* 41:990–1000. doi: 10.2134/jeq2011.0070

610 Koerner C, Renhardt U (1987) Dry matter partitioning and root length / leaf area ratios in herbaceous perennial plants
611 with diverse altitudinal distribution. *Oecologia* 74:411–418.

612 Koerselman W, Meuleman AFM (1996) The vegetation N : P ratio : a new tool to detect the nature of nutrient
613 limitation. *J Appl Ecol* 33:1441–1450.

614 Kołtowski M, Oleszczuk P (2015) Toxicity of biochars after polycyclic aromatic hydrocarbons removal by thermal

615 treatment. *Ecol Eng* 75:79–85. doi: 10.1016/j.ecoleng.2014.11.004

616 Kopittke PM, Menzies NW (2007) A Review of the Use of the Basic Cation Saturation Ratio and the “Ideal” Soil. *Soil*
617 *Sci Soc Am J* 71:259–265. doi: 10.2136/sssaj2006.0186

618 Krahulec F, Skálová H, Herben T, et al. (2001) Vegetation changes following sheep grazing in abandoned mountain
619 meadows. *Appl Veg Sci* 4:97–102. doi: 10.1111/j.1654-109X.2001.tb00239.x

620 Kuppusamy S, Thavamani P, Megharaj M, et al. (2016) Agronomic and remedial benefits and risks of applying biochar
621 to soil: Current knowledge and future research directions. *Environ Int* 87:1–12. doi: 10.1016/j.envint.2015.10.018

622 de la Rosa JM, Knicker H (2011) Bioavailability of N released from N-rich pyrogenic organic matter: An incubation
623 study. *Soil Biol Biochem* 43:2368–2373. doi: 10.1016/j.soilbio.2011.08.008

624 Lehmann J, Gaunt J, Rondon M (2006) Bio-char Sequestration in Terrestrial Ecosystems – A Review. *Mitig Adapt*
625 *Strateg Glob Chang* 11:395–419. doi: 10.1007/s11027-005-9006-5

626 Liang B, Lehmann J, Solomon D, et al. (2006) Black Carbon Increases Cation Exchange Capacity in Soils. *Soil Sci Soc*
627 *Am J* 70:1719. doi: 10.2136/sssaj2005.0383

628 Von Liebig J (1840) *Die organische chemie in ihrer anwendung auf agrikultur und physiologie*. Friedrich Vieweg,
629 Braunschweig

630 Lim TJ, Spokas KA, Feyereisen G, Novak JM (2016) Predicting the impact of biochar additions on soil hydraulic
631 properties. *Chemosphere* 142:136–144. doi: 10.1016/j.chemosphere.2015.06.069

632 Di Lonardo S, Vaccari FP, Baronti S, et al. (2013) Biochar successfully replaces activated charcoal for in vitro culture
633 of two white poplar clones reducing ethylene concentration. *Plant Growth Regul* 69:43–50. doi: 10.1007/s10725-
634 012-9745-8

635 Makoto K, Choi D, Hashidoko Y, Koike T (2011) The growth of *Larix gmelinii* seedlings as affected by charcoal
636 produced at two different temperatures. *Biol Fertil Soils* 47:467–472. doi: 10.1007/s00374-010-0518-0

637 Mäser P, Gierth M, Schroeder JI (2002) Molecular mechanisms of potassium and sodium uptake in plants. *Plant Soil*
638 247:43–54.

639 Maurer K (2005) Natural and anthropogenic determinants of biodiversity of grasslands in the Swiss Alps. Universität
640 Basel

641 Muller S, Dutoit T, Alard D, Gréville F (1998) Restoration and Rehabilitation of Species-Rich Grassland Ecosystems
642 in France : a Review. *Restor Ecol* 6:94–101.

643 Naisse C, Girardin C, Davasse B, et al. (2014) Effect of biochar addition on C mineralisation and soil organic matter
644 priming in two subsoil horizons. *J Soils Sediments* 15:825–832. doi: 10.1007/s11368-014-1002-5

645 National Research Council (2001) *Nutrient Requirements of Dairy Cattle*, 7th revise. National Academy Press,
646 Washington, D.C.

647 Nelson DC, Flematti GR, Ghisalberti EL, et al. (2012) Regulation of seed germination and seedling growth by chemical
648 signals from burning vegetation. *Annu Rev Plant Biol* 63:107–30. doi: 10.1146/annurev-arplant-042811-105545

649 Novak J, Sigua G, Watts D, et al. (2016) Biochars impact on water infiltration and water quality through a compacted
650 subsoil layer. *Chemosphere* 142:160–167. doi: 10.1016/j.chemosphere.2015.06.038

651 Ogawa M, Okimori Y (2010) Pioneering works in biochar research , Japan. *Aust J Soil Res* 48:489–500.

652 Poeplau C, Don A (2013) Sensitivity of soil organic carbon stocks and fractions to different land-use changes across
653 Europe. *Geoderma* 192:189–201. doi: 10.1016/j.geoderma.2012.08.003

654 Poschlod P, Wallisdevries MF (2002) The historical and socioeconomic perspective of calcareous grasslands — lessons
655 from the distant and recent past. *Biol Conserv* 104:361–376.

656 Pusceddu E, Criscuoli I, Miglietta F (2013) Morphological investigation and physical characterization of ancient
657 fragments of pyrogenic carbon. *J Phys Conf Ser* 470:012003. doi: 10.1088/1742-6596/470/1/012003

658 Reid RL, Horvath DJ (1980) Soil chemistry and mineral problems in farm livestock. A review. *Anim. Feed Sci.*
659 *Technol.* 5:

660 Riedener E, Rusterholz H-P, Baur B (2013) Effects of different irrigation systems on the biodiversity of species-rich
661 hay meadows. *Agric Ecosyst Environ* 164:62–69. doi: 10.1016/j.agee.2012.09.020

662 Rombolà AG, Marisi G, Torri C, et al. (2015) Relationships between Chemical Characteristics and Phytotoxicity of
663 Biochar from Poultry Litter Pyrolysis. *J Agric Food Chem* 63:6660–6667. doi: 10.1021/acs.jafc.5b01540

664 Rondon M a., Lehmann J, Ramírez J, Hurtado M (2006) Biological nitrogen fixation by common beans (*Phaseolus*
665 *vulgaris* L.) increases with bio-char additions. *Biol Fertil Soils* 43:699–708. doi: 10.1007/s00374-006-0152-z

666 Schachtman D, Liu W (1999) Molecular pieces to the puzzle of the interaction between potassium and sodium uptake in
667 plants. *Trends Plant Sci* 4:281–287.

668 Schimmelpfennig S, Kammann C, Moser G, et al. (2015) Changes in macro- and micronutrient contents of grasses and
669 forbs following *Miscanthus x giganteus* feedstock, hydrochar and biochar application to temperate grassland.
670 *Grass Forage Sci* 70:582–599. doi: 10.1111/gfs.12158

671 Schimmelpfennig S, Müller C, Grünhage L, et al. (2014) Biochar, hydrochar and uncarbonized feedstock application to
672 permanent grassland—Effects on greenhouse gas emissions and plant growth. *Agric Ecosyst Environ* 191:39–52.
673 doi: 10.1016/j.agee.2014.03.027

674 Smith RT, Atkinson K (1975) *Techniques in pedology*. Paul Elek (Scientific Books) Ltd., London

675 Sohi S, Lopez-capel E, Krull E, Bol R (2009) Biochar , climate change and soil : A review to guide future research.

676 Solaiman ZM, Murphy D V., Abbott LK (2011) Biochars influence seed germination and early growth of seedlings.
677 *Plant Soil* 353:273–287. doi: 10.1007/s11104-011-1031-4

678 Spokas K a., Baker JM, Reicosky DC (2010) Ethylene: potential key for biochar amendment impacts. *Plant Soil*
679 333:443–452. doi: 10.1007/s11104-010-0359-5

680 Stanchi S, Freppaz M, Zanini E (2012) The influence of Alpine soil properties on shallow movement hazards,
681 investigated through factor analysis. *Nat Hazards Earth Syst Sci* 12:1845–1854. doi: 10.5194/nhess-12-1845-2012

682 Tasser E, Mader M, Tappeiner U (2003) Effects of land use in alpine grasslands on the probability of landslides. *Basic*
683 *Appl Ecol* 280:271–280.

684 Tasser E, Tappeiner U (2002) Impact of land use changes on mountain vegetation. *Appl Veg Sci* 5:173–184.

685 Tasser E, Walde J, Tappeiner U, et al. (2007) Land-use changes and natural reforestation in the Eastern Central Alps.
686 *Agric Ecosyst Environ* 118:115–129. doi: 10.1016/j.agee.2006.05.004

687 Tozer KN, Rennie GM, King WM, et al. (2013) Pasture renewal on Bay of Plenty and Waikato dairy farms : impacts on
688 pasture production and invertebrate populations post-establishment. *Proc New Zeal Grassl Assoc* 75:227–234.

689 USEPA (1996) Method 3052, Microwave assisted acid digestion of siliceous and organically based matrices. 1–20.

690 Vaccari F., Maienza A, Miglietta F, et al. (2015) Biochar stimulates plant growth but not fruit yield of processing
691 tomato in a fertile soil. *Agric Ecosyst Environ* 207:163–170. doi: 10.1016/j.agee.2015.04.015

692 Väre H, Lampinen R, Humphries C, Williams P (2003) Taxonomic Diversity of Vascular Plants in the European Alpine
693 Areas. *Alp. Biodivers. Europe*

694 Ventura M, Alberti G, Viger M, et al. (2015) Biochar stability and priming effect on SOM decomposition in two
695 European short rotation coppices. *Glob Chang Biol Bioenergy* 7:1150–1160.

696 van de Voorde TFJ, Bezemer TM, Van Groenigen JW, et al. (2014) Soil biochar amendment in a nature restoration
697 area : effects on plant productivity and community composition. *Ecol Appl* 24:1167–1177.

698 Wang J, Xiong Z, Kuzyakov Y (2015) Biochar stability in soil: meta-analysis of decomposition and priming effects.
699 *GCB Bioenergy* n/a–n/a. doi: 10.1111/gcbb.12266

700 Van Zwieten L, Rose T, Herridge D, et al. (2015) Enhanced biological N₂ fixation and yield of faba bean (*Vicia faba*
701 L.) in an acid soil following biochar addition: dissection of causal mechanisms. *Plant Soil* 395:7–20. doi:
702 10.1007/s11104-015-2427-3

703