



HAL
open science

The Uniform Rugosity Effect

Matthieu Bonnivard, Dorin Bucur

► **To cite this version:**

Matthieu Bonnivard, Dorin Bucur. The Uniform Rugosity Effect. Journal of Mathematical Fluid Mechanics, 2012, 14, pp.201 - 215. 10.1007/s00021-011-0052-3 . hal-01471639

HAL Id: hal-01471639

<https://hal.sorbonne-universite.fr/hal-01471639>

Submitted on 20 Feb 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The uniform rugosity effect

Matthieu Bonnivard

Dorin Bucur

Laboratoire de Mathématiques, CNRS UMR 5127 , Université de Savoie
Campus Scientifique, 73376 Le-Bourget-Du-Lac, France

Abstract

Relying on the effect of microscopic asperities, one can mathematically justify that viscous fluids adhere completely on the boundary of an impermeable domain. The rugosity effect accounts asymptotically for the transformation of complete slip boundary conditions on a rough surface in total adherence boundary conditions, as the amplitude of the rugosities vanishes. The decreasing rate (average velocity divided by the amplitude of the rugosities) computed on close flat layers is definitely influenced by the geometry. Recent results prove that this ratio has a uniform upper bound for certain geometries, like periodical and "almost Lipschitz" boundaries. The purpose of this paper is to prove that such a result holds for arbitrary (non-periodical) crystalline boundaries and general (non-smooth) periodical boundaries.

Keywords: rugosity effect, non-periodic boundaries, uniform decreasing of the velocity

1 Introduction

It is commonly accepted that viscous fluids adhere to rough surfaces. One mathematical explanation is based on the so called rugosity effect. For the Navier-Stokes equation, complete slip boundary conditions on a rough surface transform asymptotically in no-slip conditions as the amplitude of the rugosities vanishes, provided that the energy of the solutions is uniformly bounded and there is "enough roughness" of the oscillating boundaries. We refer the reader to the pioneering paper of Casado-Díaz, Fernández-Cara and Simon where this result is proved in the case of periodic, self-similar C^2 - boundaries [9]. In their proof, the authors prove implicitly that the rugosity effect has a uniform character, in the sense that the decreasing rate of the average velocity on a flat layer close to the boundary can be estimated uniformly.

Recent results obtained in [5, 8, 7] (see also [12, 13]) give a quite complete understanding of the rugosity effect for arbitrary boundaries. For equi-Lipschitz domains, if Ω_ε is a geometric perturbation of Ω (in the sense that the Hausdorff distance vanishes, $d_H(\Omega_\varepsilon, \Omega) \rightarrow 0$), the solutions \mathbf{u}_ε of a Stokes equation with complete slip boundary conditions in Ω_ε converge to the solution \mathbf{u} of the same Stokes equation complemented by the so called friction-driven boundary conditions (see [8]): there exists a suitable trio $\{\mu, A, V\}$ such that

- μ is a capacitary measure concentrated on $\partial\Omega$
- $\{V(x)\}_{x \in \Gamma}$ is a family of vector subspaces in \mathbb{R}^{N-1}
- A is a positive symmetric matrix function A defined on $\partial\Omega$

and formally the boundary conditions read

$$\begin{cases} \mathbf{u}(x) \in V(x) \text{ for q.e. } x \in \partial\Omega \\ [\mathbf{D}[\mathbf{u}] \cdot \mathbf{n} + \mu A \mathbf{u}] \cdot \mathbf{v} = 0 \text{ for any } \mathbf{v} \in V(x), x \in \Gamma \end{cases} \quad (1)$$

The rugosity effect holds provided that above $V(x) = \{0\}$ for a.e. $x \in \partial\Omega$, which implies that $\mathbf{u} = 0$ on $\partial\Omega$. However, assuming that the energies of the solutions $\|u_\varepsilon\|_{\mathbf{H}^1(\Omega_\varepsilon)}$ are uniformly bounded, the estimate of the average velocity decay rate is more delicate and relies on finding on an upper bound for

$$\limsup_{\varepsilon \rightarrow 0} \frac{\int_{\partial\Omega_\varepsilon} |\mathbf{u}_\varepsilon|^2 d\mathcal{H}^2}{d_H(\Omega_\varepsilon, \Omega)}. \quad (2)$$

For general situations one can not expect this number to be finite. If finite, the rugosity effect is said to be uniform (we refer to [6] for applications). All previous results in the literature providing an upper bound for (2) were given for periodic self-similar boundaries in the framework of [9], under more or less regularity assumptions. We refer to the recent paper of Březina [4], where the C^2 -regularity is weakened to *almost Lipschitz* boundaries (see the precise sense in [4]). Nevertheless, at least intuitively, less regularity of the boundary should enforce the physical rugosity effect but, of course, new technical difficulties arise.

The main purpose of the paper is to analyse general geometric perturbations of flat domains for which an upper bound can be found in (2). On the one hand, we remove any smoothness hypothesis and prove that the rugosity effect is uniform for general, continuous, periodic self-similar boundaries which are not riblets. This first result is only a technical improvement of previous results of [9, 6, 4] and involves a weak interpretation of the riblets associated to continuous boundaries (gradientless formulation). In a second step, we remove the periodicity assumption and prove that for arbitrary crystalline boundaries the rugosity effect is again uniform. This second result relies on a fine use of the Young measures to understand second order pointwise oscillations of the boundaries, i.e. the oscillations of the normal fields of *locally rescaled* domains. Uniform behaviours are in general difficult to capture for non-periodic structures. In our case, this is possible due to the very specific crystalline structure. We also refer to the paper [3] for a different result involving uniformity in a random geometric framework.

Uniform trace theorems in vector valued Sobolev spaces. The results of the paper can be interpreted as continuity estimates of trace operators in vector valued Sobolev spaces related to Poincaré type inequalities. Roughly speaking, if $\Omega \subseteq \mathbb{R}^N$ is a bounded connected Lipschitz set and $\Gamma \subseteq \partial\Omega$ is a subset of positive capacity, then there exists a constant $C > 0$ such that

$$\forall u \in H^1(\Omega), u = 0 \text{ q.e. on } \Gamma \quad \int_{\partial\Omega} u^2 d\mathcal{H}^{N-1} \leq C \int_{\Omega} |\nabla u|^2 dx. \quad (3)$$

If the set Γ is empty (or of zero capacity), the inequality is false since non zero constant functions become admissible. The constant C depends precisely on the Lipschitz character of Ω and on the size of Γ in terms of capacity, but not on the geometry of Γ . For example, if $N = 2$, for all connected Lipschitz domains satisfying a uniform cone condition and having a bounded diameter and for all connected sets Γ having a prescribed length, the constant C is uniform.

For vector valued Sobolev spaces, a trace continuity result can be obtained replacing the zero condition by the non penetration condition on Γ , provided that the set Γ is "curved" enough. Precisely, if Γ satisfies suitable *geometric* conditions, then there exists a constant $C > 0$ such that

$$\forall \mathbf{u} \in \mathbf{H}^1(\Omega), \mathbf{u} \cdot \mathbf{n} = 0 \text{ q.e. on } \Gamma \quad \int_{\partial\Omega} |\mathbf{u}|^2 d\mathcal{H}^{N-1} \leq C \int_{\Omega} |\nabla \mathbf{u}|^2 dx. \quad (4)$$

Requiring $\mathbf{u} \cdot \mathbf{n} = 0$ on Γ is much less than requiring $\mathbf{u} = 0$ on Γ so that inequality (3) cannot be applied. The curved character of Γ is crucial. For example, if Γ is flat inequality (4) fails, while (3) continue to hold. Roughly speaking, if Γ contains N capacity points such that their corresponding normals $\mathbf{n}_1, \dots, \mathbf{n}_N$ are independent, then (4) holds.

A rephrase of the purpose of the paper is to analyse the behaviour of the constant C and to give estimates of this constant with respect to the "oscillations" of Γ , in some particular situations coming from fluid dynamics. Understanding the influence of the geometry of Γ requires to zoom in the neighbourhood of every point of Γ and to handle the local amount of oscillations. In the previous works [5, 7] it was established that the constant C vanishes as soon as Γ is oscillating in a non degenerate sense. The results of this paper give estimates of the constant C with respect to the amplitude of the oscillations and prove that the constants are vanishing uniformly with respect to the amplitude in both the crystalline and periodical frameworks.

One of the main applications of this kind of inequalities is the study of the Navier- Stokes equations with Navier slip boundary conditions. The uniform estimates of the constant C in (4) are one of the key points for the extraction of suitable wall laws in [10] or in the study the incompressible limits of the Navier-Stokes-Fourier system on domains with rough boundaries [6].

2 General settings and main results

Throughout the paper, we fix the dimension of the space $N = 3$. For simplicity, and without losing generality, we shall assume that Ω is the cube $(0, 1)^2 \times (-1, 0)$. Let us denote $\varphi_\varepsilon : (0, 1)^2 \rightarrow [0, 1]$ a family of lower semi-continuous functions such that $(\varphi_\varepsilon)_\varepsilon$ converges uniformly to zero as $\varepsilon \rightarrow 0$. We introduce the geometric perturbations of Ω ,

$$\Omega_\varepsilon = \{(x_1, x_2, x_3) \mid (x_1, x_2) \in (0, 1)^2, -1 < x_3 < \varphi_\varepsilon(x_1, x_2)\}, \quad (5)$$

which are open sets approaching in a certain geometric sense Ω , as $\varepsilon \rightarrow 0$. In this case, it is convenient to replace in (2) the Hausdorff distance with the amplitude of the rugosities

$\|\varphi_\varepsilon\|_\infty$. Note that discontinuity points on φ_ε are *a priori* admissible and correspond to severe "roughness points" of the boundary. Define $D = (0, 1)^2 \times (-1, 2)$ so that

$$\Omega_\varepsilon \subset D \quad \forall \varepsilon > 0.$$

The non penetration condition. We denote $\mathbf{H}^1(\Omega_\varepsilon)$ the usual Sobolev space $H^1(\Omega_\varepsilon, \mathbb{R}^3)$. For a function $\mathbf{v} \in \mathbf{H}^1(\Omega_\varepsilon)$, the non-penetration condition $\mathbf{v} \cdot \mathbf{n}_\varepsilon = 0$ on

$$\Gamma_\varepsilon = \{x \in \partial\Omega_\varepsilon : x_3 \geq 0\},$$

has to be understood in a weak sense, as soon as the normal vector field \mathbf{n}_ε is not properly defined on Γ_ε . Precisely, if φ_ε is Lipschitz, the trace of \mathbf{v} on Γ_ε and the normal vector field \mathbf{n}_ε are pointwise defined a.e. If φ_ε is only continuous, we say that $\mathbf{v} \in \mathbf{H}^1(\Omega_\varepsilon)$ satisfies the *non-penetration condition* provided that

$$\forall \psi \in C_c^1(D) \quad \int_{\Omega_\varepsilon} [(\operatorname{div} \mathbf{v})\psi + \mathbf{v} \cdot \nabla \psi] dx = 0. \quad (6)$$

If φ_ε is lower semi-continuous, the above form of the non-penetration condition is incomplete, and special attention has to be given to discontinuity points. The following form of the non-penetration condition turns out to be equivalent to (6) as soon as the boundary is represented by a continuous graph, but captures also information at discontinuity points:

$$\forall \psi \in H^1(\Omega_\varepsilon) \quad \int_{\Omega_\varepsilon} [(\operatorname{div} \mathbf{v})\psi + \mathbf{v} \cdot \nabla \psi] dx = 0. \quad (7)$$

Rugosity effect and uniform decay rates. The rugosity effect reads: let $\mathbf{v}_\varepsilon \in \mathbf{H}^1(\Omega_\varepsilon)$ satisfy the non-penetration condition on Γ_ε , such that $(1_{\Omega_\varepsilon} \mathbf{v}_\varepsilon, 1_{\Omega_\varepsilon} \cdot \nabla \mathbf{v}_\varepsilon)$ converge weakly in $L^2(D, \mathbb{R}^{12})$ to $(1_\Omega \mathbf{v}, 1_\Omega \cdot \nabla \mathbf{v})$. Then $\mathbf{v} = 0$ on $\Gamma = (0, 1)^2 \times \{0\}$.

The rugosity effect above is said to be *uniform* if the following estimate holds:

$$\exists C > 0, \exists \varepsilon_0 > 0, \forall \varepsilon_0 > \varepsilon > 0, \forall \mathbf{v} \in \mathbf{H}^1(\Omega_\varepsilon), \mathbf{v} \cdot \mathbf{n}_\varepsilon = 0 \text{ on } \Gamma_\varepsilon \quad (8)$$

$$\int_\Gamma |\mathbf{v}|^2 d\mathcal{H}^2 \leq C \|\varphi_\varepsilon\|_\infty \int_{\Omega_\varepsilon} |\nabla \mathbf{v}|^2 dx. \quad (9)$$

Notice that in this inequality, one does not require \mathbf{v} to vanish on some part of the boundary. As a consequence of the continuity of the trace operator, estimate (9) proves that the rugosity effect holds, when $\varepsilon \rightarrow 0$.

Provided that φ_ε are equi-Lipschitz, one can replace in (8)-(9) the sum on Γ by the sum on Γ_ε

$$\int_{\Gamma_\varepsilon} |\mathbf{v}|^2 d\mathcal{H}^2 \leq C' \|\varphi_\varepsilon\|_\infty \int_{\Omega_\varepsilon} |\nabla \mathbf{v}|^2 dx, \quad (10)$$

which provides an upper bound in (2). If φ_ε were not smooth, the trace on Γ_ε is not defined. In this case, one can expect only estimates of type (9).

Main results of the paper. We first consider periodic structures generated by

$$\varphi_\varepsilon(x_1, x_2) = \varepsilon\varphi\left(\frac{x_1}{\varepsilon}, \frac{x_2}{\varepsilon}\right), \quad (11)$$

where φ is a continuous function on the two-dimensional torus in $\mathcal{T}^2 = [0, 1]^2|_{\{0,1\}}$ extended by periodicity in \mathbb{R}^2 . We give a characterization of the continuous functions φ such that (9) holds, removing the "almost Lipschitz" hypothesis considered in [4]. Loosely speaking, non-smooth boundaries should provide a stronger rugosity effect than the smooth ones! For technical reasons we restrict our study only to continuous boundaries but in the final section we shall discuss briefly boundaries with singularities generated by lower-semicontinuous functions φ .

We say φ is a riblet if there exists $(c_1, c_2) \in \mathbb{R}^2 \setminus \{0\}$ such that for every $(x_1, x_2) \in \mathbb{R}^2$ and every $h \in \mathbb{R}$,

$$\varphi(x_1 + hc_1, x_2 + hc_2) = \varphi(x_1, x_2). \quad (12)$$

If φ were differentiable, this would correspond to $\nabla\varphi \cdot (c_1, c_2) = 0$.

Theorem 2.1 (Characterization of the uniform rugosity effect) *Assume that φ is a continuous strictly positive function on \mathcal{T}^2 . The following statements are equivalent:*

- (i) *there exist $k > 0$ and $\varepsilon_0 > 0$ such that for every $\varepsilon_0 > \varepsilon > 0$ and every $\mathbf{v} \in \mathbf{H}^1(\Omega_\varepsilon)$ satisfying the non-penetration condition on Γ_ε (in its weak integral form), the following inequality holds:*

$$\int_{\Gamma} |\mathbf{v}|^2 d\mathcal{H}^2 \leq \varepsilon k \int_{\Omega_\varepsilon} |\nabla\mathbf{v}|^2 dx, \quad (13)$$

- (ii) *φ is not a riblet.*

In the case of lower-semicontinuous boundaries the characterization Theorem 2.1 can be rephrased, but provides a less clear geometric criterion for the uniform rugosity effect.

The second result of the paper is concerned with arbitrary crystalline boundaries (the periodicity assumption is removed). We prove that estimate (9) holds under a mild non-degeneracy assumption, similar to the one introduced in [5]. Let us consider a finite set $K \subset \mathbb{R}^2$ which satisfies the non-degeneracy assumption

$$\forall y_1, y_2 \in K, \quad 0 \notin [y_1, y_2]. \quad (14)$$

This condition simply avoids the creation of riblets on a crystalline structure. A function $\varphi \in W^{1,\infty}((0, 1)^2)$ is admissible provided that

$$\text{for a.e. } y \in (0, 1)^2 \quad \varphi(y) \in [0, 1] \text{ and } \nabla\varphi(y) \in K.$$

For every admissible function φ we define the crystalline boundary

$$\Gamma_\varphi = \{(x_1, x_2, \varphi(x_1, x_2)) \mid (x_1, x_2) \in (0, 1)^2\},$$

and the corresponding domain

$$\Omega_\varphi = \{(x_1, x_2, x_3) \mid (x_1, x_2) \in (0, 1)^2, \quad -1 < x_3 < \varphi(x_1, x_2)\}.$$

Theorem 2.2 (Crystalline boundaries) *There exist $k > 0$ and $\varepsilon_0 > 0$ such that inequality*

$$\int_{\Gamma} |\mathbf{v}|^2 d\mathcal{H}^2 + \int_{\Gamma_\varphi} |\mathbf{v}|^2 d\mathcal{H}^2 \leq k \|\varphi\|_\infty \int_{\Omega_\varphi} |\nabla \mathbf{v}|^2 dx \quad (15)$$

holds for every admissible function φ verifying $\varepsilon_0 > \|\varphi\|_\infty > 0$, and every $\mathbf{v} \in \mathbf{H}^1(\Omega_\varphi)$ satisfying the non-penetration condition on Γ_φ .

From a technical point of view, the uniformity of the rugosity effect is related to the analysis of the spectral abscissa of an elliptic operator in a family of infinite domains with "wildly" moving boundaries. The rugosity (Young) measures introduced in [5] provide a useful tool for dealing with the crystalline case. For our purpose, their use has to be refined in order to understand the local oscillations of the rescaled boundaries.

All results of the paper are given in \mathbf{H}^1 but they can be extended easily to $\mathbf{W}^{1,p}$ spaces. For instance, inequality (15) becomes

$$\|\mathbf{v}\|_{L^q(\Gamma)} + \|\mathbf{v}\|_{L^q(\Gamma_\varphi)} \leq k \|\varphi\|_\infty^\alpha \|\nabla \mathbf{v}\|_{L^p(\Omega_\varphi)}, \quad (16)$$

which holds for every admissible function $\varphi \in W^{1,\infty}((0,1)^2)$ such that $\varepsilon_0 > \|\varphi\|_\infty > 0$ and every $\mathbf{v} \in W^{1,p}(\Omega_\varphi, \mathbb{R}^3)$ satisfying the non-penetration condition on Γ_φ , where

$$1 < p < 3, \quad 1 < q < \frac{2}{3-p}, \quad \alpha = 1 - \frac{3}{p} + \frac{2}{q}.$$

3 Pointwise roughness and Young measures

A fine tool allowing to understand the rugosity effect is given by the general theory of Young measures (see for instance [11, 14]). For the sake of clarity, we recall the fundamental theorem of Young measures.

Theorem 3.1 (Fundamental theorem of Young measures) *Let $U \subset \mathbb{R}^n$ be an open set and $K \subset \mathbb{R}^m$ be a compact set. Consider a sequence $(f_k)_{k \in \mathbb{N}} \subset L^\infty(U, K)$. There exists a subsequence (f_{k_j}) and for a.e. $y \in U$ a Borel probability measure \mathcal{R}_y on \mathbb{R}^m such that for each $F \in C(\mathbb{R}^m)$ we have*

$$F(f_{k_j}) \rightharpoonup \overline{F} \quad \text{weakly-} * L^\infty(U),$$

where

$$\overline{F}(y) = \int_{\mathbb{R}^m} F(Z) d\mathcal{R}_y(Z) \quad \text{a.e. } y \in U.$$

We call $\{\mathcal{R}_y\}_{y \in U}$ a family of Young measures associated with the subsequence (f_{k_j}) .

Remark 3.2 We recall that the family of Young measures $\{\mathcal{R}_y\}_{y \in U}$ may be not unique for a given sequence $(f_k)_{k \in \mathbb{N}}$ and satisfies the following property:

$$\text{spt}(\mathcal{R}_y) \subset K \quad \text{a.e. } y \in U.$$

In particular, for the crystalline framework when K is finite, the support of the measures \mathcal{R}_y is discrete.

Remark 3.3 Let $(\varphi_k)_k$ be a bounded sequence in $W^{1,\infty}((0,1)^2, [0,1])$. We assume, up to renaming the indices, that there exists $\varphi \in W^{1,\infty}((0,1)^2, [0,1])$ such that

$$\varphi_k \rightarrow \varphi \quad \text{uniformly on } (0,1)^2, \quad \nabla\varphi_k \rightharpoonup \nabla\varphi \quad \text{weakly-}^* \text{ } L^\infty((0,1)^2).$$

We denote by $\{\mathcal{R}_y\}_{y \in (0,1)^2}$ a family of Young measures associated with a subsequence of $(\nabla\varphi_k)_{k \in \mathbb{N}}$. Then

$$\int_{\mathbb{R}^2} Z d\mathcal{R}_y(Z) = \nabla\varphi(y) \quad \text{a.e. } y \in (0,1)^2.$$

This is a direct consequence of the fundamental theorem of Young measures, taking for F the identity function.

Lemma 3.4 *With the notations of Remark 3.3, let us denote*

$$\Omega_k = \{(x_1, x_2, x_3) \mid (x_1, x_2) \in (0,1)^2, -1 < x_3 < \varphi_k(x_1, x_2)\},$$

and consider $\mathbf{u}_k \in \mathbf{H}^1(\Omega_k)$ satisfying the non-penetration condition

$$\mathbf{u}_k \cdot \mathbf{n}_k = 0 \quad \text{on } \Gamma_k = \{x \in \partial\Omega_k : x_3 \geq 0\}$$

converging weakly in $\mathbf{H}^1(D)$ to $\mathbf{c} = (c_1, c_2, c_3) \in \mathbb{R}^3$.

Then for every $g \in C(\mathbb{R}^2, \mathbb{R})$ we have

$$(c_1, c_2) \cdot \int_{\mathbb{R}^2} g(Z) Z d\mathcal{R}_y(Z) = c_3 \int_{\mathbb{R}^2} g(Z) d\mathcal{R}_y(Z) \quad \text{a.e. } y \in (0,1)^2.$$

Proof Let $\xi \in C(\mathbb{R}^2, \mathbb{R})$. We prove that for every $\psi \in C_c^\infty((0,1)^2, \mathbb{R})$,

$$\int_{(0,1)^2} \psi(y) \left[(c_1, c_2) \cdot \int_{\mathbb{R}^2} \sqrt{1+Z^2} \xi(Z) Z d\mathcal{R}_y(Z) - c_3 \int_{\mathbb{R}^2} \sqrt{1+Z^2} \xi(Z) d\mathcal{R}_y(Z) \right] dy = 0. \quad (17)$$

Taking into account that $\mathbf{n}_k(y, \varphi_k(y))$ is co-linear with $(-\nabla\varphi_k(y), 1)$, the non penetration condition on Γ_k yields

$$\int_{\Gamma_k} \psi(y) \xi(\nabla\varphi_k(y)) (-\nabla\varphi_k(y), 1) \cdot \mathbf{u}_k(y, \varphi_k(y)) d\mathcal{H}^2 = 0.$$

Performing a change of variables from Γ_k to Γ , we get

$$\begin{aligned} & \int_{(0,1)^2} \psi(y) \xi(\nabla\varphi_k(y)) \nabla\varphi_k(y) \cdot (u_k^1(y, \varphi_k(y)), u_k^2(y, \varphi_k(y))) \sqrt{1 + \|\nabla\varphi_k(y)\|^2} dy \\ &= \int_{(0,1)^2} \psi(y) \xi(\nabla\varphi_k(y)) u_k^3(y, \varphi_k(y)) \sqrt{1 + \|\nabla\varphi_k(y)\|^2} dy \end{aligned} \quad (18)$$

Without loss of generality, we may assume that u_k belongs to $\mathbf{H}^1(D) \cap C^\infty(D, \mathbb{R}^3)$. Then integrating on vertical lines and using the Cauchy-Schwartz inequality, we get

$$\int_{(0,1)^2} |\mathbf{u}_k(y, \varphi_k(y)) - \mathbf{u}_k(y, \varphi(y))| dy \leq \|\varphi_k - \varphi\|_\infty \|\nabla\mathbf{u}_k\|_{L^2(D)} \rightarrow 0 \quad \text{as } k \rightarrow +\infty.$$

Passing to the limit in (18), we get (17). Since ψ is arbitrary, we finish the proof choosing $\xi(Z) = \frac{g(Z)}{\sqrt{1+Z^2}}$. \square

Lemma 3.5 *Let $(\varphi_k)_k$ be a bounded sequence in $W^{1,\infty}((0,1)^2)$. Let $(\mathcal{R}_y)_{y \in (0,1)^2}$ be a family of Young measures associated to a subsequence of $(\nabla\varphi_k)_k$, and $\mathbf{N} \in K$. If*

$$\liminf_{k \rightarrow +\infty} |\{y \in (0,1)^2 \mid \nabla\varphi_k(y) = \mathbf{N}\}| > 0. \quad (19)$$

then

$$|\{y \in (0,1)^2 \mid \mathbf{N} \in \text{spt}(\mathcal{R}_y)\}| > 0.$$

Proof Assume for contradiction that

$$|\{y \in (0,1)^2 \mid \mathbf{N} \in \text{spt}(\mathcal{R}_y)\}| = 0.$$

Since $N \notin \text{spt}(\mathcal{R}_y)$ for a.e. $y \in (0,1)^2$, there exists $r > 0$ such that

$$\mathcal{R}_y(B(\mathbf{N}, r)) = 0 \quad \text{for a.e. } y \in G.$$

Let $\psi \in C_c^\infty(\mathbb{R}^2)$ such that $\text{spt}(\psi) \subset B(\mathbf{N}, r)$,

$$\psi(\mathbf{N}) = 1, \quad \psi(n) = 0 \quad \forall n \in K \setminus \{\mathbf{N}\}.$$

By definition of the Young measures, up to a subsequence,

$$\psi(\nabla\varphi_k) \rightharpoonup 0 \quad \text{weakly } * \text{ in } L^\infty((0,1)^2),$$

which yields

$$\int_{(0,1)^2} \psi(\nabla\varphi_k) dx \rightarrow 0,$$

that is,

$$\lim_{k \rightarrow +\infty} |\{y \in (0,1)^2 \mid \nabla\tilde{\varphi}_\varepsilon(y) = \mathbf{N}\}| = 0$$

in contradiction with hypothesis (19). □

4 Proof Theorem 2.1

In order to prove Theorem 2.1, we start with two general results involving lower semicontinuous boundaries.

First, we recall the following Poincaré Lemma, which does not require smoothness over all the boundary. The proof of this result is standard, summing on vertical lines.

Lemma 4.1 (Poincaré Lemma for l.s.c. boundaries) *Let $\varphi : (0,1)^2 \rightarrow [0,1]$ be lower semicontinuous. Then there exists a constant $K > 0$ depending only on φ , such that*

$$\int_{\Omega_\varphi} |\mathbf{u}|^2 dx \leq K \left(\int_{\Omega_\varphi} |\nabla \mathbf{u}|^2 dx + \int_{\Gamma} |\mathbf{u}|^2 d\mathcal{H}^2 \right) \quad \forall \mathbf{u} \in \mathbf{H}^1(\Omega_\varphi) \quad (20)$$

wher Ω_φ is the open set defined by

$$\Omega_\varphi = \{(x_1, x_2, x_3) \mid (x_1, x_2) \in (0,1)^2, -1 < x_3 < \varphi(x_1, x_2)\}.$$

Lemma 4.2 Assume that $\varphi : \mathcal{T}^2 \rightarrow [0, 1]$ is a continuous function and define φ_ε by (11). The following statements are equivalent:

(i) the only constant $\mathbf{c} \in \mathbb{R}^3$ satisfying

$$\forall \psi \in C_c^1(D) \quad \int_{\Omega_1} \mathbf{c} \cdot \nabla \psi dx = 0 \quad (21)$$

is $\mathbf{c} = 0$;

(ii) there exists $k, \varepsilon_0 > 0$ such that for every $\varepsilon_0 > \varepsilon > 0$ and every $\mathbf{v} \in \mathbf{H}^1(\Omega_\varepsilon)$ satisfying the weak form of the non-penetration condition (6), the following inequality holds:

$$\int_{\Gamma} |\mathbf{v}|^2 d\mathcal{H}^2 \leq \varepsilon k \int_{\Omega_\varepsilon} |\nabla \mathbf{v}|^2 dx. \quad (22)$$

Proof (i) \Rightarrow (ii) Considering a subdivision of $(0, 1)^2$ in squares $(S_i^\varepsilon)_i$ of size $\varepsilon \times \varepsilon$ parallel to the axes, it is enough to prove the following:

$$\int_{S_i^\varepsilon \times \{0\}} |\mathbf{v}|^2 d\mathcal{H}^2 \leq \varepsilon k \int_{V_i^\varepsilon} |\nabla \mathbf{v}|^2 dx \quad (23)$$

where

$$V_i^\varepsilon = \{(x_1, x_2, x_3) \mid (x_1, x_2) \in S_i^\varepsilon, -\varepsilon < x_3 < \varphi_\varepsilon(x_1, x_2)\}.$$

Performing a translation if necessary, we may assume that the square S_i^ε coincides with $(0, \varepsilon) \times (0, \varepsilon)$. Using the scaling function $H_\varepsilon : (x_1, x_2, x_3) \in \mathbb{R}^3 \rightarrow (\frac{x_1}{\varepsilon}, \frac{x_2}{\varepsilon}, \frac{x_3}{\varepsilon}) \in \mathbb{R}^3$, (23) is equivalent to

$$\int_{\Gamma} |\mathbf{v}|^2 d\mathcal{H}^2 \leq k \int_{\Omega_1} |\nabla \mathbf{v}|^2 dx, \quad (24)$$

where $\Omega_1 = H_\varepsilon(V_i^\varepsilon)$ is not depending on ε .

Assume for contradiction that there exists a sequence $\mathbf{v}_k \in \mathbf{H}^1(\Omega_1)$ satisfying the non-penetration condition (6) and such that $\int_{\Gamma} |\mathbf{v}_k|^2 = 1$ and $\|\nabla \mathbf{v}_k\|_{L^2(\Omega_1)} \rightarrow 0$. From Lemma 4.1 we get that $\|\mathbf{v}_k\|_{\mathbf{H}^1(\Omega_1)}$ is bounded, so there exists \mathbf{v}^* in $\mathbf{H}^1(\Omega_1)$ such that up to a subsequence,

$$\mathbf{v}_k \rightharpoonup \mathbf{v}^* \quad \text{weakly in } \mathbf{H}^1(\Omega_1).$$

As a result, $\|\nabla \mathbf{v}^*\|_{L^2(\Omega_1)} = 0$ so there exists $\mathbf{c} \in \mathbb{R}^3$ such that

$$\mathbf{v}^* = \mathbf{c} \quad \text{a.e. in } \Omega_1.$$

Using the compactness of the trace operator from $H^1(\Omega_1)$ to $L^2(\Gamma)$, we get

$$\int_{\Gamma} |\mathbf{v}^*|^2 d\mathcal{H}^2 = 1$$

which implies that $\mathbf{c} \neq 0$. This contradicts hypothesis (i) passing to the limit the non-penetration condition (6) for \mathbf{v}_k .

(ii) \Rightarrow (i) Assume for contradiction that there exists $\mathbf{c} \in \mathbb{R}^3 \setminus \{0\}$ such that (21) holds. Then the function $v \in \mathbf{H}^1(\Omega_1)$ defined by $\mathbf{v} = \mathbf{c}$ satisfies the non-penetration condition in the weak form (6), but contradicts inequality (13) for every $k > 0$. □

Proof of Theorem 2.1. (i) \Rightarrow (ii) Assume for contradiction that φ is a riblet in the sense of (12). From Lemma 4.2, it is enough to show that a constant vector field of the form $\mathbf{v} = (c_1, c_2, 0)$, with $(c_1, c_2) \in \mathbb{R}^2 \setminus \{0\}$, satisfies (21) (or (6)). This would readily imply that hypothesis (i) does not hold.

Since φ is a riblet, one may consider a sequence of smooth functions θ_n approaching uniformly φ on $\mathbb{R} \times \{0\} \times \{0\}$ and construct a riblet from θ_n in the direction $(c_1, c_2, 0)$. Then, the weak form of the non-penetration condition can be written on the domain defined by the smooth riblet associated to θ_n

$$\Theta_n = \{(x_1, x_2, x_3) \mid (x_1, x_2) \in (0, 1)^2, -1 < x_3 < \theta_n(x_1, x_2)\}.$$

Since $1_{\Theta_n} \rightarrow 1_{\Omega_1}$ in $L^1(\mathbb{R}^3)$, the passage to the limit of the non-penetration condition concludes the proof.

(ii) \Rightarrow (i) We need the following result concerning the uniqueness of riblet structures associated to a non constant, periodic and continuous function.

Lemma 4.3 *Let $\varphi \in C(\mathcal{T}^2)$ be a non-constant function, extended on \mathbb{R}^2 by periodicity. Assume there exist $(c_1, c_2, c_3), (c_1^*, c_2^*, c_3)$ in $\mathbb{R}^3 \setminus \{0\}$ such that for every $(x_1, x_2) \in (0, 1)^2$ and every $h \in \mathbb{R}$,*

$$\begin{aligned} \varphi(x_1 + hc_1, x_2 + hc_2) &= \varphi(x_1, x_2) + hc_3, \\ \varphi(x_1 + hc_1^*, x_2 + hc_2^*) &= \varphi(x_1, x_2) + hc_3. \end{aligned} \quad (25)$$

Then either $c_3 = 0$ and $(c_1, c_2), (c_1^, c_2^*)$ are co-linear, or $(c_1, c_2) = (c_1^*, c_2^*)$.*

Proof Assume that $(c_1, c_2) = \alpha(c_1^*, c_2^*)$, for some $\alpha \in \mathbb{R}$. Then, relations (25) give directly $hc_3 = h\alpha c_3$ for every $h \in \mathbb{R}$. Consequently, either $c_3 = 0$, or $\alpha = 1$.

Assume now that (c_1, c_2) and (c_1^*, c_2^*) are not co-linear. We will prove that φ is constant, in contradiction with our hypothesis. Indeed, let $(x_1, x_2) \in (0, 1)^2$. For every $\alpha, \beta > 0$ we define

$$(d_1, d_2) = \alpha(c_1, c_2) + \beta(c_1^*, c_2^*). \quad (26)$$

By formula (25) we get

$$\varphi(x_1 + d_1, x_2 + d_2) = \varphi(x_1, x_2) + (\alpha + \beta)c_3. \quad (27)$$

Since φ is continuous and periodic, and since (d_1, d_2) is arbitrary, (27) implies that $c_3 = 0$, i.e. φ is constant. \square

Proof of Theorem 2.1 (continuation). Assume for contradiction that condition (i) in Theorem 2.1 does not hold.

If φ is constant then (12) holds for every $(c_1, c_2) \in \mathbb{R}^2 \setminus \{0\}$. Assume that φ is not constant. From the proof of Lemma 4.2, there exists $\mathbf{c} = (c_1, c_2, c_3) \in \mathbb{R}^3 \setminus \{0\}$ such that (21) holds. We prove that for every $(x_1, x_2) \in (0, 1)^2$ and every $h \in \mathbb{R}$ such that $(x_1 + hc_1, x_2 + hc_2) \in (0, 1)^2$,

$$\varphi(x_1 + hc_1, x_2 + hc_2) = \varphi(x_1, x_2) + hc_3. \quad (28)$$

Let ρ be the standard mollifier in \mathbb{R}^3 . For every $\varepsilon > 0$ we define ρ_ε by

$$\rho_\varepsilon(x) = \frac{1}{\varepsilon^3} \rho\left(\frac{x}{\varepsilon}\right) \quad \forall x \in \mathbb{R}^3.$$

Let $\eta \in (0, 1)$ be a given constant. We introduce the following open subset of D :

$$D_\eta = \{x \in D \mid \text{dist}(x, \partial D) > \eta\}.$$

We will prove that for every $\varepsilon \in (0, \eta)$, every $x \in D_\eta$ and every $h \in \mathbb{R}$ such that $x + h\mathbf{c} \in D_\eta$, the following equality holds:

$$\mathbf{1}_{\Omega_1} * \rho_\varepsilon(x + h\mathbf{c}) = \mathbf{1}_{\Omega_1} * \rho_\varepsilon(x). \quad (29)$$

Let $\psi \in C_c^1(D)$ such that $\text{supp}(\psi) \subset D_\eta$, and let $\varepsilon \in (0, \eta)$. Applying (21) to $\psi * \rho_\varepsilon \in C_c^1(D)$ we get

$$\int_D \mathbf{1}_{\Omega_1} \mathbf{c} \cdot \nabla(\psi * \rho_\varepsilon) dx = 0$$

which yields

$$\int_D (\mathbf{1}_{\Omega_1} * \rho_\varepsilon) \mathbf{c} \cdot \nabla \psi dx = 0.$$

Integrating by part we get

$$\int_D \frac{\partial(\mathbf{1}_{\Omega_1} * \rho_\varepsilon)}{\partial \mathbf{c}} \psi dx = 0. \quad (30)$$

Since (30) holds for every $\psi \in C_c^1(D_\eta)$ and since $\mathbf{1}_{\Omega_1} * \rho_\varepsilon \in C^\infty(D_\eta)$, we obtain

$$\frac{\partial(\mathbf{1}_{\Omega_1} * \rho_\varepsilon)}{\partial \mathbf{c}}(x) = 0 \quad \forall x \in D_\eta.$$

Consequently (29) is proved.

Let us denote Γ_1 the upper part of $\partial\Omega_1$, namely

$$\Gamma_1 = \{(x_1, x_2, x_3) \mid (x_1, x_2) \in (0, 1)^2, x_3 = \varphi(x_1, x_2)\}.$$

We need the following result.

Lemma 4.4 *Let $\phi : \mathbb{R} \rightarrow \mathbb{R}$ be a function satisfying the following property:*

$$\exists b \in \mathbb{R}, \forall x \in \mathbb{R}, \forall y \in \mathbb{R} \quad \phi(x) > y \implies \forall h \in \mathbb{R} \quad \phi(x + h) \geq y + hb. \quad (31)$$

Then $\phi(h) = \phi(0) + bh$ for every $h \in \mathbb{R}$.

Proof On the one hand, for every $\varepsilon > 0$, we have $\phi(0) > \phi(0) - \varepsilon$, so that

$$\forall h \in \mathbb{R}, \phi(h) \geq \phi(0) - \varepsilon + hb.$$

Making $\varepsilon \rightarrow 0$, we get

$$\forall h \in \mathbb{R}, \phi(h) \geq \phi(0) + hb.$$

Assume for contradiction that there exists $x_0 \in \mathbb{R}$ such that

$$\phi(x_0) > \phi(0) + x_0 b.$$

Then, using property (31), we get for a suitable $\varepsilon > 0$

$$\phi(x_0 + h) \geq \phi(0) + x_0 b + \varepsilon + hb.$$

Taking $h = -x_0$ we get a contradiction. □

We shall prove in the sequel that for every $x \in D$ and for every $h \in \mathbb{R}$ such that $x + hc \in D$, we have

$$x \in \Omega_1 \Rightarrow x + hc \in \Omega_1 \cup \Gamma_1,$$

or equivalently that property (31) is satisfied by the restriction of φ on any line parallel to (c_1, c_2) .

Let $x \in \Omega_1$ and $h \in \mathbb{R}$ such that $x + hc \in D$. There exists $\eta > 0$ such that $x, x + hc \in D_\eta$ so (29) holds for every $\varepsilon \in (0, \eta)$. Since φ is continuous, Ω_1 is open so there exists $\eta_1 > 0$ such that $B(x, \eta_1) \subset \Omega_1$. In particular, for every $\varepsilon \in (0, \min(\eta, \eta_1))$,

$$\mathbf{1}_{\Omega_1} * \rho_\varepsilon(x) = \mathbf{1}_{\Omega_1}(x) = 1.$$

To prove that $x + hc \in \Omega_1 \cup \Gamma_1$ we argue by contradiction. Suppose that $x + hc \in D \setminus (\Omega_1 \cup \Gamma_1)$. Since φ is continuous, $D \setminus (\Omega_1 \cup \Gamma_1)$ is open, so by the same argument as above we obtain that there exists $\eta_2 > 0$ such that

$$\mathbf{1}_{\Omega_1} * \rho_\varepsilon(x + hc) = 0 \quad \forall \varepsilon \in (0, \eta_2).$$

This is a contradiction with (29) and (4).

Applying Lemma 4.4 to the function

$$h \in \mathbb{R} \rightarrow \varphi(x_1 + hc_1, x_2 + hc_2)$$

we get (28).

In order to prove that $c_3 = 0$, we reproduce the previous argument for p periods of φ . Let us fix $p \in \mathbb{N}^*$. We define the following subsets of \mathbb{R}^3 :

$$\begin{aligned} \Omega'_1 &= \{(x_1, x_2, x_3) \mid (x_1, x_2) \in (0, p)^2, -1 < x_3 < \varphi(x_1, x_2)\}, \\ D' &= \{(x_1, x_2, x_3) \mid (x_1, x_2) \in (0, p)^2, -1 < x_3 < 2\}, \\ \Gamma' &= \{(x_1, x_2, x_3) \mid (x_1, x_2) \in (0, p)^2, x_3 = 0\}. \end{aligned}$$

As above, there exists $c' = (c'_1, c'_2, c'_3)$ such that

$$\int_{\Omega'_1} c' \cdot \nabla \psi dx = 0 \quad \forall \psi \in C_c^1(D') \tag{32}$$

and

$$(c'_1)^2 + (c'_2)^2 + (c'_3)^2 = 1. \tag{33}$$

We first prove that $c' = \pm c$. By (32) we get that for every $(x_1, x_2) \in (0, p)^2$ and every $h \in \mathbb{R}$ such that $(x_1 + hc'_1, x_2 + hc'_2) \in (0, p)^2$,

$$\varphi(x_1 + hc'_1, x_2 + hc'_2) = \varphi(x_1, x_2) + hc'_3. \tag{34}$$

Since $|c'| = |c|$ it is enough to prove that c' and c are colinear. Recall the following relations:

$$\varphi(x_1 + hc_1, x_2 + hc_2) = \varphi(x_1, x_2) + hc_3, \tag{35}$$

$$\varphi(x_1 + h^*c'_1, x_2 + h^*c'_2) = \varphi(x_1, x_2) + h^*c'_3, \tag{36}$$

which hold for every $(x_1, x_2) \in (0, 1)^2$ and every $h, h^* \in \mathbb{R}$ such that $(x_1 + hc_1, x_2 + hc_2), (x_1 + h^*c'_1, x_2 + h^*c'_2) \in (0, 1)^2$.

If $c'_3 = 0$ then immediately $c_3 = 0$. Assume that $c'_3 \neq 0$. Setting $h^* = h \frac{c_3}{c'_3}$ in (36), we get that for every $(x_1, x_2) \in (0, 1)^2$ and for h small enough,

$$\varphi(x_1 + h \frac{c_3}{c'_3} c'_1, x_2 + h \frac{c_3}{c'_3} c'_2) = \varphi(x_1, x_2) + hc_3. \quad (37)$$

Lemma 4.3 and relations (35)-(37) imply that $(\frac{c_3}{c'_3} c'_1, \frac{c_3}{c'_3} c'_2)$ and (c_1, c_2) are colinear, which yields

$$c_3 (c_1 c'_2 - c'_1 c_2) = 0.$$

If $c_3 \neq 0$ by (35)-(36) we get that $(c_1, c_2) \neq 0$ and $(c'_1, c'_2) \neq 0$, so there exists $\lambda \in \mathbb{R} \setminus \{0\}$ such that

$$(c_1, c_2) = \lambda(c'_1, c'_2).$$

Denoting $h^* = \lambda h$ in (36) and using (35) we get

$$c_3 = \lambda c'_3.$$

Thus $c = \lambda c'$, with $\lambda = \pm 1$.

Finally, for every $p \in \mathbb{N}^*$ relation (34) holds with the same constant vector \mathbf{c} . Since φ is bounded, we get that $c_3 = 0$ hence φ is a riblet. \square

5 Proof of Theorem 2.2

Proof of Theorem 2.2. Since Γ_φ is a $W^{1,\infty}$ - parametrization of Γ , with bounded norm independent on φ , it is enough to prove that there exists $\varepsilon_0 > 0$ such that

$$\int_{\Gamma} |\mathbf{v}|^2 d\mathcal{H}^2 \leq k \|\varphi\|_{\infty} \int_{\Omega_\varphi} |\nabla \mathbf{v}|^2 dx \quad (38)$$

for every admissible function $\varphi \in W^{1,\infty}((0, 1)^2)$ such that $\varepsilon_0 > \|\varphi\|_{\infty} > 0$ and every $\mathbf{v} \in \mathbf{H}^1(\Omega_\varphi)$ satisfying the non-penetration condition on Γ_φ .

We fix $M > 0$. The value of M depends only on K , and will be fixed later in the proof. Let $\Gamma_M = (0, M)^2 \times \{0\}$,

$$\tilde{\varphi}(x_1, x_2) = \frac{1}{\|\varphi\|_{\infty}} \varphi(\|\varphi\|_{\infty} x_1, \|\varphi\|_{\infty} x_2) \quad \forall (x_1, x_2) \in (0, M)^2.$$

and

$$U_\varphi = \{(x_1, x_2, x_3) \mid (x_1, x_2) \in (0, M)^2, -1 < x_3 < \tilde{\varphi}(x_1, x_2)\}.$$

Using the same scaling method as in the proof of Theorem 2.1 (formally for $\varepsilon = \|\varphi\|_{\infty}$), it is enough to prove the existence of a positive constant k such that

$$\int_{\Gamma_M} |\mathbf{v}|^2 d\mathcal{H}^2 \leq k \|\nabla \mathbf{v}\|_{L^2(U_\varphi)}^2 \quad (39)$$

for every admissible function $\varphi \in W^{1,\infty}((0,1)^2)$ with $\varepsilon_0 > \|\varphi\|_\infty > 0$ and for every $\mathbf{v} \in \mathbf{H}^1(U_\varphi)$ satisfying the non-penetration condition

$$\int_{U_\varphi} [(\operatorname{div} \mathbf{v})\psi + \mathbf{v} \cdot \nabla \psi] dx = 0 \quad \forall \psi \in C_c^1((0, M)^2 \times (-1, 2)). \quad (40)$$

Assume for contradiction that there exists a sequence of admissible functions φ_k and a sequence of functions $\mathbf{v}_k \in \mathbf{H}^1(U_{\varphi_k})$ satisfying (40) such that

$$\int_{\Gamma_M} |\mathbf{v}_k|^2 d\mathcal{H}^2 = 1 \quad \text{and} \quad \|\nabla \mathbf{v}_k\|_{L^2(U_{\varphi_k})} \rightarrow 0. \quad (41)$$

Since $(\tilde{\varphi}_k)_k$ is uniformly bounded in $W^{1,\infty}((0, M)^2)$, there exists a Lipschitz-continuous function $\tilde{\varphi} \in C((0, M)^2)$ such that, up to a subsequence,

$$\tilde{\varphi}_k \rightarrow \tilde{\varphi}$$

uniformly on $(0, M)^2$ and weakly $*$ in $W^{1,\infty}((0, M)^2)$.

We now define

$$U := \{(x_1, x_2, x_3) \mid (x_1, x_2) \in (0, 1)^2, \quad -1 < x_3 < \tilde{\varphi}(x_1, x_2)\}.$$

Clearly

$$1_{U_{\varphi_k}} \rightarrow 1_U \quad \text{in } L^1(\mathbb{R}^3). \quad (42)$$

As a consequence of a lemma 4.1, there exists a constant $C > 0$ such that

$$\forall k, \quad \|\mathbf{v}_k\|_{\mathbf{H}^1(U_{\varphi_k})} \leq C.$$

Define $D_M = (0, M)^2 \times (-1, 2)$. Since the family $(U_{\varphi_k})_k$ is equi-Lipschitz, there exists a family of extension operators

$$P_k : \mathbf{H}^1(U_{\varphi_k}) \rightarrow \mathbf{H}^1(D_M)$$

and a constant $C > 0$ such that

$$\|P_k\| \leq C \quad \forall k \in \mathbb{N}.$$

As a consequence, v_k (identified to $P_k(\mathbf{v}_k)$) is uniformly bounded in $\mathbf{H}^1(D_M)$, so there exists $\mathbf{v}^* \in \mathbf{H}^1(D_M)$ such that, up to a subsequence,

$$\mathbf{v}_k \rightharpoonup \mathbf{v}^* \quad \text{weakly in } \mathbf{H}^1(D_M). \quad (43)$$

By (41), there exists $\mathbf{c} = (c_1, c_2, c_3) \in \mathbb{R}^3$ such that

$$\mathbf{v}^* = \mathbf{c} \quad \text{a.e. in } U_\varphi.$$

Using the compactness of the trace operator from $H^1(U)$ to $L^2(\Gamma_M)$, we obtain that $M^2\|\mathbf{c}\|^2 = 1$, hence $c \neq 0$.

At this point, we get from Lemma 3.4 that for every $g \in C(\mathbb{R}^2, \mathbb{R})$

$$(c_1, c_2) \cdot \int_{\mathbb{R}^2} g(Z) Z d\mathcal{R}_y(Z) = c_3 \int_{\mathbb{R}^2} g(Z) d\mathcal{R}_y(Z) \quad \text{a.e. } y \in (0, M)^2.$$

On the other hand, from Remark 3.3 we get

$$\int_{\mathbb{R}^2} Z d\mathcal{R}_y(Z) = \nabla \tilde{\varphi}(y) \quad \text{a.e. } y \in (0, M)^2,$$

while the non-penetration condition gives (in a purely geometric way, or as a consequence of Lemma 3.4 for $g \equiv 1$)

$$-(c_1, c_2) \cdot \nabla \tilde{\varphi}(y) + c_3 = 0, \quad \text{a.e. } y \in (0, M)^2. \quad (44)$$

Finally, for every $g \in C(\mathbb{R}^2, \mathbb{R})$

$$(c_1, c_2) \cdot \left[\int_{\mathbb{R}^2} g(Z) Z d\mathcal{R}_y(Z) - \int_{\mathbb{R}^2} Z d\mathcal{R}_y(Z) \int_{\mathbb{R}^2} g(Z) d\mathcal{R}_y(Z) \right] = 0, \quad \text{a.e. } y \in (0, M)^2. \quad (45)$$

If for some $y \in (0, M)^2$ the support of \mathcal{R}_y contains two independent vectors of K , say N_1 and N_2 , then we get $(c_1, c_2) = 0$. This can be proved by taking test functions g of the form

$$g(Z) = ag_1(Z) + bg_2(Z),$$

where $a, b \in \mathbb{R}$ and g_i are continuous functions equal to 1 on N_i and vanishing on $K \setminus \{N_i\}$, $i = 1, 2$. Consequently, from (44) we also have $c_3 = 0$, in contradiction with $M^2 \|c\|^2 = 1$.

We may assume that for a.e. $y \in (0, M)^2$, the support of \mathcal{R}_y contains only co-linear vectors of K . Recall, that the non degeneracy hypothesis implies that all co-linear vectors in K have the same sense. Since $\nabla \tilde{\varphi}(y) = \int_{\mathbb{R}^2} Z d\mathcal{R}_y(Z)$, there exists $\alpha_y \geq 1$ and $N_y \in K$ such that

$$\nabla \tilde{\varphi}(y) = \alpha_y N_y.$$

For simplicity we choose N_y as the shortest vector in each family of co-linear vectors in K .

If $c_3 = 0$ then from (44) all N_y coincide with some fixed vector $\mathbf{v} \in K$, otherwise (c_1, c_2) would vanish. Consequently,

$$\frac{\partial \tilde{\varphi}}{\partial \mathbf{v}}(y) \geq \|\mathbf{v}\|^2 \quad \text{a.e. } y \in (0, M)^2.$$

Since K is finite, we can fix $M > 0$ depending only on K to conclude that $\|\tilde{\varphi}\|_\infty > 1$ and get a contradiction.

Assuming now that $c_3 \neq 0$, we get from (44) that

$$-(c_1, c_2) \alpha_y N_y + 1 = 0 \quad \text{for a.e. } y \in (0, M)^2.$$

This implies that there exists a half plane containing almost every N_y . Hence, we can repeat the argument above with a suitable vector v contained in the half plane and such that $v \cdot N_y$ is bounded from below by a positive constant depending only on K . \square

6 Further remarks on singular rough boundaries.

Assume that the function $\varphi : \mathcal{T}^2 \rightarrow [0, 1]$ is only lower semicontinuous and $\varphi_\varepsilon(x_1, x_2) = \varepsilon\varphi(\frac{x_1}{\varepsilon}, \frac{x_2}{\varepsilon})$. Notice that their sub-graphs are open sets. Due to the lower semicontinuous character, singular rugosities may occur around discontinuity points. In this case, the non-penetration condition has to be weakened, in the sense of (7). Around discontinuity points of φ , testing only with $C_c^1(D)$ -functions in (6) is not enough to capture the non-penetration of a vertical obstacle.

If φ is lower semicontinuous, a suitable version of Lemma 4.2, in which the non-penetration condition (6) is replaced by (7) can be interpreted as a characterisation of the uniform rugosity effect. The main difficulty is to give a correct geometric interpretation of a riblet, similar to the one obtained in Theorem 2.1.

The following geometry, which is not a riblet in the sense of (12), does not produce a uniform rugosity effect:

$$\varphi : \mathcal{T}^2 \rightarrow \mathbb{R}, \quad \varphi(x_1, x_2) = 1 - x_1 \frac{1}{2} \delta_{\{x_2 = \frac{1}{2}\}}.$$

In fact, around the discontinuity points of φ , vertical surfaces of positive capacity have a crucial influence on the rugosity effect. This kind of situation falls out from (12), and can not be captured, from a technical point of view, by the proof of Theorem 2.1. Indeed, the convolution method used in Theorem 2.1 ignores sets of zero Lebesgue measure, even though their capacity is strictly positive.

If the uniform rugosity effect does not hold, one we can prove, using the convolution technique of Theorem 2.1, that if $x \in \Omega_1$, there exists $\varepsilon > 0$ and $\mathbf{c} = (c_1, c_2, 0) \neq 0$ such that for every $h \in \mathbb{R}$

$$|\Omega_1^c \cap B(x + h\mathbf{c}, \varepsilon)| = 0.$$

If for every $x_0 \in \mathcal{T}^2$, the value $\varphi(x_0)$ coincides with its approximate lower limit at x_0

$$\varphi(x_0) = \text{ap} \lim_{y \rightarrow x_0} \varphi(y),$$

then Lemma 4.4 can be used and φ is a ribblet. So Theorem 2.1 can be extended for lower and approximate lower semicontinuous functions. An example of such a function is

$$\varphi : \mathcal{T}^2 \rightarrow \mathbb{R}, \quad \text{generated by } \forall (x_1, x_2) \in [0, 1)^2 \quad \varphi(x_1, x_2) = x_1.$$

References

- [1] A. A. Amirat, D. Bresch, J. Lemoine, and J. Simon. Effect of rugosity on a flow governed by stationary Navier-Stokes equations. *Quart. Appl. Math.*, **59**:768–785, 2001.
- [2] A. A. Amirat, E. Climent, E. Fernández-Cara, and J. Simon. The Stokes equations with Fourier boundary conditions on a wall with asperities. *Math. Models. Methods Appl.*, **24**:255–276, 2001.

- [3] A. Basson and D. Gérard-Varet. Wall laws for fluid flows at a boundary with random roughness. *Comm. Pure Appl. Math.* 61 (2008), no. 7, 941–987.
- [4] J. Březina *Asymptotic properties of solutions to the equations of incompressible fluid mechanics* *J. Math. Fluid Mech.* (2009) online first.
- [5] D. Bucur, E. Feireisl, Š. Nečasová, and J. Wolf. On the asymptotic limit of the Navier-Stokes system on domains with rough boundaries. *J. Differential Equations*, **244**:2890–2908, 2008.
- [6] D. Bucur, E. Feireisl. The incompressible limit of the full Navier-Stokes-Fourier system on domains with rough boundaries. *Nonlinear Analysis: Real World Applications*, **10**:3203–3229, 2009.
- [7] D. Bucur, E. Feireisl, Š. Nečasová, Influence of wall roughness on the slip behaviour of viscous fluids. *Proc. Roy. Soc. Edinburgh Sect. A* **138** (2008), no. 5, 957–973.
- [8] D. Bucur, E. Feireisl, Š. Nečasová, Boundary behavior of viscous fluids: Influence of wall roughness and friction-driven boundary conditions. *Arch. Ration. Mech. Anal.* 197 (2010), 117–138.
- [9] J. Casado-Díaz, E. Fernández-Cara, and J. Simon. Why viscous fluids adhere to rugose walls: A mathematical explanation. *J. Differential Equations*, **189**:526–537, 2003.
- [10] A.L. Dalibard, and D. Gérard-Varet. Effective boundary condition at a rough surface starting from a slip condition. <http://arxiv.org/abs/1008.5030>, arXiv:1008.5030v1, 2010.
- [11] L.C. Evans, Weak convergence methods for nonlinear partial differential equations. CBMS Regional Conference Series in Mathematics, 74. American Mathematical Society, Providence, RI, 1990.
- [12] W. Jaeger and A. Mikelić. On the roughness-induced effective boundary conditions for an incompressible viscous flow. *J. Differential Equations*, **170**:96–122, 2001.
- [13] B. Mohammadi, O. Pironneau, F. Valentin, Rough boundaries and wall laws. Finite elements in fluids. *Internat. J. Numer. Methods Fluids* **27** (1998), no. 1-4, Special Issue, 169–177.
- [14] P. Pedregal, Parametrized measures and variational principles. Progress in Nonlinear Differential Equations and their Applications, 30. Birkhäuser Verlag, Basel, 1997.