

# Cysteamine re-establishes the clearance of Pseudomonas aeruginosa by macrophages bearing the cystic fibrosis-relevant F508del-CFTR mutation

Eleonora Ferrari, Romina Monzani, Valeria R. Villella, Speranza Esposito, Francesca Saluzzo, Federica Rossin, Manuela d'Eletto, Antonella Tosco,

Fabiola de Gregorio, Valentina Izzo, et al.

## ▶ To cite this version:

Eleonora Ferrari, Romina Monzani, Valeria R. Villella, Speranza Esposito, Francesca Saluzzo, et al.. Cysteamine re-establishes the clearance of Pseudomonas aeruginosa by macrophages bearing the cystic fibrosis-relevant F508del-CFTR mutation. Cell Death and Disease, 2017, 8, pp.e2544. 10.1038/cddis.2016.476 . hal-01492521

# HAL Id: hal-01492521 https://hal.sorbonne-universite.fr/hal-01492521

Submitted on 20 Mar 2017  $\,$ 

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

www.nature.com/cddis

# Cysteamine re-establishes the clearance of *Pseudomonas aeruginosa* by macrophages bearing the cystic fibrosis-relevant F508del-CFTR mutation

Eleonora Ferrari<sup>1</sup>, Romina Monzani<sup>1</sup>, Valeria R Villella<sup>1</sup>, Speranza Esposito<sup>1</sup>, Francesca Saluzzo<sup>1</sup>, Federica Rossin<sup>2</sup>, Manuela D'Eletto<sup>2</sup>, Antonella Tosco<sup>3</sup>, Fabiola De Gregorio<sup>1,3</sup>, Valentina Izzo<sup>4</sup>, Maria C Maiuri<sup>4</sup>, Guido Kroemer<sup>\*,4,5,6,7,8,9</sup>, Valeria Raia<sup>\*,3</sup> and Luigi Maiuri<sup>\*,1,10</sup>

Cystic fibrosis (CF), the most common lethal monogenic disease in Caucasians, is characterized by recurrent bacterial infections and colonization, mainly by *Pseudomonas aeruginosa*, resulting in unresolved airway inflammation. CF is caused by mutations in the gene coding for the cystic fibrosis transmembrane conductance regulator (CFTR) protein, which functions as a chloride channel in epithelial cells, macrophages, and other cell types. Impaired bacterial handling by macrophages is a feature of CF airways, although it is still debated how defective CFTR impairs bacterial killing. Recent evidence indicates that a defective autophagy in CF macrophages leads to alterations of bacterial clearance upon infection. Here we use bone marrow-derived macrophages from transgenic mice to provide the genetic proof that defective CFTR compromises both uptake and clearance of internalized *Pseudomonas aeruginosa*. We demonstrate that the proteostasis regulator cysteamine, which rescues the function of the most common F508del-CFTR mutant and hence reduces lung inflammation in CF patients, can also repair the defects of CF macrophages, thus restoring both bacterial internalization and clearance through a process that involves upregulation of the pro-autophagic protein Beclin 1 and re-establishment of the autophagic pathway. Altogether these results indicate that cysteamine restores the function of several distinct cell types, including that of macrophages, which might contribute to its beneficial effects on CF. *Cell Death and Disease* (2017) **8**, e2544; doi:10.1038/cddis.2016.476; published online 12 January 2017

Cystic fibrosis (CF), the most common lethal monogenic disease in Caucasians, is caused by mutations in the gene coding for cystic fibrosis transmembrane conductance regulator (CFTR), a 1480 amino-acid protein that functions as a chloride channel at the plasma membrane.1 Approximately 2000 mutations, most of which are disease relevant, have been identified in the CFTR gene and then categorized in six different classes according to their functional impact.<sup>2</sup> CF is a systemic disease, characterized by highly heterogeneous manifestations in distinct organs,<sup>1,3,4</sup> comprising insufficiency of the exocrine pancreas, increased electrolytes in sweat, male infertility, and, most prevalent, a chronic progressive lung disease resulting from decreased mucociliary clearance with accumulation of thick, sticky mucus, chronic inflammation, and persistent and untreatable bacterial infections and colonization, mainly by Pseudomonas aeruginosa (P. aeruginosa), Staphilococcus aureus and Burkholderia cepacia (B. cepacia).<sup>1,3,4</sup> In spite of increased survival to date, most current CF treatments are symptomatic and hence target lung inflammation and infection rather than the primary CF defect, namely the loss of CFTR function. New anti-inflammatory drugs and antibiotics are in clinical trials in CF patients.5,6

Bacterial infections and chronic lung inflammation have a pivotal role in CF lung disease and deeply influence each other, as microbial invasion leads to an uncontrolled inflammatory host-response that in turn favors unresolved bacterial infections and colonization.<sup>3,4,6</sup> Whether the lack of CFTR channel activity directly drives inflammation is still debated. Emerging evidence indicates that CFTR does not only act as a pure ion channel, but also as a hub protein that orchestrates the proteostasis network (PN) and influence multiple cellular functions.<sup>7,8</sup> The dysfunction of this hub in CF deranges major mechanisms of cell physiology, both in primary epithelial cells biopsied from F508del-CFTR homozygous patients and in mice bearing an equivalent mutation in the mouse CFTR gene (CF mice). Autophagy, a major mechanism of cytoplasmic protein turnover, is defective in CF airways, owing to tissue transglutaminase (TG2)-mediated depletion of the essential autophagy-related protein Beclin 1 (BECN1).8-12 This leads to the accumulation of the autophagic substrate SQSTM1/ p62.<sup>8–10</sup> which sequestrates misfolded ubiguitinated F508del-CFTR and major anti-inflammatory proteins, such as PPAR<sub> $\gamma$ </sub> and IK-B<sub> $\alpha$ </sub>, in intracellular aggresomes, thus sustaining a pro-inflammatory cellular environment.8-12

Received 15.8.16; revised 17.11.16; accepted 16.12.16; Edited by RA Knight

<sup>&</sup>lt;sup>1</sup>Division of Genetics and Cell Biology, San Raffaele Scientific Institute, European Institute for Research in Cystic Fibrosis, Milan 20132, Italy; <sup>2</sup>Department of Biology, University of Rome 'Tor Vergata', Rome, Italy; <sup>3</sup>Regional Cystic Fibrosis Center, Pediatric Unit, Department of Translational Medical Sciences, Federicoll University Naples 80131, Italy; <sup>4</sup>Equipe11 labellisée Ligue Nationale contrele Cancer, Centre de Recherche des Cordeliers, Paris, France; <sup>5</sup>INSERM U1138, Centre de Recherche des Cordeliers, Paris, France; <sup>6</sup>Université Paris Descartes, Paris, France; <sup>7</sup>Metabolomics and Cell Biology Platforms, Institut Gustave Roussy, Villejuif, France; <sup>8</sup>Pôlede Biologie, Hôpital Européen Georges Pompidou, AP-HP, Paris, Franceand; <sup>9</sup>Karolinska Institute, Department of Women's and Children's Health, Karolinska University Hospital, Stockholm 17176, Sweden and <sup>10</sup>SCDU of Pediatrics, Department of Health Sciences, University of Piemonte Orientale, Novara 28100, Italy

<sup>\*</sup>Corresponding author: G Kroemer or V Raia or L Maiuri, Division of Genetics and Cell Biology, San Raffaele Scientific Institute, European Institute for Research in Cystic Fibrosis, via Olgettina 58, Milan 20131, Italy. Tel: +33 1 44 27 76 67; Fax: +33 1 44 27 76 74 or Tel: +39 081 7463273; Fax: +39 081 5463933 or Tel: +39 3207981514; Fax: 39 0226434328; E-mail: kroemer@orange.fr or raia@unina.it or luigi.maiuri@gmail.com

Notably, re-establishing a proper autophagy flux by means of genetic interventions (enforced expression of BECN1 or direct depletion of SQSTM1/p62) or the inhibition of TG2 by the proteostasis regulator cysteamine, controls lung inflammation in F508del-CFTR homozygous mice.<sup>8–10</sup>

Defective bacterial handling by macrophages is a feature of CF airways.<sup>13–19</sup> CFTR is expressed by, and localizes at the plasma membrane of, human and mouse macrophages<sup>6,13–19</sup> and can also localize at the membranes of phagosomes and lysosomes.<sup>13–16</sup> Conflicting results have been reported on whether the defective bacterial killing observed in F508del-CFTR macrophages is the consequence of impaired acidification of lysosomes and/or unbalanced redox regulation due to defective channel function.<sup>17–20</sup> More recently, several authors reported that F508del-CFTR macrophages are also characterized by defective autophagy with reduced BECN1 levels and accumulation of SQSTM1/p62.<sup>21,22</sup>

Emerging therapeutic approaches aim at controlling lung inflammation and ameliorating lung function through the repair of the primary CFTR defect.<sup>6,23-25</sup> These strategies take advantage from large-scale high-throughput screening programs that led to the identification of small molecules capable of either correcting the traffic of misfolded CFTR mutants (CFTR correctors) or improving channel function (CFTR potentiators) of plasma membrane-resident CFTR mutants. A CFTR potentiator (Ivacaftor, Kalvdeco, Vertex Pharmaceutical) is available for a small fraction (5%) of CF patients bearing class III mutations (such as the G551D membrane-resident channel-dead CFTR mutant).<sup>26</sup> Moreover, for the majority (>70%) of patients bearing the most common CFTR mutation, F508del-CFTR, the FDA approved the combination of the CFTR potentiator lvacaftor with the corrector Lumacaftor (Orkambi), although this therapy is only marginally effective.27-31

Small molecules that target the deranged PN in CF (proteostasis regulators) can represent an alternative strategy to circumvent CFTR defect.7,8,32-34 We reported that a combination of two proteostasis regulators, cysteamine, an FDA-approved drug for the treatment of patients affected by cvstinosis. and the over-the-counter nutraceutical epigallocatechin-gallate (EGCG) is effective in rescuing a functional F508del-CFTR to the cell surface and in stabilizing the rescued CFTR mutant at the plasma membrane of epithelial cells. Such a combinatory treatment succeeds in increasing survival and reducing lung inflammation in vivo in F508del homozygous mice.8-10,33,34 The effects of cysteamine are mediated by its ability to restore BECN1 expression and autophagy,<sup>8-10</sup> whereas EGCG inhibits the master kinase CK2, responsible for CFTR fragmentation and degradation.33,35 These pre-clinical results paved the way for 2 clinical trials in CF patients bearing F508del-CFTR, in whom the combination treatment restored autophagy, decreased sweat chloride concentrations and increased chloride efflux in brushed nasal cells.33,34 Moreover, the treatment significantly reduced the levels of inflammatory cytokines in patients' sputum coupled with an amelioration of lung function.33,34 Cysteamine is a known mucolytic agent and reduces the viscoelasticity of CF sputum.<sup>36,37</sup> It is also endowed with bactericidal effects against P. aeruginosa and other emerging pathogens and may disrupt established biofilms or prevent their formation in CF

patients.<sup>36,37</sup> This may synergize with antibiotic treatments and help avoiding bacterial resistance.<sup>33,34</sup>

Here, we addressed the question as to whether cysteamine might have a direct effect on the microbiocidal activity of macrophages, at least part of which depends on an intact autophagic machinery.<sup>38–43</sup> We show that cysteamine can restore the impaired ability of CFTR defective macrophages in killing *P. aeruginosa* through restoring BECN1 expression and autophagy.

### Results

Cysteamine reverts defective killing of *P. aeruginosa* by BMDMs from CF mice. To determine whether cysteamine may improve the killing of P. aeruginosa upon acute infection, bone marrow-derived macrophages (BMDMs) were isolated from the femurs of 6–12 week-old CFTR homozygous mice in the FVB/129 outbred background (below referred to as *Cftr<sup>F508del/F508del*</sub>, or their wild-type (WT) littermates.<sup>21,22</sup></sup> Then the cells were inoculated with the P. aeruginosa strain PAO1 at a multiplicity of infection (MOI) of 50 bacteria/ macrophage for 10 min, washed and treated for 10 min with 100  $\mu$ g/ml gentamicin to remove any extracellular bacteria and finally lysed to determine the clonogenic potential of surviving intracellular bacteria. To monitor the clearance of internalized bacteria, BMDMs were infected and treated as described, extensively washed and then kept in culture for 30 min, 1 h, 2 h, 3 h, and up to 48 h, as reported.<sup>21,22</sup> Colony-forming units (CFU) and bacterial DNA, together with the enumeration of the residual intracellular fluorescent-labeled P. aeruginosa were determined. To measure the effective intracellular P. aeruginosa clearance by BMDMs, we calculated the percentage of living bacteria after 3 h of culture with respect to the total amount of internalized bacteria. We observed that the internalization of P. aeruginosa was significantly lower in Cftr<sup>F508del/F508del</sup> BMDMs than in WT controls (CFU,  $22.5\pm7.3$  versus  $104.50\pm50$ , P<0.01 and bacterial DNA, 0.042+0.068 versus 0.23+0.028. P<0.001. respectively) (Figures 1a-c). Next, we measured the number of CFU, the amount of bacterial DNA and the number of the residual GFP-conjugated intracellular bacteria 3 h post-infection. Although the number of live bacteria was not significantly different in BMDMs isolated from Cftr<sup>F508del/F508del</sup> mice (CFU. 18.6 ± 4.5, bacterial DNA, 0.0042 ± 0.0021) versus WT (CFU,  $20.5 \pm 6.7$ , bacterial DNA,  $0.0064 \pm 0.0052$ ) mice (P>0.5) (Figures 1d-f), the effective clearance of internalized bacteria after 3 h of culture was greatly reduced in BMDMs from Cftr<sup>F508del/F508del</sup> (8.3±13.9%) respect to WT littermates  $(63.7 \pm 12.7\%, P < 0.01)$  (Figure 1g). Moreover, viable bacteria were still detected as late as after 48 h following infection (Figure 1h). Thus, BMDMs from Cftr<sup>F508del/F508del</sup> mice exhibit both defective internalization and impaired clearance of P. aeruginosa.

To investigate whether cysteamine may modulate these events in BMDMs isolated from either WT and  $Cftr^{F508del/F508del}$  mice, the cells were pre-treated for 18 h with 250  $\mu$ M cysteamine or vehicle. Cysteamine significantly increased the internalization of *P. aeruginosa* (CFU, 42.3 ± 16.3; *P*<0.001 *versus* vehicle-treated BMDMs)

Bacterial clearance by cysteamine in CF macrophages E Ferrari et al



3

**Figure 1** Internalization and clearance of *P. aeruginosa*-O1 (PAO1) in bone marrow-derived macrophages (BMDMs) from Wild-type (WT) and *Cftt*<sup>F508del/F508del</sup> mice. (**a**-**g**) WT or *Cftt*<sup>F508del/F508del</sup> BMDMs infected with PAO1 for 10 min followed by 10 min (**a**-**c**) or 3 h (**d**-**f**) of culture in the presence of gentamicin. (**a**, **d**) Number of colony-forming units (CFU) of PAO1 in WT (n=10) and *Cftt*<sup>F508del/F508del</sup> cells (n=13) BMDMs (\*\*P<0.01, Student's *t*-test) and (**b**, **e**) q-PCR transcript level of total PAO1 DNA in WT (n=3) and *Cftt*<sup>F508del/F508del</sup> (n=3) BMDMs related to proC housekeeping gene (\*\*P<0.001, Student's *t*-test). (**c**, **f**) Immunofluorescence detection of viable intracellular bacteria. (**g**) Clearance of bacteria expressed as percentage of living bacteria after 3 h of culture respect to the total amount of internalized bacteria in WT and *Cftt*<sup>F508del/F508del</sup> BMDMs infected with PAO1-GFP at different time points. Enumeration of intracellular PAO1-GFP by immufluorescence in WT and *Cftt*<sup>F508del/F508del</sup> BMDMs. Means  $\pm$  S.D. of three independent measurements for each group of experiments. Asterisks indicate significant differences

(Figures 2a and b) and improved the clearance of internalized *P. aeruginosa* in *Cftr<sup>F508del/F508del/* BMDMs up to  $45.5 \pm 10.5\%$  (*P*<0.001 as compared with vehicle-treated mice) (Figure 2c). No effects of cysteamine were observed in WT CFTR mice (Figures 2d–f).</sup>

**Cysteamine reverts defective killing of** *P. aeruginosa* by **restoring BECN1 expression and autophagy.** We previously reported that cysteamine restores autophagy in *Cttr<sup>F508del/F508del</sup>* mice by preventing tissue TG2-mediated targeting of the master autophagy regulator BECN1 and its

sequestration in intracellular aggregates.<sup>9–11</sup> Previous studies reported that the colocalization of BECN1 and *B. cepacia* increases in F508del-CFTR macrophages upon autophagy induction.<sup>21</sup> Accordingly, we found that cysteamine treatment of F508del-CFTR BMDMs increased BECN1 protein levels and reduced SQSTM1/p62 accumulation (Figures 3a and b).

To provide the genetic proof for the involvement of BECN1 in bacterial killing of *P. aeruginosa*, BMDMs were isolated from *Becn*<sup>+/-</sup> haplo-insufficient mice on a *C57BL/6 J* background and infected with *P. aeruginosa*. Macrophages from *Becn*<sup>1+/-</sup>

Bacterial clearance by cysteamine in CF macrophages E Ferrari et al



**Figure 2** Effects of cysteamine on internalization and clearance of *P. aeruginosa*-O1 (PAO1) in bone marrow-derived macrophages (BMDMs) from wild-type (WT) and *Cftr<sup>F508del/F508del* (**a**-**c**) or WT (**d**-**f**) mice treated with 250  $\mu$ M cysteamine (n = 10) or vehicle (n = 10) for 10 min before infection followed by 10 min (**a**-**b**, **d**-**e**) or 3 h (**c**, **f**) of gentamicin culture. *P. aeruginosa* internalization expressed as number of CFUs of PAO1 (**a**, **d**) or enumerated by immunofluosescence (**b**, **e**) after 10 min of gentamicin culture (\*\*\**P* < 0.001 *versus* vehicle; Student's *t*-test); (**c**, **f**) clearance of *P. aeruginosa* expressed as percentage of living bacteria after 3 h of culture respect to the total amount of internalized bacteria (\*\*\**P* < 0.001 *versus* treatment with vehicle; Student's *t*-test). Means ± S.D. of three independent measurements for each group of experiments. Asterisks indicate significant differences</sup>

haplo-insufficient mice efficiently internalized *P. aeruginosa* (CFU, 62.6  $\pm$  9.7) as compared with WT (*P*>0.5). In contrast, they manifested defective bacterial clearance with a percentage of clearance of 32+7.6% (*P*<0.05) (Figures 4a–c). This suggests a critical role of BECN1 and autophagy in the clearance of the internalized bacteria but not in the earliest steps of *P. aeruginosa* uptake. Notably, cysteamine was not capable of increasing the intracellular clearance of *P. aeruginosa* (*P*>0.5) in BMDMs from *Becn1*<sup>+/-</sup> upon infection (Figure 4c).

To unveil the role of BECN1 in a F508del-CFTR defective context,  $Cft^{F508del/+}$  female mice were backcrossed with  $Becn1^{+/-}$  mice to obtain Becn1 haplo-insufficient F508del heterozygous mice (abbreviated  $Cft^{F508del/+}/Becn1^{+/-}$ ) at the first generation. These  $Cftr^{F508del/+}/Becn1^{+/-}$  animals were crossbred to obtain Becn1 haplo-insufficient F508del

homozygous mice (abbreviated *Cftr*<sup>F508del/F508del/Becn1<sup>+/-</sup>).<sup>34</sup> In BMDMs from *Cftr*<sup>F508del/F508del/Becn1<sup>+/-</sup> mice, both *P. aeruginosa* uptake (CFU, 16.3±5.1) and clearance (CFU, 13±3.6) were greatly reduced as compared with either WT or their *Becn1<sup>+/-</sup>* and were similar to those in *Cftr*<sup>F508del/F508del</sup> mice (Figures 4d–f). Notably, cysteamine showed negligible effects on either uptake and clearance of *P. aeruginosa* by *Cftr*<sup>F508del/F508del</sup>/*Becn1<sup>+/-</sup>* BMDMs (*P*>0.5) (Figures 4d–f). These results indicate that cysteamine enhances bacterial clearance of *P. aeruginosa* through restoring BECN1 expression and autophagy.</sup></sup>

**Cysteamine reverts defective killing of** *P. aeruginosa* by **rescuing a functional CFTR protein.** Cysteamine is effective in either rescuing and stabilizing misfolded F508del-CFTR mutant at the PM.<sup>9,10</sup> Notably, cysteamine reduces

Λ

Bacterial clearance by cysteamine in CF macrophages E Ferrari et al



Figure 3 Effects of cysteamine on autophagy markers in bone marrow-derived macrophages (BMDMs). Wild-type (WT) or *Cftr<sup>F508del/F508del*</sup> BMDMs treated for 18 h with cysteamine or vehicle. Representative immunoblots with anti-BECN1 (**a**) and anti-SQSTM1/p62 (**b**) (top) and densitometric measurements (bottom) in BMDMs from one mice per group. Mean ± S.D. of three independent measurements (\*\**P* < 0.01 *versus* vehicle-treated mice; ANOVA). Asterisks indicate significant differences

lung inflammation *in vivo* through re-establishing CFTR function either in *Cftr*<sup>F508del/F508del</sup> mice or in CF patients bearing class II CFTR mutations.<sup>9,10,33,34</sup> To determine whether the effects of cysteamine on bacterial killing are secondary to its ability to rescue CFTR function, BMDMs were isolated from mice in which the gene coding for CFTR has been knocked out by homologous recombination (i.e., *Cftr*<sup>-/-</sup> mice). Both bacterial internalization (CFU, 15.5 ± 2.88) and intracellular clearance of *P. aeruginosa* (6.08 ± 4.18% of clearance) were similar to those observed in *Cftr*<sup>F508del/</sup><sup>F508del</sup> mice (*P*>0.5) (Figures 5a–c). However, the treatment with cysteamine failed to restore the uptake (CFU, 17.66 ± 4.5) or clearance (CFU, 13.33 ± 2.88) of *P. aeruginosa* by *Cftr*<sup>-/-</sup> BMDMs (Figures 5a–c), contrasting with its significant beneficial effects on BMDMs from *Cftr*<sup>F508del/F508del</sup> mice.

To provide the proof-of-concept that the beneficial effects of cysteamine on P. aeruginosa killing rely on the presence of a rescuable CFTR mutant, heterozygous Cftr<sup>F508del/+</sup>males were backcrossed with heterozygous  $Cftr^{+/-}$  females to obtain F508del/null CFTR heterozygous mice (abbreviated Cftr<sup>F508del</sup>/<sup>-</sup>). BMDMs were infected and treated with cysteamine in the presence or absence of the autophagy inhibitor 3-methyl-adenine (3-MA) before infection. BMDMs from Cftr<sup>F508del</sup>/<sup>-</sup> mice behaved similarly to Cftr<sup>F508del</sup>/F508del mice. in thus far that cysteamine was effective in restoring P. aeruginosa uptake (22.3 ± 7.02 CFU in controls: and  $41 \pm 4.35$ , after cysteamine treatment; P<0.05) or clearance  $(16.5\pm6.28$  of clearance in controls and  $38.02\pm6.45\%$  of after cysteamine treatment, P<0.05), unless 3-MA was also added to the culture system (CFU,  $25 \pm 7.2$  and  $14.05 \pm 1.85\%$ of clearance) (Figures 5d-f). These results provide the genetic proof that cysteamine restores bacterial killing through rescuing a functional F508del-CFTR and hence re-establishing BECN1-dependent autophagic flux. Accordingly, the addition of the CFTR inhibitor 172 (CFTR<sub>inh172</sub>) to the system

during cysteamine treatment, significantly reduced the beneficial effects of cysteamine on the internalization and clearance of *P. aeruginosa* in both  $Cftr^{F508del/F508del}$  and  $Cftr^{F508del/-}$  mice (Figures 5d–f).

Cysteamine decreases the production of inflammatory cytokines induced by P. aeruginosa in BMDMs. To explore the possibility that cysteamine may control the increased production of inflammatory cytokines by BMDMs from Cftr<sup>F508del/F508del</sup> mice following P. aeruginosa infection, we measured the levels of IL-1 $\beta$  and TNF- $\alpha$  in culture supernatants from BMDMs collected after 24 h following infection. Pre-treatment with cysteamine reduced the levels of pro-inflammatory cytokines as compared to untreated control cells (Figures 6a and b). These beneficial effects of cysteamine were abrogated if 3-MA (Figures 6a and b) or wortmannin (Supplementary Figure 1) were added to the culture system. No effect of cysteamine on cytokine levels was observed in BMDMs from Cftr<sup>F508del/F508del</sup>/Becn1<sup>+/-</sup> mice (Figures 6c and d). Hence, haploinsufficiency of Becn1 curtails the anti-inflammatory effects of cysteamine on Cftr<sup>F508del/F508del</sup> BMDMs.

### Discussion

Persistent infections by *P. aeruginosa* indicate dismal prognosis for CF patients.<sup>44</sup> Although chronic bacterial adaptation to the CF lung environment can favor colonization and antibiotic resistance, intrinsic host–macrophage defects compromise the capacity of CF phagocytes to properly fight bacterial challenges,<sup>13–20</sup> thus favouring chronic infection. The mechanisms underlying defective bacterial killing in CF are not completely understood. Conflicting results have been reported on mouse and human CF macrophages, which were either monocyte-derived or lung-resident,<sup>13–20</sup>

Bacterial clearance by cysteamine in CF macrophages E Ferrari et al



Figure 4 Cysteamine does not restore defective killing of *P. aeruginosa* by bone marrow-derived macrophages (BMDMs) from *Cftr<sup>F508del/F508d*</sup>

likely because the latter are conditioned by the proinflammatory and micro-anaerobiotic environment of CF airways. Bacterial killing is a complex mechanism resulting from a sequential cascade of events, which depend on bacterial strain, environmental conditions<sup>13–20</sup> together with the ability of the host phagocytes to properly internalize and clear bacteria.<sup>13–20</sup>

CFTR is expressed by human and mouse macrophages.<sup>18,19</sup> However, whether and how defective CFTR has a direct role in bacterial killing is still debated. Conflicting results were reported on whether defective CFTR impairs the acidification of the lysosome or deranges the clearance of *P. aeruginosa* via redoxdependent processes.<sup>17–20</sup> More recently, impaired autophagy has been involved in the defective killing of *B. cepacia* by CF macrophages.<sup>21,22</sup> Autophagy is a general mechanism of survival that cells adopt to cope with stress through targeting cytosolic or recently internalized proteins to lysosomal degradation.<sup>38–43</sup> Defective autophagy was first discovered as a major consequence of defective CFTR in bronchial epithelial cells<sup>9,10,45</sup> and subsequently described in CF macrophages.<sup>21,22</sup> Depletion of SQSTM1/p62 from F508del-CFTR macrophages *in vitro* improves autophagy, increases the colocalization of *B. cepacia* with the master autophagy protein BECN1 and reduces intracellular *B. cepacia* survival.<sup>21,22</sup> We have previously described that defective autophagy in CF is due to TG2-mediated sequestration of BECN1, thus reducing the BECN1-dependent generation of phosphatidyl-inositol-3-phosphate (*PI3P*) that is important for increasing autophagic flux.<sup>9,10,45,46</sup>

Our results provide genetic proof for the role of CFTR and autophagy in orchestrating the sequential cascade of events following the acute infection of *P. aeruginosa* in BMDMs (which are non-conditioned phagocytes). We used a panoply of established and newly generated transgenic mice bearing different CFTR defects, such as  $Cftr^{-/-}$  mice (which lack any rescuable CFTR protein) or mice bearing F508del-CFTR in homozygosis ( $Cftr^{F508del/F508del}$  mice) or in combination with a null CFTR allele ( $Cftr^{F508del/-}$  mice). Although murine models of CF do not recapitulate the clinical manifestations of CF lung disease, CF mice manifest an inflammatory lung

Bacterial clearance by cysteamine in CF macrophages E Ferrari et al



**Figure 5** Cysteamine reverts defective killing of *P. aeruginosa* through rescuing CFTR. Bone marrow-derived macrophages (BMDMs) from *Cftr<sup>F508del/F508del</sup>* (n=3) (a=c), *Cftr<sup>-/-</sup>* (n=3) (a=c), and Cftr<sup>F508del/-</sup> (n=3) (d=f), and infected with *P. aeruginosa*-O1 (PAO1) for 10 min followed by 10 min (a, d) or 3 h (b=c and e=f) of culture in the presence of gentamicin. (a=b, d=e) number of CFU of PAO1; (c, f) clearance of bacteria expressed as percentage of living bacteria after 3 h of culture respect to the total amount of internalized bacteria. BMDMs from *Cftr<sup>-/-</sup>* (n=3) (a=c) and Cftr<sup>F508del/-</sup> (d=f) mice treated with 250  $\mu$ M cysteamine or vehicle in the presence or absence of 3-methyl-adenine (3-MA) (3 mM) and/or CFTR<sub>inh172</sub> (20  $\mu$ M) (d=f) and then infected with PAO1 for 10 min followed by 10 min (a, d) or 3 h (b=c and e=f) of gentamicin culture. \*P<0.05, \*\*P<0.01, Student's *t*-test. Means  $\pm$  S.D. of three independent measurements for each group of experiments. Asterisks indicate significant differences

phenotype coupled to hyper-responsiveness to bacteria or microbial products and are widely used to test candidate drugs and their mechanism of action.47 In addition, we used autophagy-defective Becn1+/- mice as well as Cft/F508del/F508del mice in a background of BECN1 haploinsufficiency (*Cftr*<sup>F508del/F508del</sup>/*Becn*1<sup>+/-</sup> mice). First, we analyzed the early steps following acute infection. As soon as 10 min following infection, CFTR competent BMDMs are able to internalize bacteria, whereas BMDMs isolated from either  $Cftr^{-/-}$  or  $Cftr^{F508del/F508del}$  or  $Cftr^{F508del/-}$  mice exhibited a major internalization defect. Notably, the CFU enumerated after 3 h of incubation following infection, were similar in WT and CF mice. However, the percentage of clearance of internalized bacteria was impaired in BMDMs from CF with distinct genotypes (Cftr<sup>-/-</sup>, Cftr<sup>F508del</sup>/-, mice CftrF508del/F508del), indicating that CFTR defective macrophages are unable to properly clear internalized bacteria. The defect of *P. aeruginosa* internalization was not secondary to defective BECN1 function or reduced autophagy as it was not observed in Becn1+/- mice that, however, manifested reduced intracellular clearance of internalized bacteria to

approximately a half of the values of their autophagycompetent littermates. Accordingly, when  $Cftr^{F508del/F508del}$ were backcrossed in a BECN1 haplo-insufficient background  $(Cftr^{F508del/F508del}/Becn1^{+/-}$  mice), the response of BMDMs to the acute *P. aeruginosa* infection was similar to that observed in  $Cftr^{F508del/F508del}$  mice.

Cysteamine was effective in restoring BMDMs dysfunction from either *Cftr*<sup>F508del/F508del</sup> and *Cftr*<sup>F508del/-</sup>, but not *Cftr*<sup>-/-</sup> transgenic mice. This supports the hypothesis that cysteamine exerts its beneficial effects on CF through restoring a functional F508del-CFTR, in agreement with our previous data in transgenic mice and in CF patients bearing class II CFTR mutations. In contrast, cysteamine failed to have any beneficial effect of *P. aeruginosa* uptake or clearance by BMDMs when CFTR protein was not rescuable, namely in *Cftr*<sup>-/-</sup> BMDMs. Moreover, cysteamine was not effective on *Cftr*<sup>F508del/F508del</sup>/Becn1<sup>+/-</sup> BMDMs as well as on BMDMs from *Cftr*<sup>F508del/F508del</sup> that were simultaneously treated with the autophagy inhibitor 3-MA. Altogether these results indicate that cysteamine can rescue a WT like phenotype in CF macrophages

Bacterial clearance by cysteamine in CF macrophages E Ferrari et al



**Figure 6** Cysteamine controls *P. aeruginosa*-induced increase of TNF $\alpha$  and IL-1 $\beta$  levels in bone marrow-derived macrophages (BMDMs) from *Cttr<sup>F508del/F508del</sup>* mice. BMDMs from *Cttr<sup>F508del/F508del</sup>* (n=3) (**a** and **b**) and *Cttr<sup>F508del/F508del</sup>/Becn1<sup>+/-</sup>* (n=3) (**c** and **d**) mice treated with 250  $\mu$ M cysteamine in the presence or absence of 3-methyladenine (3-MA) (3 mM) (**a** and **b**) and then infected with *P. aeruginosa*-O1 (PAO1) for 10 min followed by culture with gentamicin up to 24 h. Quantitation of TNF $\alpha$  (**a**, **c**) and IL-1 $\beta$ (**b**, **d**) levels in culture supernatants by ELISA. \*\*\**P*<0.001, °°°*P*<0.001, Student's *t*-test. Means  $\pm$  S.D. of three independent measurements for each group of experiments. Asterisks indicate significant differences

through restoring autophagy, exactly as it does in CF epithelial cells.

Unresolved inflammation is a major feature of CF lungs. Defective CFTR generates a dysregulation of the innate immune response at the mucosal sites facing airway and gut lumen.<sup>1,3,4</sup> This leads to an unbalanced response to either cell-autonomous or luminal stressors, as it is the case of bacterial infections in airways. Such a constitutive, yet inefficient, activation of the innate immune response increases the susceptibility of CF patients to manifest recurrent and unresolved bacterial infections that further increase lung inflammation.

Therefore, defective *P. aeruginosa* internalization coupled to excessive and unbalanced activation of mucosal immune response is a puzzling feature of CF airways. The mechanisms responsible for *P. aeruginosa* internalization in different cell types and tissues are complex and not completely understood. CFTR has a key role in the formation of an

internalization platform that requires cytoskeleton integrity and the cooperation of the essential scaffold protein Caveolin 1 (CAV-1).<sup>15</sup> CF macrophages fail to enhance CAV-1 expression in response to LPS<sup>15</sup> and also show defective compartimentalization of heme-oxygenase 1 (HO-1),48 a regulator of redox balance induced by stress conditions including infections, to the cell surface where the CAV-1/HO-1 complex acts as negative regulator of TLR-4 signaling.<sup>15,48,49</sup> Thus, defective CFTR compromises major mechanisms of bacterial internalization by either mouse and human macrophages, while sustaining an inappropriate inflammatory response. These puzzling features of CF macrophages may explain how even low bacterial challenges may result in an exaggerated inflammatory response due to defective elimination of internalized bacteria coupled to sustained TLR signaling.

In conclusion, our study provides genetic proof on how the interplay between defective CFTR and impaired autophagy

8

compromises the ability of CF macrophages to fight P. aeruginosa infection. Our results offer a new perspective for the treatment of *P. aeruginosa* infection in CF patients. as they provide the proof-of-concept for the beneficial effect of cysteamine in reverting macrophage dysfunction in CF. Recurrent and chronic bacterial infections represent a major problem in the treatment of CF patients and entail the need of long-term and expensive antibiotic treatments with the risk of adverse reactions and antibiotic resistance. Cvsteamine is a repurposed drug, with a known safety profile. Combining the effects of cysteamine on CFTR rescue and lung inflammation with the ability to improve bacterial clearance by macrophages, as we describe in this study, might result in long-term clinical benefit to CF patients bearing the F508del-CFTR mutation. At this point, we surmise that this speculation warrants further clinical evaluation by investigating primary macrophages from cysteamine-treated CF patients.

#### Materials and Methods

Mice and treatments. CF mice homozygous for the F508del-CFTR in the FVB/129 outbred background (Cftrtm<sup>1EUR</sup>, F508del, FVB/129, abbreviated Cftr<sup>F50Bdel/F50Bdel</sup>) were obtained from Bob Scholte, Erasmus Medical Center Rotterdam, The Netherlands, CF coordinated action program EU FP6 LSHMCT-2005-018932.<sup>50</sup> Transgenic KO Cftr mice (B6.129P2-KOCftrtm<sup>1UNC</sup>, abbreviated Cftr-/-), were purchased from The Jackson Laboratory (Bar Harbor, ME, USA). The heterozygous *Cftr<sup>F508del/+</sup>* males were backcrossed with the heterozygous Cftr<sup>+/-</sup> females to obtain F508del/null CFTR heterozygous mice (abbreviated Cftr<sup>F508del/-).34</sup> Cftr<sup>F508del/+</sup> female mice were backcrossed to the C57BL/6J background Becn1<sup>+/-</sup> male mice (generous gift from Beth Levine, Center for Autophagy Research, Department of Internal Medicine, UT Southwestern Medical Center, Dallas, USA and Francesco Cecconi, University of Tor Vergata, Rome, Italy) to obtain at the first generation Becn1 haplo-insufficient F508del heterozygous mice (abbreviated  $Cftr^{F508del/+}/Becn^{1+/-}$ ). These mice  $Cftr^{F508del/+}/Becn^{1+/-}$  were crossbred to obtain Becn1 haplo-insufficient F508del homozygous mice (abbreviated Cftr<sup>F508del/F508del</sup>/Becn<sup>1+/-</sup>).<sup>34</sup> The newly generated Cftr<sup>F508del/-</sup> and the Cftr<sup>F508del/F508del</sup>/Becn1<sup>+/-</sup> were housed at the San Raffaele Scientific Institute SOPF animal house (Milan, Italy). These mice were provided with a special food, consisting of an equal mixture of SRM-A (Arie Blok, Woerden, The Netherlands) and Teklad 2019 (Harlan Laboratories, San Pietro al Natisone, Udine, Italy) and water acidified to pH 2.0 with HCl and containing 60 g/l PEG 3350, 1.46 g/l NaCl, 0.745 g/l KCl, 1.68 g/l NaHCO, and 5.68 g/l Na2SO4. Newborn mice were genotyped by cutting a small piece of tail 12 days after birth. DNA was extracted by digesting tails with Direct PCR Lysis Reagent (Viagen, CA, USA) and 1 mg/ml Proteinase K overnight at 56 °C. For Cftr<sup>F508del/-</sup> two PCR reactions were performed. For the *Cftr<sup>F508del/F508del</sup>* mutation, thermocycling consisted of an initial polymerase activation step at 95 °C for 5 min, amplification was performed with 30 cycles of 95 °C for 1 min, 52 °C for 1 min and 72 °C for 1 min with a final extension at 72 °C for 2 min; for the  $Cftr^{-/-}$  mutation thermocycling consisted of an initial polymerase activation step at 94 °C for 3 min, amplification was performed with 30 cycles of 94 °C for 30 s, 57 °C for 30 s and 72 °C for 30 s with a final extension at 72 °C for 10 min. For  $Cthr^{508del/F508del/Becn1^{+/-}}$ , after analyzing the Cftr<sup>F508del/F508del</sup> mutation, the Becn1<sup>+/-</sup> thermocycling consisted of an initial polymerase activation step at 95 °C for 4 min, amplification was performed with 30 cycles of 95 °C for 1 min, 62 °C for 1 min and 72 °C for 1.30 min with a final extension at 72 °C for 10 min. Mice were anesthetized with Avertine (tribromoethanol, 250 mg/kg. Sigma Aldrich, Milan, Italy, T48402) and then killed and bone marrow was collected. All the procedures in mice were approved by the local Ethics Committee for Animal Welfare (IACUC No. 713) and were carried out in strict respect of European and National regulations.

**Bacteria.** *P. aeruginosa* strain PAO1 or PAO1-GFP (kindly provided by EM Bruscia, Yale University School of Medicine) was used for these studies. Bacteria were kept frozen at -80 °C in 10% glycerol. Bacteria were thawed and grown in broth LB (Sodium Chloride, Sigma Aldrich, bacto tryptone, BD yeast extract, BD) overnight. The concentration of bacteria was determined by spectrophotometric reading at 600 nm, as reported.<sup>51</sup>

**Isolation and culture of BMDMs.** BMDMs were isolated from the femurs of 6- to 12-wk-old mice and cultured as previously described.<sup>21,22</sup> After overnight culture, the non-adherent cells were differentiated for 7 days in 20 ng/ml recombinant M-CSF (ConnStem Inc., CT, USA). After 7 days, cells were detached and characterized by flow cytometry (F4-80+/MAC-1+ population). We obtained  $1-3 \times 10^7$  mature macrophages for each mice.

**BMDMs treatments.** Macrophages were treated for 18 h with medium or cysteamine, (250  $\mu$ M, Sigma Aldrich, M9768), with or without the autophagy inhibitor 3-MA (3 mM, Sigma Aldrich, M9281) or wortmannin (0.5  $\mu$ M Sigma Aldrich, W1628) or the CFTR inhibitor (CFTR<sub>inh-172</sub>) (20  $\mu$ M, Sigma Aldrich, C2992) and then infected in the presence or absence of drugs. The cells were infected, with PAO1 or PAO-1GFP at a MOI of 50 bacteria/macrophage for 10 min and then were washed and treated with gentamicin (100  $\mu$ g/ml, Sigma Aldrich, G1397) for 10 min and then lysed or extensively washed and kept in culture up to 3 h or 24 h. The cells were lysed with a solution 0.2% triton x-100 (Biorad, Milan, Italy 1610407) and then plated in agar petridish in presence of carbenicillin (100  $\mu$ g/ml, Sigma Aldrich, C1613) overnight at 37 °C. CFU were enumerated as desctibed.<sup>21,22</sup> In addition, the enumeration of the residual intracellular *R aeruginosa* clearance by BMDMs, was determined by calculating the percentage of living bacteria after 3 h of culture with respect to the total amount of internalized bacteria.

**Immunofluorescence.** The microscopy samples were analyzed with an Axivert 200LSM5 Meta microscope (Zeiss, NY, USA). Two million BMDMs were cultured on 12 mm glass cover slips in 24-well tissue culture plates and infected synchronously with PAO1-GFP at an MOI of 50 bacteria/macrophage. The BMDMs were fixed with PFA 4%. All experiments were performed in at least triplicate.

**Immunoblot analysis.** Western blot analysis was performed as previously described<sup>9,10</sup> with antibodies against the following proteins: BECN1, 1:1000, (Abcam, Cambridge, UK Ab55878),  $\beta$ -actin 1:1000, (Cell Signaling Inc., Danvers, MA, USA, 4970), SQSTM1/p62 1:1000, (Sigma Aldrich, MABN130). The densitometric analysis was performed by Image J software and each data point was expressed as the mean  $\pm$  S.D. of independent experiments.

**Real-time PCR.** Total RNA was extracted with the RNeasy Mini Kit (Qiagen, Milan, Italy, 74104). The mRNA was reverse transcribed with a SuperScriptTM III First Strand Synthesis System (Promega, Milan, Italy, A5001). Quantitative RT-PCR was performed with an iCycler iQ Multicolor Real-Time PCR Detector (BioRad) with iQ TM SYBR Green supermix (Five Prime, 2900217). The relative amounts of mRNA were calculated by using the comparative Ct method. Real-time RT-PCR analyses were executed for evaluating the efficiency of expression. Thermocycling consisted of an initial polymerase activation step at 98 °C for 5 min, and amplification was performed with 35 cycles of 95 °C for 15 s, 68 °C for 10 s, and 72 °C for 20 s with data acquisition at this stage and the reaction finished by the built in melt curve. Expression levels of genes were normalized to the housekeeping gene proC in the same sample.

The sequence of PAO1 was: lasB forward 5'-ATGAACGAGGCGTTCTCCG-3' and reverse 5'-GTTGTACACGCCGCTGGAGT-3'.

Housekeeping gene proC forward 5'-CAGGCCGGGCAGTTGCTGTC-3' and reverse 5'-GGTCAGGCGCGAGGCTGTCT-3'.

**Cytokine assessment.** Levels of IL-1 $\beta$  and TNF- $\alpha$  in culture supernatants from BMDMs collected 24 h following infection were assessed by means of standard ELISA kits (R&D Systems, Minneapolis, MN, USA), according to the manufacturer's instructions, as reported.<sup>33,34</sup> Samples were read in triplicate at 450 nm in Microplate Reader (BioRad) using Microplate Manager 5.2.1 software. Values were normalized to protein concentration evaluated by Bradford analysis.

**Statistical analysis.** Data are reported as arithmetic mean  $\pm$  S.D. Data distribution was analyzed for normality and comparisons between groups for statistical difference were done using Student's two tailed *t*-test or the one-way ANOVA, as indicated. Significant differences are indicated in the figures. All experiments were performed at least three independent times with similar results. Data were analyzed using SPSS 20 software. Statistical significance was defined as *P*-value of <0.05.

### **Conflict of Interest**

VR, LM, and GK are listed as inventors on a patent application describing the use of cysteamine for the treatment of CF.

Acknowledgements. We thank Dr Bob Scholte, Erasmus Medical Center Rotterdam, The Netherlands, who provided Cftr<sup>tm1EUR</sup> (F508del (FVB/129) mice (European Economic Community European Coordination Action for Research in Cystic Fibrosis program EU FP6 SHM-CT-2005-018932), Dr Beth Levine, Center for Autophagy Research, Department of Internal Medicine, UT Southwestern Medical Center, Dallas, USA, and Dr Francesco Cecconi, University of Tor Vergata, Rome, Italy, who provided C57BL/6J background Becn1+/-mice, Dr EM Bruscia, Yale University School of Medicine, who provided GFP-labelled P. aeruginosa. This study was supported by The European Institute for Research in Cystic Fibrosis (IERFC) non-profit foundation, Regional Cystic Fibrosis Associations of Campania, Sicilia, Lazio, Puglia, Piemonte (to Dr L Maiuri, Dr Raia), E-Rare (Rescue CFTR preclinic) (to Dr L Maiuri and Dr. Kroemer); Telethon (#GGP12128) (to Dr L Maiuri, Dr Raia, Dr MC Maiuri), Agence National de la Recherche (ANR) - Projetsblancs; ANR under the frame of E-Rare-2. the ERA-Net for Research on Rare Diseases: Association pour la recherche sur le cancer (ARC); Cancéropôle lle-de-France; Institut National du Cancer (INCa); Fondation Bettencourt-Schueller; Fondation de France; Fondation pour la Recherche Médicale (FRM); the European Commission (ArtForce); the European Research Council (ERC); the LabEx Immuno-Oncology; the SIRIC Stratified Oncology Cell DNA Repair and Tumor Immune Elimination (SOCRATE); the SIRIC Cancer Research and Personalized Medicine (CARPEM); and the Paris Alliance of Cancer Research Institutes (PACRI) (all to Dr Kroemer).

- 1. Ratjen F, Doring G. Cystic fibrosis. Lancet 2003; 361: 681-689.
- De Boeck K, Zolin A, Cuppens H, Olesen HV, Viviani L. The relative frequency of CFTR mutation classes in European patients with cystic fibrosis. J Cyst Fibrosis 2014; 13: 403–409.
- O'Sullivan BP, Freedman SD. Cystic fibrosis. *Lancet* 2009; **373**: 1891–1904.
   Ratjen FA. Cystic fibrosis: pathogenesis and future treatment strategies. *Respir Care* 2009; **54**: 595–605.
- 5. Pittman JE, Ferkol TW. The evolution of cystic fibrosis care. *Chest* 2015; **148**: 533–427.
- De Boeck K, Amaral MD. Progress in therapies for cystic fibrosis. Lancet Resp Med 2016; 4: 662–674.
- Amaral MD, Balch WE. Hallmarks of therapeutic management of the cystic fibrosis functional landscape. J Cyst Fibros 2015: 14: 687–699.
- Esposito S, Tosco A, Villella VR, Raia V, Kroemer G, Maiuri L. Manipulating proteostasis to repair the F508del-CFTR defect in cystic fibrosis. *Mol Cell Pediatr* 2016; 3: 13.
- Luciani A, Villella VR, Esposito S, Brunetti-Pierri N, Medina D, Settembre C et al. Defective CFTR induces aggresome formation and lung inflammation in cystic fibrosis through ROS-mediated autophagy inhibition. Nat Cell Biol 2010; 12: 863–875.
- Luciani A, Villella VR, Esposito S, Gavina M, Russo I, Silano M et al. Targeting autophagy as a novel strategy for facilitating the therapeutic action of potentiators on DF508 cystic fibrosis transmembrane conductance regulator. Autophagy 2012; 8: 1657–1672.
- Maiuri L, Luciani A, Giardino I, Raia V, Villella VR, D'Apolito M et al. Tissue transglutaminase activation modulates inflammation in cystic fibrosis via PPARgamma down-regulation. J Immunol 2008; 180: 7697–7705.
- Luciani A, Villella VR, Vasaturo A, Giardino I, Raia V, Pettoello-Mantovani M *et al.* SUMOylation of tissue transglutaminase as link between oxidative stress and inflammation. *J Immunol* 2009; **183**: 2775–2784.
- Bruscia EM, Zhang PX, Ferreira E, Caputo C, Emerson JW, Tuck D et al. Macrophages directly contribute to the exaggerated inflammatory response in cystic fibrosis transmembrane conductance regulator – / – mice. Am J Respir Cell Mol Biol 2009; 40: 295–304.
- Bruscia EM, Zhang PX, Satoh A, Caputo C, Medzhitov R, Shenoy A et al. Abnormal trafficking and degradation of TLR4 underlie the elevated inflammatory response in cystic fibrosis. J Immunol 2011; 186: 6990–6998.
- Bajmoczi M, Gadjeva M, Alper SL, Pier GB, Golan DE. Cystic fibrosis transmembrane conductance regulator and caveolin-1 regulate epithelial cell internalization of Pseudomonas aeruginosa. *Am J Physiol Cell Physiol* 2009; **297**: C263–C277.
- Deriy LV, Gomez EA, Zhang G, Beacham DW, Hopson JA, Gallan AJ et al. Disease causing mutations in the cystic fibrosis transmembrane conductance regulator determine the functional responses of alveolar macrophages. J Biol Chem 2009; 284: 35926–35938.
- Zhang Y, Li X, Grassme H, Do'ring G, Gulbins E. Alterations in ceramide concentration and pH determine the release of reactive oxygen species by Cftr-deficient macrophages on infection. J Immunol 2010; 184: 5104–5111.

- Del Porto P, Cifani N, Guarnieri S, Di Domenico EG, Mariggio MA, Spadaro F et al. Dysfunctional CFTR alters the bactericidal activity of human macrophages against Pseudomonas aeruginosa. PLoS One 2011; 6: e19970.
- Painter RG, Valentine VG, Lanson NA Jr, Leidal K, Zhang Q, Lombard G et al. CFTR expression in human neutrophils and the phagolysosomal chlorination defect in cystic fibrosis. *Biochemistry* 2006; 45: 10260–10269.
- Di A, Brown ME, Deriy LV, Li C, Szeto FL et al. CFTR regulates phagosome acidification in macrophages and alters bactericidal activity. Nat Cell Biol 2006; 8: 933–944.
- Abdulrahman BA, Khweek AA, Akhter A, Caution K, Kotrange S et al. Autophagy stimulation by rapamycin suppresses lung inflammation and infection by Burkholderia cenocepacia in a model of cystic fibrosis. *Autophagy* 2011; 7: 1359–1370.
- Abdulrahman BA, Khwk AA, Akhter A, Caution K, Tazi M, Hassan H *et al.* Depletion of the ubiquitin-binding adaptor molecule SQSTM1/p62 from macrophages harboring cftr DF508 mutation improves the delivery of Burkholderia cenocepacia to the autophagic machinery. *J Biol Chem* 2013; 288: 2049–2058.
- Amaral MD. Novel personalized therapies for cystic fibrosis: treating the basic defect in all patients. J Intern Med 2015; 277: 155–166.
- Quon BS, Rowe SM. New and emerging targeted therapies for cystic fibrosis. BMJ 2016; 352: i859.
- Farinha CM, Matos P. Repairing the basic defect in cystic fibrosis one approach is not enough. FEBS J 2016; 283: 246–264.
- Harrison MJ, Murphy DM, Plant BJ. Ivacaftor in a G551D homozygote with cystic fibrosis. N Engl J Med 2013; 369: 1280–1282.
- Wainwright CE, Elborn JS, Ramsey BW, Marigowda G, Huang X, Cipolli M et al. TRAFFIC and TRANSPORT Study Groups Lumacaftor-Ivacaftor in patients with cystic fibrosis homozygous for Phe508del CFTR. N Engl J Med 2015; 373: 220–231.
- Boyle AMP, Bell SC, Konstan MW, McColley SA, Rowe SM, Rietschel E et al. VX09-809102 study group. CFTR corrector (lumacaftor) and a CFTR potentiator (ivacaftor) for treatment of patients with cystic fibrosis who have a phe508del CFTR mutation: a phase 2 randomised controlled trial. *Lancet Respir Med* 2014; 2:527–538.
- 29. Davis PB. Another beginning for cystic fibrosis therapy. N Engl J Med 2015; 373: 274-276.
- Jones AM, Barry PJ. Lumacaftor/ivacaftor for patients homozygous for Phe508del-CFTR: should we curb our enthusiasm? *Thorax* 2015; 70: 615–616.
- Amaral MD, Farinha C. Rescuing mutant. CFTR: a multi-task approach to a better outcome in treating cystic fibrosis. *Curr Pharm Des* 2013; 19: 3497–3508.
- Hutt DM, Herman D, Rodrigues AP, Noel S, Pilewski JM, Matteson J et al. Reduced histone deacetylase 7 activity restores function to misfolded CFTR in cystic fibrosis. Nat Chem Biol 2010; 6: 25–33.
- De Stefano D, Villella VR, Esposito S, Tosco A, Sepe A, De Gregorio F et al. Restoration of CFTR function in patients with cystic fibrosis carrying the F508del-CFTR mutation. *Autophagy* 2014; 10: 2053–2074.
- Tosco A, De Gregorio F, Esposito S, De Stefano D, Sana I, Ferrari E et al. A novel treatment of cystic fibrosis acting on-target: cysteamine plus epigallocatechin gallate for the autophagy-dependent rescue of class II-mutated CFTR. Cell Death Differ 2016; 8: 1380–1393.
- Venerando A, Franchin C, Cant N, Cozza G, Pagano MA, Tosoni K et al. Detection of phospho-sites generated by protein kinase CK2 in CFTR: mechanistic aspects of Thr1471 phosphorylation. PloS One 2013; 8: e74232.
- Devereux G, Fraser-Pitt D, Robertson J, Devlin E, Mercer D, O'Neil D et al. Cysteamine as a future intervention in cystic fibrosis against current and emerging pathogens: a patient-based ex vivo study confirming its antimicrobial and mucoactive potential in sputum. *Ebio Med* 2015; 2: 1507–1512.
- Charrer C, Rodger J, Robertson A, Kowalczuk N, Shand D, Fraser-Pitt D *et al.* Cysteamine (Lynovex), a novel mucoactive antimicrobial & antibiofilm agent for the treatment of cystic fibrosis. *Orphanet J Rare Dis* 2014; 9: 189.
- 38. Yang Z, Klionsky DJ. Eaten alive: a history of macroautophagy. Nat Cell Biol 2010; 12: 814–822.
- Mizushima N, Yoshimori T, Ohsumi Y. The role of Atg proteins in autophagosome formation. *Annu Rev Cell Dev Biol* 2011: 27: 07–32.
- Kroemer G, Mariño G, Levine B. Autophagy and the integrated stress response. *Mol Cell* 2010: 40: 280–293.
- Codogno P, Mehrpour M, Proikas-Cezanne T. Canonical and non-canonical autophagy: variations on a common theme of self-eating? *Nat Rev Mol Cell Biol* 2012; 13: 7–12.
- Levine B, Mizushima N, Virgin HW. Autophagy in immunity and inflammation. *Nature* 2011; 469: 323–335.
- Sridhar S, Botbol Y, Macian F, Cuervo AM. Autophagy and disease: always two sides to a problem. J Pathol 2012; 226: 255–273.
- Talwalkar JS, Murray TS. The approach to pseudomonas aeruginosa in cystic fibrosis. Clin Chest Med 2016. 37: 69–81.
- 45. Villella V R, Esposito S, Bruscia E M, Maiuri M C, Raia V, Kroemer G, Maiuri L. Targeting the intracellular environment in cystic fibrosis: restoring autophagy as a novel strategy to circumvent the CFTR defect. *Front Pharmacol* 2013; 21: 1.
- Villella VR, Esposito S, Bruscia EM, Vicinanza M, Cenci S, Guido S et al. Disease-relevant proteostasis regulation of cystic fibrosis transmembrane conductance regulator. *Cell Death Differ* 2013; 20: 1101–1115.

- Lavelle GM, White MM, Browne N, McElvaney NG, Reeves EP. Animal models of cystic fibrosis pathology: phenotypic parallels and divergences. *Biomed Res Int* 2016; 2016: 5258727.
- Zhang PX, Murray TS, Villella RV, Ferrari E, Esposito S, D'Souza A et al. Reduced caveolin-1 promotes hyper-inflammation due to abnormal heme oxygenase-1 localizationin LPS challenged macrophages with dysfunctional CFTR. J Immunol 2013; 190: 5196–5206.
- Abuarqoub H, Foresti R, Green CJ, Motterlini R. Heme oxygenase-1mediates the anti-inflammatory actions of 29-hydroxychalcone in RAW 264.7 murine macrophages. *Am J Physiol Cell Physiol* 2006; 290: C1092–C1099.
- vanDoorninck JH, French PJ, Verbeek E, Peters RH, Morreau H, Bijman J et al. A mouse model for the cystic fibrosis delta F508 mutation. EMBO J 1995; 14: 4403–4411.
- Kukavica I, Bragonzi A, Paroni M, Winstanley C, Sanschagrin F, O'Toole GA *et al. In vivo* growth of Pseudomonas aeruginosa strains PAO1 and PA14 and the hypervirulent strain LESB58 in a rat model of chronic lung infection. *J Bacteriol* 2008; **190**: 2804–2813.

**Cell Death and Disease** is an open-access journal published by *Nature Publishing Group*. This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/

11

© The Author(s) 2017

Supplementary Information accompanies this paper on Cell Death and Disease website (http://www.nature.com/cddis)