



HAL
open science

Self-organized pattern formation of run-and-tumble chemotactic bacteria: Instability analysis of a kinetic chemotaxis model

Benoît Perthame, Shugo Yasuda

► To cite this version:

Benoît Perthame, Shugo Yasuda. Self-organized pattern formation of run-and-tumble chemotactic bacteria: Instability analysis of a kinetic chemotaxis model. 2017. hal-01494963v1

HAL Id: hal-01494963

<https://hal.sorbonne-universite.fr/hal-01494963v1>

Preprint submitted on 24 Mar 2017 (v1), last revised 12 Jan 2018 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Self-organized pattern formation of run-and-tumble chemotactic bacteria: Instability analysis of a kinetic chemotaxis model

Benôit PERTHAME^{a,*}, Shugo YASUDA^{b,**}

^a*Sorbonne Universités, UPMC Univ Paris 06, Laboratoire Jacques-Louis Lions UMR CNRS 7598, Université Paris-Diderot, Inria de Paris, F75005 Paris*

^b*Graduate School of Simulation Studies, University of Hyogo, Kobe 650-0047, Japan*

Abstract

Collective motion of chemotactic bacteria as *E. Coli* relies, at the individual level, on a continuous reorientation by runs and tumbles. It has been established that the length of run is decided by a stiff response to temporal sensing of chemical cues along the pathway. We analyze consequences on self-organized pattern formation resulting from modulation of tumbling frequency with stiff response. Thanks to both analytical arguments and numerical simulations, we show that the stationary homogeneous state of population density is destabilized in realistic ranges of parameters. A remarkable property is that the unstable frequencies remain bounded, as it is the case in Turing instability. These theoretical developments are in accordance with several biological observations.

Keywords: Self-organization, Stiff response, Population growth, Kinetic reaction-transport equation, Instability analysis, Monte Carlo method

*Email: benoit.perthame@upmc.fr

**Email: yasuda@sim.u-hyogo.ac.jp

1. Introduction

Collective motion of chemotactic bacteria as *E. Coli* relies, at the individual level, on a continuous reorientation by runs and tumbles sensing extracellular chemoattractants produced by themselves [1, 2, 3, 4]. It has been established that the length of run is decided by a stiff response to temporal sensing of chemical cues along pathway, i.e., bacteria reduce their tumbling frequency and extend the run length as they sense an increase in concentrations of chemoattractants along the pathway. Thus, the modulation of tumbling frequency in the stiff response is an essential mechanism for bacterial communities self-organization.

This paper is concerned with the pattern formation of the population density of run-and-tumble chemotactic bacteria as *E. Coli*. We investigate the emergence of self-organized pattern formation stemming from a modulation of tumbling frequency in chemotactic response at an individual level. In order to investigate the multiscale nature in self-organized pattern formation, we use a mesoscopic model, i.e., a kinetic reaction-transport equation for the chemotactic bacteria coupled with a reaction-diffusion equation for the chemoattractants, in which the microscopic dynamic properties such as tumbling rate, modulation in stiff response, and proliferation (division/death) rate are involved at the individual level.

We consider the following three main ingredients in the pattern formation: (i) the random run-and-tumble motion bacteria, where the bacteria run linearly with a constant speed when rotating their flagella in counter-clockwise direction, but occasionally change the running directions (tumbling) when rotating their flagella in clockwise direction; (ii) the chemotactic response to

the spatiotemporal variation of the concentration of chemoattractant along the pathway of bacterium, which generates the biased random motion searching for the higher-concentration region of chemoattractant; and (iii) the division/death of bacteria, where the population-growth rate depends on the local population density of bacteria. The kinetic transport equation considered in this paper describe all these ingredients at the microscopic (individual) level. The pattern formations in the chemotaxis with population growth have been investigated at the macroscopic level by the Keller-Segel type equations [5, 6, 7]; for example, in [8, 9, 10, 11], the pattern formations induced by the properties of chemotaxis, i.e., so-called *chemotaxis-induced instability*, are demonstrated both theoretically and numerically. This paper is also concerned with the chemotaxis-induced instability, but which is based on the kinetic transport equation. The kinetic approach has a significant advantage in studying the multiscale mechanism and mathematical hierarchy between the individual dynamics and macroscopic phenomena.

The kinetic approach to describe the chemotaxis of run-and-tumble bacteria was first proposed in [12, 13] and then further has been developed toward involving the spatiotemporal variation of the chemoattractant along the pathway of bacterium [15], the internal dynamics of the cellular states [14], and the multi-cellular interactions [16, 17]. The mathematical foundations for the kinetic chemotaxis model have been strengthened involving the mathematical hierarchy between the kinetic and continuum equations and the existence of solution for the kinetic chemotaxis equation [19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32]. The numerical methods to solve the kinetic chemotaxis equations have been also developed in [24, 36, 37, 38, 39, 40]. The

use of the kinetic chemotaxis model is also advanced due to the development of experimental technologies, which allow experimentalists to measure the individual velocities and turning angles in the collective motions of bacteria and give access to time scale measurements. For example, in [33, 34, 35], the advantage of the kinetic modeling is demonstrated by the comparison of the numerical and experimental results.

In this paper, we carry out the linear instability analysis of a kinetic chemotaxis equation coupled with a diffusion-reaction equation for the chemoattractant. The kinetic chemotaxis equation involves a population-growth term, which depends on the local population density of bacteria, as well as a chemotactic response function in the integral kernel, which depends on the spatiotemporal variation of the chemoattractant along the pathway of each bacterium. We obtain a linear instability condition for which the stationary homogeneous state of the population density of bacteria becomes linearly unstable and where periodic patterns are generated. We also numerically demonstrate the pattern formations by Monte Carlo simulations in which we vary the parameters involved in the linear instability condition. A theoretical foundation for the Monte Carlo method is also presented.

2. Main result

Since the observation of the run and tumble movement of bacteria [33, 35], the kinetic chemotaxis equation has been proposed as an accurate description [13, 14, 15]. Here we also consider a growth term.

$$\partial_t f(t, \mathbf{x}, \mathbf{v}) + \mathbf{v} \cdot \nabla f = \frac{1}{k} \left\{ \frac{1}{4\pi} \int_V K[D_t \log S|_{\mathbf{v}'}] f(\mathbf{v}') d\Omega(\mathbf{v}') - K[D_t \log S|_{\mathbf{v}}] f(\mathbf{v}) \right\} + P[\rho] f(\mathbf{v}). \quad (1)$$

Here $f(t, \mathbf{x}, \mathbf{v})$ is the microscopic population density of bacteria with a velocity $\mathbf{v} \in V$ at a position $\mathbf{x} \in \mathbb{R}^3$ and a time $t \geq 0$. The velocity space of \mathbf{v} , V is the surface of the unit ball, i.e., $|\mathbf{v}| = 1$ and $\Omega(\mathbf{v})$ is the unit measure on V . $S(t, \mathbf{x})$ is the concentration of the chemical attractant and $\rho(t, \mathbf{x})$ is the macroscopic population density of the bacteria, which are calculated as, respectively,

$$-d\Delta S(t, \mathbf{x}) + S(t, \mathbf{x}) = \rho(t, \mathbf{x}), \quad (2)$$

$$\rho(t, \mathbf{x}) = \frac{1}{4\pi} \int_V f(t, \mathbf{x}, \mathbf{v}) d\Omega(\mathbf{v}), \quad (3)$$

where d is the diffusion constant. In Eq. (1), $K[X]$ is the tumbling kernel (which involves the chemotactic responses of bacteria as defined in Eq. (5)), $D_t X|_{\mathbf{v}}$ is the material derivative, i.e., $D_t X|_{\mathbf{v}} = \partial_t X + \mathbf{v} \cdot \nabla X$, and $P[\rho]$ is the population-growth rate of bacteria which is dependent on the local population density ρ as

$$P[\rho] > 0, \quad \text{for } 0 < \rho < 1, \quad (4a)$$

$$P[\rho] < 0, \quad \text{for } \rho > 1, \quad (4b)$$

$$P[\rho] \simeq 1 - \rho, \quad \text{for } \rho \simeq 1. \quad (4c)$$

Thus, the bacteria may divide when the local population density is lower than unity and the new born bacteria have the same velocities as the parents, but they may die when the local population density is larger than the unity.

We remark that the existence of the traveling wave in the kinetic transport equations with population growth (but without chemotactic responses, i.e., $K[X] = \text{const.}$) are proved in [44, 45, 46], where new born particles may choose their velocities according to a prescribed equilibrium velocity distribution function. In this paper, we use the simplest population-growth

model among those for which the existence of the traveling wave is proved mathematically and the logistic population-growth term is recovered in the continuum limit.

The tumbling kernel $K[X]$ in Eq. (1) is a decreasing function of X and we choose it as

$$K[X] = 1 - F[X], \quad (5)$$

where $F[X]$ is a smooth function which satisfies the following properties,

$$F[0] = 0, \quad (6a)$$

$$\frac{dF[X]}{dX} > 0, \quad (6b)$$

$$F[X] \rightarrow \pm\chi \text{ as } X \rightarrow \pm\infty. \quad (6c)$$

Here, $F[X]$ represents the chemotactic response of the bacteria, say the response function, and χ represents the amplitude of modulation in the chemotactic response and takes a constant value between $0 \leq \chi < 1$.

In Eqs. (1)–(4), all quantities are nondimensionalized by the following characteristic quantities; the characteristic time t_0 is defined as $t_0 = |(\frac{d\tilde{P}}{d\rho})_{\rho=1}|^{-1}$, where \tilde{P} represents the population-growth rate in the dimensional form, the characteristic length L_0 is defined as $L_0 = t_0 V_0$, where V_0 is a constant speed of the bacteria. The nondimensional parameter k is defined as $k = 1/(t_0 \psi_0)$, where ψ_0 is a mean tumbling frequency of the bacteria. The population density is scaled by that in the uniform stationary state ρ_0 and the concentration of the chemoattractant is scaled by $(a_0/b_0)\rho_0$, where a_0 is the production rate of chemoattractant by the bacteria and b_0 is the degradation rate of chemoattractant.

It is easily seen that Eqs. (1)–(4) have a constant uniform solution with $f(t, \mathbf{x}, \mathbf{v}) = 1$, $S(t, \mathbf{x}) = 1$, and $\rho(t, \mathbf{x}) = 1$. In our main result, this uniform solution gives the Turing-like instability [47, 48]. That is, the uniform solution is linearly unstable if the stiffness of the response function $F'[0]$ is sufficiently large as

$$\frac{F'[0]}{k} > \inf_{\lambda} \left[1 + \frac{k}{\frac{k\lambda}{\arctan(k\lambda)} - 1} \right] (1 + d\lambda^2). \quad (7)$$

In addition, the unstable eigenmodes are bounded, i.e., no high frequency oscillations occur and obtain patterns.

Figure 1 shows the linear instability diagram. We numerically calculate the minimum values of the right-hand side of Eq. (7) with variation in the Fourier mode λ for given values of k and d . It is seen that as increasing the stiffness of the response function $F'[0]$, the stationary homogeneous state with $f = S = 1$ is destabilized (the *chemotaxis-induced* instability). The critical lines increase monotonically as the diffusion coefficient d increases, so that the stationary homogeneous state is more destabilized when the diffusion coefficient d becomes smaller. It is also seen that as decreasing k , the critical line decreases but converges to that for the continuum equation, which will be given in Eq. (34), in the limit of $k \rightarrow 0$. The stationary homogeneous state is more likely destabilized as decreasing k .

3. Linear instability analysis

3.1. Linearization

It is easily seen that Eq. (1) has the uniform solution, $f(t, \mathbf{x}, \mathbf{v}) = S(t, \mathbf{x}) = 1$. We carry out the linear instability analysis about this uniform solution.




Figure 1: The diagram of the kinetic instability obtained by Eq. (7). The linear instability takes place when the stiffness of the response function $F'[0]/k$ exceeds the critical line of each k . The numerical simulations are performed for the marks A, B, C, and D with several values of k . See Table 1.

We consider a small perturbation around the uniform solution as

$$f(t, \mathbf{x}, \mathbf{v}) = 1 + g(\mathbf{x}, \mathbf{v})e^{\mu t}, \quad (8a)$$

$$S(t, \mathbf{x}) = 1 + S_g(\mathbf{x})e^{\mu t}, \quad (8b)$$

$$\rho(t, \mathbf{x}) = 1 + \rho_g(\mathbf{x})e^{\mu t}, \quad (8c)$$

where μ is a constant which distinguishes stability ($\text{Re}(\mu) < 0$) and instability ($\text{Re}(\mu) > 0$). From Eqs. (4)–(6), we can linearize the population-growth rate $P[\rho]$ and tumbling kernel $K[D_t \log S|_{\mathbf{v}}]$ in Eq. (1) as, respectively,

$$P[1 + \rho_g e^{\mu t}] = -\rho_g e^{\mu t}, \quad (9)$$

and

$$K[D_t \log(1 + S_g e^{\mu t})|_{\mathbf{v}}] = 1 - F'[0](\mu S_g + \mathbf{v} \cdot \nabla S_g)e^{\mu t}. \quad (10)$$

Thus, Eq. (1) is linearized as

$$\begin{aligned} & k(\mu g(\mathbf{x}, \mathbf{v}) + \mathbf{v} \cdot \nabla g(\mathbf{x}, \mathbf{v}))e^{\mu t} \\ &= \frac{1}{4\pi} \int_V (1 - F'[0](\mu S_g(\mathbf{x}) + \mathbf{v}' \cdot \nabla S_g(\mathbf{x}))e^{\mu t} + g(\mathbf{x}, \mathbf{v}')e^{\mu t})d\Omega(\mathbf{v}') \\ & - (1 - F'[0](\mu S_g(\mathbf{x}) + \mathbf{v} \cdot \nabla S_g(\mathbf{x}))e^{\mu t} + g(\mathbf{x}, \mathbf{v})e^{\mu t}) - k\rho_g(\mathbf{x}), \end{aligned} \quad (11a)$$

$$\begin{aligned} & k\mu g(\mathbf{x}, \mathbf{v}) + k\mathbf{v} \cdot \nabla g(\mathbf{x}, \mathbf{v}) \\ &= \frac{1}{4\pi} \int_V g(\mathbf{x}, \mathbf{v}')d\Omega(\mathbf{v}') + F'[0]\mathbf{v} \cdot \nabla S_g(\mathbf{x}) - g(\mathbf{x}, \mathbf{v}) - k\rho_g(\mathbf{x}). \end{aligned} \quad (11b)$$

By taking the Fourier transform of Eqs. (2) and (11), we obtain

$$\hat{g}(\boldsymbol{\lambda}, \mathbf{v}) = \frac{1 - k + i\frac{F'[0]}{1+d|\mathbf{v}|^2}\boldsymbol{\lambda} \cdot \mathbf{v}}{1 + k\mu + ik\boldsymbol{\lambda} \cdot \mathbf{v}}\hat{\rho}_g(\boldsymbol{\lambda}). \quad (12)$$

Hereafter, i represents the imaginary unit, $\boldsymbol{\lambda} \in \mathbb{R}$ the wave vector, and $\hat{a}(\boldsymbol{\lambda})$ the Fourier transform of the function $a(\mathbf{x})$ as $\hat{a}(\boldsymbol{\lambda}) = \int_{\mathbb{R}} a(\mathbf{x}) e^{-i\boldsymbol{\lambda} \cdot \mathbf{x}} d\mathbf{x}$. By integrating the above equation as to \mathbf{v} on V , we obtain an equation for $\hat{\rho}_g(\boldsymbol{\lambda})$ as

$$\hat{\rho}_g(\boldsymbol{\lambda}) = \frac{1}{2} \int_{-1}^1 \frac{1 - k + i \frac{F'[0]\lambda v}{1+d\lambda^2}}{1 + k\mu + ik\lambda v} dv \hat{\rho}_g(\boldsymbol{\lambda}), \quad (13a)$$

$$\hat{\rho}_g(\boldsymbol{\lambda}) = \frac{1}{2} \int_{-1}^1 \frac{\left(1 - k + i \frac{F'[0]\lambda v}{1+d\lambda^2}\right) (1 + k\mu_1 - ik\lambda(\mu_2 + v))}{(1 + k\mu_1)^2 + k^2\lambda^2(\mu_2 + v)^2} dv \rho_g(\boldsymbol{\lambda}), \quad (13b)$$

where μ_1 is the real part of μ , i.e., $\mu_1 = \text{Re}(\mu)$, and μ_2 is defined as $\mu_2 = \text{Im}(\mu)/\lambda$. Here we write $\lambda = |\boldsymbol{\lambda}|$.

Thus, in order to obtain a non-trivial solution of $\hat{\rho}_g(\boldsymbol{\lambda})$, the following equations must be simultaneously satisfied, i.e.,

$$\int_{-1}^1 \frac{(1 - k)(1 + k\mu_1) + \frac{F'[0]k\lambda^2 v}{1+d\lambda^2}(\mu_2 + v)}{(1 + k\mu_1)^2 + k^2\lambda^2(\mu_2 + v)^2} dv = 2, \quad (14)$$

and

$$\int_{-1}^1 \frac{(1 - k)k\lambda(\mu_2 + v) - \frac{F'[0]\lambda v}{1+d\lambda^2}(1 + k\mu_1)}{(1 + k\mu_1)^2 + k^2\lambda^2(\mu_2 + v)^2} dv = 0. \quad (15)$$

Further, Eqs. (14) and (15) are analytically calculated as, respectively,

$$\begin{aligned} \left(\alpha - \frac{\beta}{\xi}\right) [\arctan(\xi(\mu_2 + 1)) - \arctan(\xi(\mu_2 - 1))] \\ - \mu_2 \beta \log \left(1 + \frac{4\mu_2}{\xi^{-2} + (\mu_2 - 1)^2}\right) = 2 - 2\beta, \end{aligned} \quad (16)$$

and

$$\begin{aligned} \mu_2 \beta [\arctan(\xi(\mu_2 + 1)) - \arctan(\xi(\mu_2 - 1))] \\ + \frac{1}{2} \left(\alpha - \frac{\beta}{\xi}\right) \log \left(1 + \frac{4\mu_2}{\xi^{-2} + (\mu_2 - 1)^2}\right) = 0, \end{aligned} \quad (17)$$

where

$$\alpha = \frac{1 - k}{k\lambda}, \quad (18a)$$

$$\beta = \frac{F'[0]}{k(1 + d\lambda^2)}, \quad (18b)$$

$$\xi = \frac{k\lambda}{1 + k\mu_1}. \quad (18c)$$

Note that Eqs. (16) and (17) are symmetric as to the sign of μ_2 and $\mu_2=0$ always satisfies Eq. (17) irrespective of the values of α , β , and ξ . The eigenvalue $\mu (= \mu_1 + i\mu_2\lambda)$ is obtained by solving Eqs. (16) and (17) simultaneously, and the sign of the growing rate μ_1 determines the instability of the uniform solution of the kinetic equation, Eq. (1).

Before we consider the instability condition, we first verify that Eq. (16) is never satisfied as $\lambda \rightarrow \infty$. This is explained as follows.

We consider the first term of the left hand side of Eq. (16). In Eq. (18), α and β vanishes as $\lambda \rightarrow \infty$ while the limiting values of ξ and μ_2 are unknown. In the case that ξ converges to a finite value or diverges as $\lambda \rightarrow \infty$, the first term of the L.H.S of Eq. (16) vanishes because the first factor vanishes while the second factor is bounded. On the other hand, if ξ converges to zero as $\lambda \rightarrow \infty$, the second factor of the first term of the L.H.S of Eq. (16) is estimated as

$$\begin{aligned} |\arctan(\xi(\mu_2 + 1)) - \arctan(\xi(\mu_2 - 1))| &= \left| \arctan\left(\frac{2\xi}{1 + \xi^2(\mu_2^2 - 1)}\right) \right| \\ &< \left| \arctan\left(\frac{2\xi}{1 - \xi^2}\right) \right| \rightarrow |2\xi + \mathcal{O}(\xi^2)|, \end{aligned} \quad (19)$$

so that the first term of the L.H.S of Eq. (16) vanishes as $\lambda \rightarrow \infty$. Thus, the first term of the L.H.S of Eq. (16) vanishes as $\lambda \rightarrow \infty$ regardless the limiting values of ξ and μ_2 . However, the second term of the L.H.S of Eq. (16) is

always non-positive while the R.H.S of Eq. (16) converges to 2 as $\lambda \rightarrow \infty$. Thus, Eq. (16) cannot be satisfied in the limit of large λ whatever the limiting values of ξ and μ_2 take, so that the eigenmode does not exist in the limit of large oscillation.

3.2. Instability condition

In following of the text, we assume the imaginary part of eigenvalue is zero, i.e., $\mu_2 = 0$, and consider a linear instability condition, i.e., $\mu_1 > 0$, under this assumption. Because $\mu_2 = 0$ satisfies Eq. (17), we only consider Eq. (16) with $\mu_2=0$, i.e.,

$$(\alpha\xi - \beta) \frac{\arctan(\xi)}{\xi} = 1 - \beta. \quad (20)$$

We can write μ_1 as a function of ξ , i.e.,

$$\mu_1 = \frac{\lambda}{\xi} - \frac{1}{k}, \quad (21)$$

so that μ_1 takes a positive value if and only if $0 < \xi < k\lambda$. Thus, the instability condition, i.e., $\mu_1 > 0$, is equivalent to the condition that Eq. (20) has a solution with $0 < \xi < k\lambda$.

First, we consider Eq. (20) with $\beta = 1$, which is obtained at $\lambda = \sqrt{\frac{1}{d}(F'[0]/k - 1)}$ when $F'[0]/k \geq 1$. It is immediate that Eq. (20) gives $\mu_1 = -1$, so that $\beta = 1$ does not hold the instability condition. (We also note that $\mu_1 = -1$ is also obtained as $\lambda=0$.)

For $\beta \neq 1$, we consider the condition that the intersection of the following functions,

$$\phi(\xi) = \frac{\arctan(\xi)}{\xi}, \quad (22)$$

and

$$\psi(\xi) = \frac{1 - \beta}{\alpha\xi - \beta}, \quad (23)$$

exists in $0 < \xi < k\lambda$. Note that $\phi(\xi)$ is monotonically decreasing for $\xi \geq 0$ and $\phi(0) = 1$ and $\phi(+\infty) = 0$, while the behavior of $\psi(\xi)$ depends on the coefficients α and β




Figure 2: An auxiliary figure for the analysis below Eqs. (22) and (23). In figure (a), $\alpha = 1$ and $\beta = 1.1$ is used for Case I. In figure (b), $\alpha = 5$ and $\beta = 0.5$ is used for Case II and $\alpha = -5$ and $\beta = 2$ is used for Case IV. The vertical dotted lines show the asymptotic lines $\xi = \beta/\alpha$ for each case.

Case I: $\alpha > 0$ and $\beta > 1$

The coefficient of $(\xi - \beta/\alpha)^{-1}$ in Eq. (23) is negative, so that $\psi(\xi)$ takes positive values only when ξ is smaller than the asymptotic line, $\xi = \beta/\alpha$. See also Fig. 2(a). Thus, we only consider the intersection of $\phi(\xi)$ and $\psi(\xi)$ in $0 < \xi < \beta/\alpha$. It is found that $\psi(\xi)$ monotonically increases in $0 < \xi < k\lambda$ because the asymptotic line, $\xi = \beta/\alpha$, is always larger than $\xi = k\lambda$, because $\frac{\beta}{\alpha} > \frac{1}{\alpha} = \frac{k\lambda}{1-k} > k\lambda$. It is also found that $\psi(\xi)$ is always smaller than $\phi(\xi)$ at $\xi = 0$, because $\psi(0) = 1 - \frac{1}{\beta} < 1$. Thus, the condition that $\phi(\xi)$ and $\psi(\xi)$

intersect in $0 < \xi < k\lambda$ is given by

$$\phi(k\lambda) < \psi(k\lambda). \quad (24)$$

Case II: $\alpha > 0$ and $0 < \beta < 1$

The coefficient of $(\xi - \beta/\alpha)^{-1}$ in Eq. (23) is positive, so that $\psi(\xi)$ takes positive values only when $\xi > \beta/\alpha$. It is obvious that if the asymptotic line $\xi = \beta/\alpha$ is larger than or equal to $\xi = k\lambda$, $\phi(\xi)$ and $\psi(\xi)$ does not intersect in $0 < \xi < k\lambda$. See also Fig. 2(b). Thus, we only consider the case $\beta/\alpha < k\lambda$, in which $\psi(\xi)$ is positive and monotonically decreases in $\beta/\alpha < \xi < k\lambda$. However, $\psi(\xi)$ is always larger than the unity in this range. This is explained as follows.

From the condition $\beta/\alpha < k\lambda$ and Eq. (18a), we obtain

$$0 < \frac{k}{1-\beta} < 1. \quad (25)$$

Since $\psi(\xi)$ monotonically decreases in $\beta/\alpha < \xi < k\lambda$, we get

$$\psi(\xi) > \psi(k\lambda) = \frac{1}{1 - \frac{k}{1-\beta}} > 1. \quad (26)$$

Thus, we can conclude that $\phi(\xi)$ and $\psi(\xi)$ does not intersect in $0 < \xi < k\lambda$ in this case.

Case III: $\alpha = 0$, i.e., $k = 1$.

In this case, $\psi(\xi) = 1 - \frac{1}{\beta}$ is constant, so that the condition that $\phi(\xi)$ and $\psi(\xi)$ intersect in $0 < \xi < k\lambda$ is given by Eq. (24).

Case IV: $\alpha < 0$ and $\beta > 1$.

The coefficient of $(\xi - \beta/\alpha)^{-1}$ in Eq. (23) is positive and the asymptotic line $\xi = \beta/\alpha$ is in the negative region, so that $\psi(\xi)$ takes finite positive values

and monotonically decreases in $0 < \xi < k\lambda$. See also Fig. 2(b). In addition, $\psi(\xi)$ is smaller than $\phi(\xi)$ at $\xi = 0$, because $\psi(0) = 1 - 1/\beta < 1$. Thus, the condition $\phi(\xi)$ and $\psi(\xi)$ intersect in $0 < \xi < k\lambda$ is given by Eq. (24).

Case V: $\alpha < 0$ and $0 < \beta < 1$.

The coefficient of $(\xi - \beta/\alpha)^{-1}$ in Eq. (23) is negative and the asymptotic line $\xi = \beta/\alpha$ is in the negative region, so that $\psi(\xi)$ is always negative in $0 < \xi < k\lambda$. Thus, $\phi(\xi)$ and $\psi(\xi)$ does not intersect in $0 < \xi < k\lambda$.

In summary, the condition that $\phi(\xi)$ and $\psi(\xi)$ intersect in $0 < \xi < k\lambda$ is solely given by Eq. (24). Thus, the instability condition of the perturbation with mode λ for Eqs. (1)–(4) is written as

$$\frac{F'[0]}{k} > \left[1 + \frac{k}{\frac{k\lambda}{\arctan(k\lambda)} - 1} \right] (1 + d\lambda^2). \quad (27)$$

We also remark that the auxiliary condition $\beta > 1$, i.e., $0 < \lambda < \sqrt{(F'[0]/k - 1)/d}$, is automatically satisfied under the above condition. Thus, when $F'[0]/k$ exceeds the minimum of the right-hand side of Eq. (27) with variation in $0 < \lambda < \sqrt{(F'[0]/k - 1)/d}$, the stationary homogeneous state in the population density becomes linearly unstable. Furthermore, stationary periodic patterns are generated because no unstable eigenmodes exist in the limit of large λ .

3.3. Continuum limit

We introduce a small parameter ε and scale k , x , and d in Eqs. (1) and (2) as follows,

$$k = \varepsilon^2, \quad x = \varepsilon \hat{x}, \quad d = \varepsilon^2 \hat{d}. \quad (28)$$

We also suppose the gradient of the response function is scaled as

$$F'[0] = \varepsilon^2 \hat{F}'[0]. \quad (29)$$

Then, by the asymptotic analysis of small ε for Eqs. (1) and (2), we obtain the following continuum equations in the limit $\varepsilon \rightarrow 0$,

$$\partial_t \rho_0 + \partial_{\hat{x}} (U[\log S_0] \rho_0) = \frac{1}{3} \partial_{\hat{x}\hat{x}} \rho_0 + P[\rho_0], \quad (30)$$

and

$$-d \partial_{\hat{x}\hat{x}} S_0 + S_0 = \rho_0, \quad (31)$$

where $U[\log S]$ is defined as

$$U[\log S_0] = \int_0^1 v \hat{F}[v \partial_{\hat{x}} \log S_0] dv. \quad (32)$$

When we carry out the linear instability analysis for Eqs. (31)–(32) as in the previous subsection, we obtain the growth rate μ_1 as

$$\mu_1 = -1 + \frac{\hat{F}'[0] \hat{\lambda}^2}{3(1 + d \hat{\lambda}^2)} - \frac{1}{3} \hat{\lambda}^2, \quad (33)$$

where $\hat{\lambda}$ is the Fourier variable as to \hat{x} , i.e., $\hat{\lambda} = \varepsilon \lambda$. Eq. (33) takes the maximum at $\hat{\lambda}^2 = (\sqrt{\hat{F}'[0]} - 1)/d$, and the sign of the maximum value of the growth rate μ_1 determines the linear stability. Thus, the instability condition for Eqs. (30)–(32) is written as

$$\hat{F}'[0] > (1 + \sqrt{3d})^2. \quad (34)$$

The above equation is consistent with the instability condition of the Keller-Segel system obtained earlier by Nadin, *et. al.*, Ref. [10]. We also note that Eq. (34) can be obtained by expanding Eq. (7) as to ε under the same scaling of Eq. (28) and taking the limit $\varepsilon \rightarrow 0$.

4. Numerical analysis

The numerical simulations are performed for Eqs. (1)–(3) with the uniform initial density, i.e., $f(0, x, v) = S(0, x) = 1$, and periodic boundary condition in the one-dimensional interval $x = [0, L]$, i.e., $f(t, 0, v) = f(t, L, v)$ and $S(t, 0) = S(t, L)$. The kinetic equation Eq. (1) is solved by the Monte Carlo (MC) method, which has been recently developed in Ref. [40], coupled with the finite volume (FV) scheme for Eq. (2). The details of the MC method and FV scheme is presented in Sec. 5. In the MC simulations, the one-dimensional interval L is set as $L=100$ and divided into 2000 cubic lattice boxes with a side length $\Delta x=0.05$. The 1×10^7 simulation particles are used initially as a total in the whole lattice boxes, and they are distributed randomly into each lattice box. See also Fig. 5. The time-step size Δt is set as $\Delta t=5 \times 10^{-3}$. In the previous study, it has been verified that these simulation parameters produce accurate numerical solutions. Eq. (2) is discretised using the FV scheme on the uniform lattice mesh system with a mesh interval Δx and solved implicitly at each time step.

For the population-growth rate $P[\rho]$ and response function $F[X]$, which satisfies Eq. (4) and Eq. (6), respectively, we consider

$$P[\rho] = 1 - \rho, \quad (35)$$

and

$$F[X] = \chi \tanh\left(\frac{X}{\delta}\right). \quad (36)$$

Here, χ and δ^{-1} represent the amplitude of modulation and stiffness of the response function, respectively. Note that for Eq. (36), $F'[0]$ is given as $F'[0] = \chi/\delta$. The values of parameters χ , δ , and d are set so as to correspond

to the marks A, B, C in Fig. 1 with four values of k , i.e., $k=0.1, 1, 2$, and 10 . More specifically, we set the parameters as $(\chi/\sqrt{k}, \sqrt{k}\delta, d/k)=(0.5, 0.05, 1)$ for A, $(0.5, 0.0625, 1)$ for B, $(0.5, 0.0625, 0.7)$ for C, and $(0.5, 0.1, 1)$ for D. We perform the MC simulations for the parameters listed in Table 1 and investigate the compatibility and sharpness of the kinetic instability condition Eq. (7) numerically. Table 1 also shows the prediction of the linear instability by Eq. (7). For example, the parameter set C with $k = 1.0$ is slightly above the critical line in Fig. 1, so that the kinetic instability condition Eq. (7) affirms the occurrence of periodic pattern formation. However, for the parameter set C with $k = 2.0$, which is slightly below (but very close to) the critical line in Fig. 1, we cannot confirm neither the linear instability nor homogeneous state theoretically because Eq. (7) is a sufficient condition of the linear instability. However, the MC simulation can numerically demonstrate how sharply the kinetic instability condition Eq. (7) can predict the pattern formations.

Figure 3 shows the time progress of the population density ρ . It is obviously seen that the results for the black squares in Table 1 exhibit the stationary periodic patterns after some transient period. On the other hand, those for the white squares in Table 1 do not show any distinct patterns. In order to quantify the patterns, we also calculate the power spectra of the Fourier transforms of population density profiles between $t=[400,500]$. Figure 4 shows the results of the power spectra. Here, we take the averages of the snap shots of power spectra obtained at $t=[400,500]$ with a time interval 4. The power spectrum for the continuum equations Eqs. (30)–(32) is calculated from the snap shot of the population density at $t = 200$. For the




Figure 3: The time progress of the population densities. Figure (a) and (b) correspond to the parameter set A with $k=1.0$ and 2.0 , respectively, Figure (c) and (d) to the parameter set B with $k=0.1$ and 1.0 , respectively, and Figure (e) and (f) to the parameter set C with $k=1.0$ and 2.0 , respectively. See also Table 1.

k	A(1, 0.5, 0.05)	B(1, 0.5, 0.0625)	C(0.7, 0.5, 0.0625)	D(1, 0.5, 0.1)
0.1	—	■	—	—
1.0	■	□	■	□
2.0	□	—	□	—

Table 1: The parameter sets with which the Monte Carlo simulations are performed. The parameter sets A, B, C, and D corresponds to the marks A, B, C, and D in Fig. 1, respectively. The values of the round brackets show the values of d/k , χ/\sqrt{k} , and $\sqrt{k}\delta$, i.e., $(d/k, \chi/\sqrt{k}, \sqrt{k}\delta)$. The black squares satisfy the kinetic instability condition Eq. (7) while white squares do not satisfy it.




Figure 4: The power spectra of the Fourier transform of population density profiles, $|\hat{\rho}(\lambda)|^2/k$. Figure (a) shows the results for the parameter sets A with $k = 1.0$, B with $k = 0.1$, and C with $k = 1.0$ (the black squares ■) in Table 1 and figure (b) shows those for A with $k = 2.0$, B with $k = 1.0$, C with $k = 2.0$, and D with $k = 1$ (the white squares □) in Table 1. In the figure (a), the result of the continuum equations Eqs. (30)–(32) with the parameter set B in Table 1 (the dotted line) is also included.

parameter sets shown as the black squares ■ in Table 1 (See Fig. 4(a)), steep peaks are observed and the first peaks appear in $0 < \hat{\lambda} < \sqrt{(\hat{F}'[0] - 1)/\hat{d}}$ for each parameter set. The second and third peaks also appear as the non-linear effects although they are much smaller than the first peaks. The power spectra decreases as the wave number $\hat{\lambda}$ decreases from the first-peak position, while they neither grow nor damp at the large wave number, so that a plateau regime appears at the large wave numbers. The peaks of the power spectrum for the continuum equations with the parameter set B coincide with those for the parameter set B with $k = 0.1$. However, no plateau regime appears for the continuum equations. This result is compatible with the fact that there are no eigenmodes obtained in the linearized kinetic equation.

For the parameter sets shown as the white squares □ in Table 1 (See Fig. 4(b)), the behaviors of power spectra are similar to those for the black squares in Table 1, i.e., Fig. 4(a), except the peak behaviors. In Fig. 4(b), we cannot see the steep first peaks as seen in Fig. 4(a). The small peaks of A, B, and C in Fig. 4(b) are even smaller than the second peaks appearing in the figure (a). For the parameter set C, we cannot see any peaks. Thus, we cannot observe the stationary periodic patterns evidently in Fig. 3 for the parameter sets shown as white squares in Table 1.

The parameter set A and B are very close to the critical lines for $k=2.0$ and $k=0.1$ in Fig. 1, respectively. (The parameter set A is only slightly lower and B is only slightly upper than each critical line.) However, the result of the parameter set B shows the stationary periodic pattern evidently but the result of the parameter set A does not show it. These numerical results demonstrate that the critical lines in the instability diagram Fig. 1 can predict




Figure 5: The geometry in the Monte Carlo method. The concentrations of chemoattractant are calculated by a finite-volume method on the uniform cubic lattice boxes. The Monte Carlo particles are distributed in each lattice boxes. The periodic boundary conditions are considered both for the MC particles and chemoattractant in this paper.

sharply the occurrence of the linear instability and pattern formations.

5. Monte Carlo method

The motions of the chemotactic bacteria are simulated by using the Monte Carlo particles which follows the process described by the kinetic chemotaxis equation Eq. (1) coupled with the reaction-diffusion equation for the chemoattractant Eq. (2). The one-dimensional space interval $x \in [0, L]$ is divided into I cubic lattices with a uniform side length Δx , i.e., $L = I\Delta x$. See also Fig. 5. The reaction-diffusion equation Eq. (2) is implicitly solved on the uniform lattice system by using a finite volume scheme as

$$-\frac{d}{\Delta x^2}(S_{i+1}^n - 2S_i^n + S_{i-1}^n) + S_i^n = \rho_i^n, \quad (i = 0, \dots, I-1), \quad (37)$$

where S_i^n and ρ_i^n are the concentration of chemical attractant and population density of bacteria in the i th lattice site $[i\Delta x, (i+1)\Delta x] \times \Delta x^2$ at a time $t = n\Delta t$, respectively. Here we also set $S_{-1}^n = S_{I-1}^n$ and $S_I^n = S_0^n$ according to the periodic condition. Hereafter the superscript represents the time-step

number, the subscript without parenthesis represents the lattice-site number, and the subscript in parenthesis represents the index of each MC particle. The population density of bacteria ρ_i^n is calculated from the number of the MC particles involved in the i th lattice site, M_i^n , as

$$\rho_i^n = M_i^n / M, \quad (38)$$

where M is the number of MC particles involved in one lattice site in the reference state, i.e, $M = N/I$, where N is the total number of MC particles in the initial state.

The MC simulation is conducted using the following steps. Hereafter, the position and velocity of the l th particle are expressed as $\mathbf{r}_{(l)}^n$ and $\mathbf{v}_{(l)}^n$, respectively.

0. At $\hat{t} = 0$, MC particles are distributed according to the initial density. In each lattice site, MC particles are distributed uniformly at random positions and their velocities \mathbf{v} are determined by the probability density $f_i^0(\mathbf{v})/(4\pi\rho_i^0)$.
1. Particles move with their velocities for a duration $\Delta\hat{t}$:

$$\mathbf{r}_{(l)}^{n+1} = \mathbf{r}_{(l)}^n + \mathbf{v}_{(l)}^n \Delta\hat{t} \quad (l = 1, \dots, N^n), \quad (39)$$

where N^n is the total number of simulation particles at a time step n , i.e., $N^n = \sum_{i=0}^{I-1} M_i^n$. The particles that move beyond the boundaries are inserted at the opposite boundaries according to the periodic boundary conditions.

2. At each lattice site, the macroscopic population densities ρ_i^{n+1} and concentrations of chemical cues S_i^{n+1} ($i = 0, \dots, I - 1$) are calculated by Eqs. (38) and (37), respectively.
3. The tumbling of each particle is calculated using the scattering kernel in Eq. (1). The tumbling of the l th particle may occur with a probability

$$\frac{\Delta t}{k} K[D_t \log S_{(l)}^{n+1}], \quad (40)$$

where $D_t \log S_{(l)}$ represents the temporal variation of the chemical cue experienced by the l th MC particle along the pathway, and is defined by the following forward difference,

$$D_t \log S_{(l)}^n = \frac{\log S(t^n, \mathbf{r}_{(l)}^n) - \log S(t^n - \Delta t, \mathbf{r}_{(l)}^n - \mathbf{v}_{(l)}^n \Delta t)}{\Delta t}. \quad (41)$$

The local concentration of chemical cue at the position of the l th MC particle is calculated by using linear interpolation between the neighboring lattice sites, i.e.,

$$\log S(t^n, \mathbf{r}_{(l)}^n) = \log S_i^n + \frac{\log S_{i+1}^n - \log S_{i-1}^n}{2\Delta x} (r_{x(l)}^n - x_{i+\frac{1}{2}}). \quad (42)$$

This generates a chemoattractant gradient and MC particles that stay at the same lattice site after a single time step can sense the chemoattractant gradient along their pathways.

For the particle that is judged to tumble, say the l_t th particle, a new velocity after the tumbling, $\mathbf{v}_{(l_t)}^{n+1}$, is determined randomly as,

$$v_x = 1 - 2U_1, \quad v_y = \sqrt{1 - v_x^2} \cos(2\pi U_2), \quad v_z = \sqrt{1 - v_x^2} \sin(2\pi U_2). \quad (43)$$

Here U_1 and U_2 are the uniform random variables between 0 and 1.

4. The divisions/deaths are judged for all MC particles. The division (or death) occurs with a probability $|P[\rho_i^n]|\Delta t$, if $|P[\rho_i^n]|$ is positive (or negative), where ρ_i^n is the local population density at the lattice site where each MC particle is involved. For a particle that is judged to undergo division, e.g., the l th particle, a new particle with the same velocity $\mathbf{v}_{(l)}$ is created at a random position within the same lattice site. The numbers of MC particles involved in each lattice site are counted, M_i^{n+1} ($i = 0, \dots, I - 1$), and the total number of simulation particles is updated as N^{n+1} .
5. Return to step 1 with the obtained $\mathbf{r}_{(l)}$, $\mathbf{e}_{(l)}$, $S_{(l)}$ ($l=1, \dots, M$) at the new time step.

Weak formulation The overall procedure corresponds to the first-order time difference equation of the kinetic chemotaxis equation in the weak formulation. We consider the following moment equation,

$$\begin{aligned}
& \langle \Phi(\mathbf{x}, \mathbf{v}), f(t + \Delta t, \mathbf{x}, \mathbf{v}) \rangle = \langle \Phi(\mathbf{x}, \mathbf{v}), f(t, \mathbf{x}, \mathbf{v}) - \Delta t \mathbf{v} \cdot \nabla f(t, \mathbf{x}, \mathbf{v}) \rangle \\
& + \langle \Phi(\mathbf{x}, \mathbf{v}), \frac{1}{4\pi k} \int_V K[D_t \log S|\mathbf{v}'] f(t, \mathbf{x}, \mathbf{v}') d\Omega(\mathbf{v}') - \frac{1}{k} K[D_t \log S|\mathbf{v}] f(t, \mathbf{x}, \mathbf{v}) \rangle \Delta t \\
& + \langle \Phi(\mathbf{x}, \mathbf{v}), P[\rho(t, \mathbf{x})] f(t, \mathbf{x}, \mathbf{v}) \rangle \Delta t, \tag{44}
\end{aligned}$$

where $\Phi(\mathbf{x}, \mathbf{v})$ is an arbitrary smooth function which vanishes outside the computational domain on \mathbf{x} .

Here $\langle \cdot, \cdot \rangle$ defines the integration of the arbitrary functions $a(\mathbf{x}, \mathbf{v})$ and $b(\mathbf{x}, \mathbf{v})$ as

$$\langle a(\mathbf{x}, \mathbf{v}), b(\mathbf{x}, \mathbf{v}) \rangle = \int_{V, \mathbb{R}} a(\mathbf{x}, \mathbf{v}) b(\mathbf{x}, \mathbf{v}) d\mathbf{x} d\Omega(\mathbf{v}). \tag{45}$$

We consider the functions f^A , f^B , and f^C which are determined, respectively, as

$$\langle \Phi, f^A \rangle = \langle \Phi(\mathbf{x}, \mathbf{v}), f(t, \mathbf{x}, \mathbf{v}) - \Delta t \mathbf{v} \cdot \nabla f(t, \mathbf{x}, \mathbf{v}) \rangle, \quad (46a)$$

$$\begin{aligned} \langle \Phi, f^B \rangle = & \langle \Phi(\mathbf{x}, \mathbf{v}), \frac{1}{4\pi k} \int_V K[D_t \log S|_{\mathbf{v}}] f(t, \mathbf{x}, \mathbf{v}') d\Omega(\mathbf{v}') \\ & - \frac{1}{k} K[D_t \log S|_{\mathbf{v}}] f(t, \mathbf{x}, \mathbf{v}) \rangle \Delta t, \end{aligned} \quad (46b)$$

$$\langle \Phi, f^C \rangle = \langle \Phi(\mathbf{x}, \mathbf{v}), P[\rho(t, \mathbf{x})] f(t, \mathbf{x}, \mathbf{v}) \rangle \Delta t. \quad (46c)$$

Then, $f(t + \Delta t, \mathbf{x}, \mathbf{v})$ is obtained by the sum of three functions:

$$f(t + \Delta t, \mathbf{x}, \mathbf{v}) = f^A + f^B + f^C. \quad (47)$$

In the MC method, the microscopic population density f at the n th time step is approximated as

$$f^n(\mathbf{x}, \mathbf{v}) = \frac{4\pi(\Delta x)^3}{M} \sum_{l=1}^{N^n} \delta(\mathbf{x} - \mathbf{r}_{(l)}^n) \delta(\mathbf{v} - \mathbf{v}_{(l)}^n). \quad (48)$$

By substituting Eq. (48) into Eq. (46a) we obtain the following equation:

$$\begin{aligned} \langle \Phi, f^A \rangle &= \sum_{l=1}^{N^n} \Phi(\mathbf{r}_{(l)}^n, \mathbf{v}_{(l)}^n) + \Delta t \mathbf{v} \cdot \nabla \Phi(\mathbf{r}_{(l)}^n, \mathbf{v}_{(l)}^n) \\ &= \sum_{l=1}^{N^n} \Phi(\mathbf{r}_{(l)}^n + \mathbf{v}_{(l)}^n \Delta t, \mathbf{v}_{(l)}^n) + \mathcal{O}(\Delta t^2). \end{aligned} \quad (49)$$

It is easily seen that the function f^A corresponds to the distribution that is obtained by the moving process, say \mathcal{S}_1 in the above MC method, i.e.,

$$\mathcal{S}_1\{f^n\} = f^A, \quad (50)$$

where $\mathcal{S}_1\{ \ }$ represents the operator of the process 1 in the MC method.

The tumbling process and division/death process in the MC method, say \mathcal{S}_3 and \mathcal{S}_4 respectively, are performed independently in each lattice site. Thus, we consider the processes in a fixed lattice site, say the i th lattice site, with using a test function written as $\Phi(\mathbf{x}, \mathbf{v}) = \phi_i(\mathbf{x})\psi(\mathbf{v})$, where $\phi_i(\mathbf{x})$ is an arbitrary smooth function which vanishes outside the i th lattice site. Then, by substituting Eq. (48) into Eqs. (46b) and (46c) we obtain the following equations:

$$\begin{aligned} \langle \phi_i(\mathbf{x})\psi(\mathbf{v}), f^B \rangle = & \frac{\Delta t}{kM} \sum_{m=1}^{M_i^n} K [D_t \log S_{(m)}^n] \phi_i(\mathbf{r}_{(m)}^n) (\bar{\psi} - \psi(\mathbf{v}_{(m)}^n)) \\ & + \mathcal{O}(\Delta t^2), \end{aligned} \quad (51)$$

$$\begin{aligned} \langle \phi_i(\mathbf{x})\psi(\mathbf{v}), f^C \rangle = & \frac{\Delta t}{M} \sum_{m=1}^{M_i^n} P[\rho(\mathbf{r}_{(m)}^n)] \phi_i(\mathbf{r}_{(m)}^n) \psi(\mathbf{v}_{(m)}^n), \\ = & \frac{\Delta t P[\rho_i^n]}{M} \sum_{m=1}^{M_i^n} \bar{\phi}_i \psi(\mathbf{v}_{(m)}^n) + \mathcal{O}(\Delta t \Delta x), \end{aligned} \quad (52)$$

where the subscript m ($m = 1, \dots, M_i^n$) counts the MC particles involved in the i th lattice site at the n th time step. Here $D_t \log S_{(m)}^n$ is defined in Eq. (41), $\bar{\psi} = \frac{1}{4\pi} \int \psi(\mathbf{v}) d\Omega(\mathbf{v})$, and $\bar{\phi}_i = \phi(\mathbf{x}_{i+\frac{1}{2}})$ where $\mathbf{x}_{i+\frac{1}{2}}$ represents the center of the i th lattice site. In the derivation of Eq. (51), we approximate the tumbling kernel as

$$K[D_t \log S|_{\mathbf{v}}] = K \left[\frac{\log S(t, \mathbf{x}) - \log S(t - \Delta t, \mathbf{x} - \mathbf{v}\Delta t)}{\Delta t} \right] + \mathcal{O}(\Delta t). \quad (53)$$

In the tumbling process \mathcal{S}_3 , the particle which creates the tumbling, say the m_t th particle, changes its velocity randomly as $\mathbf{v}_{m_t}^n \rightarrow \mathbf{v}_{m_t}^{n+1} = \mathbf{u}_t$, where the random velocity \mathbf{u}_t is given by Eq. (43). We now suppose that M_t

particles make the tumbling in the i th lattice site, then the microscopic population density f^n in the i th lattice site changes as

$$\mathcal{S}_3\{f^n\} = \frac{1}{M} \left\{ \sum_{\substack{m=1 \\ m \neq m_t}}^{M_i^n} \delta(\mathbf{x} - \mathbf{r}_{(m)}^n) \delta(\mathbf{v} - \mathbf{v}_{(m)}^n) + \sum_{t=1}^{M_t} \delta(\mathbf{x} - \mathbf{r}_{(m_t)}^n) \delta(\mathbf{v} - \mathbf{u}_t) \right\}, \quad (54)$$

and the moment is written as

$$\begin{aligned} & \langle \phi_i(\mathbf{x}) \psi(\mathbf{v}), \mathcal{S}_3\{f^n\} \rangle \\ &= \frac{1}{M} \left\{ \sum_{m=1}^{M_i^n} \phi_i(\mathbf{r}_{(m)}^n) \psi(\mathbf{v}_{(m)}^n) + \sum_{t=1}^{M_t} \phi_i(\mathbf{r}_{(m_t)}^n) [\psi(\mathbf{u}_t) - \psi(\mathbf{v}_{(m_t)}^n)] \right\}. \end{aligned} \quad (55)$$

We introduce the stochastic variables Z_m and Z which are defined as

$$Z_m = \begin{cases} \phi_i(\mathbf{r}_{(m)}) [\psi(\mathbf{u}) - \psi(\mathbf{v}_{(m)})], & \text{(if tumbling),} \\ 0, & \text{(otherwise),} \end{cases} \quad (56)$$

and

$$Z = \frac{1}{M} \sum_{m=1}^{M_i^n} Z_m. \quad (57)$$

If the tumbling occurs for M_t particles, the realized value of Z is written as

$$\hat{Z} = \frac{1}{M} \sum_{t=1}^{M_t} \phi_i(\mathbf{r}_{m_t}) [\psi(\mathbf{u}_t) - \psi(\mathbf{v}_{m_t})] = \langle \phi_i(\mathbf{r}) \psi(\mathbf{v}), \mathcal{S}_3\{f^n\} - f^n \rangle. \quad (58)$$

Here we use Eq. (55). On the other hand, the expected value of Z_m is written as

$$E_p(Z_m) = \phi_i(\mathbf{r}_m) [\bar{\psi} - \psi(\mathbf{v}_{(m)})] \frac{\Delta t}{k} K[D_t \log S_{(m)}], \quad (59)$$

and the expected value of Z is written by the moment Eq. (51) as

$$E_p(Z) = \langle \phi_i(\mathbf{x}) \psi(\mathbf{v}), f^B \rangle + \mathcal{O}(\Delta t^2). \quad (60)$$

From Eqs. (58) and (60), it is seen that the microscopic population density obtained by the tumbling process on f^n , $\mathcal{S}_3\{f^n\}$ is approximated by the sum of f^n and f^B under the assumption of the law of large numbers, i.e., $\hat{Z} \simeq E_p(Z)$,

$$\mathcal{S}_3\{f^n\} = f^n + f^B. \quad (61)$$

In the division/death process \mathcal{S}_4 , the particles in the i th lattice site may divide (or die) if the local macroscopic population density ρ_i is smaller (or larger) than unity. In the divisions, where $P[\rho_i] > 0$, the particle creates a new particle with the same velocity at a random position within the same lattice site. Thus, when the divisions occur for M_c particles for the distribution Eq. (48), the microscopic population density in the i th lattice site changes as

$$\begin{aligned} \mathcal{S}_4\{f^n\} = & \\ \frac{1}{M} & \left\{ \sum_{m=1}^{M_i^n} \delta(\mathbf{x} - \mathbf{r}_{(m)}^n) \delta(\mathbf{v} - \mathbf{v}_{(m)}^n) + \sum_{c=1}^{M_c} \delta(\mathbf{x} - (\mathbf{x}_{i+\frac{1}{2}} + \Delta x \mathbf{w}_c)) \delta(\mathbf{v} - \mathbf{v}_{(m_c)}^n) \right\}, \end{aligned} \quad (62)$$

where \mathbf{w} is a random vector whose components are uniform random numbers in $[-\frac{1}{2}, \frac{1}{2}]$. The moment is written as

$$\begin{aligned} & \langle \phi_i(\mathbf{x}) \psi(\mathbf{v}), \mathcal{S}_4\{f^n\} \rangle \\ & = \frac{1}{M} \left\{ \sum_{m=1}^{M_i^n} \phi_i(\mathbf{r}_{(m)}^n) \psi(\mathbf{v}_{(m)}^n) + \sum_{c=1}^{M_c} \phi_i(\mathbf{x}_{i+\frac{1}{2}} + \Delta x \mathbf{w}_c) \psi(\mathbf{v}_{(m_c)}^n) \right\}. \end{aligned} \quad (63)$$

In deaths, where $P[\rho_i] < 0$, the particle is just removed from the lattice site. Thus, when the deaths occurs for M_c particles in the i th lattice site, the microscopic population density obtained after the death process and its

moment are written as replacing the sign of the second term with minus and $\mathbf{x}_{i+\frac{1}{2}} + \Delta x \mathbf{v}$ with $\mathbf{r}_{(m)}^n$ in Eqs. (62) and (63), respectively. Hence, the moment for the population density after the division/death process \mathcal{S}_4 can be written as

$$\begin{aligned} & \langle \phi_i(\mathbf{x})\psi(\mathbf{v}), \mathcal{S}_4\{f^n\} \rangle \\ &= \frac{1}{M} \left\{ \sum_{m=1}^{M_i^n} \phi_i(\mathbf{r}_{(m)}^n)\psi(\mathbf{v}_{(m)}^n) + \text{sign}(P[\rho_i]) \sum_{c=1}^{M_c} \bar{\phi}_i\psi(\mathbf{v}_{(m_c)}^n) \right\} + \mathcal{O}\left(\frac{M_c}{M}\Delta x\right). \end{aligned} \quad (64)$$

We remark that M_c/M is estimated as $M_c/M \sim \mathcal{O}(\Delta t)$.

We introduce the stochastic variable Ξ_m and Ξ which are defined as

$$\Xi_m = \begin{cases} \text{sign}(P[\rho_i])\bar{\phi}_i\psi(\mathbf{v}_{(m)}), & \text{(if division/death),} \\ 0 & \text{(otherwise),} \end{cases} \quad (65)$$

and

$$\Xi = \frac{1}{M} \sum_{m=1}^{M_i^n} \Xi_m. \quad (66)$$

If the divisions(or deaths) occur for M_c particles, the realized value of Ξ is written as

$$\hat{\Xi} = \frac{\text{sign}(P[\rho_i])}{M} \sum_{c=1}^{M_c} \bar{\phi}_i\psi(\mathbf{v}_{(m_c)}^n) = \langle \phi_i(\mathbf{x})\psi(\mathbf{v}), \mathcal{S}_4\{f^n\} - f^n \rangle + \mathcal{O}\left(\frac{M_c}{M}\Delta x\right). \quad (67)$$

On the other hand, the expected value of Ξ is written by Eq. (52) as

$$E_p(\Xi) = \frac{\Delta t P[\rho_i]}{M} \sum_{m=1}^{M_i^n} \bar{\phi}_i\psi(\mathbf{v}_{(m)}^n) = \langle \phi_i(\mathbf{x})\psi(\mathbf{v}), f^C \rangle + \mathcal{O}(\Delta t \Delta x). \quad (68)$$

Thus, the microscopic population density obtained by the division/death process on f^n , $\mathcal{S}_4\{f^n\}$ is approximated by the sum of f^n and f^C under the

assumption of the law of large numbers, i.e., $\hat{\Xi} \simeq E_p(\Xi)$,

$$\mathcal{S}_4\{f^n\} = f^n + f^C. \quad (69)$$

In the MC simulation, the processes \mathcal{S}_1 , \mathcal{S}_3 , and \mathcal{S}_4 are successively conducted. For example, the process \mathcal{S}_2 is performed on the distribution obtained by the process \mathcal{S}_1 on f^n ;

$$\mathcal{S}_3\{\mathcal{S}_1\{f^n\}\} = \mathcal{S}_3\{f^A\} = f^A + f^{B'}, \quad (70)$$

where $f^{B'}$ is obtained by replacing f^n with f^A in Eq. (46b). However, the difference of f^A and f^n is $\mathcal{O}(\Delta t)$, so that $f^{B'}$ can be replaced with f^B within the difference of $\mathcal{O}(\Delta t^2)$. Similarly, it is seen that the microscopic population density obtained by the successive three processes of \mathcal{S}_1 , \mathcal{S}_3 , and \mathcal{S}_4 approximates the microscopic population density at the next time step which satisfies the weak formulation Eq. (44) within the difference of $\mathcal{O}(\Delta t^2)$, i.e.,

$$\mathcal{S}_4\{\mathcal{S}_3\{\mathcal{S}_1\{f^n\}\}\} = f^A + f^B + f^C + \mathcal{O}(\Delta t^2) = f^{n+1} + \mathcal{O}(\Delta t^2). \quad (71)$$

6. Critical analysis and concluding remarks

Departing from a classical kinetic chemotaxis model of bacterial motion [13, 14, 15], we have discovered a linear instability condition Eq. (7). The stationary homogeneous state of the macroscopic population density becomes linearly unstable and stationary periodic patterns are generated. It is also found that no eigenmodes exist in the large-oscillation limit in the linearized kinetic chemotaxis equation. MC simulations for the kinetic chemotaxis equation coupled with the diffusion-reaction equation are performed with changing the parameters involved in the linear instability condition as shown

ψ_0	3.0 [s]
t_0	4500 [s]
V_0	25 [$\mu\text{m/s}$]
\tilde{d}	8×10^{-6} [cm^2/s]
$\tilde{\delta}^{-1}$	20 [s]
χ	0.2

Table 2: Experimental values for mean tumbling frequency ψ_0 , doubling time t_0 , running speed V_0 , diffusion coefficient of chemoattractant \tilde{d} , and stiffness $\tilde{\delta}^{-1}$ and modulation χ in chemotactic response obtained in Ref. [33].

in Fig. 1. The numerical results demonstrate that the obtained linear instability condition is compatible and even sharply predict the occurrence of the periodic pattern formations. See Fig. 3. The power spectra of the macroscopic population density show the plateau regime at the large wave numbers, where the perturbations neither grow nor damp irrespective of the linear instability condition. See Fig. 4 This observation is compatible with the fact that no eigenmodes exist in the large oscillation limit in the linearized kinetic chemotaxis equation.

The obtained instability condition Eq. (7) includes three control parameters, i.e., k , d/k , and $F'[0]/k$, which are written in the dimensional form as

$$k = 1/(\psi_0 t_0), \quad d/k = \tilde{d}\psi_0/(V_0^2 t_0 b_0), \quad F'[0]/k = \chi \tilde{\delta}^{-1} \psi_0. \quad (72)$$

Here ψ_0 is the mean tumbling frequency, V_0 is the running speed of bacteria, b_0 is the degradation rate of chemoattractant, and t_0 is the characteristic time which corresponds to the doubling time in cell division for Eq. (35).

See also the paragraph above Eq. (7). \tilde{d} and $\tilde{\delta}^{-1}$ are the diffusion coefficient of chemoattractant and stiffness of response function, respectively, in the dimensional form. The values of control parameters are estimated from experimental data in Ref. [33] (See Table 2) as $k = 7.5 \times 10^{-5}$, $d/k = 3.8 \times (t_0 b_0)^{-1}$, and $F'[0]/k = 12$. It is difficult to measure the degradation rate of chemoattractant b_0 in experiments. In some references [35, 51, 52], the value of b_0 is estimated as $b_0 = 4 \times 10^{-3} \sim 5 \times 10^{-2}$ from the comparison between experimental and numerical results, so that we may estimate d/k as $d/k \lesssim 1$. Thus, from our analysis, we can expect the stationary homogeneous state becomes destabilized and pattern formation occurs for chemotactic bacteria. Furthermore, for example, in Ref. [2], it is argued that the pattern formation is suppressed by reduction of chemotactic sensitivity. This argument is also consistent with the obtained instability condition. Although it remains to be assessed how quantitatively the present instability condition explains the experimental results in terms of pattern formation, the present study convinces us that the self-organized pattern formation occurs due to the modulation of stiff response in chemotaxis in a realistic range of parameters.

However our study is based on a simplified model where the adaptation dynamics in chemotactic response to the extracellular environment is ignored and replaced by a instantaneous material derivative of chemoattractant along the pathway of bacterium. The time scale of adaptation, say τ_M , is larger than the inverse of tumbling rate ψ_0^{-1} but is much smaller than the doubling time t_0 , i.e., $\psi_0^{-1}/t_0 < \tau_M/t_0 \ll 1$ [49, 50]. Thus, the adaptation dynamics may be significant for the pattern formation. In order to consider the adaptation dynamics the memory or internal state has to be taken into account in

the tumbling kernel. [14, 28, 30] The present MC method has an important advantage in extending toward the adaptation dynamics of internal states because the MC method employs the particle-based method. In a forthcoming paper, we extend the MC method toward this direction and consider this issue.

Acknowledgements

This study was financially supported by JSPS KAKENHI Grant Number 15KT0110 and 16K17554 and Institut Henri Poincaré RIP program.

- [1] J. Adler, “Chemotaxis in bacteria”, *Annu. Rev. Biochem.* **44**, 341 (1975).
- [2] E. O. Budrene and H. C. Berg, “Complex patterns formed by motile cells of *Escherichia coli*”, *Nature* **349**, 630 (1991).
- [3] E. O. Budrene and H. C. Berg, “Dynamics of formation of symmetrical patterns by chemotactic bacteria”, *Nature* **376**, 49 (1995).
- [4] H. C. Berg, *E. Coli in Motion* (Springer, Berlin, 2003).
- [5] E. F. Keller and L. A. Segel, “Model for Chemotaxis”, *J. Theor. Biol.* **30**, 225 (1971).
- [6] E. F. Keller and L. A. Segel, “Traveling bands of chemotactic bacteria: a theoretical analysis”, *J. Theor. Biol.* **30**, 235 (1971).
- [7] M. J. Tindall, P. K. Maini, S. L. Porter, and J. P. Armitage, “Overview of mathematical approaches used to model bacterial chemotaxis II: Bacterial populations”, *Bull. Math. Biol.* **70**, 1570–1607 (2008).

- [8] P. K. Maini, M. R. Myerscough, K. H. Winters, and J. D. Murray, “Bifurcating spatially heterogeneous solutions in a chemotaxis model for biological pattern generation”, *Bull. Math. Biol.* **53**, 701–719 (1991).
- [9] M. Mimura and T. Tsujikawa, “Aggregating pattern dynamics in a chemotaxis model including growth”, *Physica A* **230**, 499–543 (1996).
- [10] G. Nadin, B. Perthame, and L. Ryzhik, “Traveling waves for the Keller-Segel system with Fisher birth terms”, *Interface free bound.* **10**, 517–538 (2008).
- [11] K. J. Painter and T. Hillen, “Spatio-temporal chaos in a chemotaxis model”, *Physica D* **240**, 363–375 (2011).
- [12] W. Alt, “Biased random walk models for chemotaxis and related diffusion approximations”, *J. Math. Biol.* **9**, 147 (1980).
- [13] H. G. Othmer, S. R. Dunbar, and W. Alt, “Models of dispersal in biological systems”, *J. Math. Biol.* **26**, 263 (1988).
- [14] R. Erban and H. G. Othmer, “From individual to collective behavior in bacterial chemotaxis”, *SIAM J. Appl. Math.* **65**, 361 (2004).
- [15] Y. Dolak and C. Schmeiser, “Kinetic models for chemotaxis: Hydrodynamic limits and spatio-temporal mechanisms”, *J. Math. Biol.* **51**, 595 (2005).
- [16] N. Bellomo, A. Bellouquid, J. Nieto, and J. Soler, “Multicellular biological growing systems: Hyperbolic limits towards macroscopic description”, *Math. Model Methods Appl. Sci.* **17**, 1675–1693 (2007).

- [17] N. Bellomo, N. K. Li, and P. K. Maini, “On the foundations of cancer modelling: selected topics, speculations, and perspectives”, *Math. Model Methods Appl. Sci.* **18**, 593–646 (2008).
- [18] B. Perthame, *Transport Equations in Biology* (Birkhäuser Verlag, Basel, 2007).
- [19] T. Hillen and H. G. Othmer, “The diffusion limit of transport equations derived from velocity-jump processes”, *SIAM J. Appl. Math.* **61**, 751 (2000).
- [20] H. Othmer and T. Hillen, “The diffusion limit of transport equations. II. Chemotaxis equations”, *SIAM J. Appl. Math.* **62**, 1222 (2002).
- [21] F. A.C.C. Chalub, P. Markowich, B. Perthame, and C. Schmeiser, “Kinetic Models for chemotaxis and their drift-diffusion limits”, *Monatsh. Math.* **142**, 123 (2004).
- [22] N. Bournaveas, V. Calvez, S. Gutiérrez and B. Perthame, “Global existence for a kinetic model of chemotaxis via dispersion and Strichartz estimates”, *Commun. Part. Diff. Eq.* **33**, 79–95 (2008).
- [23] N. Bournaveas and V. Calvez, “Global existence for the kinetic chemotaxis model without pointwise memory effects, and including internal variables”, *Kinet. Relat. Mod.* **1**, 29–48 (2008).
- [24] N. Vauchelet, “Numerical simulation of a kinetic model for chemotaxis”, *Kinet. Relat. Mod.* **3**, 501–528 (2010).

- [25] F. James and N. Vauchelet, “Chemotaxis: from kinetic equations to aggregate dynamics”, *Nonlinear Differ. Equ. Appl.* **20**, 101 (2013).
- [26] G. Si, M. Tang, and X. Yang, “A pathway-based mean-field model for *E. coli* chemo-taxis: mathematical derivation and keller-segel limit”, *Multiscale Model Simul* **12**, 907–926 (2014).
- [27] L. Almeida, C. Emako, N. Vauchelet, “Existence and diffusive limit of a two-species kinetic model of chemotaxis”, *Kinet. Relat. Models* **8**, 359 (2015).
- [28] C. Xue, “Macroscopic equations for bacterial chemotaxis: integration of detailed biochemistry of cell signaling”, *J. Math. Biol.* **70**, 1 (2015).
- [29] J. Liao, “Global solution for a kinetic chemotaxis model with internal dynamics and its fast adaptation limit”, *J. Differ. Equ.* **259**, 6432–6458 (2015).
- [30] B. Perthame, M. Tang, and N. Vauchelet, “Derivation of the bacterial run-and-tumble kinetic equation from a model with biochemical pathway”, *J. Math. Biol.* **73**, 1161 (2016).
- [31] V. Calvez, “Chemotactic waves of bacteria at the mesoscale”, [arXiv:1607.00429](https://arxiv.org/abs/1607.00429), (2016).
- [32] S. Mischler and Q. Weng, “On a linear runs and tumbles equation”, *Kinetic and Related Models* **10**, 799–822 (2017).
- [33] J. Saragosti, V. Calvez, N. Bournaveas, B. Perthame, A. Buguin, and

- P. Silberzan, “Directional persistence of chemotactic bacteria in a traveling concentration wave”, *PNAS* **108**, 16235 (2011).
- [34] G. Rosser, A. G. Fletcher, D. A. Wilkinson, J. A. de Beyer, C. A. Yates, J. P. Armitage, P. K. Maini, and R. E. Baker, “Novel methods for analysing bacterial tracks reveal persistence in *Rhodobacter sphaeroides*”, *PLoS Comput. Biol.* **9**, e1003276 (2013).
- [35] C. Emako, C. Gayraud, A. Buguin, L. Almeida, and N. Vauchelet, “Traveling pulses for a two-species chemotaxis model”, *PLoS Comput. Biol.* **12**, e1004843 (2016).
- [36] M. Rousset and G. Samaey, “Individual-based models for bacterial chemotaxis in the diffusion asymptotics”, *Math. Models Methods Appl. Sci.* **23**, 2005 (2013)
- [37] M. Rousset and G. Samaey, “Simulating individual-based models of bacterial chemotaxis with asymptotic variance reduction”, *Math. Models Methods Appl. Sci.* **23**, 2155 (2013).
- [38] C. Yang and F. Filbet, “Numerical simulations of kinetic models for chemotaxis”, *SIAM J. Scientific Computing* **36**, B348 (2014)
- [39] J. Ritter, A. Klar, and F. Schneider, “Partial-moment minimum-entropy models for kinetic chemotaxis equations in one and two dimensions”, *J. Comput. Appl. Math.* **306**, 300–315 (2016).
- [40] S. Yasuda, “Monte Carlo simulation for kinetic chemotaxis model: An application to the traveling population wave”, *J. Comput. Phys.* **330**, 1022–1042 (2017).

- [41] R. A. Fisher, “The advance of advantageous genes”, *Ann. Eugenics* **65**, 335–369 (1937).
- [42] A. N. Kolmogorov, I.G. Petrovsky, and N.S. Piskunov, “Étude de l’équation de la diffusion avec croissance de la quantité de matière et son application à un problème biologique”, *Moskow. Univ. Math. Bull.* **1**, 1–25 (1937).
- [43] R. Eftimie, “Hyperbolic and kinetic models for self-organized biological aggregations and movement: a brief review”, *J. Math. Biol.* **65**, 35–75 (2012).
- [44] K. Hadeler, “Reaction transport systems in biological modelling”, *Math. Comput. Model.* **31**, 75–81 (2000).
- [45] H. Schwetlick, “Travelling fronts for multidimensional nonlinear transport equations”, *Ann. Inst. Henri Poincaré, Analyse non linéaire* **17**, 523–550 (2000).
- [46] E. Bouin, V. Calvez, and G. Nadin, “Propagation in a kinetic reaction-transport equation: Travelling waves and accelerating”, *Arch. Rational Mech. Anal.* **217**, 571–617 (2015).
- [47] A. M. Turing, “The chemical basis of morphogenesis”, *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **237**, 37–72 (1952).
- [48] B. Perthame, *Parabolic equations in biology*, Springer (2015).
- [49] D. A. Clark and L. C. Grant, “The bacterial chemotactic response

reflects a compromise between transient and steady-state behavior”, PNAS **102**, 9150–9155 (2005).

- [50] X. Zhu, G. Si, N. Deng, Q. Ouyang, T. Wu, Z. He, L. Jiang, C. Luo, and Y. Tu, “Frequency-dependent *Escherichia coli* chemotaxis behavior”, Phys. Rev. Lett. **108**, 128101 (2012).
- [51] H. Salman, A. Zilman, C. Loverdo, M. Jeffroy, and A. Libchaber, “Solitary modes of bacterial culture in a temperature gradient”, Phys. Rev. Lett. **97**, 118101 (2006).
- [52] J. Saragosti, V. Calvez, N. Bournaveas, A. Buguin, P. Silberzan, and B. Perthame, “Mathematical description of bacterial traveling pulses”, PLoS Comput. Biol. **6**, e1000890 (2010).