



HAL
open science

Hyperpolarization of Frozen Hydrocarbon Gases by Dynamic Nuclear Polarization at 1.2 K

Basile Vuichoud, Estel Canet, Jonas Milani, Aurélien Bornet, David Baudouin, Laurent Veyre, David Gajan, Lyndon Emsley, Anne Lesage, Christophe Copéret, et al.

► **To cite this version:**

Basile Vuichoud, Estel Canet, Jonas Milani, Aurélien Bornet, David Baudouin, et al.. Hyperpolarization of Frozen Hydrocarbon Gases by Dynamic Nuclear Polarization at 1.2 K. *Journal of Physical Chemistry Letters*, 2016, 7 (16), pp.3235 - 3239. 10.1021/acs.jpcllett.6b01345 . hal-01498671

HAL Id: hal-01498671

<https://hal.sorbonne-universite.fr/hal-01498671>

Submitted on 4 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Hyperpolarization of Frozen Hydrocarbon Gases by Dynamic Nuclear Polarization at 1.2 K

Basile Vuichoud ^{*a}, Estel Canet ^{a,b,c}, Jonas Milani ^a, Aurélien Bornet ^a, David Baudouin ^d,
Laurent Veyre ^d, David Gajan ^e, Lyndon Emsley ^a, Anne Lesage ^e, Christophe Copéret ^f, Chloé
Thieuleux ^d, Geoffrey Bodenhausen ^{a,b,c}, Igor Koptug ^{b,c,g,h}, and Sami Jannin ^{*a}

^a*Institut des Sciences et Ingénierie Chimiques, Ecole Polytechnique Fédérale de Lausanne (EPFL), Batochime, CH-1015 Lausanne, Switzerland.*

^b*Département de Chimie, Ecole Normale Supérieure, PSL Research University, UPMC Univ Paris 06, CNRS, Laboratoire des Biomolécules (LBM), 24 Rue Lhomond, 75005 Paris, France.*

^c*Sorbonnes Universités, UPMC Univ Paris 06, Ecole Normale Supérieure, CNRS, Laboratoires des Biomolécules (LBM), Paris, France.*

^d*Université de Lyon, Institut de Chimie de Lyon, LC2P2, UMR 5265 CNRS-CPE Lyon-UCBL, CPE Lyon, 43 Bvd du 11 Novembre 1918, 69100 Villeurbanne, France.*

^e*Université de Lyon, Institut des Sciences Analytiques, UMR 5280, CNRS, Université Lyon 1, ENS Lyon-5, rue de la Doua, 69100 Villeurbanne, France.*

^f*ETH Zürich, Department of Chemistry and Applied Biosciences, Vladimir-Prelog-Weg 1-5/10, 8093 Zürich, Switzerland.*

^g*International Tomography Center, SB RAS, 3A Institutskaya St., Novosibirsk, 630090, Russia.*

^h*Novosibirsk State University, Pirogova St. 2, Novosibirsk, 630090, Russia.*

AUTHOR INFORMATION

Corresponding Authors

* Basile Vuichoud

Ecole Polytechnique Fédérale de Lausanne, Institut des Sciences et Ingénierie Chimiques

1015 Lausanne, Switzerland.

E-Mail: basile.vuichoud@epfl.ch

Tel: (+41) 21-693-94-27

Homepage: <http://lrmb.epfl.ch>

* Sami Jannin

Ecole Polytechnique Fédérale de Lausanne, Institut des Sciences et Ingénierie Chimiques

1015 Lausanne, Switzerland.

E-Mail: sami.jannin@epfl.ch


Tel: (+41) 21-693-97-24

Homepage: <http://lrmb.epfl.ch>

ABSTRACT

We report a simple and general method for the hyperpolarization of condensed gases by DNP. The gases are adsorbed in the pores of mesoporous silica matrices known as HYPSOs (HYper Polarizing SOLids) that have paramagnetic polarizing agents covalently bound at the surface of the mesopores. DNP is performed at low temperatures and moderate magnetic fields ($T = 1.2$ K and $B_0 = 6.7$ T). Frequency-modulated microwave irradiation is applied close to the electron spin resonance frequency ($f = 188.3$ GHz), and the electron spin polarization of the polarizing agents of HYPSO is transferred to the nuclear spins of the frozen gas. A proton polarization as high as $P(^1\text{H}) = 70\%$ can be obtained, which can be subsequently transferred to ^{13}C in natural abundance by cross-polarization, yielding up to $P(^{13}\text{C}) = 27\%$ for ethylene.

TOC GRAPHICS



Optical pumping^{1,2} is an established method for hyperpolarizing noble gases such as ^{129}Xe and ^3He ,^{3, 4} offering polarization levels close to unity, albeit with limited throughputs.⁵ Applications range from material science to biomedical research, in particular to lung imaging by MRI.^{6, 7} Unfortunately, optical pumping is restricted to a few noble gases, and molecular gases have never been hyperpolarized so far. Molecules with double or triple carbon-carbon bonds can be hydrogenated with parahydrogen by heterogeneous catalysis to produce hyperpolarized gaseous reaction products,⁸⁻¹⁰ though with polarization levels well below unity. Dissolution dynamic nuclear polarization (D-DNP)¹¹ has become a method of choice for the hyperpolarization of a broad range of small molecules in solution. The use of D-DNP to polarize ^{129}Xe ¹²⁻¹⁴ and $^{15}\text{N}_2\text{O}$ ¹⁵ has been demonstrated. In this study, we propose a general method to extend the use of D-DNP to virtually any condensed gas.

Compared to optical pumping or hydrogenation with parahydrogen, D-DNP requires rather complex instrumentation (low temperature cryostats, superconducting magnets, microwave sources, etc.) D-DNP has found applications to various fields of chemistry such as drug discovery¹⁶⁻¹⁹ and metabolomics.^{20, 21} One can detect the presence of tumors in patients by real-time imaging of metabolic conversion rates after intravenous injection of hyperpolarized metabolites such as ^{13}C -enriched pyruvate.²² D-DNP usually consists in three steps: *(i)* preparing a frozen sample containing the molecules of interest that are mixed with paramagnetic polarizing agents (PAs) in solutions that become glassy upon freezing. *(ii)* Polarizing the sample at low temperatures in moderate fields ($T = 1.2\text{ K}$ and $B_0 = 3.35\text{ T}$ in many laboratories; in our case $B_0 = 6.7\text{ T}$) by applying suitable microwave irradiation, *(iii)* rapidly melting the frozen hyperpolarized sample with superheated water, and transferring the solution to an NMR or MRI machine.

The DNP sample preparation is a critical step, which is traditionally performed as follows: (i) the polarizing agents are homogeneously dissolved in a solution together with a substance to be hyperpolarized, (ii) the solution is subsequently frozen, and (iii) the resulting sample is visually inspected and discarded if it has a "milky" appearance. When a transparent glassy state cannot be achieved, the formation of crystalline domains inevitably leads to the aggregation of the polarizing agents. Such samples are not suitable for DNP. Obtaining a glassy frozen sample often requires the addition of a glassing agent such as glycerol. This is critical to ensure a good DNP efficiency. To achieve hyperpolarization of ^{129}Xe by D-DNP,¹² the condensed gas was mixed with a co-solvent like isobutanol doped with TEMPO as polarizing agent. This was achieved under relatively high pressure (*ca.* 1 MPa) and at low temperatures (*ca.* 200 K) to ensure that both xenon and isobutanol were in a liquid state and could be homogeneously mixed. The solution was subsequently rapidly frozen by plunging into a liquid nitrogen bath, which resulted in a homogeneous glassy matrix, suitable for DNP. This sample preparation is difficult to reproduce since it may lead to the formation of Xe clusters.^{14 15} Furthermore, this method cannot easily be generalized to other gases. Our attempts to use a similar approach for ethylene have failed so far. More specifically, we have successfully mixed liquid ethylene with ethanol and TEMPO, but this mixture did not produce a homogeneous glass upon freezing. The frozen mixture was characterized by a non-uniform color distribution, clearly indicating segregation of the components, and the proton polarization could not be enhanced by DNP.

We therefore propose an alternative route for polarizing condensed gases by using recently developed HYper Polarizing SOLids (HYPSOs).^{23, 24} These mesoporous hybrid organo-silica materials contain surface-bound polarizing agents (TEMPO here) that are homogeneously distributed on the surface of the mesopores (see Fig 1A). HYPSOs were initially developed for

polarizing metabolites, since they can easily be filtered out after dissolution, resulting in pure hyperpolarized solutions without contaminants. We show here that HYPsOs can be also used for polarizing gases such as propane, butane and ethylene. In this preliminary study, we simply condensed these gases using first and third generations of HYPsO materials (HYPsO-1²⁴ and HYPsO-3²⁵), and we polarized the frozen impregnated materials using our state-of-the-art DNP polarizer.^{26, 27} We have designed and built a simple system for impregnating HYPsOs with controlled amounts of gas (Fig. 1B). Here, 50 mL of a mixture of *n*-butane, isobutane and propane from a gas canister (“butane refill gas” for lighters, sold by Migros) was condensed with 10.2 mg of HYPsO-1. The loading capacity of HYPsO-1 being 1.8 mL/g, only about 8.9 mg of the liquefied gas were expected to enter the pores (Fig. 1C).²⁴ The sample was then rapidly frozen in our DNP polarizer^{26, 27} at 4.2 K, before lowering the temperature to 1.2 K. A similar procedure was applied for 50 mL of ethylene (Carbagas) with 22.5 mg of HYPsO-3²⁵ precooled in a bath of liquid ethanol at 159 K. The loading capacity of the cubic three-dimensional HYPsO-3 is 0.47 mL/g, hence only 6 mg of liquid ethylene were expected to enter the pores. The samples were polarized at $B_0 = 6.7$ T and $T = 1.2$ K^{26, 27}, using a saddle-coil probe doubly tuned for $f(^1\text{H}) = 285.3$ MHz and $f(^{13}\text{C}) = 71.73$ MHz. The sample was irradiated at 188.29 GHz (negative polarization) with $P_{\mu\text{w}} = 87.5$ mW using triangular microwave frequency modulation²⁸ with a frequency $f_{\text{mod}} = 10$ kHz over a range $\Delta f_{\text{mod}} = \pm 50$ MHz. The build-up curves of the polarization $P(^1\text{H})$ were monitored with 1° pulses applied every 5 s. Cross-polarization^{29,30} from ^1H to ^{13}C was performed with multiple adiabatic contacts as described elsewhere,³¹ and the polarization $P(^{13}\text{C})$ was measured with 5° pulses every 37.5 s for the mixture of *n*-butane, isobutane and propane, and every 75 s for ethylene. After DNP, the initial volume of condensed gas (ca. 50 mL) could be fully recovered by connecting the sample cup to an empty syringe while the sample was still frozen, and

gently warming-up to room temperature. Such an amount of undiluted gas would be sufficient for various applications, for instance to follow a chemical reaction in a batch or a continuous flow reactor. Similar studies dealing with D-DNP of ^{129}Xe yielded comparable volumes of gas.¹²⁻¹⁴




Figure 1. (A) Transmission electron microscopy image of mesoporous HYPSO-1 taken with a Philips CM30 TEM operating at 300 kV. In the hexagonal structure, the red dots indicate the polarizing agents derived from TEMPO. (B) System used to impregnate

HYPSO materials with condensed gases equipped with (a) a flask containing the HYPSO material which can be cooled to liquid nitrogen temperature. A vacuum is established in the entire system by opening valves V1, V2, V3 and V4 and pumping with a rotary pump connected to (b). A manometer (c) monitors the pressure. With valves V1 and V2 closed, valves V3 and V4 are opened and the control valve VC is opened gently to let a desired volume of gas escape from the gas tank (d) and fill the syringe (e). Finally, V4 is closed and V2 is opened to let the gas adsorb onto the HYPSO powder. An overpressure valve (f) ensures safety. (C) By decreasing the temperature in (a) to 159 K using an ethanol – liquid N₂ bath and increasing the pressure, the gas is forced to enter the pores of HYPSO.

Figure 2 shows the build-up of the proton polarization in a mixture of *n*-butane, isobutane and propane at 1.2 K. A remarkable $P(^1\text{H}) = 53\%$ could be achieved in 1000 s with a build-up time constant $\tau_{\text{DNP}}(^1\text{H}) = 228 \pm 4$ s. The build-up of the polarization $P(^{13}\text{C})$ (Figure 3) was boosted by cross-polarization (CP) every 5 min to transfer the ^1H polarization to ^{13}C . For the mixture of *n*-butane, isobutane and propane, an average polarization $P(^{13}\text{C}) = 9\%$ was achieved after 20 min. For ethylene, the build-up at 1.2 K reaches $P(^1\text{H}) = 70\%$ with $\tau_{\text{DNP}}(^1\text{H}) = 1384 \pm 13$ s. Two cross-polarization (CP) contacts were performed (after 10 and 20 min) to transfer the polarization from ^1H to ^{13}C , to achieve $P(^{13}\text{C}) = 27\%$ for ethylene.




Figure 2. Build-up of the proton polarization in a mixture of *n*-butane, isobutane and propane adsorbed on HYPISO-1 at 1.2 K to $P(^1\text{H}) = 53$ % with a time constant $\tau_{\text{DNP}} = 228 \pm 4$ s.




Figure 3. Build-up of natural-abundance ^{13}C polarization in a mixture of *n*-butane, isobutane and propane adsorbed on HYPISO-1 at 1.2 K with 5 cross-polarization (CP) contacts every 5 min, leading to $P(^{13}\text{C}) = 9$ %.

This initial study demonstrates that hydrocarbon gases such as propane, butane and ethylene can be condensed in porous HYPISO materials and readily polarized by DNP. With these encouraging results in hand, we are currently trying to develop strategies for collecting the hyperpolarized hydrocarbon gases as a gas, a pure liquid, a solution in a solvent, or a solid, while preserving the hyperpolarization. In this respect, gases represent a greater challenge than liquids since the predominance of the spin-rotation mechanism leads to fairly short T_1 relaxation times.³²⁻

³⁶ There are a number of ways to significantly extend the lifetimes of non-equilibrium spin states in gases, such as increasing the pressure or mixing with a heavier buffer gas.^{32,36} By simply liquefying gases one can extend the lifetime of the polarization; the $T_1(^1\text{H})$ of liquid ethylene at 169 K is ca. 20 s.³⁷ Pure liquids could be produced by melting the frozen gas (for example with an IR laser).³⁸ Finally, one can use long-lived spin states.^{19, 39-43} In particular, at low magnetic fields, non-equilibrium spin states with the lifetimes as long as 5-6 s have been reported for propane gas,^{44,45} 100 s for ethylene-d₂ dissolved in perdeuterated hexane⁴⁶ and 1000 s for ethylene gas.³⁴ As demonstrated earlier,⁴⁷ a non-equilibrium spin state corresponding to a singlet order can be directly populated by D-DNP. Typically, for a polarization $P(^1\text{H}) = 0.7$ that was achieved in this work, for a pair of $I=1/2$ nuclei the singlet-triplet imbalance (i.e., the difference between the population of the singlet state and the average population of the three triplet states, which is proportional to $[P(^1\text{H})]^2 = 0.49$) is 49% of its maximum possible value, with the latter achieved when the singlet state is fully depleted ($[P(^1\text{H})]^2 = 1$).

The ability to produce hyperpolarized gases with lifetimes as long as 5-1000 s is promising for the development of a number of novel applications. In particular, catalytic transformations such as hydrogenation, oxidative and non-oxidative dehydrogenation, hydroformylation, oxidation, metathesis and other reactions are of primary importance in modern chemical industry.⁴⁸ Such reactions can unlock long-lived spin states produced by DNP and make them observable by NMR. In most cases, a much deeper insight into the mechanisms of industrially important processes, including synthesis of various hydrocarbons, production of fuels, polymerization, etc., is needed to achieve any significant progress in improving reaction conversion and selectivity and reducing its environmental impact. This requires detailed mechanistic studies of these reactions and calls for the implementation of novel advanced methods and instrumental techniques. Most

hyperpolarized NMR studies of catalytic processes are currently limited to the use of parahydrogen in the homogeneous catalytic hydrogenation of unsaturated hydrocarbons in solution.⁴⁹⁻⁵² Yet the vast majority of industrial catalytic processes use heterogeneous rather than homogeneous catalysis. The extension of parahydrogen-based studies to heterogeneous catalysis, while successful,⁸⁻¹⁰ faces some major limitations and is still naturally restricted to reactions that involve H₂. The use of hyperpolarized gases to study chemical reactions is particularly promising in this context and has the potential to significantly advance hyperpolarization-based mechanistic NMR studies of catalysis and other areas of research.

Notes

The authors declare no competing financial interests.

ACKNOWLEDGMENTS

Igor Koptyug acknowledges the Ecole Normale Supérieure for a visiting professorship and the Russian Science Foundation (Project No. 14-35-00020) for financial support. The authors thank Anto Barisic and Dr. Pascal Miéville at EPFL for valuable technical assistance, and André Fattet for building the system to impregnate HYPSON with gases. This work was supported by the Swiss National Science Foundation (SNSF), the Ecole Polytechnique Fédérale de Lausanne (EPFL), ETH Zürich, the Swiss Commission for Technology and Innovation (CTI), Bruker BioSpin Switzerland AG, the French CNRS, Pulsalys in Lyon, and the European Research Council (ERC Advanced Grant No. 339754 "Dilute para-water", and ERC Advanced Grant No. 320860).

REFERENCES

- (1) Kastler A. Optical Methods of Atomic Orientation. *J. Opt. Soc. Am.*, **1957**, *47*, 460-465.
- (2) Walker T.; Happer W. Spin-exchange optical pumping of noble-gas nuclei. *Rev. Mod. Phys.*, **1997**, *69*, 629-642.
- (3) Raftery D.; Long H.; Meersmann T.; Grandinetti P. J.; Reven, L.; Pines A. High-field of Adsorbed Xenon Polarized by Laser Pumping. *Phys. Rev. Lett.*, **1991**, *66*, 584-587.
- (4) Colegrove F. D.; Franken P. A.; Optical Pumping of Helium in the 3S_1 Metastable State. *Phys. Rev.*, **1960**, *119*, 680-690.
- (5) Ruset I. C.; Ketel S.; Hersman F. W. Optical Pumping System Design for Large Production of Hyperpolarized ^{129}Xe . *Phys. Rev. Lett.*, **2006**, *96*, 053002.
- (6) Hyperpolarized Xenon-129 Magnetic Resonance: Concepts, Production, Techniques and Applications. Meersmann T. and Brunner E., Eds., RSC Publishing, Cambridge, UK, 2015.
- (7) Cherubini A.; Bifone A.; Hyperpolarized xenon in biology. *Prog. Nucl. Magn. Reson. Spectrosc.*, **2003**, *42*, 1-30.
- (8) Kovtunov K. V.; Zhivonitko V. V.; Skovpin I. V.; Barskiy D. A.; Koptyug I. V.; Parahydrogen-induced polarization in heterogeneous catalytic processes. *Top. Curr. Chem.*, **2013**, *338*, 123-180.
- (9) Kovtunov K. V.; Barskiy D. A.; Coffey A. M.; Truong M. L.; Salnikov O. G.; Khudorozhkov A. K.; Inozemtseva E. A.; Prosvirin I. P.; Bukhtiyarov V. I.; Waddell K. W.; Chekmenev E. Y.; Koptyug I. V. High-resolution 3D proton MRI of hyperpolarized gas enabled by parahydrogen and Rh/TiO₂ heterogeneous catalyst. *Chem. Eur. J.*, **2014**, *20*, 11636-11639.
- (10) Zhivonitko V. V.; Telkki V. -V.; Koptyug I. V. Characterization of Microfluidic Gas Reactors Using Remote-Detection MRI and Parahydrogen-Induced Polarization. *Angew. Chem. Int. Ed.*, **2012**, *51*, 8054-8058.

- (11) Ardenkjaer-Larsen J. H.; Fridlund B.; Gram A.; Hansson G.; Hansson L.; Lerche M. H.; Servin R.; Thaning M.; Golman K. Increase in signal-to-noise ratio of > 10,000 times in liquid-state NMR. *Proc. Nat. Aca. Sci.*, **2003**, *100*, 10158-10163.
- (12) Comment A.; Jannin S.; Hyacinthe J. N.; Miéville P.; Sarkar R.; Ahuja P.; Vasos P. R.; Montet X.; Lazeyras F.; Vallée J.-P.; Hautle P.; Konter J. A.; van den Brandt B.; Ansermet J. –Ph.; Gruetter R.; Bodenhausen G. Hyperpolarizing Gases via Dynamic Nuclear Polarization and Sublimation. *Phys. Rev. Lett.*, **2010**, *105*, 018104.
- (13) Capozzi A.; Roussel C.; Comment A.; Hyacinthe J.-N. Optimal glass-forming solvent brings sublimation dynamic nuclear polarization to ¹²⁹Xe hyperpolarization biomedical imaging standards. *J. Phys. Chem. C*, **2015**, *119*, 5020.
- (14) Kuzma N. N.; Pourfathi M.; Kara H.; Manasseh P.; Ghosh R. K.; Ardenkjaer-Larsen J. H.; Kadlecsek S. J.; Rizi R. R. Cluster formation restricts dynamic nuclear polarization of xenon in solid mixtures. *J. Chem. Phys.*, **2012**, *137*, 104508.
- (15) Ghosh R. K.; Kuzma N. N.; Kadlecsek S. J.; Rizi R. R. Versatile pulse sequence device to conserve hyperpolarization for NMR and MRI studies, *Magn. Reson. Med.*, **2015**, *75*, 182.
- (16) Lee Y.; Zeng H. F.; Mazur A.; Wegstroth M.; Carlomagno T.; Reese M.; Lee D.; Becker S.; Griesinger C.; Hilty C. Hyperpolarized Binding Pocket Nuclear Overhauser Effect for Determination of Competitive Ligand Binding. *Angew. Chem. Int. Ed.*, **2012**, *51*, 5179-5182.
- (17) Lee Y.; Zeng H. F.; Ruedisser S.; Gossert A. D.; Hilty C. Nuclear magnetic resonance of hyperpolarized fluorine for characterization of protein-ligand interactions. *J. Am. Chem. Soc.*, **2012**, *134*, 17448-17451.
- (18) Chappuis Q.; Milani J.; Vuichoud B.; Bornet A.; Gossert A. D.; Bodenhausen G.; Jannin S. Hyperpolarized Water to Study Protein-Ligand Interactions. *J. Phys. Chem. Lett.*, **2015**, *6*, 1674-1678.
- (19) Buratto R.; Bornet A.; Milani J.; Mammoli D.; Vuichoud B.; Salvi N.; Singh M.; Laguerre A.; Passemard S.; Gerber-Lemaire S.; Jannin S.; Bodenhausen G. Drug screening boosted by hyperpolarized long-lived states in NMR. *ChemMedChem*, **2014**, *9*, 2509-2515.

- (20) Meier S.; Jensen P. R.; Karlsson M.; Lerche M. H. Hyperpolarized NMR Probes for Biological Assays. *Sensors*, **2014**, *14*, 1576-1597.
- (21) Dumez J.-N.; Milani J.; Vuichoud B.; Bornet A.; Lalande-Martin J.; Tea I.; Yon M.; Maucourt M.; Deborde C.; Moing A.; Frydman L.; Bodenhausen G.; Jannin S.; Giraudeau P. Hyperpolarized NMR of plant and cancer cell extracts at natural abundance. *Analyst*, **2015**, *140*, 58605863.
- (22) Nelson S. J.; Kurhanewicz J.; Vigneron D. B.; Larson P. E. Z.; Harzstark A. L.; Ferrone M.; van Criekinge M.; Chang J. W.; Bok R.; Park I.; Reed G.; Carvajal L.; Small E. J.; Munster P.; Weinberg V. K.; Ardenkjaer-Larsen J. H.; Chen A. P.; Hurd R. E.; Odegaardstuen L.-I.; Robb F. J.; Tropp J.; Murray J. A. Metabolic imaging of patients with prostate cancer using hyperpolarized [1-¹³C]pyruvate. *Science Translational Medicine*, **2013**, *5*, 198ra108.
- (23) Gajan D.; Schwarzwälder M.; Conley M. P.; Grüning W. R.; Rossini A. J.; Zagdoun A.; Lelli M.; Yulikov M.; Jeschke G.; Sauvée C.; Ouari O.; Tordo P.; Veyre L.; Lesage A.; Thieuleux C.; Emsley L.; Copéret C. Solid-Phase Polarization Matrixes for Dynamic Nuclear Polarization from Homogeneously Distributed Radicals in Mesostuctured Hybrid Silica Materials. *J. Am. Chem. Soc.*, **2013**, *135*, 15459-15466.
- (24) Gajan D.; Bornet A.; Vuichoud B.; Milani J.; Melzi R.; van Kalkerem H. A.; Veyre L.; Thieuleux C.; Conley M. P.; Grüning W. R.; Schwarzwälder M.; Lesage A.; Copéret C.; Bodenhausen G.; Emsley L.; Jannin S. Hybrid polarizing solids for pure hyperpolarized liquids through dissolution dynamic nuclear polarization. *P. Natl. Acad. Sci. USA*, **2014**, *111*, 14693-14697.
- (25) Baudouin D.; van Kalkerem H. A.; Bornet A.; Vuichoud B.; Veyre L.; Cavallès M.; Schwarzwälder M.; Liao W.-C.; Gajan D.; Bodenhausen G.; Emsley L.; Lesage A.; Jannin S.; Copéret C.; Thieuleux C. Cubic Three-Dimensional Hybrid Silica Solids for Nuclear Hyperpolarization. *Chem. Sci.* **2016**, DOI: 10.1039/C6SC02055K.
- (26) Comment A.; van den Brandt B.; Uffmann K.; Kurdzesau F.; Jannin S.; Konter J. A.; Hautle P.; Wenckebach W. T. H.; Gruetter R.; van der Klink J. J. Design and Performance of

a DNP Prepolarizer Coupled to a Rodent MRI Scanner. *Concepts Magn. Reson. B*, **2007**, *31B*, 255-269.

(27) Jannin S.; Bornet A.; Melzi R.; Bodenhausen G. High field dynamic nuclear polarization at 6.7 T: Carbon-13 polarization above 70% within 20 min. *Chem. Phys. Lett.*, **2012**, *549*, 99-102.

(28) Bornet A.; Milani J.; Vuichoud B.; Perez Linde A. J.; Bodenhausen G. Microwave frequency modulation to enhance Dissolution Dynamic Nuclear Polarization. *Chem. Phys. Lett.*, **2014**, *602*, 63-67.

(29) Hartmann S. R.; Hahn E. L. Nuclear Double Resonance in the Rotating Frame. *Phys. Rev.* **1962**, *128*, 2042-2053.

(30) Pines A.; Gibby M. G.; Waugh J. S. Proton-Enhanced Nuclear Induction Spectroscopy. A Method for High Resolution NMR of Dilute Spins in Solids. *J. Chem. Phys.*, **1972**, *56*, 1776-1777.

(31) Bornet A.; Melzi R.; Linde A. J. P.; Hautle P.; van den Brandt B.; Jannin S.; Bodenhausen G. Boosting Dissolution Dynamic Nuclear Polarization by Cross Polarization. *J. Phys. Chem. Lett.*, **2013**, *4*, 111-114.

(32) Bloom M.; Lipsicias M.; Muller B. H. Proton spin-lattice relaxation in polyatomic gases. *Canadian Journal of Physics*, **1961**, *39*, 1093-1109.

(33) Jameson C. J.; Jameson A. K.; Smith N. C.; Hwang J. K.; Zia T. Carbon-13 and proton spin relaxation in methane in the gas phase. *J. Phys. Chem.*, **1991**, *95*, 1092-1098.

(34) Zhivonitko V. V.; Kovtunov K. V.; Chapovsky P. L.; Koptug I. V. Nuclear spin isomers of ethylene: enrichment by chemical synthesis and application for NMR signal enhancement. *Angew. Chem. Int. Ed.*, **2013**, *52*, 13251-13255.

- (35) Hirasaki G. J.; Lo S.-W.; Zhang Y. NMR properties of petroleum reservoir fluids. *Magn. Reson. Imaging*, **2003**, *21*, 269-277.
- (36) Gas Phase NMR. Jackowski K. and Jaszunski M., Eds. 2016, Royal Society of Chemistry, Cambridge, UK, 2016.
- (37) Trappeniers N. J.; Ligthart F. A. S. Proton spin relaxation in ethylene. A new high pressure phase. *Chem. Phys. Lett.*, **1973**, *19*, 465-470.
- (38) Joo C.-G.; Hu K.-N.; Bryant J. A.; Griffin R. G. In situ temperature jump high-frequency dynamic nuclear polarization experiments: enhanced sensitivity in liquid-state NMR spectroscopy. *J. Am. Chem. Soc.*, **2006**, *128*, 9428-9432.
- (39) Stevanato G.; Hill-Cousins J. T.; Hakansson P.; Roy S. S.; Brown L. J.; Brown R. C. D.; Pileio G.; Levitt M. H. A nuclear singlet lifetime of more than one hour in room-temperature solution. *Angew. Chem. Int. Ed.*, **2015**, *54*, 3740-3743.
- (40) Levitt M. H. Singlet nuclear magnetic resonance. *Annu. Rev. Phys. Chem.*, **2012**, *63*, 89-105.
- (41) Vasos P. R.; Comment A.; Sarkar R.; Ahuja P.; Jannin S.; Ansermet J.-P.; Konter J. A.; Hautle P.; Van Den Brandt B.; Bodenhausen G. Long-lived states to sustain hyperpolarized magnetization. *Proc. Natl. Acad. Sci. USA*, **2009**, *106*, 18469-18473.
- (42) Stevanato G.; Roy S. S.; Hill-Cousins J.; Kuprov I.; Brown L. J.; Brown R. C. D.; Pileio G.; Levitt M. H. Long-lived nuclear spin states far from magnetic equivalence. *Phys. Chem. Chem. Phys.*, **2015**, *17*, 5913-5922.

- (43) Feng Y.; Theis T.; Wu T.-L.; Claytor K.; Warren W. S. Long-lived polarization protected by symmetry. *J. Chem. Phys.*, **2014**, *141*, 134307.
- (44) Kovtunov K. V.; Truong M. L.; Barskiy D. A.; Koptug I. V.; Coffey A. M.; Waddell K. W.; Chekmenev E. Y. Long-lived spin states for low-field hyperpolarized gas MRI. *Chem. Eur. J.*, **2014**, *20*, 14629-14632.
- (45) Kovtunov K. V.; Truong M. L.; Barskiy D. A.; Salnikov O. G.; Bukhtiyarov V. I.; Coffey A. M.; Waddell K. W.; Koptug I. V.; Chekmenev E. Y. Propane-d₆ heterogeneously hyperpolarized by parahydrogen. *J. Phys. Chem. C*, **2014**, *118*, 28234-28243.
- (46) Zhou Z.; Claytor K.; Warren W. S.; Theis T. Accessing long lived ¹H states via ²H couplings. *J. Magn. Reson.*, **2016**, *263*, 108-115.
- (47) Tayler M. C. D.; Marco-Rius I.; Kettunen M. I.; Brindle K. M.; Levitt M. H.; Pileio G. Direct enhancement of nuclear singlet order by dynamic nuclear polarization. *J. Am. Chem. Soc.*, **2012**, *134*, 7668-7671.
- (48) Ullmann's Encyclopedia of Industrial Chemistry, 2012, Wiley-VCH Verlag GmbH & Co. KGaA.
- (49) Natterer J.; Bargon J. Parahydrogen induced polarization. *Progr. NMR Spectr.*, **1997**, *31*, 293-315.
- (50) Canet D.; Aroulanda C.; Mutzenhardt P.; Aime S.; Gobetto R.; Reineri F. Para-hydrogen enrichment and hyperpolarization. *Concepts Magn. Reson.*, **2006**, *28A*, 321-330.

(51) Duckett S. B.; Wood N. J. Parahydrogen-based NMR methods as a mechanistic probe in inorganic chemistry. *Coord. Chem. Rev.*, **2008**, *252*, 2278-2291.

(52) Koch A.; Bargon J. In situ NMR observation of mono- and binuclear rhodium dihydride complexes using parahydrogen-induced polarization. *Inorg. Chem.*, **2001**, *40*, 533-539.